

# Molecular evidence of Pleistocene bidirectional faunal exchange between Europe and the Near East: the case of the bicoloured shrew (*Crocidura leucodon*, Soricidae)

S. DUBEY,\* J.-F. COSSON,† V. VOHRALÍK,‡ B. KRYŠTUFEK,§ E. DIKER¶ & P. VOGEL\*

\*Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland

†Centre de Biologie et de Gestion des Populations (CBGP), INRA-EFPA, Campus International de Baillarguet, Montferrier/Lez cedex, France

‡Department of Zoology, Charles University, Viničná, Prague, Czech Republic

§Science and Research Centre of Koper, University of Primorska, Garibaldijeva, Koper, Slovenia

¶Department of Biology, Faculty of Art and Sciences, Trakya University, Edirne, Turkey

## Keywords:

apolipoprotein B gene;  
*Crocidura leucodon*;  
cytochrome *b* gene;  
phylogeography.

## Abstract

We sequenced 1077 bp of the mitochondrial cytochrome *b* gene and 511 bp of the nuclear Apolipoprotein B gene in bicoloured shrew (*Crocidura leucodon*, Soricidae) populations ranging from France to Georgia. The aims of the study were to identify the main genetic clades within this species and the influence of Pleistocene climatic variations on the respective clades. The mitochondrial analyses revealed a European clade distributed from France eastwards to north-western Turkey and a Near East clade distributed from Georgia to Romania; the two clades separated during the Middle Pleistocene. We clearly identified a population expansion after a bottleneck for the European clade based on mitochondrial and nuclear sequencing data; this expansion was not observed for the eastern clade. We hypothesize that the western population was confined to a small Italo-Balkan refugium, whereas the eastern population subsisted in several refugia along the southern coast of the Black Sea.

## Introduction

The impact of Pleistocene climatic fluctuations on European biota is well documented. Studies of mitochondrial DNA markers reveal general patterns involving southern refugia and northern recolonization routes, as a modification of genetic diversity of terrestrial biota throughout the Holarctic (Hewitt, 2000, 2004a,b). Populations were isolated in different glacial refugia by barriers such as mountains and seas, leading to various recolonization patterns (e.g. Taberlet *et al.*, 1998; Hewitt, 1999). Widely accepted refugia include the Iberian Peninsula in the west and the Italo-Balkan region in the east (e.g. Thorpe, 1984; Ferris *et al.*, 1993, 1998; Dumolin-Lapegue *et al.*, 1997; Santucci *et al.*, 1998; Taberlet *et al.*, 1998;

Dubey *et al.*, 2006); however, a number of phylogeographical studies of various taxa have revealed an additional pattern of colonization of Europe by populations originating from eastern areas such as the Caucasus, southern Urals and western Asia (Cooper *et al.*, 1995; Bilton *et al.*, 1998; Nesbo *et al.*, 1999; Palme & Vendramin, 2002; Seddon *et al.*, 2002; Michaux *et al.*, 2004; Culling *et al.*, 2006; Dubey *et al.*, 2006). This pattern of post-glacial recolonization appears to be more common than previously suspected and probably affected a large range of taxa.

With regard to population dynamics, few studies have detailed the impact of Pleistocene climatic oscillations, e.g. the dating of population expansions and identifying differences in patterns among clades of different geographical origin (Michaux *et al.*, 2004; Brändli *et al.*, 2005; Culling *et al.*, 2006; Dubey *et al.*, 2006; Koch *et al.*, 2006; Marmi *et al.*, 2006; Vörös *et al.*, 2006; Dubey *et al.*, 2007a). In the same way, the majority of previous studies are based solely on mitochondrial DNA, whereas the inclusion of uniparentally and biparentally inherited markers present in nuclear genomes can enhance our

Correspondence: Sylvain Dubey, Département d'Ecologie et d'Evolution, Bâtiment du Biophore, Université de Lausanne, 1015 Lausanne, Switzerland.

Tel.: +41 21 692 4163; fax: +41 21 692 4165;

e-mail: sylvain.dubey@unil.ch

Address from September 2007: Shine Laboratory, School of Biological Sciences, University of Sydney, Sydney, NSW 2006, Australia.

understanding of population history (Brändli *et al.*, 2005; Dubey *et al.*, 2006).

In the present study, we focused on the bicoloured shrew, *Crocidura leucodon* (Hermann, 1780), which is a strictly western Eurasian species, distributed from France to central and southern Europe to Turkey and Georgia. This species seems to be an interesting candidate for a phylogeographical study, as Vogel *et al.* (2003), based on only two samples, from Switzerland and Turkey, noted a substantial Kimura 2-paramters genetic distance of 3.8% between them for the cytochrome *b* gene. In addition, Poulakakis *et al.* (2005) analysed three Greek samples (from Peloponnesus and Lesbos) obtained from owl pellets, based on the same gene, promoting such material as usable for phylogeographical studies, without giving any information concerning the distances between haplotypes. Nevertheless, the intraspecific genetic distances were considered low.

We tested (i) in which way the Bosphorus strait has isolated the European and Near East populations and (ii) in which way the Pleistocene climatic fluctuations have had different impacts on the western and eastern populations, such as bottleneck effects. To resolve these questions, we analysed sequence data from mitochondrial and nuclear markers (cytochrome *b* and apolipoprotein B) and considered the molecular clock.

## Materials and methods

### Sampling


We analysed 60 samples of *C. leucodon* collected from France to Georgia (Fig. 1 and Table 1), three other Eurasian crocidurine taxa, and as an out-group a soricine, *Sorex minutus*. This set of samples (Table 1)

included material from the collections of Lausanne (IZEA), Switzerland; Prague (DZCU), Czech Republic and Ljubljana (PMS), Slovenia. Some sequences were taken from Ohdachi *et al.* (2004); Fontanillas *et al.* (2005) and Dubey *et al.* (2006). We also used samples from Poulakakis *et al.* (2005) that were treated separately.

### DNA extraction and amplification of cytochrome *b* and apolipoprotein B genes

Samples (livers) from IZEA collection were first frozen in the field in liquid nitrogen and kept for several years at  $-70^{\circ}\text{C}$  before being stored in ethanol until DNA extraction. Samples from the other collections were directly stored in ethanol. DNA extraction was carried out using the QIA Amp DNA Mini Kit (Qiagen, Valencia, PA, USA). Double-stranded DNA amplifications of the mitochondrial cytochrome *b* gene (*cyt-b*) were performed with the primer pairs L14724/H15149, C1/C2, C3/H15915 and L14724/H15915 (Irwin *et al.*, 1991; Dubey *et al.*, 2006). Amplification of the apolipoprotein B (*ApoB*) nuclear genes was performed using the primer pairs ApoBf/ApoBr (Dubey *et al.*, 2007b). Amplification conditions for the *ApoB* consisted of 40 cycles of 45 s denaturation at  $94^{\circ}\text{C}$ , 45 s annealing at  $50^{\circ}\text{C}$  and 90 s extension at  $72^{\circ}\text{C}$ ; for the *cyt-b*, the conditions of Dubey *et al.* (2006) were used.

The PCR products were checked on a 1% agarose gel and then purified using the QIAquick PCR Purification Kit (QIAGEN) following the manufacturer's instructions. DNA sequencing was performed in a total volume of  $10\ \mu\text{L}$  containing 1–3  $\mu\text{L}$  of amplified PCR product, 1  $\mu\text{L}$  of  $10\ \mu\text{M}$  primer, and 4  $\mu\text{L}$  of ABI PRISM<sup>TM</sup> Dye Terminator 1 (Perkin-Elmer, Boston, MA, USA). Sequence reactions were visualized on an ABI 3100 genetic analyser (Applied Biosystems, Foster City, CA, USA).



**Fig. 1** (a) Location of analysed samples, tentative distribution of mitochondrial clades (grey areas), allelic frequencies of nuclear genes (cheese), and nucleotide diversity of mitochondrial and nuclear gene; (b,c) Observed (solid line) and expected (dotted-line) mismatch distributions for a sudden expansion of the western and eastern clades, respectively.

**Table 1** Details of samples used in this study: species, samples location, collection, location on map (Fig. 1), *cyt-b* Id. Code, accession number of *cyt-b* sequences, and *ApoB* alleles.

Species	Samples location	Coll. code	Number on the map	<i>Cyt-b</i> Id. code	Accession ( <i>cyt-b</i> ) and <i>ApoB</i> allele
<i>Sorex minutus</i>	Champmartin, CH	IZEA 7622	/	/	DQ630379/-
<i>Crocidura suaveolens</i>	Fulophasa, HU	Dubey <i>et al.</i> (2006)	/	/	AY843451/-
<i>Crocidura russula</i>	Vaud, CH	Fontanillas <i>et al.</i> (2005)	/	/	AY769264/-
<i>Crocidura lasiura</i>	Ussurisk, RU	Ohdachi <i>et al.</i> (2004)	/	/	AB077071/-
<i>Crocidura leucodon</i>	Visp, CH	IZEA 7553	1	CH1	DQ994744/A1
<i>Crocidura leucodon</i>	Réchy, CH	IZEA 7552	1	CH2	DQ994745/A1
<i>Crocidura leucodon</i>	Brigerbad, CH	IZEA 2951	1	CH3	DQ994747/-
<i>Crocidura leucodon</i>	Raron, CH	IZEA 7532	1	CH4	DQ994746/-
<i>Crocidura leucodon</i>	Réchy, CH	IZEA 5590	1	CH9	DQ994794/-
<i>Crocidura leucodon</i>	Brigerbad, CH	IZEA 7526	1	/	-/A1
<i>Crocidura leucodon</i>	Gordevio, CH	IZEA 5965	2	CH5	DQ994748/-
<i>Crocidura leucodon</i>	Gordevio, CH	IZEA 5963	2	CH6	DQ994749/A1
<i>Crocidura leucodon</i>	Quartino, CH	IZEA 5455	2	CH7	DQ994750/A1
<i>Crocidura leucodon</i>	Grison, CH	IZEA 9013	3	CH8	DQ994785/A1
<i>Crocidura leucodon</i>	Karlsruhe, DE	IZEA 5441	4	DE5	DQ994795/A1
<i>Crocidura leucodon</i>	Rendsburg, DE	IZEA 7835	5	DE1	DQ994761/-
<i>Crocidura leucodon</i>	Rendsburg, DE	IZEA 7836	5	DE2	DQ994762/-
<i>Crocidura leucodon</i>	Rendsburg, DE	IZEA 7837	5	DE3	DQ994763/A1
<i>Crocidura leucodon</i>	Rendsburg, DE	IZEA 7838	5	DE4	DQ994764/-
<i>Crocidura leucodon</i>	Mignouillard, FR	IZEA 9006	6	FR4	DQ994780/A1
<i>Crocidura leucodon</i>	Mignouillard, FR	IZEA 9002	6	FR5	DQ994781/A1
<i>Crocidura leucodon</i>	Chapelle, FR	IZEA 9003	7	FR2	DQ994779/A1
<i>Crocidura leucodon</i>	Chapelle, FR	IZEA 9005	7	FR3	DQ994782/A1
<i>Crocidura leucodon</i>	St-Martin, FR	IZEA 9008	8	FR1	DQ994783/A1
<i>Crocidura leucodon</i>	St-Etienne, FR	IZEA 9009	9	FR6	DQ994784/A1
<i>Crocidura leucodon</i>	Vercelli, IT	IZEA 7948	10	IT1	DQ994767/-
<i>Crocidura leucodon</i>	Vercelli, IT	IZEA 7946	10	IT2	DQ994768/-
<i>Crocidura leucodon</i>	Vercelli, IT	IZEA 7952	10	IT3	DQ994769/-
<i>Crocidura leucodon</i>	Serramazoni, IT	IZEA 5662	11	IT4	DQ994770/A1
<i>Crocidura leucodon</i>	Serramazoni, IT	IZEA 5649	11	IT5	DQ994771/-
<i>Crocidura leucodon</i>	Serramazoni, IT	IZEA 5663	11	IT6	DQ994772/A1
<i>Crocidura leucodon</i>	Piacenza, IT	IZEA 7517	12	IT7	DQ994773/A1
<i>Crocidura leucodon</i>	Bratislava, SK	IZEA 5728	13	SK1	DQ994765/-
<i>Crocidura leucodon</i>	Bratislava, SK	IZEA 5723	13	SK2	DQ994766/A1
<i>Crocidura leucodon</i>	Lesbos, GR	IZEA 4153	14	GR1	DQ994777/A1
<i>Crocidura leucodon</i>	Lesbos, GR	IZEA 3929	14	GR2	DQ994778/A1
<i>Crocidura leucodon</i>	Lesbos, GR	Poulakakis <i>et al.</i> , 2005	14	-	AY452165/-
<i>Crocidura leucodon</i>	Burgas, BG	IZEA 8059	15	BG1	DQ994774/A1/A2
<i>Crocidura leucodon</i>	Burgas, BG	IZEA 8060	15	BG2	DQ994775/A1
<i>Crocidura leucodon</i>	Burgas, BG	IZEA 8063	15	BG3	DQ994776/A1
<i>Crocidura leucodon</i>	Burgas, BG	IZEA 8059	15	-	-/A1/A2
<i>Crocidura leucodon</i>	Alazani, GE	IZEA 23629	16	GE1	DQ994756/A1/A4
<i>Crocidura leucodon</i>	Alazani, GE	IZEA 23635	16	GE2	DQ994757/A4
<i>Crocidura leucodon</i>	Alazani, GE	IZEA 23613	16	GE3	DQ994758/A1
<i>Crocidura leucodon</i>	Dusheti, GE	IZEA 2880	17	GE4	DQ994760/A1/A3
<i>Crocidura leucodon</i>	Cakalli, TR	IZEA 6064	18	TR1	DQ994751/A1
<i>Crocidura leucodon</i>	Rize, TR	IZEA 6045	19	TR2	DQ994752/-
<i>Crocidura leucodon</i>	Rize, TR	IZEA 6049	19	TR3	DQ994753/A1
<i>Crocidura leucodon</i>	Rize, TR	IZEA 6047	19	TR4	DQ994754/A1
<i>Crocidura leucodon</i>	Rize, TR	IZEA 6046	19	TR5	DQ994755/A1
<i>Crocidura leucodon</i>	Altindere, TR	IZEA 6039	20	TR6	DQ994759/A1
<i>Crocidura leucodon</i>	Altindere, TR	IZEA 6038	20	TR7	DQ994796/-
<i>Crocidura leucodon</i>	Altindere, TR	IZEA 6044	20	-	-/A1
<i>Crocidura leucodon</i>	Yelibeli, Kar., TR	DZCU TU-1179	23	TR8	DQ994787/A1
<i>Crocidura leucodon</i>	Çiğlikara, Ant., TR	DZCUTU-1195	24	TR9	DQ994786/A1
<i>Crocidura leucodon</i>	Karlovy Vary, CZ	DZCU 01400	25	-	-/A1
<i>Crocidura leucodon</i>	Karlovy Vary, CZ	DZCU 01399	25	CZ1	DQ994788/A1

Table 1 (Continued).

Species	Samples location	Coll. code	Number on the map	Cyt- <i>b</i> Id. code	Accession (cyt- <i>b</i> ) and Apob allele
<i>Crocidura leucodon</i>	Karlovy Vary, CZ	DZCU 01398	25	CZ2	DQ994789/A1
<i>Crocidura leucodon</i>	Mt. Cer, SE	PMS 7391	26	SE1	DQ994790/A1
<i>Crocidura leucodon</i>	Cataloi, RO	IZEA 8160	27	RO1	DQ994791/A1
<i>Crocidura leucodon</i>	Slobozia, RO	IZEA 8170	28	RO2	DQ994792/A1
<i>Crocidura leucodon</i>	Vrhnik, SI	PMS Slo 1	29	SI1	DQ994793/A1
<i>Crocidura leucodon</i>	Katrançi-Biga, TR	2003.131	30	TR10	EF417543/A1
<i>Crocidura leucodon</i>	Özbek, İzmir, TR	2003.199	31	TR11	EF417544/A1
<i>Crocidura leucodon</i>	Terzialan-Çan., TR	2003.217	32	TR12	EF417545/A1
<i>Crocidura leucodon</i>	Parnitha, GR	Poulakakis <i>et al.</i> , 2005	21	–	AY452166/-
<i>Crocidura leucodon</i>	Stymfalia, GR	Poulakakis <i>et al.</i> , 2005	22	–	AY452176/-

BG, Bulgaria; CZ, Czech Republic; DE, Germany; FR, France; GE, Georgia; GR, Greece; HU, Hungary; IT, Italy; RU, Russia; SE, Serbia; SK, Slovak Republic; SI, Slovenia; CH, Switzerland; TR, Turkey.

### Phylogenetic methods

Nucleotide sequences of *cyt-b* and *ApoB* genes were edited with Sequence Navigator (Parker, 1997) and manually aligned. Two methods of phylogenetic analyses were carried out for *cyt-b*, using PAUP\* version 4.0b10 PPC (Swofford, 1998). Tests were conducted on the complete fragment (1077 bp), all codon positions were used, and trees were rooted using sequences of *S. minutus* (DQ630379). A Neighbour Joining (NJ) tree was constructed using the general time reversible (GTR; Rodriguez *et al.*, 1990) genetic distance, which was selected previously using MODELTEST 3.06 according to the protocol of Posada & Crandall (1998). The Parsimony analyses (MP) were performed using the following options: heuristic search, stepwise-addition of sequences, 10 replicates of random addition of taxa and TBR branch swapping (Swofford, 1998); all codon positions were equally weighted. Bootstrap support values were obtained with 1000 pseudo-replicates and 10 random replicates of stepwise-addition sequences. Fast maximum likelihood (ML) heuristic searches and bootstrap analyses (1000 replicates) were performed using PHYL (Guindon & Gascuel, 2003) with a GTR model.

### Nucleotide diversity and genetic structure

The nucleotide diversities ( $\pi$ ) of *cyt-b* and *ApoB* and the population genetic structure for *cyt-b* (analysis of molecular variance, AMOVA) were estimated using Arlequin version 2.0 Software (Schneider, 2000). AMOVA was performed at two different hierarchical levels: among clades and within clades.

### Molecular clock

The molecular clock hypothesis was tested following the method of Posada & Crandall (1998). Estimation of divergence time from the molecular data was performed according to the calibration developed for *cyt-b* in

Soricidae by Fumagalli *et al.* (1999), based on an estimate of 20 Myr for the split between the Soricinae and the Crocidurinae and on the number of third-position transversions observed. Nevertheless, it could not be used directly because of the low number of third-position transversions observed at a specific level within the family. Thus, we used the calibration of Dubey *et al.* (2006), based on the previous study that allowed a better estimate because of the use of a ML distance including all the codon positions with a divergence rate of 0.057 ML distance Myr<sup>-1</sup> (95% CI: 0.044–0.070).

### Isolation with migration

We used the 'isolation with migration' (or IM, Nielsen & Wakeley, 2001) model to date the divergence between populations of both sides of the Marmara Sea within each clade. As suggested by Hey (2007), we started the simulations using a burn-in of 100 000 steps followed by a 1/2 h run with maximum values for theta, *m* and *t* arbitrary set to 10 and the -J1 run option. From there, appropriate values for priors were chosen. In each following run, plots of parameter trend lines were systematically consulted for assessing how well the Markov chain was exploring the parameter space. Convergence by the Markov chain simulations was assessed by monitoring three independent chains, and by assessing the autocorrelation of parameter values over the course of each run. Individual simulations were run for 10 million updates or more. For each of the demographic parameters we recorded the marginal density. The peaks of the resulting distributions were taken as estimates of the parameters and the 90% highest posterior density (HPD) interval was taken for the credibility intervals. Estimates with IM are scaled by the overall neutral mutation rate per gene per generation. We assigned an inheritance scalar of 0.25 as usual for mtDNA and assumed 1 year per generation. To convert parameter estimates to time scale units, we used the average rate of evolution for the *cyt-b* gene in *Crocidura* species from

Dubey *et al.* (2006), corresponding to  $3.07E-5$  (95% CI:  $2.37E-5$ – $3.77E-5$ ) mutation events/locus/year.

### Expansion time

To test the hypothesis of recent population growth from a low-diversity founder population within the different clades, several tests were performed for *cyt-b*. We used three methods implemented in Arlequin version 2.0 (Schneider, 2000). The first method, Fu's (1997)  $F_S$  statistic, tests the probability of having no fewer than the number of observed alleles in the sample given that  $\theta$  (heterozygosity per sites) =  $\pi$ . This statistic tends to be negative when there is an excess of recent mutations (or rare alleles). The second method, Tajima's (1989)  $D$  statistic, tests the null hypothesis that two estimates of the neutral mutation parameter, one derived from the average number of pairwise nucleotide differences and the other based on the number of segregating sites in the sample, are equal. In the third test, pairwise mismatch distributions among individuals were plotted and tested for goodness-of-fit to a model of sudden expansion using parametric bootstrapping with 1000 replicates (Schneider & Excoffier, 1999). Expansion time after the bottleneck was estimated from the mismatch distribution ( $\tau$ ) (Rogers, 1995) and uncorrected distances ( $p$ ). Evolutionary rate for uncorrected ( $p$ ) distance was estimated using the molecular clock developed by Fumagalli *et al.* (1999) with a divergence rate of  $0.061$  uncorrected distance ( $p$ )  $\text{Myr}^{-1}$  (95% CI:  $0.054$ – $0.069$ ).

## Results


### Cytochrome *b* gene

The 60 *C. leucodon* samples showed 41 different haplotypes of 1077 bp and contained 366 variable sites, of which 218 were parsimony-informative. No insertions or deletions were observed. As the three phylogenetic methods gave identical arrangements of the main branches, the relationship between haplotypes is given only for the ML analysis in Fig. 2.

The *C. leucodon* samples formed a monophyletic unit (all bootstrap values of 100%), well differentiated from the other Eurasian species (Fig. 2). Within this unit, two major and strongly supported clades were found (all bootstrap values  $\geq 80\%$ ).

Clade I (western clade) included European samples from the Czech Republic, France, Germany, Greece (Lesbos), Italy, Serbia, Slovakia, Slovenia, Switzerland, and samples from north-western Anatolia, Turkey. Clade II (eastern clade) included the European samples from Bulgaria and Romania and the samples from Georgia and Anatolia (with the exception of westernmost Anatolia).

AMOVA showed that the majority of mtDNA variation (84.09%) is distributed between the two clades of *C. leucodon*; only a small percentage of this variation



**Fig. 2** Phylogeny of the 1077 bp *cyt-b* fragment analysed with maximum likelihood, using the general time reversible model of substitution. Values in branches are indices of support for the major branches for maximum likelihood (ML), maximum parsimony (MP) and distance (NJ) analyses (percentage of 1000 replications). Codes are as in Table 1.

(15.91%) is observed within clades. Mean pairwise GTR distance between clades is 3.94%. The mean pairwise GTR distances within clades and nucleotide diversities were 0.65% and 0.057 for the western clade, respectively, and 0.71% and 0.059 for the eastern clade.

Three additional sequences from Lesbos, Stymfalia, and Parnitha (Greece) described by Poulakakis *et al.* (2005) belong to the western clade, but were not included in our phylogenetic analyses. The analysis of these data (AY452166, AY452176, AY452165) revealed an unexpected result. The mean GTR distance between our samples and the Poulakakis *et al.* samples from Lesbos was 2.8%. In comparing the sequences, we observed a large number of mutations at the beginning and end of their *cyt-b* sequence. When the last 13 bp of their sequence were deleted, the mean GTR distance fell to 0.89%, and this was further reduced to 0.42% by deleting the first 48 bp of their sequence (see Table 2 for the details of the mutations observed at the beginning and the end of the sequences of Poulakakis *et al.*). In addition, the mean GTR distance between the

**Table 2** Example of mutations observed between some of our samples and the samples of Poulakakis *et al.* (2005; \*Lesvos, AY452165; Parnitha, AY452166; Stymfalia, AY452176).

Samples	Example of mutations observed between sequences																			
SK2, Bratislava	TC	TGC	TTA	ATT	GCA	CAA	ATC	CTA	ACA	GGA	TTA	TTC	CTA	GCC	ATA	CAC	TAT	ACA	[53]	
IT5, Serramazoni	..	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[53]
TR10, Katranci	..	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[53]
GR2, Lesvos	..	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	..T	[53]
GR1, Lesvos	..	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[53]
Lesvos*	NN	NNN	NNN	NNN	NNN	...	...	...	...	...	...	..T	...	...	...	...	...	...	..T	[53]
Parnitha*	NN	NNN	NNN	NNN	NNN	...	...	..G	...	...	...	..T	...	...	...	...	...	...	...	[53]
Stymfalia*	..	...	...	...	...	...	...	..T	...	...	C..	..T	...	...	...	...	...	...	...	[53]
TR6, Altindere	..	...	...	...	...	...	...	T..	...	...	C..	...	T..	...	...	...	...	...	...	[53]
GE1, Alazani	..	...	...	...	...	...	...	T..	...	...	C..	...	...	...	...	...	...	...	...	[53]
SK2, Bratislava	TCT	GAT	ACT	ACA	ACA	GCT	TTC	TCC	TCC	GTA	ACC	CAT	ATT	TGC	CGA	GAT	GTA	AAT	[107]	
IT5, Serramazoni	..A	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[107]
TR10, Katranci	..A	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[107]
GR2, Lesvos	..A	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[107]
GR1, Lesvos	..A	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[107]
Lesvos*	..A	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[107]
Parnitha*	..A	...	..C	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[107]
Stymfalia*	..A	...	..C	...	...	...	...	...	...	...	...	...	...	..T	...	...	...	...	...	[107]
TR6, Altindere	..A	...	...	...	...	...	...	...	...	..A	...	...	...	...	...	...	...	...	...	[107]
GE1, Alazani	..A	...	...	...	...	...	...	...	...	..A	...	...	...	...	...	...	...	...	...	[107]
SK2, Bratislava	TA	GGA	TAT	GTT	CTT	CCC	TGA	GGT	CAA	ATA	TCA	TTT	TGA	GGT	GCA	ACA	GTA	AT	[332]	
IT5, Serramazoni	..	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[332]
TR10, Katranci	..	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[332]
GR2, Lesvos	..	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[332]
GR1, Lesvos	..	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[332]
Lesvos*	..	...	...	...	...	...	..A	...	...	...	..C	...	..G	..T	G..	NNN	NN	...	[332]	
Parnitha*	..	...	...	...	...	...	..A	...	...	...	..C	...	..G	..T	G..	NNN	NN	...	[332]	
Stymfalia*	..	...	...	...	...	...	..A	...	...	...	..C	...	..G	..T	G..	AGT	T.	...	[332]	
TR6, Altindere	..	..G	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[332]
GE1, Alazani	..	..G	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	[332]

Only the mutations including the first 107 bp and the last 52 bp of the Poulakakis sequences are shown. The samples SK2, IT5, TR10, GR1 and GR2 (this study) belong to the western mitochondrial clade, and the samples TR6 and GE1 from the eastern mitochondrial clade.


Poulakakis *et al.* samples from Lesvos and those from continental Greece was 2.15%, whereas the mean distance between our samples from Lesbos and the Poulakakis *et al.* samples was 6.22%; this value is still much greater than the mean GTR distance between all of our sequences from this clade (0.65%) and the mean GTR distance between the two major clades (western and eastern) obtained in our analyses (3.94%, Fig. 2).

A likelihood ratio test led to the acceptance of the molecular clock hypothesis for the whole sample ( $\chi^2_{43} = 53.44$ ,  $P = 0.13$ ) –Ln Likelihood values with and without the molecular clock assumption are 4169.79 and 4143.07, respectively, for the best trees. On the basis of the calibration of Fumagalli *et al.* (1999) and Dubey *et al.* (2006), we estimated the divergence time between the western and eastern clade to be 0.691 Myr (95% CI: 0.510–0.980; Fig. 2).

The IM model has six demographic parameters but we were particularly interested in three of them:  $t$ , the time of population splitting (in generation) in the past, and  $m1$  and  $m2$ , the gene-flow rates per gene copy per generation. Once priors were correctly defined, we ran three

runs for each clade. The different runs revealed unambiguous marginal posterior probability distributions of the parameters in each clade. The migration parameters revealed a peak at the lower limit of resolution in both directions (from East to West and vice versa) in both clades, as expected given that the Marmora Sea is a non-negotiable barrier to dispersal once filled with water (during interglacials). We hereafter interpret the locations of these peaks as being at zero and then simulated three more runs for each clade fixing  $m1$  and  $m2$  to zero for estimating the time of the splits. The marginal posterior probability distribution of  $t$  revealed a sharp peak at 0.525 and 1.785 for the Western and the Eastern clade respectively (Fig. 3). When converted to a scale of years, the divergence times between both sides of the Marmara Sea were estimated to be 17 100 years (90% HPD interval: 1700–40 600) for the Western clade, and 58 200 years (90% HPD interval: 16 100–109 000) for the Eastern one.

We observed a nonsignificant  $P$ -value for the mismatch distribution test of goodness-of-fit for the western clade (*Harpending's Raggedness index* = 0.02,  $P = 0.28$ )



**Fig. 3** Marginal posterior probability ( $p$ ) distribution of the divergence time parameter ( $t$ ) in year, from the isolation with migration model, between populations situated western and eastern of the Bosphorus Strait within the Western and Eastern Clade, and estimated time for the splits with the 90% highest posterior density (HPD) interval.

and a significant  $P$ -value for the eastern clade (*Harpending's Raggedness index* = 0.12,  $P = 0.03$ ; Table 2). The frequency of the mean pairwise difference between haplotypes showed a bell-shaped distribution for the western clade, contrasting with the more complex distribution obtained for the eastern clade (Fig. 1).  $F_S$  statistics and Tajima's  $D$  were significant ( $F_S = -10.68$ ,  $P < 0.001$ ;  $D = -1.48$ ,  $P = 0.04$ ; Table 3) for the western clade but not significant for the eastern ( $F_S = -2.02$ ,  $P = 0.16$ ;  $D = -0.67$ ,  $P = 0.26$ ); hence, we inferred a scenario of expansion for the western population and nonexpansion for the eastern population.

The timing of expansions was estimated from the mismatch distribution according to the method proposed

by Rogers (1995; Fig. 1b,c). The  $\tau$  value for the western population was 6.71 (95% CI: 4.19–8.39). Assuming no saturation of uncorrected distances ( $p$ ), as shown in Dubey *et al.* (2006), the distance was 0.0612 per million years (95% CI: 0.054–0.069). With a generation time of 1 year, the population expansion time was estimated to be 51 900 years (95% CI: 28 100–72 100).

### Apolipoprotein B gene

The 44 analysed samples showed four different *ApoB* alleles of 511 bp, named A1–A4 (Genbank accession: EF011555–EF011558), all the alleles differing from each others by only one mutation at the positions 62, 96 and 144 of our sequence alignment. Consequently, the different alleles of heterozygous samples were easily determined, as only one mutation was observed between alleles. The western mitochondrial clade samples were all homozygous for the allele A1 (Table 1). In contrast, 13 A1 homozygotes, two A1/A2 compound heterozygotes, one A1/A3 compound heterozygote, one A1/A4 compound heterozygote and one A4 homozygote were found in the eastern mitochondrial clade. This means that nucleotide diversities were 0.0000 for the western clade and 0.0006 for the eastern clade.

## Discussion

### Quality of samples

The integration of our *cyt-b* sequences with those of Ohdachi *et al.* (2004); Fontanillas *et al.* (2005) and Dubey *et al.* (2006), analysed from preserved tissues, showed the expected genetic distances between samples. In contrast, the material of *C. leucodon* from owl pellets analysed by Poulakakis *et al.* (2005) showed GTR distances with our *C. leucodon* samples that were much higher than expected, even between samples from the same island (Lesbos; mean GTR distance: 2.8%). A distance that fell to 0.42% by deleting the first and the last bp of their sequence from Lesbos. Moreover, no nonsynonymous mutations were observed in the *cyt-b* gene for our *C. leucodon* samples, whereas mutations at the end of the Poulakakis *et al.* (2005) sequences resulted in two amino acid substitu-

**Table 3** Number of samples (Ns) and number of haplotypes (Nh) within the western and eastern mitochondrial bicoloured shrew clades, genetic diversity, nucleotide diversity, mean pairwise differences between haplotypes, Goodness-of-fit test probability, Fu's  $F_S$  test probability, Tajima's  $D$  test probability, estimated  $\tau$  value and expansion time for western clade.

Clade (Ns, Nh)	Nucleotide div./ mean pairwise diff.	Goodness-of-fit test		Fu's $F_S$ test		Tajima's test			Expansion time
		Rag. Index	$P$	$F_S$	$P$	Tajima's $D$	$P$	$\tau$	
Western (34, 23)	0.0057/6.21	0.02	0.28	-10.68	< 0.001	-1.48	0.04	6.80 (95% CI: 4.19–8.39)	51 900 years BP (95% CI: 28 100–72 100)
Eastern (16, 11)	0.0059/6.31	0.12	0.03	-2.02	0.16	-0.67	0.26	7.47 (95% CI: 3.71–10.85)	–

tions; thus, these mutations appear to be highly suspect. These mutations may in fact be artefacts of DNA sequencing as a result of the poor template quality obtained from the owl pellets (Taberlet & Fumagalli, 1996; Waits & Paetkau, 2005). This reinforces the value of repeating the DNA extraction and/or the analysis for each noninvasive sample several times in order to validate the sequencing result (Waits & Paetkau, 2005). This means that the affirmation of Poulakakis *et al.* (2005) concerning the use of their samples for phylogenetic and phylogeographical studies of small mammals has to be questioned.

### Biogeography

Based on our mitochondrial phylogenetic analyses, *C. leucodon* populations are divided in two main mitochondrial clades (mean GTR distance between clades: 3.94%; Figs 1a and 2). The first clade (western clade) includes samples from western and central Europe, and unexpectedly, two samples from Lesbos Island and three from north-western Anatolia, to which the Greek samples (Lesbos, Parnitha and Stymfalia) from Poulakakis *et al.* (2005; results not shown) should be added. The second clade (eastern clade) includes samples from Bulgaria, Romania, Anatolia (with the exception of westernmost Anatolia) and Georgia. These two clades could represent the chromosomal differences observed between the Georgian samples and those from the Czech Republic and Lesbos that were detected in the karyotype analyses undertaken by Biltueva *et al.* (2001).

The separation between these two mitochondrial lineages of *C. leucodon* occurred in the Middle Pleistocene, 0.691 Myr (95% CI: 0.510–0.980), the period immediately following the Günz glacial events (790 000–950 000 years BP). This suggests the isolation of populations by submergence of the Bosphorus Strait (between the Black Sea and the Marmora Sea) with increasing sea levels following the glacial period.

Sedimentological and palaeontological evidence reveals that the Bosphorus Strait has alternatively submerged and emerged since the Middle Pleistocene before being completely submerged from the Mid-Late Holocene (Kerey *et al.*, 2004). Consequently, the fact that we observed a lack of clear structure between samples situated on either side of the Bosphorus Strait indicates that south-eastern Europe was probably colonized by the eastern mitochondrial lineages during a recent land bridge connection between Europe and the Near East (Late Pleistocene), a hypothesis confirmed by the migration analyses that estimated this event of vicariance to the Upper Pleistocene (58 200 years BP; 90% HPD: 16 100–109 000). Based on similar results, the colonization of western Anatolia by the western mitochondrial lineage occurred during the same period (17 100 years BP; 90% HPD: 1700–40 600; Late Pleistocene–Holocene). Thus, this strait appears to be a permeable biogeographic

barrier for *C. leucodon*. This permeability has already been demonstrated in classical zoogeography for various species (e.g. Hosey, 1982) on the basis of unidirectional migrations from Europe to Anatolia (Hosey, 1982; Filippucci & Simson, 1996; Kryštufek, 2002). However, our study demonstrates for the first time a Late Pleistocene–Holocene bidirectional exchange of two different conspecific lineages between Europe and the Near East, whereas previous studies have only revealed unidirectional colonization.

The last climatic fluctuations of the Upper Pleistocene (126 000–11 500 years BP; Ogg, 2004) had contrasting impacts on the western and eastern mitochondrial clades. For the western clade, the bell-shaped curve of the mismatch distributions (Fig. 1b) of the *cyt-b* gene 0 indicates an expansion following a bottleneck 50 800 years ago (95% CI: 28 100–72 100). This finding reveals that the last glacial maximum of the Pleistocene (22 000 years BP) had a moderate impact on these small mammal populations. Conversely, no sign of expansion was detected for the eastern mitochondrial clade of *C. leucodon*, which is consistent with the absence of nuclear polymorphism in the western clade ( $\pi$ : 0.0000), whereas four alleles are present in the eastern ( $\pi$ : 0.0006).

Thus, the European population appears to have been reduced to a small number of individuals during the last glaciations, probably confined to a small Italo-Balkan refugium. Conversely, the eastern clade appears to have persisted with a greater population size and probably in several refugia around the Black Sea. This pattern has also been suggested for mammals and plants (Michaux *et al.*, 2004; Heuertz *et al.*, 2006; Kučera *et al.*, 2006), based on a higher genetic diversity observed in Anatolian populations compared with European ones. Moreover, these results are also supported by palynological data that indicate that open wooded cover in this area persisted through the full glacial condition of the Pleistocene (Tarasov *et al.*, 2000).

### Conclusion

Based on molecular data, we confirmed that the Bosphorus Strait was a permeable barrier. This finding has been suggested previously by the colonization of western Anatolia by European populations of the lesser white-toothed shrew in the Lower Pleistocene (*C. suaveolens*; Dubey *et al.*, 2006) and more recently by the hedgehog (Filippucci & Simson, 1996; Kryštufek, 2002). Moreover, we provide evidence that European populations of the bicoloured shrew may be the source of recent West Anatolian populations.

We also found a marked difference in population history between the two divergent mitochondrial lineages of bicoloured shrew, suggesting that Pleistocene climatic variations have more strongly reduced the genetic diversity of the European population than that

of the Near East population. We hypothesize that this could be a general pattern for fauna and flora, as other studies comparing these geographical areas have reported similar differences (Michaux *et al.*, 2004; Heuertz *et al.*, 2006; Kučera *et al.*, 2006).

## Acknowledgments

We thank N. Di Marco for laboratory facilities, C. Ohayon for scientific and linguistic advice and M. Fontaine for noticeable advice in using IM software. We are grateful to P. Borkenhagen, P. Brunnet-Lecompte, H. Brünner, F. Catzeflis, P. Müller, R. Parapanov, C. Vogel and M. Zaitsev for tissues samples, and to A. Andreeva, S. Uzunov (Bulgaria), C. Kurtonur (Turkey), D. Murariu (Romania) for integrating us (PV) in local fieldwork for trapping shrews. The study was partly sponsored by GACR (grant 206/05/2334) and MSMT CR (project 0021620828).

## References

- Bilton, D.T., Mirol, P.M., Mascheretti, S., Fredga, K., Zima, J. & Searle, J.B. 1998. Mediterranean Europe as an area of endemism for small mammals rather than a source for northwards postglacial colonization. *Proc. R. Soc. Lond., Ser. B Biol. Sci.* **265**: 1219–1226.
- Biltueva, L.S., Rogatcheva, M.B., Perelman, P.L., Borodin, P.M., Oda, S.I., Koyasu, K., Harada, M., Zima, J. & Graphodatsky, A.S. 2001. Chromosomal phylogeny of certain shrews of the genera *Crocidura* and *Suncus* (Insectivora). *J. Zool. Syst. Evol. Res.* **39**: 69–76.
- Brändli, L., Handley, L.J., Vogel, P. & Perrin, N. 2005. Evolutionary history of the greater white-toothed shrew (*Crocidura russula*) inferred from analysis of mtDNA, Y and X chromosome markers. *Mol. Phylogenet. Evol.* **37**: 832–844.
- Cooper, S.J.B., Ibrahim, K.M. & Hewitt, G.M. 1995. Postglacial expansion and genome subdivision in the European Grasshopper *Chorthippus parallelus*. *Mol. Ecol.* **4**: 49–60.
- Culling, M.A., Janko, K., Boron, A., Vasil'ev, V.P., Cote, I.M. & Hewitt, G.M. 2006. European colonization by the spined loach (*Cobitis taenia*) from Ponto-Caspian refugia based on mitochondrial DNA variation. *Mol. Ecol.* **15**: 173–190.
- Dubey, S., Zaitsev, M., Cosson, J.-F., Abdulkadier, A. & Vogel, P. 2006. Pliocene and Pleistocene diversification and multiple refugia in a Eurasian shrew (*Crocidura suaveolens* group). *Mol. Phylogenet. Evol.* **38**: 635–647.
- Dubey, S., Cosson, J.-F., Magnanou, E., Vohralík, V., Benda, P., Frynta, D., Hutterer, R., Vogel, V. & Vogel, P. 2007a. Mediterranean populations of the lesser white-toothed shrew (*Crocidura suaveolens* group): an unexpected puzzle of Pleistocene survivors and prehistoric introductions. *Mol. Ecol.* in press.
- Dubey, S., Salamin, N., Ohdachi, S.D., Barriere, P. & Vogel, P. 2007b. Molecular phylogenetics of shrews (Mammalia: Soricidae) reveal timing of transcontinental colonizations. *Mol. Phylogenet. Evol.* **44**: 126–137.
- Dumolin-Lapegue, S., Demesure, B., Fineschi, S., LeCorre, V. & Petit, R.J. 1997. Phylogeographic structure of white oaks throughout the European continent. *Genetics* **146**: 1475–1487.
- Ferris, C., Oliver, R.P., Davy, A.J. & Hewitt, G.M. 1993. Native oak chloroplasts reveal an ancient divide across Europe. *Mol. Ecol.* **2**: 337–344.
- Ferris, C., King, R.A., Vainola, R. & Hewitt, G.M. 1998. Chloroplast DNA recognizes three refugial sources of European oaks and suggests independent eastern and western immigrations to Finland. *Heredity* **80**: 584–593.
- Filippucci, M.G. & Simson, S. 1996. Allozyme variation and divergence in Erinaceidae (Mammalia, Insectivora). *Isr. J. Zool.* **42**: 335–345.
- Fontanillas, P., Depraz, A., Giorgi, M.S. & Perrin, N. 2005. Nonshivering thermogenesis capacity associated to mitochondrial DNA haplotypes and gender in the greater white-toothed shrew, *Crocidura russula*. *Mol. Ecol.* **14**: 661–670.
- Fu, Y.X. 1997. Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics* **147**: 915–925.
- Fumagalli, L., Taberlet, P., Stewart, D.T., Gielly, L., Hausser, J. & Vogel, P. 1999. Molecular phylogeny and evolution of *Sorex* shrews (Soricidae: Insectivora) inferred from mitochondrial DNA sequence data. *Mol. Phylogenet. Evol.* **11**: 222–235.
- Guindon, S. & Gascuel, O. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* **52**: 692–704.
- Heuertz, M., Carnevale, S., Fineschi, S., Sebastiani, F., Hausman, F., Paule, L. & Vendramin, G.G. 2006. Chloroplast DNA phylogeography of European ashes, *Fraxinus* sp. (Oleaceae): roles of hybridization and life history traits. *Mol. Ecol.* **15**: 2131–2140.
- Hewitt, G.M. 1999. Post-glacial re-colonization of European biota. *Biol. J. Linn. Soc.* **68**: 87–112.
- Hewitt, G.M. 2000. The genetic legacy of the quaternary ice ages. *Nature* **405**: 907–913.
- Hewitt, G.M. 2004a. Genetic consequences of climatic oscillations in the Quaternary. *Phil. Trans. R. Soc. Lond. Ser. B.* **359**: 183–195.
- Hewitt, G.M. 2004b. The structure of biodiversity – insights from molecular phylogeography. *Front. Zool.* **1**: 4.
- Hey, J. 2007. *Using the IM program*, IM Documentation. [http://lifesci.rutgers.edu/~hey/lab/ProgramsandData/Programs/IM/Using\\_IM\\_3\\_5\\_2007.pdf](http://lifesci.rutgers.edu/~hey/lab/ProgramsandData/Programs/IM/Using_IM_3_5_2007.pdf).
- Hosey, G.R. 1982. The Bosphorus land-bridge and mammal distributions in Asia Minor and the Balkans. *Säugetierkd. Mitt.* **30**: 53–62.
- Irwin, D.M., Kocher, T.D. & Wilson, A.C. 1991. Evolution of the cytochrome *b* gene of Mammals. *J. Mol. Evol.* **32**: 128–144.
- Kerey, I.E., Meric, E., Kelling, G., Brenner, R.A. & Dogan, A.U. 2004. Black Sea-Marmara Sea Quaternary connections: new data from the Bosphorus, Istanbul, Turkey. *Paleogeogr. Paleoclimatol.* **204**: 277–295.
- Koch, M.A., Kiefer, C., Ehrlich, D., Vogel, J., Brochmann, C. & Mummenhoff, K. 2006. Three times out of Asia Minor: the phylogeography of *Arabis alpina* L. (Brassicaceae). *Mol. Ecol.* **15**: 825–839.
- Kryštufek, B. 2002. Cranial variability in the eastern hedgehog *Erinaceus concolor* (Mammalia: Insectivora). *J. Zool. (London)* **258**: 365–373.
- Kučera, J., Lihová, J. & Marhold, K. 2006. Taxonomy and phylogeography of *Cardamine impatiens* and *C. pectinata* (Brassicaceae). *Bot. J. Linn. Soc.* **152**: 169–195.
- Marmi, J., Lopez-Giraldez, F., MacDonald, D.W., Calafell, F., Zholnerovskaya, E. & Domingo-Roura, X. 2006. Mitochond-

- rial DNA reveals a strong phylogeographic structure in the badger across Eurasia. *Mol. Ecol.* **15**: 1007–1020.
- Michaux, J.R., Libois, R., Paradis, E. & Filippucci, M.G. 2004. Phylogeographic history of the yellow-necked fieldmouse (*Apodemus flavicollis*) in Europe and in the Near and Middle East. *Mol. Phylogenet. Evol.* **32**: 788–798.
- Nesbo, C.L., Fosshem, T., Vollestad, L.A. & Jakobsen, K.S. 1999. Genetic divergence and phylogeographic relationships among European perch (*Perca fluviatilis*) populations reflect glacial refugia and postglacial colonization. *Mol. Ecol.* **8**: 1387–1404.
- Nielsen, R. & Wakeley, J. 2001. Distinguishing migration from isolation: a Markov chain Monte Carlo approach. *Genetics* **158**: 885–896.
- Ogg, J.G. 2004. Status of divisions of the International Geologic Time Scale. *Lethaia* **37**: 183–199.
- Ohdachi, S.D., Iwasa, M.A., Nesterenko, V.A., Abe, H., Masuda, R. & Haberl, W. 2004. Molecular phylogenetics of *Crocidura* shrews (Insectivora) in east and central Asia. *J. Mammal* **85**: 396–403.
- Palme, A.E. & Vendramin, G.G. 2002. Chloroplast DNA variation, postglacial recolonization and hybridization in hazel, *Corylus avellana*. *Mol. Ecol.* **11**: 1769–1780.
- Parker, S.R., 1997. Sequence Navigator. Multiple sequence alignment software. *Methods Mol. Biol.* **70**: 145–154.
- Posada, D. & Crandall, K.A. 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* **14**: 817–818.
- Poulakakis, N., Lymberakis, P., Paragamian, K. & Mylonas, M. 2005. Isolation and amplification of shrew DNA from barn owl pellets. *Biol. J. Linn. Soc.* **85**: 331–340.
- Rodriguez, R., Oliver, J.L., Marin, A. & Medina, J.R. 1990. The general stochastic model of nucleotide substitution. *J. Theor. Bio.* **142**: 485–501.
- Rogers, A.R. 1995. Genetic-evidence for a Pleistocene population explosion. *Evolution* **49**: 608–615.
- Santucci, F., Emerson, B.C. & Hewitt, G.M. 1998. Mitochondrial DNA phylogeography of European hedgehogs. *Mol. Ecol.* **7**: 1163–1172.
- Schneider, I. 2000. Software programs for DNA sequence analysis. *Genet. Eng. News* **20**: 8.
- Schneider, S. & Excoffier, L. 1999. Estimation of past demographic parameters from the distribution of pairwise differences when the mutation rates vary among sites: Application to human mitochondrial DNA. *Genetics* **152**: 1079–1089.
- Seddon, J.M., Santucci, F., Reeve, N. & Hewitt, G.M. 2002. Caucasus Mountains divide postulated postglacial colonization routes in the white-breasted hedgehog, *Erinaceus concolor*. *J. Evol. Biol.* **15**: 463–467.
- Swofford, D.L. 1998. PAUP\*. Phylogenetic Analysis Using Parsimony (\* and other Methods). Version 4.0b1. Sinauer Associates, Sunderland, MA, USA.
- Taberlet, P. & Fumagalli, L. 1996. Owl pellets as a source of DNA for genetics studies of small mammals. *Mol. Ecol.* **5**: 301–305.
- Taberlet, P., Fumagalli, L., Wust-Saucy, A.G. & Cosson, J.F. 1998. Comparative phylogeography and postglacial colonization routes in Europe. *Mol. Ecol.* **7**: 453–464.
- Tajima, F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* **123**: 585–595.
- Tarasov, P.E., Volkova, V.S., Webb, T., Guiot, J., Andreev, A.A., Bezusko, L.-G., Bezusko, T.V., Bykova, G.V., Dorofeyuk, N.I., Kvavadze, E.V., Osipova, I.M., Panova, N.K. & Sevastyanov, D.V. 2000. Last glacial maximum biomes reconstructed from pollen and plant macrofossil data from northern Eurasia. *J. Biogeogr.* **27**: 609–620.
- Thorpe, R.S. 1984. Primary and secondary transition zones in speciation and population differentiation – a phylogenetic analysis of range expansion. *Evolution* **38**: 233–243.
- Vogel, P., Cosson, J.F. & Jurado, L.F.L. 2003. Taxonomic status and origin of the shrews (Soricidae) from the Canary Islands inferred from a mtDNA comparison with the European *Crocidura* species. *Mol. Phylogenet. Evol.* **27**: 271–282.
- Vörös, J., Alcobendas, M., Martinez-Solano, I. & Garcia-Paris, M. 2006. Evolution of *Bombina bombina* and *Bombina variegata* (Anura: Discoglossidae) in the Carpathian Basin: a history of repeated mt-DNA introgression across species. *Mol. Phylogenet. Evol.* **38**: 705–718.
- Waits, L.P. & Paetkau, D. 2005. Noninvasive genetic sampling tools for wildlife biologists: a review of applications and recommendations for accurate data collection. *J. Wildl. Manage.* **69**: 1419–1433.

Received 14 February 2007; revised 28 April 2007; accepted 2 May 2007