

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Are HDL receptors really located where we think they are in the liver?

Authors: Rützi S, Widmann C

Journal: Current opinion in lipidology

Year: 2016 Aug

Volume: 27

Issue: 4

Pages: 424-5

DOI: 10.1097/MOL.0000000000000322

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

Are HDL receptors really located where we think they are in the liver?

Sabine Rütli and Christian Widmann

Department of Physiology, Lausanne University, Switzerland.

Word counts: 509

Address correspondence: Christian Widmann, Department of Physiology, Rue du Bugnon 7, 1005 Lausanne, Switzerland, Phone: +41 21 692 5123, Fax: +41 21 692 5505, E-mail: Christian.Widmann@unil.ch.

The central role played by high density lipoproteins (HDLs) in reverse cholesterol transport has been studied in details over the past years. One of the many functions of HDLs is to facilitate the uptake of cholesterol from cells of peripheral organs and to deliver it to the liver where it is transferred to the bile for excretion. Scavenger receptor class B type 1 (SRB1), which is abundantly expressed in the liver, participates in the selective hepatic cholesterol uptake from mature HDL particles [1]. A second recently discovered pathway involves HDL particle uptake by the P2Y13 purinergic receptor [2]. How SRB1 binds to HDLs and unload its cholesterol cargo remains to be precisely determined, as different mechanisms have been proposed, implicating or not HDL particle endocytosis [3, 4].

The experiments studying the implication of SRB1 in reverse cholesterol transport have so far always considered that this occurs in hepatocytes. In our current understanding of reverse cholesterol transport, liver sinusoidal endothelial cells (LSECs) act as a hedgerow separating blood circulation from the space of Disse (the extracellular space located between endothelial cells and hepatocytes), allowing HDLs to passively move from the circulation to the hepatocytes where their cholesterol cargo is processed [5]. Given the known high capacity of LSECs to scavenge many different particles, Ganesan and colleagues evaluated the participation of these cells in HDL metabolism in a study recently published in Scientific Reports [6]. Using two different approaches (high-resolution confocal microscopy of ultrathin section of mouse liver and flow cytometry of purified cell populations), they convincingly conclude that LSECs express substantially more SRB1 than hepatocytes. They explain the discrepancy between their findings and what has been described so far in the literature by technical limitations of earlier approaches.

If hepatocytes really express almost no SRB1, how do HDLs enter hepatocytes to be metabolized and how is cholesterol eventually secreted into the bile? Different routes are possible. Ganesan and colleagues do not exclude that a very small amount of

SRB1, not detectable with current technologies, is present on hepatocytes and mediates the interaction with HDLs. Alternatively, other HDL receptors might mediate the uptake of HDLs in hepatocytes, such as CD36 or P2Y13 [2, 7]. Another possibility is that HDLs captured by the SRB1-bearing LSECs is made available to the hepatocytes by a mechanism involving transcytosis. Similar mechanisms have been observed in other endothelial cells [8, 9]. Whether HDLs that bind to SRB1 expressed on LSECs transcytose as intact molecules or whether unloading of cholesterol already happens in LSECs and only cholesterol is then transferred to hepatocytes remains to be investigated.

Mice and humans are quite different in their lipoprotein metabolism [10]. In mice, cholesterol is mainly transported by HDLs whereas in humans LDLs fulfil this function. This difference comes from the presence of the cholesterol ester transfer protein, present in humans and absent in mice, which allows the transfer of cholesterol from HDLs to LDLs. These species differences highlight the importance of conducting a study on human samples similar to the one mentioned here on SRB1 expression in hepatocytes and LSECs.

Acknowledgements

None

Financial support

C.W. is supported by grants from the Swiss National Science Foundation (no. 31003A_160181/1, CRSII3_154420, and IZLSZ3_148907/1).

Conflicts of interest

There are no conflicts of interest.

REFERENCES

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

[1] Kent AP, Stylianou IM. Scavenger receptor class B member 1 protein: hepatic regulation and its effects on lipids, reverse cholesterol transport, and atherosclerosis. *Hepatic medicine : evidence and research* 2011; 3:29-44.

[2] Fabre AC, Malaval C, Ben Addi A *et al.* P2Y13 receptor is critical for reverse cholesterol transport. *Hepatology* 2010; 52:1477-1483.

[3] Trigatti BL, Krieger M, Rigotti A. Influence of the HDL receptor SR-BI on lipoprotein metabolism and atherosclerosis. *Arteriosclerosis, thrombosis, and vascular biology* 2003; 23:1732-1738.

[4] Silver DL, Wang N, Xiao X, Tall AR. High density lipoprotein (HDL) particle uptake mediated by scavenger receptor class B type 1 results in selective sorting of HDL cholesterol from protein and polarized cholesterol secretion. *The Journal of biological chemistry* 2001; 276:25287-25293.

[5] Sorensen KK, Simon-Santamaria J, McCuskey RS, Smedsrod B. Liver Sinusoidal Endothelial Cells. *Comprehensive Physiology* 2015; 5:1751-1774.

[6] • Ganesan LP, Mates JM, Cheplowitz AM *et al.* Scavenger receptor B1, the HDL receptor, is expressed abundantly in liver sinusoidal endothelial cells. *Scientific reports* 2016; 6:20646.

The authors of this study provide compelling evidence that SRB1 is more abundantly expressed in liver sinusoidal endothelial cells than in hepatocytes, challenging the current view of how HDL particles are captured by the liver.

[7] Brundert M, Heeren J, Merkel M *et al.* Scavenger receptor CD36 mediates uptake of high density lipoproteins in mice and by cultured cells. *Journal of lipid research* 2011; 52:745-758.

[8] Rohrer L, Ohnsorg PM, Lehner M *et al.* High-density lipoprotein transport through aortic endothelial cells involves scavenger receptor BI and ATP-binding cassette transporter G1. *Circulation research* 2009; 104:1142-1150.

[9] Lim HY, Thiam CH, Yeo KP *et al.* Lymphatic vessels are essential for the removal of cholesterol from peripheral tissues by SR-BI-mediated transport of HDL. *Cell metabolism* 2013; 17:671-684.

[10] • Lee-Rueckert M, Escola-Gil JC, Kovanen PT. HDL functionality in reverse cholesterol transport - Challenges in translating data emerging from mouse models to human disease. *Biochimica et biophysica acta* 2016; 1861:566-583.

This review provides a detailed up-to-date comparison and critical analysis of the various steps of reverse cholesterol transport in mouse and man.