



Disponible en ligne sur

ScienceDirect  
www.sciencedirect.com

Elsevier Masson France

EM|consulte  
www.em-consulte.com



Original article

# A rare condition: Spontaneous subarachnoid haemorrhage due to spinal hemangioblastoma: Report of 2 cases and review of the literature



C. Tuleasca<sup>a,b,c,d,e,\*</sup>, S. Knafo<sup>e</sup>, P. David<sup>e,i</sup>, S. Richard<sup>f,g,i</sup>, C. Adam<sup>h,i</sup>, N. Aghakhani<sup>e,i</sup>, F. Parker<sup>e,i</sup>

<sup>a</sup> Department of clinical neurosciences, Neurosurgery service and Gamma knife centre, Lausanne university hospital (CHUV), Lausanne, Switzerland

<sup>b</sup> Faculty of biology and medicine (FBM), university of Lausanne (Unil), Lausanne, Switzerland

<sup>c</sup> Signal processing laboratory (LTS 5), École polytechnique fédérale de Lausanne (EPFL), Lausanne, Switzerland

<sup>d</sup> Faculté de médecine, Sorbonne université, Paris, France

<sup>e</sup> Service de neurochirurgie, Assistance publique–Hôpitaux de Paris, hôpitaux universitaires Paris-Sud, centre hospitalier universitaire de Bicêtre, Paris, France

<sup>f</sup> Génétique oncologique EPHE, PSL research university, Paris, France

<sup>g</sup> Faculté de médecine Paris-Sud, université Paris-Saclay, Inserm UMR 1186, Gustave-Roussy, Villejuif, France

<sup>h</sup> Laboratoire de neuropathologie, GHU Paris-Sud – hôpital Bicêtre, Le Kremlin-Bicêtre, France

<sup>i</sup> PREDIR National rare adult cancer reference network: INCA, Assistance publique–Hôpitaux de Paris, GHU Paris-Sud – hôpital Bicêtre, Le Kremlin-Bicêtre, France

## ARTICLE INFO

### Article history:

Received 19 March 2020

Received in revised form 13 May 2020

Accepted 24 May 2020

Available online 28 August 2020

### Keywords:

Hemangioblastomas

Subarachnoid haemorrhage

Spinal

Von Hippel–Lindau

## ABSTRACT

**Introduction.** – Subarachnoid haemorrhage (SAH), secondary to spinal hemangioblastoma (HBL), is extremely rare, with only a few case reports to date. We report the experience of our reference centre for spinal tumours and Von Hippel–Lindau (VHL) disease in patients with spinal HBL presenting with SAH. We further performed a systematic review of the literature.

**Methods.** – We report two cases. A systematic search was performed using the PubMed, Embase and Cochrane databases, with no limit for publication date. Inclusion criteria were: patients with HBL presenting with SAH, with or without VHL. The systematic review retrieved only 10 studies, including 16 patients.

**Results.** – In our centre, the first case concerned radicular HBL at D12 level, presenting with spinal and brain SAH. The patient underwent uneventful microsurgical *en bloc* resection. Postoperative course was normal. The second case concerned HBL with SAH at the cervico-medullary junction, with rapidly fatal course. The systematic review revealed female predominance, at a median age of 40 years, with HBL predominantly located at cervical level, common preoperative symptoms being headache and signs of meningeal irritation.

**Conclusions.** – In conclusion, spinal HBL is an extremely rare cause of SAH. The systematic review found putative risk factors: female gender, age 40–50 years, cervical location, and median size 2 cm. Diagnosis can be difficult when presentation mimics intracerebral SAH. We advocate early surgical removal. The risk of rapidly fatal course, in case of major haemorrhage, needs to be borne in mind.

© 2020 The Authors. Published by Elsevier Masson SAS. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Hemangioblastomas (HBL) are highly vascularised lesions of the central nervous system (CNS), most frequently within the

cerebellum and spinal cord [1–3]. They account for approximately 2–3% of primary spinal cord tumours [4]. They are considered benign (World Health Organisation grade I) [5]. However, they can lead to significant morbidity and mortality through their mass effect, with specific clinical manifestation depending on the anatomical location [6]. In most cases, HBL is sporadic (70 to 80% [7]), although 20–30% of cases are related to a dominantly inherited genetic familial cancer syndrome known as von Hippel–Lindau (VHL) [6,8]. Thirteen per cent of HBLs occur in the spinal canal [9].

\* Corresponding author at: Centre hospitalier universitaire de Lille, hôpital Roger-Salengro; centre hospitalier universitaire de Vaudois, Neurosurgery Service and Gamma Knife Centre, rue du Bugnon 44–46, BH-08, CH-1011 Lausanne, Switzerland.

E-mail addresses: [constantin.tuleasca@gmail.com](mailto:constantin.tuleasca@gmail.com), [constantin.tuleasca@chuv.ch](mailto:constantin.tuleasca@chuv.ch) (C. Tuleasca).

Spinal diseases are an extremely rare cause of spontaneous sub-arachnoid haemorrhage (SAH), accounting for less than 1% of all non-traumatic cases [10]. Haemorrhage secondary to spinal HBL is extremely rare, with currently only a few reports.

The present report describes our experience as a reference centre for spinal tumours and VHL disease, regarding HBL presenting with SAH. We detail clinical course, surgical management (if such was the case) and postoperative assessment. We further performed a systematic review of the literature for publications describing patients with HBL presenting with SAH, with or without surgical exploration, together with clinical and radiological outcomes. The purpose was to investigate potential factors that could help distinguish this rare clinical entity.

## 2. Methods

### 2.1. Patient population and selection

This was a retrospective, single-centre observational study. All patients had been diagnosed and treated in our reference centre for spinal neoplasms, vascular disorders and VHL disease (CHU de Bicêtre, France), over a period of 30 years. The local review board of the Bicêtre University Hospital was notified.

Inclusion criteria were: patients harbouring an HBL at the level of the spine and craniocervical junction who presented with SAH confirmed by anatomopathological diagnosis (of the same or another lesion in the context of VHL). Exclusion criteria were: refusal or inability to understand and sign the informed consent form.

After cross matching between the surgeons' (CT, FB) and the pathologist's patients list (CA), 2 cases were included for analysis.

### 2.2. Systematic review of the literature

#### 2.2.1. Literature search

We performed a systematic search using the PubMed, Embase and Cochrane databases, with no limit for dates. The PubMed database was searched using the following word combinations in the "title" or "title/abstract" item: ("hemangioblastoma" AND "haemorrhage" AND "bleeding"), ("hemangioblastoma" AND "bleed" AND "subarachnoid") etc. We also used the Google Scholar and Google search engines to expand our list of studies, including Abstracts; for final analysis, only peer-reviewed papers were included.

#### 2.2.2. Study selection

Inclusion criteria were patients:

- with HBL;
- who presented with SAH;
- notwithstanding their VHL status.

Exclusion criteria were: non-English-language studies and articles reporting other type of haemorrhage (i.e., intralesional).

Titles and Abstracts were screened by two different people (CT, FP). Potentially relevant articles were selected for full-text screening evaluation, independently performed by 3 investigators (CT, NA, FP). Discrepancies were resolved by the corresponding and senior authors (CT, FP).

#### 2.2.3. Data extraction

Ten studies were finally selected, including 16 patients (Table 1). The study flowchart is shown in Supplementary Fig. S1. Study characteristics, publication year, sample size, pre- and postoperative symptoms, and lesion location and size were extracted.

## 3. Results

### 3.1. Case series

#### 3.1.1. Illustrative case 1

A 64 year-old male was diagnosed with VHL syndrome at the age of 46 years, with pancreatic neuroendocrine tumours and renal tumours. He underwent surgery for cerebellar HBL at another institution 18 years previously and resection of a left cerebellar HBL one year before presentation. Multiple spinal HBLs were already diagnosed, at levels C3, C4–5, T4, T12 and L1–L2.

One year after the second posterior fossa surgery, he presented with acute headache, urinary retention and mild progressively reversible paraparesis. Spinal MRI revealed SAH at T12–L1: i.e., the location of a known HBL (Fig. 1A1). Recent MRI scans (18 days and 11 months beforehand) further documented the SAH. Spinal digital subtraction angiography (DSA) revealed the anterior spinal artery at the left T11 level. He underwent a brain CT scan, showing SAH at the sylvian fissure and intraventricular haemorrhage (Fig. 1A2). Brain DSA revealed no arteriovenous malformation (AVM) and no vasospasm. MRI confirmed the presence of the known lesion (Fig. 1B).

In standard D12–L1 laminectomy, dural exposure revealed arachnoid webs spanning both segments around the lesion, together with SAH (Fig. 1C). After arachnoid dissection, no other HBL than the one located along the D12 rootlets was noted. Following dissection of the posterior rootlet attached to the HBL, the artery and veins were coagulated and monobloc resection was performed. Postoperative course was uneventful, and MRI confirmed complete microsurgical resection (Fig. 1D).

#### 3.1.2. Illustrative case 2

A 61 year-old female was diagnosed with VHL syndrome at the age of 23. She had surgical history of pheochromocytoma, and also a pancreatic neuroendocrine tumour and pancreatic and renal cyst. She presented with multiple central nervous system HBLs, including a growing cerebellar HBL for which she had refused surgery, and an HBL at the medulla oblongata (Fig. 2A).

She initially presented with obstructive hydrocephalus, for which she underwent ventriculocisternostomy, and went on to undergo microsurgical resection of the cerebellar HBL. External ventricular drainage was performed postoperatively due to posterior fossa edema and acute hydrocephalus.

She further presented acute spontaneous SAH due to the HBL of the medulla oblongata (Fig. 2B), 17 days after the posterior fossa surgery, clinically manifesting with tetraplegia, respiratory failure and swallowing disorder, revealed during her stay in our intensive care unit.

She died several days later after a multidisciplinary decision to stop further treatment.

### 3.2. Systematic review

The systematic review retrieved 10 case series, including 16 patients. A vast majority were females (female to male sex ratio: 10/6; Table 1). Three had VHL syndrome.

Median age was 39 years (mean 42.9, range 27–71).


The most common symptoms were headache and signs of meningeal irritation.

CT usually confirmed the SAH, the vast majority of which were cervical (9 out of 16). Syrinx was rarely present. Median size was 2 cm (mean 2.6, range 1–5).


When detailed postoperative status was reported, it was normal in most cases, even in patients with major preoperative

**Table 1**  
Systematic review of the literature.

Publication	Number of patients	Age	Sex	Location	Syrinx (yes/no)	Preoperative symptom	Preoperative sign	Preoperative status	Confirmation of SAH	Size (cm)	Postoperative status
Djindjan (1978) [13]	1	30	M	Upper lumbar C1	No	Acute low back pain	Acute low back pain	Coma	Intraoperative	2	–
Kormos (1980) [19]	1	31	F	C1	Yes	Paresthesias left side	Meningeal irritation	Normal	LP	2	Normal
Hille (1986) [20]	1	37	M	C2	No	Left upper limb paresthesias	Meningeal irritation	Normal	LP	4	Normal
Neumann (1989) [6]	2.1.	27	F	C1–C2	No	Headaches	Meningeal irritation	Normal	LP	2	–
	2.2.	53	F	T10–T11	No	Weakness	Motor weakness	Motor deficit	LP	1	–
	2.3.	71	F	C2–C4	No	Brachialgia	Sensory and motor deficit	Sensory and motor deficit	CT	4	–
Cerejo (1990) [10]	1	36	F	C1	No	Headaches, confusion	Meningeal irritation	Normal	CT	1.5	Normal
Yu (1994) [21]	2.1.	31	F	T6–T7	No	Neck pain	Acute lower limb weakness	Acute paraplegia	LP	5	Paraplegia
	2.2. (VHL)	28	M	T7–T8	No	Lower back electric shock like	Lower limb weakness	Paraplegia	Intraoperative/MR	–	Paraplegia
Cervoni (1995) [22]	1	20	F	C3	No	None	paresthesias	Normal	CT	2	Normal
Minami (1998) [23]	1	48	M	C0	No	Severe headache	None	Slight hypoesthesia	CT	1.5	Normal
Irie (1998) [24]	1	41	M	C5–C6	No	Nuchal pain	Nuchal rigidity	Normal	CT	1	Normal
Bertis (2003) [25]	1 (VHL)	47	F	C0–C1	No	Severe headache	Right paresis	Right paresis	CT	1	–
Gläsker (2005) [14]	4.1.	27	F	C1–C2	No	Asymptomatic	None	Normal	CT	3	–
	4.2. (VHL)	47	F	C4–C5	No	Asymptomatic	None	Normal	CT	4.5	–
	4.3.	53	F	T10–T11	No	Motor weakness	Motor weakness	Motor deficit	CT	1.5	–
	4.4.	71	F	C2–C4	No	Brachialgia	Brachialgia	Sensorimotor deficit	CT	4	–
Nishimura (2012) [12]	1 (VHL)	56	M	L2–L3	No	Lumbar pain	–	Transient paraparesis	CT (brain SAH)	3	Normal
						Neck pain					
						Headache					
Current series (2019)	1 (VHL)	64	M	T12	No	Headache	Paraparesis, urinary retention	Transient paraparesis and urinary retention	CT, MRI	1.7	Normal
	2 (VHL)	61	F	C0–C2	No	Motor weakness	Tetraplegia	Tetraplegia	CT	1.5	Dead (no surgery)



**Fig. 1.** Illustrative case 1: A1. Spinal MRI at different time-points before surgery, with subsequent haemorrhage at symptom onset. A2. Brain CT at symptom onset with SAH and intraventricular haemorrhage. B. Preoperative MRI (18 days after symptom onset) including, from left to right, sagittal and axial T2 sequences and sagittal and axial gadolinium-injected sequences. C. Intraoperative images. D. Preoperative images (first two on the left) and postoperative images (last two on the right).



**Fig. 2.** Illustrative case 2: A. Before SAH, and B. after SAH.

symptoms. However, as in our series, severe complications sometimes occurred, including death.

#### 4. Discussion

The present study detailed the course of two patients harbouring HBL, exceptionally associated with SAH. One concerned radicular HBL at D12, and the second was located at the cervico-medullary junction, with rapidly fatal course. The systematic review revealed female predominance (unlike in Von Hippel–Lindau in general, where the ratio is 1/1), a median age of 39 years, with cervical location, and common preoperative symptoms being headache and signs of meningeal irritation.

Owing to their vascular histology, the risk of haemorrhage, whether spontaneous, intraoperative or postoperative, is well

known in HBL. However, the most common symptoms of spinal HBL are related to progressive spinal cord compression [4]. The most common symptom of SAH is sudden headache, while the most common sign is meningeal irritation. For HBL presenting with SAH, spinal cord or root deficits are sometimes observed. When haemorrhage arises in the cervical region, bleeding may extend intracranially, and clinical differential diagnosis versus intracranial SAH may be difficult [11]. Symptom severity usually depends on severity of bleeding. Absence of pre-SAH neurological symptoms and signs makes diagnosis of HBL difficult. To date, only one case of brain SAH in a patient harbouring lumbar HBL was reported [12], and was surprisingly similar to the present case n° 1.

Arteriovenous malformation is the most frequent etiology of spinal SAH, followed by spinal cord tumour (mostly ependymoma), then neurinoma, paraganglioma and dissemination of malignant

spinal tumour [13]. Hemangioblastoma is an extremely rare case of spinal SAH. In a recent study, the risk of spontaneous haemorrhage per person per year in spinal HBL was 0.0024 [14], while the risk of SAH was even lower (0.06%) [15]. The lesion size associated with spontaneous haemorrhage is much greater. Moreover, it is important preoperatively to differentiate HBL from other types of tumour, as treatment strategies differ.

Before the advent of MRI, HBL was diagnosed on digital subtraction angiography (DSA) [16]. MRI is now the gold standard for HBL diagnosis [17]. The classical aspect is a well-circumscribed mass, often associated with a cyst. An associated syrinx was reported in more than 60% of intramedullary lesions [9], but seems rarely associated with HBL presenting with SAH. The SAH might be further explained by the superficial location of medullary HBLs, whereas cerebellar HBLs tend to bleed intraparenchymally [14].

Tumour size is considered by some authors to be implicated in spontaneous and/or postoperative bleeding [14]. In our systematic review, the median size was 2 cm. This might be explained by the high blood flow through these lesions. Some authors postulated pathophysiological mechanisms similar to those in AVM, and that partial transmission of arterial pressure to the venous side, together with the high flow, ultimately causes structural changes that lead to vascular vulnerability [14].

Nowadays, complete removal with low morbidity is possible with microsurgery [18]. Early bipolar coagulation of the nutrient artery of spinal cord HBL is the essential surgical step that facilitates resection and reduces surgical risk.

The present study had several limitations: Firstly, its retrospective nature and small number of cases; and secondly, in the systematic review, some cases may be missing, in spite of our attention, particularly if published as short and undetailed cases within much larger series.

## 5. Conclusion

In conclusion, spinal HBL is an extremely rare cause of SAH, usually sporadic, and rarely occurring in a context of VHL. The systematic review including 16 cases revealed a majority of female patients, in their forties, with cervical lesions with a median size of 2 cm. Haemorrhagic spinal HBL might lead to diagnostic difficulties when presentation mimics intracerebral SAH. We advocate early surgical removal. The possibility of rapidly fatal course in case of major haemorrhage should be borne in mind.

## Ethical statement

The authors confirm that they have read the journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

## Ethical approval

This study was approved by the review board of the Assistance publique–Hôpitaux de Paris. Written informed consent was obtained.

## Funding

Constantin Tuleasca gratefully acknowledges receipt of a “Young Researcher in Clinical Research Grant” (“Jeune Chercheur en Recherche Clinique”) from the University of Lausanne (UNIL), Faculty of Biology and Medicine (FBM) and the Lausanne University Hospital (CHUV).

## Disclosure of interest

Dr Tuleasca is a scientific advisor for Elekta Instruments, AB, Sweden. The present research project did not benefit from other sources of funding with the one exception of the university hospital. The other authors declare that they have no competing interest.

## Acknowledgments

Constantin Tuleasca gratefully acknowledges the University of Lausanne (Unil), Faculty of Biology and Medicine (FBM) for the receipt of a grant “Jeune Chercheur en Recherche Clinique”.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.neuchi.2020.05.008>.

## References

- [1] Cristante L, Herrmann HD. Surgical management of intramedullary hemangioblastoma of the spinal cord. *Acta Neurochir (Wien)* 1999;141:333–9 [discussion 339–40].
- [2] Joaquim AF, et al. Surgical management of intradural extramedullary tumours located anteriorly to the spinal cord. *J Clin Neurosci* 2012;19:1150–3.
- [3] Joaquim AF, et al. Intramedullary hemangioblastomas: surgical results in 16 patients. *Neurosurg Focus* 2015;39:E18.
- [4] Yasargil MG, et al. The microsurgical removal of intramedullary spinal hemangioblastomas. Report of twelve cases and a review of the literature. *Surg Neurol* 1976;3:141–8.
- [5] Richard S, et al. Central nervous system hemangioblastomas, endolymphatic sac tumours, and von Hippel–Lindau disease. *Neurosurg Rev* 2000;23:1–22 [discussion 23–4].
- [6] Neumann HP, et al. Hemangioblastomas of the central nervous system. A 10-year study with special reference to von Hippel–Lindau syndrome. *J Neurosurg* 1989;70:24–30.
- [7] Lonser RR, et al. Prospective natural history study of central nervous system hemangioblastomas in von Hippel–Lindau disease. *J Neurosurg* 2014;120:1055–62.
- [8] Wanebo JE, et al. The natural history of hemangioblastomas of the central nervous system in patients with von Hippel–Lindau disease. *J Neurosurg* 2003;98:82–94.
- [9] Browne TR, Adams RD, Roberson GH. Hemangioblastoma of the spinal cord. Review and report of five cases. *Arch Neurol* 1976;33:435–41.
- [10] Cerejo A, et al. Spinal cord hemangioblastoma with subarachnoid haemorrhage. *Neurosurgery* 1990;27:991–3.
- [11] Luxon LM, Harrison MJ. Subarachnoid haemorrhage and papilledema due to a cervical neurilemmoma. Case report. *J Neurosurg* 1978;48:1015–8.
- [12] Nishimura Y, et al. Intra-extradural dumbbell-shaped hemangioblastoma manifesting as subarachnoid haemorrhage in the cauda equina. *Neurol Med Chir (Tokyo)* 2012;52:659–65.
- [13] Dijindjian M, et al. Subarachnoid haemorrhage due to intraspinal tumours. *Surg Neurol* 1978;9:223–9.
- [14] Glasker S, Van Velthoven V. Risk of haemorrhage in hemangioblastomas of the central nervous system. *Neurosurgery* 2005;57:71–6 [discussion 71–6].
- [15] Gluf WM, Dailey AT. Hemorrhagic intramedullary hemangioblastoma of the cervical spinal cord presenting with acute-onset quadriplegia: case report and review of the literature. *J Spinal Cord Med* 2014;37:791–4.
- [16] Di Chiro G, et al. Digital subtraction angiography and dynamic computed tomography in the evaluation of arteriovenous malformations and hemangioblastomas of the spinal cord. *J Comput Assist Tomogr* 1982;6:655–70.
- [17] Chu BC, et al. MR findings in spinal hemangioblastoma: correlation with symptoms and with angiographic and surgical findings. *AJNR Am J Neuroradiol* 2001;22:206–17.
- [18] Giammattei L, et al. Surgical resection of medulla oblongata hemangioblastomas: outcome and complications. *Acta Neurochir (Wien)* 2016;158:1333–41.
- [19] Kormos RL, Tucker WS, Bilbao JM, Gladstone RM, Bass AG. Subarachnoid hemorrhage due to a spinal cord hemangioblastoma: case report. *Neurosurgery* 1980;6:657–60.
- [20] Van Hille PT, Abbott RJ, Cameron MM, Holland IM, Loizou LA. Recurrent spontaneous subarachnoid haemorrhage due to spinal haemangioblastoma. *J Neurol Neurosurg Psychiatr* 1987;50:639–40.
- [21] Yu JS, Short MP, Schumacher J, Chapman PH, Harsh GR. Intramedullary hemorrhage in spinal cord hemangioblastoma. Report of two cases. *J Neurosurg* 1994;81:937–40.

- [22] Cervoni L, Franco C, Celli P, Fortuna A. Spinal tumors and subarachnoid hemorrhage: pathogenetic and diagnostic aspects in 5 cases. *Neurosurg Rev* 1995;18:159–62.
- [23] Minami M, Hanakita J, Suwa H, Suzui H, Fujita K, Nakamura T. Cervical hemangioblastoma with a past history of subarachnoid hemorrhage. *Surg Neurol* 1998;49:278–81.
- [24] Irie K, Kuyama H, Nagao S. Spinal cord hemangioblastoma presenting with subarachnoid hemorrhage. *Neurologia medico-chirurgica* 1998;38:355–8.
- [25] Berlis A, Schumacher M, Spreer J, Neumann HP, van Velthoven V. Subarachnoid haemorrhage due to cervical spinal cord haemangioblastomas in a patient with von Hippel-Lindau disease. *Acta neurochirurgica* 2003;145:1009–13 [Discussion 1013].