

Serveur Académique Lausannois SERVAL [serval.unil.ch](http://serval.unil.ch)

## Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

**Title:** Altered Sleep Homeostasis in Rev-erb $\alpha$  Knockout Mice.

**Authors:** Mang GM, La Spada F, Emmenegger Y, Chappuis S, Ripperger JA, Albrecht U, Franken P

**Journal:** Sleep

**Year:** 2016 Mar 1

**Issue:** 39

**Volume:** 3

**Pages:** 589-601

**DOI:** 10.5665/sleep.5534

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

# Altered sleep homeostasis in *Rev-erba* knockout mice

Géraldine M. Mang<sup>1</sup> Ph.D., Francesco La Spada<sup>1</sup> Ph.D., Yann Emmenegger<sup>1</sup>, Sylvie Chappuis<sup>2</sup> Ph.D., Jürgen A. Ripperger<sup>2</sup> Ph.D., Urs Albrecht<sup>2§</sup> Ph.D., and Paul Franken<sup>1§</sup> Ph.D.

<sup>1</sup>Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland

<sup>2</sup>Faculty of Science, Department of Biology, Biochemistry, University of Fribourg, Fribourg, Switzerland

<sup>§</sup>Share last authorship

Corresponding author:

*Paul Franken*

*Center for Integrative Genomics*

*University of Lausanne*

*1015 Lausanne, Switzerland*

*e-mail: paul.franken@unil.ch*

*phone/fax: +41 21 6923972 / -65*

## Disclosure

This study was performed at the Universities of Lausanne and Fribourg, Switzerland, and supported by the Swiss National Science Foundation (SNF n°130825 and 136201 to PF and SNF n°146265 to UA) and the states of Vaud supporting GMM, YE, and PF and Fribourg supporting JAR and UA.

## Conflict of Interest

No conflict of interest has to be declared.

## Subtitle

Loss of *Rev-erba* alters sleep homeostasis

## Abstract

**Study objectives:** The nuclear receptor REV-ERB $\alpha$  is a potent, constitutive transcriptional repressor critical for the regulation of key circadian and metabolic genes. Recently, REV-ERB $\alpha$ 's involvement in learning, neurogenesis, mood, and dopamine turnover was demonstrated suggesting a specific role in central nervous system functioning. We have previously shown that the brain expression of several core clock genes, including *Rev-erba*, is modulated by sleep loss. We here test the consequences of a loss of REV-ERB $\alpha$  on the homeostatic regulation of sleep.

**Design:** Sleep recordings during baseline, sleep deprivation, and recovery. Gene expression measurements after sleep deprivation.

**Setting:** Mouse sleep laboratory.

**Participants:** Male *Rev-erba* knockout (KO) mice and their wild type (WT) littermates.

**Intervention:** Sleep deprivation.

**Measurements and Results:** Although baseline sleep/wake duration was remarkably similar, KO mice showed an advance of the sleep/wake distribution relative to the light-dark cycle. After sleep onset in baseline and after sleep deprivation, both EEG delta power (1-4Hz) and sleep consolidation were reduced in KO mice indicating a slower increase of homeostatic sleep need during wakefulness. This slower increase might relate to the smaller increase in theta and gamma power observed in the waking EEG prior to sleep onset under both conditions. Indeed, the increased theta activity during wakefulness predicted delta power in subsequent NREM sleep. Lack of *Rev-erba* increased *Bmal1*, *Npas2*, *Clock*, and *Fabp7* expression, confirming the direct regulation of these genes by REV-ERB $\alpha$  also in the brain.

**Conclusion:** Our results add further proof to the notion that clock genes are involved in sleep homeostasis. Because accumulating evidence directly links REV-ERB $\alpha$  to dopamine signaling the altered homeostatic regulation of sleep reported here are discussed in that context.

**Keywords:** Clock genes, mood, depression, neurogenesis, Process S, slow wave activity.

## Introduction

The timing and quality of sleep are controlled by the interaction of a homeostatic process, that tracks sleep need as a function of the previous sleep/wake history, and a circadian process that ensures the appropriate timing of sleep relative to the daily light-dark alternation <sup>1,2</sup>. Although these two processes seem functionally and neuro-physiologically distinct, at the molecular level, several core components of the circadian timing system were found to also play a role in maintaining proper sleep homeostasis (for review see <sup>3,4</sup>).

The molecular circadian oscillator consists of positive and negative elements. In mammals the positive elements comprise of CLOCK, NPAS2, and BMAL1, with CLOCK/NPAS2:BMAL1 heterodimers driving the transcription of many target genes including that of the *Period* (*Per1, -2*) and *Cryptochrome* (*Cry1, -2*) genes (reviewed in <sup>5</sup>). PER:CRY protein complexes suppress CLOCK/NPAS2:BMAL1-mediated transcription, including their own, thereby constituting the negative elements in this core feedback loop. Additional interactions between these core clock genes at the level of transcription, translocation back into the nucleus, and post-translational modifications add further complexity and stability to the circuit. One important auxiliary feedback loop involves the orphan nuclear receptor REV-ERB $\alpha$ , which binds to specific sequences in the promoters of all three positive elements <sup>6,7</sup> and inhibits their transcription <sup>8-11</sup>. In turn, *Rev-erba* expression is directly regulated by CLOCK/NPAS2:BMAL1-mediated transcription <sup>8,12</sup>. Given REV-ERB $\alpha$ 's ability to reset the phase of the molecular clock oscillation and to entrain central and peripheral clocks, it has been suggested to function as a synchronizing 'hinge' of the clock gene machinery <sup>13</sup> enabling it to act as 'gatekeeper' in coordinating the circadian metabolic response <sup>14</sup>.

The REV-ERB $\alpha$  protein, encoded by the *Nr1d1* (*nuclear receptor subfamily 1, group D, member 1*) gene, regulates gene transcription through binding to specific DNA sequences comprising an A/T-rich flank followed by AGGTCA in the promoter region of target genes <sup>15,16</sup>. REV-ERB $\alpha$  is part of the ligand-binding receptor family, but lacks the carboxy-terminal tail of the ligand-binding domain required for co-activation and thus necessary for transcriptional activation <sup>17</sup>. As a consequence, upon activation by its ligand, heme, REV-ERB $\alpha$  is a constitutive repressor of gene expression <sup>16,18-21</sup>.

Beside its important role in the generation of circadian rhythms, REV-ERB $\alpha$  is implicated in the regulation of various metabolic pathways including adipocyte

differentiation, gluconeogenesis, bile acid synthesis, and heme and cholesterol homeostasis<sup>20,22-24</sup>. Consistent with its function, *Rev-erba* is highly expressed in ~~metabolic~~ tissues with high rates of metabolism such as adipose tissue, liver, skeletal muscle, and brain<sup>25,26</sup>. *Rev-erba* deletion impacts the expression of many metabolic genes, in particular genes involved in lipid metabolic pathways<sup>12</sup>. The administration of REV-ERB $\alpha$  agonists in mice decreased diet-induced obesity by reducing fat content and improving hyperglycemia and dyslipidaemia<sup>27</sup>. The expression of *Rev-erba* can be directly activated through the peroxisome proliferator-activated receptors (PPARs) that play essential roles in energy metabolism and are themselves expressed in a circadian manner (for review see<sup>28</sup>). REV-ERB $\alpha$  is a heme sensor and crucial for heme homeostasis thereby further tightening the link between metabolic and circadian physiology<sup>16,19-21</sup>.

In the context of sleep homeostasis, we have previously shown that the expression of *Rev-erba* in the forebrain of mice is decreased after sleep deprivation<sup>29</sup>. We hypothesize that *Rev-erba* could act as an integrator of both energy demand and sleep pressure. With the aim to establish such role, we evaluated sleep, the EEG, and cortical gene expression under baseline and sleep deprivation conditions, in *Rev-erba* knockout (KO) mice and their wild-type (WT) littermate controls.

## Material & methods

### ***Animals and housing conditions***

*Rev-erba* KO mice were kindly provided by Ueli Schibler (University of Geneva, Geneva, Switzerland) and maintained on a mixed 129/Sv x C57BL6 background. In these mice, exons 3 and 4, encoding the DNA binding domain, and part of exons 2 and 5 of the *Rev-erba* gene, were replaced by an in-frame lacZ allele and a PGK-neo gene by homologous recombination in 129/SV ES cells, resulting in the absence of the transcript and protein<sup>8</sup>. Wild type (WT) littermates were used as control animals. Mice were individually housed in polycarbonate cages (31×18×18 cm) in a temperature and humidity controlled room (25°C, 50-60% respectively) and a 12h light/12h dark cycle (lights on at 9 am, 70-90 lux). Animals had access to food and water *ad libitum*. All experiments were approved by the Ethical Committee of the State of Vaud Veterinary Office, Switzerland.

### ***EEG/EMG implantation***

At the age of 9 to 14 weeks, 13 KO and 9 WT male mice were implanted with EEG and EMG electrodes under deep Xylazine/Ketamine anesthesia as previously described <sup>30</sup>. Briefly, six gold-plated screws (diameter 1.1 mm) were screwed bilaterally into the skull, over the frontal and parietal cortices. Two served as EEG electrodes and the remaining four anchored the electrode connector assembly. As EMG electrodes two gold wires were inserted into the neck musculature. The EEG and EMG electrodes were soldered to a connector and cemented to the skull. Animals were allowed to recover from surgery during 5 - 7 days before they were connected to the recording cables in their home cage. A minimum of 6 days were allowed for habituation to the cable and the experimental room prior to the experiments. Mice were 11 - 16 weeks old at the time of experiment and age did not differ between genotypes (t-test,  $p = 0.33$ ).

### ***Experimental Protocols and Data acquisition***

EEG and EMG signals were recorded continuously for 72 h. The recording started at light onset; i.e., Zeitgeber Time (ZT) 0 or ZT0. During the first 48 h, mice were left undisturbed and these two days were considered as baseline. Starting at ZT0 of day 3, animals were sleep deprived by gentle handling as described previously <sup>31</sup> during 6 hours (ZT0 – 6). The remaining 18 h of day 3 were considered as recovery.

The analog EEG and EMG signals were amplified (2000 x) and filtered and then digitized at 2 kHz and subsequently down sampled to 200 Hz and stored. The EEG was subjected to a discrete Fourier transformation yielding power spectra (range: 0 – 100 Hz; frequency resolution: 0.25 Hz; time resolution: consecutive 4-sec epochs; window function: Hamming). Hardware (EMBLA) and software (Somnologica-3) were purchased from Medcare Flaga (EMBLA, Thornton, USA).

### ***Determination of behavioral states***

Offline, the animal's behavior was visually classified as "Wakefulness", "REM sleep", or "NREM sleep" for consecutive 4-sec windows based on the EEG and EMG signals as previously described <sup>30</sup>. Wakefulness was characterized by EEG activity of mixed frequency and low amplitude. Muscle tone was present and variable. NREM sleep was defined by synchronous activity in the delta frequency (1 - 4 Hz) and low and stable muscle tone. REM sleep was

characterized by regular theta oscillations (6 - 9 Hz) and muscle atonia with occasional twitches. Four-sec epochs containing EEG artifacts were marked according to the state in which they occurred and excluded from EEG spectral analysis. As the sleep/wake state of epochs in which EEG artifacts occurred could still be determined, they were included in the analysis of time-spent-asleep and -awake.

### **Data analysis**

Analysis of the time course of time spent in each sleep/wake state was performed on 1-, 12-, and 24h values. For the 12h and 24h analysis, the two baseline days were averaged, as no significant differences between the two days were observed [~~3-way repeated measures analysis of the variance (rANOVA)~~ 3-way rANOVA, factors 'day', 'time', and 'genotype';  $p > 0.14$  for factor 'day' and all its interactions]. To further quantify the altered baseline distribution of sleep/wake states in KO mice, hourly values were accumulated ~~over baseline~~ and genotype differences calculated to assess the times at which the two distributions deviated.

Sleep/wake state quality was assessed by analyzing the spectral content of the EEG. To account for inter-individual differences in overall EEG power, EEG spectra of the three sleep/wake states were expressed as a percentage of an individual reference value calculated as the total EEG power across all frequencies considered (0.75 – 45.0 Hz) and sleep/wake states. This reference value was weighted so that for all animals the relative contribution of the three sleep/wake states to this reference value was equal, according to <sup>32</sup>. Theta peak frequency in wakefulness and REM sleep was calculated by determining the frequency at which maximum power density in the theta frequency range (5 – 10 Hz) was reached.

Effects of sleep deprivation were assessed by analyzing EEG delta power, sleep fragmentation, and time spent asleep. Time course analysis of EEG delta power (i.e., the mean EEG power density in the 1 - 4 Hz range in NREM sleep) during baseline and after sleep deprivation was performed as described previously <sup>33</sup>. Briefly, the recording was divided into sections to which an equal number of 4-sec epochs scored as NREM sleep contributed (i.e., percentiles). The baseline light periods were divided into twelve such sections; the baseline and recovery dark periods into six. The recovery light (ZT6-12) period was divided into eight sections. The ~~number of~~ percentiles per recording period ~~was~~ chosen to assure that sufficient and similar numbers of NREM sleep epochs contributed to each interval necessary to obtain meaningful mean delta power values. Delta power values were normalized by

expressing all values relative to the mean value reached in the last 4 h of the main rest periods (i.e., ZT8 - 12) when delta power is minimal during baseline consistent with the fact that delta power is thought to reflect homeostatic sleep need which is lowest at the end of the major rest period. Besides its time course, delta power levels reached immediately after sleep onset in baseline and after sleep deprivation were determined separately. To assure that the same number of 4-sec epochs contributed, delta power was averaged over the first 20 min (300 4-sec epochs) scored as NREM sleep. Sleep onset was determined as the entry into the first episode of NREM sleep lasting > 1 min and not interrupted by two or more 4-sec epochs of wakefulness. This first episode was determined from light onset in the baseline days and from the end of the sleep deprivation for the recovery period.

The effect of 6h sleep deprivation on time spent in and NREM and REM sleep was assessed by calculating the recovery-baseline difference in sleep time for 6h intervals; i.e., the first 6 h of recovery still in the light period (ZT6 - 12) and two 6h intervals in the dark (ZT12 - 18 and ZT18 - 24). The effect of sleep deprivation on NREM sleep spectra and sleep fragmentation was assessed by contrasting EEG spectra and the number of short awakenings (waking bouts lasting 4 consecutive 4s-epochs or less; i.e., <16 s, expressed per hour of NREM sleep;<sup>31</sup>) during the first 3 hours of recovery sleep after sleep onset (ZT6 - 9) from the values observed during the last 4 hours of the baseline light periods (ZT8 - 12), a period during which homeostatic sleep need (and EEG delta power) is lowest and sleep fragmentation highest. Similarly, NREM sleep spectra and the number of brief awakenings in the first 3 hours after sleep onset (ZT0 - 3) in the two baseline days was calculated and contrasted to the same reference.

### ***Cortical gene expression analysis***

Nine KO and 10 WT male mice were used to assess the effect of genotype and sleep deprivation on cortical gene expression. In each genotype, half the mice were submitted to a 6h sleep deprivation (ZT0 – 6), and the other half was left undisturbed and used as control. At ZT6, both groups of mice were sacrificed, the cerebral cortex extracted, and immediately flash frozen in liquid nitrogen. Samples were stored at -80°C. RNA was extracted and purified using the RNeasy Lipid Tissue Mini Kit 50 (QIAGEN, Hombrechtikon, Switzerland) according to manufacturer's instructions. RNA quantity (NanoDrop ND-1000 spectrophotometer; Thermo Scientific, Wilmington, NC, USA) and quality (Agilent 2100 bioanalyzer chips; Agilent



technologies, Basel, Switzerland) was measured and verified. 1000 ng of purified RNA were reverse-transcribed in 20 µl using random hexamers and Superscript II reverse transcriptase (Invitrogen, Life Technologies, Europe, Zug, Switzerland) according to standard procedures. The cDNA was diluted 10 times and 2 µl were amplified in a 10 µl TaqMan reaction in technical triplicates on a ABI PRISM HT 7900 detection system (Applied Biosystems, Life Technologies, Europe, Zug, Switzerland). Cyclor conditions were 50°C 2 min, 95°C 10 min, and 45 cycles at 95°C 15 s and 60°C 1 min. To quantify the RNA expression level, specific forward and reverse primers and probes were used (Table S2). Gene expression levels were normalized to 4 reference genes (*Gapdh*, *Tbp*, *Rsp9*, and *Eef1a1*) using Qbase<sup>PLUS</sup> software (Biogazelle, Zwijnaarde, Belgium) except for *Fabp7* which was expressed relative to *Gapdh* only. The fold change indicative of the relative gene expression are based on the mean of three biological replicates in relation to control samples.

### **Statistics and analysis tools**

TMT Pascal Multi-Target5 software (Framework Computers, Inc., Brighton, MA, USA) was used to manage the data, SigmaPlot V10.0 (Systat Software Inc., Chicago, IL, USA) for graphics, and SAS V9.2 (SAS Institute Software Inc., Cary, NC, USA) or Sigmastat V3.5. (Systat Software, Chicago, IL, USA) for statistical analysis. To assess the effect of genotype on the sleep/wake distribution, EEG power spectra, and time course of delta power, two- or three-way ~~repeated measures analysis of the variance (rANOVAs)~~ were performed. Significant effects and interactions were decomposed using post-hoc Tukey's HSD and t-tests. Genotype differences in the light/dark amount, accumulation, fragmentation of sleep/wake states, and in EEG delta power at sleep onset, as well as the cortical gene expression were evaluated using t-tests. Statistical significance was set to  $p = 0.05$  and results are reported as mean  $\pm$  standard error of the mean (SEM) or of the difference (SED).

## **Results**

### ***The sleep/wake distribution is advanced in Rev-erba knockout mice***

Similar to WT mice, *Rev-erba* KO mice were mostly asleep during the light period and mostly awake during the dark period (almost 2/3 of the time in the respective 12h periods; Figure 1, Table S1). Nevertheless, the distribution of the three sleep/wake states over the 24h day

importantly differed between the two genotypes (Figure 1). Differences were observed mainly at the time encompassing the light-to-dark transition when KO mice spent more time awake than their WT littermates; i.e., in the 2 h prior and 3 h following this transition (ZT10 - 15; Figure 1A). The dynamics of the accumulation of time spent awake over the baseline days summarizes this effect and demonstrates that by ZT15 KO mice accrued ca. 1 h extra wakefulness compared to WT mice (Figure 1B). Interestingly, this gain in wakefulness was rapidly lost over the subsequent 3 h (ZT15 - 18) resulting in almost identical 24h values for wakefulness (Figure 1B, Table S1). Similar, albeit opposite results were observed for NREM and REM sleep (analyses not shown).

Because of this redistribution, KO mice spent significantly more time awake in the 12h light period compared to WT and more time asleep during the 12h dark period, although the latter effect did not reach significance levels ( $p = 0.07$ ; Table S1). As a result, the differences between time-spent-asleep (or –awake) in the 12h dark and 12h light periods, sometimes used to estimate the amplitude of the diurnal sleep/wake distribution, was significantly reduced in KO mice for all three sleep/wake states (Table S1). To further quantify the effect of the earlier wake onset on this reduced diurnal amplitude, the 12h periods over which amplitude was calculated were systematically shifted at 1min increments (Figure 1C). Advancing the 12h periods by  $65 \pm 14$  min yielded the highest diurnal amplitude in KO mice while for WT mice maximal amplitude was already reached with a zero shift ( $+3 \pm 8$  min; Figure 1C). This suggests that in KO mice the 12h dark period does not adequately cover the active period. Moreover, the shift with which a zero amplitude was obtained (i.e., the time of day that divides the 24h day into halves with equal sleep time) was advanced by  $1.8 \pm 0.3$  h relative to WT mice (Figure 1C). Nevertheless, the maximum dark-light amplitude was still significantly lower in KO mice (Figure 1C;  $p = 0.01$ , t-test) indicating that besides an earlier onset other aspects of the sleep/wake distribution, such as the less pronounced main waking bout (analysis not shown), differed between the two genotypes.

### ***Evidence of altered sleep homeostasis in *Rev-erba* knockout mice***

To assess the consequence of a lack of *Rev-erba* on sleep homeostasis mice were challenged with a 6h sleep deprivation. We found that several aspects of the response to sleep deprivation were altered including time spent asleep, sleep continuity, and the levels of EEG delta power reached.

Both genotypes responded to the sleep deprivation by sleeping more than in the corresponding periods in baseline with state-specific recovery dynamics; i.e., whereas extra NREM sleep was already obtained within the first 6h of recovery (i.e., during the last 6 h of the light period; ZT6 - 12), recovery of REM sleep was deferred to the following dark period (Figure 2). However, for both sleep states, *Rev-erba* KO mice obtained less extra sleep in the 1<sup>st</sup> 6 h of the recovery dark period when WT mice accrued most of the extra time spent in NREM and REM sleep (ZT12 - 18; Figure 2). During this time interval, KO mice did not significantly gain any NREM sleep ( $3 \pm 6$  min) whereas WT mice obtained  $35 \pm 7$  min extra NREM sleep compared to that expressed in corresponding baseline hours (Figure 2). KO mice also gained less extra REM sleep ( $7 \pm 2$  min vs.  $13 \pm 1$  min in WT mice; Figure 2B). Despite these different recovery dynamics, at the end of the 18h recovery period, the deficit in sleep gained in KO mice no longer significantly differed from WT (NREM sleep:  $41 \pm 10$  min and  $57 \pm 10$  min; REM sleep:  $14 \pm 3$  and  $20 \pm 3$  min, for KO and WT respectively;  $p = 0.24$  and  $0.11$  for NREM and REM sleep, respectively, t-tests).

The general time course of the changes in EEG delta power was similar in the two genotypes; i.e., delta power increased during periods when waking prevails (i.e., the dark or active period), was highest immediately after light onset and decreased over the remainder of the light (or rest) period, and sleep deprivation resulted in an increase in EEG delta power in subsequent NREM sleep (Figure 3A). This increase was, however, significantly smaller in KO mice. Likewise, EEG delta power at sleep onset in baseline was lower in KO mice compared to WT mice (Figure 3A, B). High relative levels of EEG delta power during NREM sleep are often accompanied by a lower number of brief awakenings indicating deeper and more consolidated sleep<sup>34</sup>. Accordingly, sleep deprivation resulted in a pronounced reduction of the number of brief awakenings in both genotypes (Figure 3C) compared to the baseline reference. Consistent with the smaller increase in EEG delta power, the decrease in the number of brief awakenings interrupting sleep in the first 3 h after sleep onset in baseline and after sleep deprivation was smaller in KO mice (rANOVA,  $p < 0.01$ ; Figure 3C). Post-hoc testing revealed, however, that the genotype effect in recovery sleep did not reach the 5% significance level ( $p = 0.01$  and  $0.06$ , for baseline and recovery sleep, respectively). After sleep onset in baseline, WT mice showed a reduced number of brief awakenings to the number observed at the end of the baseline light periods (ZT8 - 12), while in KO mice no difference from this baseline reference was observed (Figure 3C). In keeping with the analysis of EEG delta power, as

reference for the effects on sleep fragmentation the number of brief awakenings occurring at the end of the baseline light periods was used (see Methods). This reference did not differ between genotypes (Figure S1).

To further investigate whether the genotype differences in EEG delta power reported above were specific to the delta frequency range and to homeostatic sleep need we analyzed the spectral composition of the NREM sleep EEG. Spectra did not differ between genotypes when averaged over the entire 48h baseline (Figure S2A). Similarly, in the last 4h hours of the baseline light periods, when sleep need is considered to be lowest, no spectral differences were observed (Figure 4; dashed lines labeled 5 and 6 in panels A and B). In contrast, when sleep need was high; i.e., at the start of the baseline light periods and after sleep deprivation, NREM sleep spectra in *Rev-erb $\alpha$*  KO mice did deviate from WT and the frequency bins in which power density differed in baseline and after sleep deprivation largely overlapped (Figure 4). Thus under both conditions, NREM sleep EEG power density between 2.5 and 4.75 Hz was significantly lower in KO mice.

Although the largest effects of sleep deprivation on the NREM sleep EEG concern the delta frequencies, significant recovery-to-baseline differences can be observed over a larger frequency range<sup>35,36</sup>. In the current experiment, power density in frequencies up to 20 Hz were increased compared to baseline (Figure 4C). In addition, we found evidence that EEG activity in the low gamma range (32 - 45 Hz) was decreased. Although, the shape of the differential spectral profiles as well as the frequency ranges that were significantly affected by the sleep deprivation were very similar for the two genotypes, the magnitude of these changes was smaller in *Rev-erb $\alpha$*  KO mice. The frequency range for which the sleep deprivation effect was significantly smaller was again limited to the delta frequencies (1.0 - 3.75 Hz; Figure 4C).

The results concerning the genotype difference in the increase in EEG delta power suggest that when awake, homeostatic sleep need accumulates at a slower rate in mice lacking *Rev-erb $\alpha$* . Given the published relationship between; e.g., theta activity in the waking EEG and delta activity in subsequent NREM sleep<sup>37,38</sup>, we also quantified the spectral composition of waking EEG. When analyzed over the 48h baseline, EEG activity during wakefulness showed higher power in the higher delta frequencies (1.5 - 5.5 Hz), frequencies between 11 and 20 Hz, and the low gamma range (33 - 42 Hz; Figure S2). The first two of these three frequency ranges were similarly affected in REM sleep (Figure S2). Moreover, the prevailing theta frequency

during wakefulness, but not that of REM sleep, was slower in *Rev-erb $\alpha$*  KO mice. Slower theta oscillations in the wakefulness EEG were also observed during the sleep deprivation (Figure S2B).

We next focused on the waking EEG in the 3 h immediately preceding sleep onset in baseline and during the last 3 h of the sleep deprivation in an attempt to identify those frequency components that could have contributed to the delta power differences observed in subsequent NREM sleep. In the last 3 h of both the baseline dark periods (ZT21 - 24) and of the sleep deprivation (ZT3 - 6) the waking EEG showed clear theta activity in WT mice (Figure 5A). The appearance of a distinct theta peak in the waking EEG resulted from the combination of a decrease in high delta activity (3 - 6 Hz) and an increase in theta activity (6.5 - 11 Hz; Figure 5B). In addition, in both conditions, EEG activity in the high gamma range (40 - 90 Hz) significantly increased compared to that observed in the last 4h of the baseline light periods which served as reference (Figure 5B, Figure S4). Although also in KO mice activity in the theta and high gamma frequency ranges was increased both in the baseline dark period and the sleep deprivation, the peak in theta activity during the baseline dark was less distinct because high delta activity remained elevated (Figure 5A, B). Moreover, the increase in theta power was smaller in both baseline and sleep deprivation when mice lack *Rev-erb $\alpha$* . Correlation analyses showed that this increase in theta power predicted the increase in EEG delta power in subsequent NREM sleep (Figure 5C). In the sleep deprivation condition also the increase in high gamma activity was smaller in the KO mice.

### ***The molecular consequences of lack of *Rev-erb $\alpha$* in the cerebral cortex***

To test whether the lack of *Rev-erb $\alpha$*  also affects molecular markers of sleep need, we investigated the cortical expression of genes that are known to be reliably modulated by sleep deprivation; i.e., *Homer1 $\alpha$* , *Sgk1*, *Dbp*, and *Per2* (e.g. <sup>39</sup>). *Homer1 $\alpha$* , *Sgk1*, and *Per2* were up-regulated by sleep deprivation and *Dbp* expression down-regulated although post-hoc testing showed that this decrease reached significance in WT mice only (Figure 6). We previously found that also *Npas2* expression increases after sleep deprivation<sup>29</sup> but in the current experiment ~~only in KO mice~~ this increase was significant only in KO mice. These sleep-deprivation induced changes in cortical gene expression did, however, not statistically differ between the two genotypes (Figure 6).

Next we assessed the expression of genes known to be directly regulated by *Rev-erba*, to test whether *Rev-erba* has a similar role on these targets in the cerebral cortex. The expression levels of *Bmal1*, *Npas2*, *Clock*, and the REV-ERB $\alpha$  target *Fabp7* (*fatty acid binding protein 7*)<sup>40</sup> were all significantly up-regulated in the cortex of *Rev-erba* KO mice compared to WT (Figure 6); that of *Fabp7* more than 7-fold. The overall level of *Dbp* and *Per2* in the KO was also increased, although for *Per2* it did not reach significance levels ( $p < 0.06$ , t-test). These results confirm that *Rev-erba* directly, or indirectly in the case of *Dbp* and *Per2*, controls the expression of clock genes also in the cortex.

## Discussion

### ***REV-ERB $\alpha$ impacts sleep homeostasis***

In the current study we identified a robust sleep homeostatic phenotype in mice lacking *Rev-erba*. Significantly lower values for the electrophysiological correlate of homeostatic sleep need, EEG delta power, were reached after periods of prolonged wakefulness in KO mice compared to their WT littermates. This phenotype was observed both after enforced and spontaneous periods of wakefulness arguing against the possible influence of other variables than extended wakefulness that usually accompany a sleep deprivation protocol such as; e.g., increased stimulation, locomotion, and stress. This genotype difference in the NREM sleep EEG was specific to the delta frequencies and appeared only at times when sleep need was high. The observed differences do therefore not represent a general EEG phenomena but concern only those frequencies thought to reflect homeostatic sleep need. Moreover, the less pronounced increase in sleep consolidation, another, non-EEG variable reflecting homeostatic sleep need<sup>34</sup>, confirmed that sleep immediately after sleep onset was less 'deep'.

Together these findings argue for a slower accumulation of sleep need when *Rev-erba* KO mice are awake. Correlates of the homeostatic sleep process can also be quantified in the waking EEG, the theta content of which is known to increase over the course of extended waking periods (e.g.<sup>34,38,41</sup>). Moreover, in the rat, levels of theta activity during wakefulness and/or time spent exploring, during which theta activity is prominent, predict delta power in subsequent NREM sleep<sup>38,42</sup> suggesting that these two EEG activities not only gauge the same underlying homeostatic process but that wakefulness with higher theta activity causes delta power to be higher during the sleep that follows. We here confirmed this relationship in the mouse. *Rev-erba* KO mice displayed a smaller increase in theta activity during wakefulness in

the 3 h preceding sleep onset that predicted the lower delta power immediately after sleep onset. We extended these theta findings to the 35 - 90 Hz or gamma frequencies, the activity in which ~~in the hippocampal formation~~ is closely coupled to theta oscillations in the hippocampal formation<sup>43,44</sup>. In both genotypes EEG activity in the gamma band was increased relative to the values reached when sleep need is lowest but, like for theta activity, this increase was less pronounced in the KO mice although the frequencies in which significant differences were observed concerned the 71 - 78 Hz range only. Lack of *Rev-erba* may thus lead to deficits in either engaging in waking behaviors rich in theta activity such as exploratory behavior, and/or in the recruitment of neuronal populations contributing to these oscillations. The significantly slower theta frequency we observed during wakefulness constitutes additional evidence for a role of *Rev-erba* in modulating theta oscillations of hippocampal origin. Of interest in this context is the fact that lack of *Rev-erba* leads to increased adult neurogenesis in the hippocampus<sup>40</sup>, which may affect the network properties contributing to theta oscillations<sup>45</sup>.

### ***REV-ERBa, sleep homeostasis, and dopamine***

Impaired hippocampal function in *Rev-erba* KO mice has been demonstrated for a number of hippocampal-dependent behaviors<sup>40,46</sup>. This impairment was accompanied by increased dopamine turnover<sup>46,47</sup>. The slower hippocampal theta oscillations we observed in *Rev-erba* KO mice are consistent with the theta slowing observed when dopamine tone is increased as observed in *Dopamine transporter (Dat)* KO mice<sup>48</sup>. In *Rev-erba* KO mice the increased dopamine turnover was due to an up-regulation of *Tyrosine hydroxylase (Th)*, the rate-limiting enzyme in dopamine production<sup>46,47</sup>. *Rev-erba* can directly repress *Th* expression in competition with the *Nuclear receptor-related 1 protein (Nurr1)*, another nuclear receptor and key transcriptional activator of *Th* and other elements of the dopaminergic system<sup>47,49</sup>. Of immediate interest for our current study is the observation that altered dopamine levels, resulting from altered *Dat* activity, were found to be associated with a change in the homeostatic sleep rebound both in flies and humans<sup>50-52</sup>.

Other clock genes have been implicated in the regulation of dopamine levels in the brain<sup>47,53,54</sup>. Such role might be direct, like REV-ERBa's repression of *Th* expression discussed above and ~~for the NPAS2:BMAL1's transcriptional activation circadian control~~ of the dopamine degrading enzyme *monoamine oxidase A (Maoa)*<sup>54</sup>, or indirect through the effects of other

clock genes on the expression of e.g., *Rev-erba*. For instance, CLOCK/NPAS2 $\beta$ -BMAL1 heterodimers induce *Rev-erba* transcription while REV-ERB $\alpha$ , in turn, represses the expression of these three transcription factors<sup>6-8,11</sup>. We here confirm that also in the cerebral cortex removing the repression provided by REV-ERB $\alpha$  increases the expression of *Bmal1*, *Npas2*, and *Clock*. Consistent with this up-regulation we found the CLOCK/NPAS2 $\beta$ -BMAL1 targets *Dbp* and *Per2* to be upregulated as well. PER2, in turn, not only provides negative feedback to CLOCK/NPAS2 $\beta$ -BMAL1 induced transcription but, in addition, coordinates the action of various nuclear receptors, including REV-ERB $\alpha$  and NURR1, via protein-protein interactions<sup>55</sup>. Thus REV-ERB $\alpha$  as part of a complex network of interacting transcriptional regulators, seems central in changing dopamine turnover.

The link between dopamine levels and sleep homeostasis is intriguing and could also have contributed to the profound sleep homeostatic phenotype we observed in *Rev-erba* KO mice. This relationship is, however, not straightforward; the mutations in *Dat* activity referred to above led to increased dopamine post-synaptically due to compromised dopamine re-uptake which was associated with increased homeostatic sleep rebounds<sup>50-52</sup>, while in *Rev-erba* KO mice the reduced inhibition of *Th* expression led to increased dopamine levels pre-synaptically and, as we show here, was associated with a decreased homeostatic sleep rebound. Further illustrating this complexity is the fact that REV-ERB $\alpha$  can directly inhibit the expression of the *Dopamine D3 receptor*<sup>56</sup>. The activation of this receptor is thought to be inhibitory, reducing novelty seeking behaviors, and to reduce dopamine through post-synaptic negative feedback<sup>57</sup>.

#### ***REV-ERB $\alpha$ and circadian organization of overt behavior***

Although *Rev-erba* is important for the circadian molecular circuitry and for setting the phase of circadian rhythms in peripheral tissues<sup>12</sup>, its lack only modestly affects rhythms in overt behaviors. *Rev-erba* KO mice do maintain circadian organization of locomotor activity under constant conditions albeit with a significantly shorter free-running period<sup>8,12</sup>. The earlier onset of the main waking period under the entrained conditions of our experiment is consistent with a shorter endogenous period although the ca. 24 min shortening of the period seems insufficient to account for the >1 h advance of sleep/wake distribution. The inducible depletion of both *Rev-erba* and its homolog *Rev-erb $\beta$*  does lead to a profound disruption of circadian behavior<sup>12</sup> pointing to a functional redundancy between the two at least for this



phenotype. Whether also the sleep homeostatic phenotype becomes more pronounced in double KO mice remains to be determined.

In addition to altered EEG delta power and sleep fragmentation after sleep deprivation, we also observed a distinctly different pattern for the recovery of sleep time lost during the sleep deprivation to which the advance of the sleep/wake distribution might have contributed. In both genotypes, maximal levels of wakefulness are reached within the first 6 h of the dark period under baseline conditions. After the sleep deprivation, it was during these 6 h that WT mice adapt their behavior to allow for recovery sleep gaining 35 min of NREM sleep. KO mice conspicuously maintain their baseline sleep/wake pattern and did not increase NREM sleep time above basal levels. As a consequence of the extra NREM sleep in the recovery dark period, EEG delta power in WT mice, exceptionally, reached lower levels than KO mice. The inflexibility of *Rev-erba* KO mice to mount the appropriate behavioral sleep response is reminiscent to that observed in mice lacking the clock gene *Npas2*<sup>58</sup>. Also *Npas2* KO mice have intact circadian behavior but a poor homeostatic response to food or sleep deprivation<sup>59</sup>.

#### ***Sleep deprivation, mood, metabolism, and clock genes***

Strong, bidirectional links exist between sleep and mood disorders, perhaps best illustrated by the amazingly rapid, albeit short lasting, antidepressant effects of sleep deprivation in depression (reviewed in<sup>60</sup>). These beneficial effects of sleep deprivation combined with the observation made in some, but not all studies, that EEG delta power is reduced in major depressive disorder (reviewed in<sup>61</sup>), led to the hypothesis that sleep need accumulates at a slower rate during wakefulness and is causally involved in the dysregulation of mood<sup>62</sup>. Abnormal dynamics in waking theta activity over the course of a sleep deprivation further suggest abnormal sleep homeostasis in depressed patients<sup>63</sup>. Both the altered theta activity in wakefulness and the altered delta activity in NREM sleep we here describe for the KO mice are reminiscent of these EEG changes associated with mood disorders.

Sleep homeostasis and mood disorders could also be linked at the molecular level. Many studies reported on the role of clock genes in mood regulation<sup>64,65</sup> and our own work revealed a bidirectional relationship between clock genes and sleep homeostasis<sup>66</sup>, further illustrated here by the altered homeostatic regulation of sleep in *Rev-erba* KO mice and by the effects of sleep deprivation on the cortical expression of the clock genes *Npas2* and *Per2*, and

the clock-controlled transcription factor *Dbp*. We previously showed that like *Dbp*, *Rev-erba* was decreased by sleep deprivation<sup>29</sup>. We also quantified two transcripts known to be reliably upregulated sleep deprivation; i.e., *Homer1a* and *Sgk1*. Although the expression of both was indeed increased in the current study, their increase did not differ between genotypes pointing to a dissociation between EEG and molecular markers of sleep need.

Among the clock genes, REV-ERB $\alpha$  seems to play a central role in the regulation of mood as evidenced by; e.g., a mania-like behavior in KO mice<sup>40,47,67</sup> and, perhaps, the altered sleep homeostasis in the current study. The involvement of REV-ERB $\alpha$  in mood might, in part, relate to its role in dopaminergic signaling discussed above. Other likely candidate pathways concern REV-ERB $\alpha$ 's role in adult hippocampal neurogenesis<sup>40</sup>, which has also been linked to mood disorders<sup>68</sup>, and its well-established role in metabolism and circadian rhythms<sup>65</sup>. One important mediator of the increased adult hippocampal neurogenesis is FABP7<sup>40,69</sup>, a direct target of REV-ERB $\alpha$ <sup>40</sup>. Because of the large, 7.5-fold over-expression of *Fabp7* in the brain that we observed (see Figure 6 and<sup>40</sup>), FABP7 might also be involved in mediating other phenotypes observed in *Rev-erba* KO mice such as anxiety and memory deficits<sup>69-71</sup>. Concerning the sleep homeostatic phenotype, overexpression of the mouse *Fabp7* or its fly homologue *Fabp-B* in the fruit fly *Drosophila melanogaster* resulted in decreased sleep duration and consolidation<sup>70</sup>, the *Drosophila* correlates of reduced homeostatic sleep need<sup>51,72</sup>. Whether FABP7 plays a role in the homeostatic regulation of sleep in mammals has yet to be determined.

Several groups have investigated the role of the core clock genes in sleep homeostasis by studying the effects of sleep deprivation on EEG delta power in mice carrying targeted disruptions of single or a combination of clock genes<sup>3,4</sup>. The first core clock genes for which altered sleep-wake dependent dynamics of EEG delta power was demonstrated were the *Cryptochromes*<sup>73</sup>. *Cry1,2* double KO mice displayed a more rapid build-up of homeostatic sleep need during wakefulness resulting in overall higher levels of EEG delta power during baseline, explaining the smaller relative increase in EEG delta power after sleep deprivation. In contrast, as in our *Rev-erba* KO mice, evidence for a slower build rate was obtained in *Bmal1* and *Npas2* knock-out mice<sup>58,74</sup>, while in *Clock*-mutant mice no differences were reported<sup>75</sup>. Sleep homeostasis was also assessed in *Per1* and *Per2* single and double mutant KO mice but with inconsistent outcomes<sup>76,77</sup>. These disparate results do not support a simple, unifying mechanism through which the clock gene circuitry, as a whole, alters the dynamics of the sleep

homeostat but rather suggest that factors such as tissue/cell type-specificity and yet to be identified modifiers associated with differences in genetic background play important roles as has been demonstrated for circadian-related phenotypes <sup>5</sup>.

### **Conclusion**

REV-ERB $\alpha$ 's role in the control of sleep homeostasis is of particular interest because both the molecule and the process are tightly linked to metabolism as well as circadian rhythms (for reviews see <sup>3,14,78,79</sup>). REV-ERB $\alpha$ 's activity as a transcriptional repressor is modulated by cellular redox state through altered binding of its endogenous ligand, heme <sup>80,81</sup>. In the context of our sleep homeostatic phenotype, *Rev-erba* could thus act as a sensor of the metabolic imbalance imposed at the neuronal level by periods of extended wakefulness in keeping with our proposal that clock genes not only set time-of-day, but in the cerebral cortex, can also be used to keep track of and respond to time-spent-awake <sup>3,79</sup>. A recent study demonstrated that synthetic agonists targeting both REV-ERB proteins leads to a strong, immediate reduction in sleep time when administrated at ZT6 <sup>82</sup>, implying that pharmacologically targeting of REV-ERB $\alpha$  could be useful in the treatment sleep disorders.

### **Acknowledgements**

We are grateful for the expert technical support provided by Ashot Sargsyan and Antoinette Hayoz and all that helped with the sleep deprivations.

### **References**

1. Borbely AA. A two process model of sleep regulation. *Hum Neurobiol* 1982;1(3):195-204.
2. Daan S, Beersma DG, Borbely AA. Timing of human sleep: recovery process gated by a circadian pacemaker. *Am J Physiol* 1984;246(2 Pt 2):R161-83.
3. Franken P. A role for clock genes in sleep homeostasis. *Curr Opin Neurobiol* 2013;23(5):864-72.
4. Mang GM, Franken P. Genetic Dissection of Sleep Homeostasis. *Curr Top Behav Neurosci*. 2013 Dec 14 ed: Springer Berlin Heidelberg, 2014.
5. Lowrey PL, Takahashi JS. Genetics of circadian rhythms in Mammalian model organisms. *Adv Genet* 2011;74:175-230.
6. Crumbley C, Burriss TP. Direct regulation of CLOCK expression by REV-ERB. *PLoS One* 2011;6(3):e17290.
7. Crumbley C, Wang Y, Kojetin DJ, Burriss TP. Characterization of the core mammalian clock component, NPAS2, as a REV-ERB $\alpha$ /ROR $\alpha$  target gene. *J Biol Chem* 2010;285(46):35386-92.

8. Preitner N, Damiola F, Lopez-Molina L, et al. The orphan nuclear receptor REV-ERB $\alpha$  controls circadian transcription within the positive limb of the mammalian circadian oscillator. *Cell* 2002;110(2):251-60.
9. Ueda HR, Chen W, Adachi A, et al. A transcription factor response element for gene expression during circadian night. *Nature* 2002;418(6897):534-9.
10. Sato TK, Panda S, Miraglia LJ, et al. A functional genomics strategy reveals Rora as a component of the mammalian circadian clock. *Neuron* 2004;43(4):527-37.
11. Akashi M, Takumi T. The orphan nuclear receptor ROR $\alpha$  regulates circadian transcription of the mammalian core-clock Bmal1. *Nat Struct Mol Biol* 2005;12(5):441-8.
12. Cho H, Zhao X, Hatori M, et al. Regulation of circadian behaviour and metabolism by REV-ERB $\alpha$  and REV-ERB $\beta$ . *Nature* 2012;485(7396):123-7.
13. Mazzocchi G, Cai Y, Liu S, et al. REV-ERB $\alpha$  and the clock gene machinery in mouse peripheral tissues: a possible role as a synchronizing hinge. *Journal of biological regulators and homeostatic agents* 2012;26(2):265-76.
14. Duez H, Staels B. Rev-erb $\alpha$ : an integrator of circadian rhythms and metabolism. *Journal of applied physiology* 2009;107(6):1972-80.
15. Harding HP, Lazar MA. The monomer-binding orphan receptor Rev-Erb represses transcription as a dimer on a novel direct repeat. *Molecular and cellular biology* 1995;15(9):4791-802.
16. Burris TP. Nuclear hormone receptors for heme: REV-ERB $\alpha$  and REV-ERB $\beta$  are ligand-regulated components of the mammalian clock. *Mol Endocrinol* 2008;22(7):1509-20.
17. Renaud JP, Harris JM, Downes M, Burke LJ, Muscat GE. Structure-function analysis of the Rev-erbA and RVR ligand-binding domains reveals a large hydrophobic surface that mediates corepressor binding and a ligand cavity occupied by side chains. *Mol Endocrinol* 2000;14(5):700-17.
18. Yin L, Lazar MA. The orphan nuclear receptor Rev-erb $\alpha$  recruits the N-CoR/histone deacetylase 3 corepressor to regulate the circadian Bmal1 gene. *Mol Endocrinol* 2005;19(6):1452-9.
19. Raghuram S, Stayrook KR, Huang P, et al. Identification of heme as the ligand for the orphan nuclear receptors REV-ERB $\alpha$  and REV-ERB $\beta$ . *Nat Struct Mol Biol* 2007;14(12):1207-13.
20. Yin L, Wu N, Curtin JC, et al. Rev-erb $\alpha$ , a heme sensor that coordinates metabolic and circadian pathways. *Science* 2007;318(5857):1786-9.
21. Estall JL, Ruas JL, Choi CS, et al. PGC-1 $\alpha$  negatively regulates hepatic FGF21 expression by modulating the heme/Rev-Erb( $\alpha$ ) axis. *Proc Natl Acad Sci U S A* 2009;106(52):22510-5.
22. Duez H, van der Veen JN, Duhem C, et al. Regulation of bile acid synthesis by the nuclear receptor Rev-erb $\alpha$ . *Gastroenterology* 2008;135(2):689-98.
23. Le Martelot G, Claudel T, Gatfield D, et al. REV-ERB $\alpha$  participates in circadian SREBP signaling and bile acid homeostasis. *PLoS Biol* 2009;7(9):e1000181.
24. Delezie J, Dumont S, Dardente H, et al. The nuclear receptor REV-ERB $\alpha$  is required for the daily balance of carbohydrate and lipid metabolism. *Faseb J* 2012;26(8):3321-35.
25. Balsalobre A, Damiola F, Schibler U. A serum shock induces circadian gene expression in mammalian tissue culture cells. *Cell* 1998;93(6):929-37.
26. Torra IP, Tsibulsky V, Delaunay F, et al. Circadian and glucocorticoid regulation of Rev-erb $\alpha$  expression in liver. *Endocrinology* 2000;141(10):3799-806.
27. Solt LA, Wang Y, Banerjee S, et al. Regulation of circadian behaviour and metabolism by synthetic REV-ERB agonists. *Nature* 2012;485(7396):62-8.
28. Chen L, Yang G. PPARs Integrate the Mammalian Clock and Energy Metabolism. *PPAR research* 2014;2014:653017.
29. Mongrain V, Hernandez SA, Pradervand S, et al. Separating the contribution of glucocorticoids and wakefulness to the molecular and electrophysiological correlates of sleep homeostasis. *Sleep* 2010;33(9):1147-57.
30. Mang GM, Franken P. Sleep and EEG Phenotyping in Mice. *Current Protocols in Mouse Biology* 2012;2:55-74.
31. Franken P, Malafosse A, Tafti M. Genetic determinants of sleep regulation in inbred mice. *Sleep* 1999;22(2):155-69.

32. Franken P, Malafosse A, Tafti M. Genetic variation in EEG activity during sleep in inbred mice. *Am J Physiol* 1998;275(4 Pt 2):R1127-37.
33. Franken P, Chollet D, Tafti M. The homeostatic regulation of sleep need is under genetic control. *J Neurosci* 2001;21(8):2610-21.
34. Franken P, Tobler I, Borbely AA. Sleep homeostasis in the rat: simulation of the time course of EEG slow-wave activity. *Neuroscience Letters* 1991;130(2):141-4.
35. Borbely AA, Baumann F, Brandeis D, Strauch I, Lehmann D. Sleep deprivation: effect on sleep stages and EEG power density in man. *Electroencephalography and clinical neurophysiology* 1981;51(5):483-95.
36. Kopp C, Petit JM, Magistretti P, Borbely AA, Tobler I. Comparison of the effects of modafinil and sleep deprivation on sleep and cortical EEG spectra in mice. *Neuropharmacology* 2002;43(1):110-8.
37. Huber R, Esser SK, Ferrarelli F, Massimini M, Peterson MJ, Tononi G. TMS-induced cortical potentiation during wakefulness locally increases slow wave activity during sleep. *PLoS One* 2007;2(3):e276.
38. Vyazovskiy VV, Tobler I. Theta activity in the waking EEG is a marker of sleep propensity in the rat. *Brain research* 2005;1050(1-2):64-71.
39. Curie T, Mongrain V, Dorsaz S, Mang GM, Emmenegger Y, Franken P. Homeostatic and circadian contribution to EEG and molecular state variables of sleep regulation. *Sleep* 2013;36(3):311-23.
40. Schnell A, Chappuis S, Schmutz I, et al. The nuclear receptor REV-ERB $\alpha$  regulates *Fabp7* and modulates adult hippocampal neurogenesis. *PLoS One* 2014;9(6):e99883.
41. Cajochen C, Brunner DP, Krauchi K, Graw P, Wirz-Justice A. Power density in theta/alpha frequencies of the waking EEG progressively increases during sustained wakefulness. *Sleep* 1995;18(10):890-4.
42. Huber R, Tononi G, Cirelli C. Exploratory behavior, cortical BDNF expression, and sleep homeostasis. *Sleep* 2007;30(2):129-39.
43. Penley SC, Hinman JR, Sabolek HR, Escabi MA, Markus EJ, Chrobak JJ. Theta and gamma coherence across the septotemporal axis during distinct behavioral states. *Hippocampus* 2012;22(5):1164-75.
44. Sirota A, Montgomery S, Fujisawa S, Isomura Y, Zugaro M, Buzsaki G. Entrainment of neocortical neurons and gamma oscillations by the hippocampal theta rhythm. *Neuron* 2008;60(4):683-97.
45. Rangel LM, Quinn LK, Chiba AA, Gage FH, Aimone JB. A hypothesis for temporal coding of young and mature granule cells. *Front Neurosci* 2013;7:75.
46. Jager J, O'Brien WT, Manlove J, et al. Behavioral changes and dopaminergic dysregulation in mice lacking the nuclear receptor Rev-erb $\alpha$ . *Mol Endocrinol* 2014;28(4):490-8.
47. Chung S, Lee EJ, Yun S, et al. Impact of circadian nuclear receptor REV-ERB $\alpha$  on midbrain dopamine production and mood regulation. *Cell* 2014;157(4):858-68.
48. Dzirasa K, Santos LM, Ribeiro S, et al. Persistent hyperdopaminergia decreases the peak frequency of hippocampal theta oscillations during quiet waking and REM sleep. *PLoS One* 2009;4(4):e5238.
49. Zetterstrom RH, Solomin L, Jansson L, Hoffer BJ, Olson L, Perlmann T. Dopamine neuron agenesis in *Nurr1*-deficient mice. *Science* 1997;276(5310):248-50.
50. Greenspan RJ, Tononi G, Cirelli C, Shaw PJ. Sleep and the fruit fly. *Trends Neurosci* 2001;24(3):142-5.
51. Pfeiffenberger C, Allada R. *Cul3* and the BTB adaptor *insomniac* are key regulators of sleep homeostasis and a dopamine arousal pathway in *Drosophila*. *PLoS Genet* 2012;8(10):e1003003.
52. Holst SC, Bersagliere A, Bachmann V, Berger W, Achermann P, Landolt HP. Dopaminergic role in regulating neurophysiological markers of sleep homeostasis in humans. *J Neurosci* 2014;34(2):566-73.

53. Parekh PK, Ozburn AR, McClung CA. Circadian clock genes: Effects on dopamine, reward and addiction. *Alcohol* 2015.
54. Hampp G, Ripperger JA, Houben T, et al. Regulation of monoamine oxidase A by circadian-clock components implies clock influence on mood. *Current Biology* 2008;18(9):678-83.
55. Schmutz I, Ripperger JA, Baeriswyl-Aebischer S, Albrecht U. The mammalian clock component PERIOD2 coordinates circadian output by interaction with nuclear receptors. *Genes & development* 2010;24(4):345-57.
56. Ikeda E, Matsunaga N, Kakimoto K, et al. Molecular mechanism regulating 24-hour rhythm of dopamine D3 receptor expression in mouse ventral striatum. *Mol Pharmacol* 2013;83(5):959-67.
57. Holmes A, Lachowicz JE, Sibley DR. Phenotypic analysis of dopamine receptor knockout mice; recent insights into the functional specificity of dopamine receptor subtypes. *Neuropharmacology* 2004;47(8):1117-34.
58. Franken P, Dudley CA, Estill SJ, et al. NPAS2 as a transcriptional regulator of non-rapid eye movement sleep: genotype and sex interactions. *Proc Natl Acad Sci U S A* 2006;103(18):7118-23.
59. Dudley CA, Erbel-Sieler C, Estill SJ, et al. Altered patterns of sleep and behavioral adaptability in NPAS2-deficient mice. *Science* 2003;301(5631):379-83.
60. Benedetti F. Antidepressant chronotherapeutics for bipolar depression. *Dialogues Clin Neurosci* 2012;14(4):401-11.
61. Frey S, Birchler-Pedross A, Hofstetter M, et al. Challenging the sleep homeostat: sleep in depression is not premature aging. *Sleep Med* 2012;13(7):933-45.
62. Borbely AA, Wirz-Justice A. Sleep, sleep deprivation and depression. A hypothesis derived from a model of sleep regulation. *Hum Neurobiol* 1982;1(3):205-10.
63. Plante DT, Goldstein MR, Landsness EC, et al. Altered overnight modulation of spontaneous waking EEG reflects altered sleep homeostasis in major depressive disorder: a high-density EEG investigation. *Journal of affective disorders* 2013;150(3):1167-73.
64. McClung CA. Circadian genes, rhythms and the biology of mood disorders. *Pharmacol Ther* 2007;114(2):222-32.
65. Albrecht U. Circadian clocks and mood-related behaviors. *Handbook of experimental pharmacology* 2013(217):227-39.
66. Franken P. A role for clock genes in sleep homeostasis. *Current opinion in neurobiology* 2013;23(5):864-72.
67. Yin L, Wang J, Klein PS, Lazar MA. Nuclear receptor Rev-erb $\alpha$  is a critical lithium-sensitive component of the circadian clock. *Science* 2006;311(5763):1002-5.
68. Samuels BA, Hen R. Neurogenesis and affective disorders. *Eur J Neurosci* 2011;33(6):1152-9.
69. Matsumata M, Inada H, Osumi N. Fatty acid binding proteins and the nervous system: Their impact on mental conditions. *Neuroscience research* 2014.
70. Gerstner JR, Vanderheyden WM, Shaw PJ, Landry CF, Yin JC. Fatty-acid binding proteins modulate sleep and enhance long-term memory consolidation in *Drosophila*. *PLoS One* 2011;6(1):e15890.
71. Shimamoto C, Ohnishi T, Maekawa M, et al. Functional characterization of FABP3, 5 and 7 gene variants identified in schizophrenia and autism spectrum disorder and mouse behavioral studies. *Human molecular genetics* 2014;23(24):6495-511.
72. Andretic R, Shaw PJ. Essentials of sleep recordings in *Drosophila*: moving beyond sleep time. *Methods in enzymology* 2005;393:759-72.
73. Wisor JP, O'Hara BF, Terao A, et al. A role for cryptochromes in sleep regulation. *BMC Neuroscience* 2002;3:20.
74. Laposky A, Easton A, Dugovic C, Walisser J, Bradfield C, Turek F. Deletion of the mammalian circadian clock gene BMAL1/Mop3 alters baseline sleep architecture and the response to sleep deprivation. *Sleep* 2005;28(4):395-409.
75. Naylor E, Bergmann BM, Krauski K, et al. The circadian clock mutation alters sleep homeostasis in the mouse. *J Neurosci* 2000;20(21):8138-43.

76. Kopp C, Albrecht U, Zheng B, Tobler I. Homeostatic sleep regulation is preserved in mPer1 and mPer2 mutant mice. *Eur J Neurosci* 2002;16(6):1099-106.
77. Shiromani PJ, Xu M, Winston EM, Shiromani SN, Gerashchenko D, Weaver DR. Sleep rhythmicity and homeostasis in mice with targeted disruption of mPeriod genes. *Am J Physiol Regul Integr Comp Physiol* 2004;287(1):R47-57.
78. Benington JH, Heller HC. Restoration of brain energy metabolism as the function of sleep. *Prog Neurobiol* 1995;45(4):347-60.
79. Albrecht U. Circadian rhythms and sleep--the metabolic connection. *Pflugers Arch* 2012;463(1):23-30.
80. Gupta N, Ragsdale SW. Thiol-disulfide redox dependence of heme binding and heme ligand switching in nuclear hormone receptor rev-erb $\beta$ . *J Biol Chem* 2011;286(6):4392-403.
81. Carter EL, Ragsdale SW. Modulation of nuclear receptor function by cellular redox poise. *Journal of inorganic biochemistry* 2014;133:92-103.
82. Banerjee S, Wang Y, Solt LA, et al. Pharmacological targeting of the mammalian clock regulates sleep architecture and emotional behaviour. *Nature communications* 2014;5:5759.

## Figure Legends

**Figure 1. *Rev-erba* knockout mice have an earlier increase in wakefulness relative to dark onset. A)** Mean ( $\pm$  1 SEM) hourly values of Wakefulness, NREM sleep, and REM sleep over 48h baseline. Genotype significantly affected the time course for all three states while the two baseline days did not differ [3-way rANOVA, factors genotype ('G': KO, WT), day ('D': 1, 2), and hour ('H': 1 - 24); Wakefulness: 'G' p = 0.94, 'D' p = 0.19, 'H' p < 0.001, 'GxD' p = 0.33, 'GxH' p < 0.001, 'DxH' p = 0.56; NREMS: 'G' p = 0.35, 'D' p = 0.12, 'H' p < 0.001, 'GxD' p = 0.19, 'GxH' p < 0.001, 'DxH' p = 0.46; REMS: 'G' p = 0.22, 'D' p = 0.13, 'H' p < 0.001, 'GxD' p = 0.18, 'GxH' p < 0.001, 'DxH' p = 0.87]. **B)** Upper curve: Mean ( $\pm$  1 SEM) accumulation of time spent awake calculated at 1h increments. The two baseline days were averaged. Lower curve: Mean differences (KO-WT;  $\pm$  1 SED) in accumulated values represented in the upper curve. **C)** Amplitudes of the difference in time spent awake between the two 12h periods. The start time of the 12h periods was shifted at 1min increments with 0 shift corresponding to the actual dark onset. For each 1min shift the amplitude was calculated. Circles with bi-directional error bars indicated the mean ( $\pm$  1 SEM) of the individual amplitude with 0 shift (symbols labeled '1'; same values as in Table S1) or at the shift yielding maximal amplitude (symbols labeled '2'). All panels, knockout (KO, n=13) and wild type (WT, n=8) mice are indicated with black lines and symbols and grey line and open symbols, respectively. Grey areas represent 12h dark

periods. Red squares in A) and B) indicate hours with significant genotype differences (post-hoc t-tests,  $p < 0.05$ ).

**Figure 2. *Rev-erba* knockout mice recover less sleep between ZT12 and ZT18 in recovery from sleep deprivation compared to their wild type littermates.** Recovery-baseline differences in NREM sleep and REM sleep time spent in three 6h intervals (i.e., during the light from ZT6-12, during the dark from ZT12-18 and ZT18-24). Indicated are the mean ( $\pm 1$  SEM) differences for KO (black bars,  $n = 13$ ) and WT (white bars,  $n = 8$ ) mice. Red stars indicate significant genotype differences (t-tests,  $p < 0.05$ ); blue stars significant recovery-baseline differences (paired t-tests,  $p < 0.05$ ).

**Figure 3. EEG delta power after sleep onset in baseline and after sleep deprivation is lower in the absence of *Rev-erba*.** **A)** Mean ( $\pm 1$  SEM) EEG delta power (upper curves) and NREM sleep amount (lower curves) during 48h baseline, 6h sleep deprivation (SD; grey shaded area), and 18h recovery. The time course of the changes in EEG delta power during baseline differed significantly between KO and WT (3-way rANOVA, factors genotype ('G'  $p < 0.001$ ) and hour ('H'  $p < 0.001$ ), 'GxH'  $p < 0.001$ ; analyses of average of 2 baselines). Also in recovery the time course of both EEG delta power (2-way rANOVA, 'G'  $p = 0.0012$ , 'H'  $p < 0.001$ , 'GxH'  $p < 0.001$ ) and of the hourly NREM sleep values (2-way rANOVA, 'G'  $p = 0.69$ , 'H'  $p < 0.001$ , 'GxH'  $p < 0.001$ ) were affected by genotype. Intervals in which EEG delta power and NREM sleep in KO mice significantly differed from WT are indicated by red squares (post-hoc t-tests,  $p < 0.05$ ). For baseline NREM sleep statistics see Figure 1A. **B)** Mean ( $\pm 1$  SEM) level of EEG delta power in the first 20 min of NREM sleep after light onset in baseline (average of two days) and after the SD. Both KO and WT have a significant increase in EEG delta power after sleep onset in baseline and after sleep deprivation compared to their lowest baseline levels (=100%; t-tests,  $p < 0.05$ ) but the levels reached in KO mice were lower in both conditions (red stars; t-tests,  $p < 0.05$ ). **C)** Recovery-baseline differences ( $\pm 1$  SEM) in the number of brief awakenings (i.e., waking bouts lasting 16 sec or less) interrupting sleep during the first 3 h of the light period in baseline (Baseline) and during the 3 first hours at sleep onset after sleep deprivation (recovery), contrasted to the number obtained during the last 4 h of the baseline light period (= 0 level). The decrease in number of brief awakenings was smaller in KO mice (2-way rANOVA, factors genotype ( $p = 0.01$ ) and day ( $p < 0.001$ ), interaction  $p = 0.44$ ). Red stars mark



significant genotype differences (post-hoc t-tests;  $p < 0.05$ ). Note that in recovery, genotype difference reached  $p = 0.06$ . Also note that while brief awakenings after recovery sleep onset was significantly lower than in the last 4 h of the baseline light periods for both genotypes (post-hoc t-tests;  $p < 0.001$ ), after baseline sleep onset, they were reduced in WT mice only (post-hoc t-tests,  $p < 0.001$  and  $p = 0.98$  for WT and KO, respectively).

**Figure 4. Genotype differences in NREM sleep EEG spectra are specific to the delta frequencies activity and appear only when sleep need is high. A)** Mean EEG spectral profiles in NREM sleep at times when sleep need is assumed to be high; i.e., in the first 3 h after sleep onset in baseline [left panel; lines labeled 3 and 4 for WT (grey) and KO (black), resp.] and after sleep deprivation [right panel, recovery, lines labeled '1' and '2' for WT (grey) and KO (blue), resp.]. For comparison, spectral profiles of NREM sleep in the last 4 h of baseline, which served as reference for the analysis of EEG delta power (see Figure 3), when sleep need is lowest, are depicted [left panel, dashed lines, labeled '5' and '6' for KO (black) and WT (grey), resp.]. **B)** Genotype differences for the spectral profiles in panels A. Mean spectral KO/WT ratios for recovery (blue line labeled '2/1') and baseline (black line; '4/3') were remarkably similar and significant differences concerned higher delta frequencies only for both conditions (2.50 - 4.75 Hz; post-hoc t-tests,  $p < 0.05$ , horizontal lines at the bottom), while no significant genotype differences were observed when sleep need is low (dashed black line labeled '5/6'). **C)** The effect of sleep deprivation on NREM sleep spectra in panel A. Analysis of spectral recovery-to-baseline (Rec/Bsl) ratios in KO (blue line labeled '2/5') and WT (grey; '1/6') mice demonstrated that sleep loss increased EEG power density in frequencies  $< 21$  Hz, while suppressing EEG activity in the low gamma range ( $> 32$  Hz) in both genotypes (blue and grey lines at bottom). Although shape and the affected frequency ranges were highly similar, both the increases and the decreases in EEG power density were smaller in KO mice. Significant genotype differences were, however, highly frequency specific and limited to the delta frequency range (1.0 - 3.75 Hz) and in one frequency bin within the low gamma band (42.5 Hz).

**Figure 5. Mean EEG spectral profiles in wakefulness preceding the times at which NREMS EEG delta power differed between genotypes. A)** Mean spectral profiles in the 3 h before sleep onset in baseline (ZT21 - 24; left panel) and the last 3 h of the sleep deprivation (ZT3 - 6; right panel; KO black lines, WT grey lines). EEG power density was expressed as a percentage

of individual total EEG power (see Methods and FigureS1C). **B)** Effects of prolonged wakefulness on the spectral composition of the waking EEG was assessed by comparing the EEG spectra in panels A to the waking spectra obtained in the last 4 h of baseline (ZT8 - 12) when sleep need is lowest (see Figure S5 for a genotype comparison of the spectral EEG profiles at this time). Mean profiles of spectral ratios (last 3 h / baseline ZT8-12 reference) reached in last 3 h of the baseline dark periods (left panel) and the last 3 h of the sleep deprivation (right panel) were similar and significant differences concerned a decrease in EEG power in high delta frequencies (3 - 6 Hz) and increases in the theta (6.5 - 11 Hz) and gamma (40 - 90 Hz) frequency ranges (horizontal lines at bottom of panel; WT grey, KO black; post-hoc paired t-tests,  $p < 0.05$ ). These differences were, however, larger in sleep deprivation than in baseline (analyses not shown) and larger in WT than KO mice (red horizontal line at bottom of panel, post-hoc t-tests  $p < 0.05$ ). **C)** The increase in EEG theta power in wakefulness as quantified in panel B, predicted the level of EEG delta power reached in NREM sleep after sleep onset as quantified in Figure 2B. The changes in theta power were averaged over the 2Hz frequency range for which significant genotype differences were observed (8.5 - 10.5 Hz). For calculation of EEG delta power values see Fig. 2B. Lines represent the linear correlation through all KO (black line:  $\text{delta} = 38 \pm 9 \times \text{theta} + 125 \pm 8$ ;  $R^2=0.45$ ,  $p = 0.0002$ ,  $n = 26$ ) and WT (grey line:  $\text{delta} = 50 \pm 15 \times \text{theta} + 133 \pm 18$ ;  $R^2 = 0.45$ ,  $p = 0.0047$ ,  $n = 16$ ) data points (i.e., Bsl + Rec). Slope and offset did not differ between genotypes. Black symbols: KO ( $n = 13$ ); grey symbols WT ( $n = 8$ ); circles baseline (Bsl); triangles recovery (Rec).

**Figure 6. Cortical expression of several genes is affected by the lack of *Rev-erba* and sleep deprivation.** Mean (1 SEM) levels of mRNA expression in the cortex of WT ( $n=10$ ) and KO ( $n=9$ ) mice, submitted to either a 6h sleep deprivation from ZT0-6 (SD, black bars) or allowed to sleep *ad lib* (control, white bars). Normalized expression levels (see Methods) are expressed relative to the WT control group (set to 1.0). Red stars indicate a significant effect of the sleep deprivation within genotypes ( $p < 0.05$ ; post-hoc t-tests), while blue stars indicate genotype differences [2-way ANOVA, factors genotype ('G') and sleep condition ('SD'); *Homer1a*: 'G'  $p = 0.99$ , 'SD'  $p < 0.001$ ; 'GxSD'  $p = 0.19$ ; *Sgk1*: 'G'  $p = 0.77$ , 'SD'  $p < 0.001$ , 'GxSD'  $p = 0.68$ ; *Fabp7*: 'G'  $p < 0.001$ ; 'SD'  $p = 0.98$ , 'GxSD'  $p = 0.91$ ; *Bmal1*: 'G'  $p < 0.001$ , 'SD'  $p = 0.42$ , 'GxSD'  $p = 0.36$ ; *Npas2*: 'G'  $p = 0.004$ , 'SD'  $p = 0.003$ , 'GxSD'  $p = 0.13$ ; *Clock*: 'G'  $p = 0.02$ , 'SD'  $p = 0.18$ , 'GxSD'  $p$

= 0.73; *Dbp*: 'G' < 0.001, 'SD'  $p = 0.02$ , 'GxSD'  $p = 0.48$ ; *Per2*: 'G'  $p = 0.06$ ; 'SD'  $p = 0.003$ ; 'GxSD'  $p = 0.84$ ].






Figure 1




Figure 2





Figure 3




Figure 4




Figure 5






Figure 6

## Supplemental Information

Mang *et al.* Altered sleep homeostasis in *Rev-erb $\alpha$*  knockout mice

**Table S1. Time spent asleep and awake in baseline.**


|             | 12h light |         |              | 12h dark |          |          | 24h      |          |          |
|-------------|-----------|---------|--------------|----------|----------|----------|----------|----------|----------|
|             | KO        | WT      | <i>P</i>     | KO       | WT       | <i>p</i> | KO       | WT       | <i>p</i> |
| <b>Wake</b> | 280 (6)   | 249 (7) | <b>0.003</b> | 452 (11) | 484 (11) | 0.07     | 732 (13) | 733 (15) | 0.82     |
| <b>NREM</b> | 372 (7)   | 390 (6) | 0.09         | 233 (11) | 203 (9)  | 0.06     | 606 (13) | 593 (11) | 0.97     |
| <b>REM</b>  | 67 (3)    | 75 (3)  | 0.11         | 33 (2)   | 30 (2)   | 0.23     | 101 (3)  | 105 (4)  | 0.87     |

|             | 12h dark- 12h light |           |              |
|-------------|---------------------|-----------|--------------|
|             | KO                  | WT        | <i>P</i>     |
| <b>Wake</b> | 172 (12)            | 235 (10)  | <b>0.002</b> |
| <b>NREM</b> | -139 (12)           | -187 (12) | <b>0.02</b>  |
| <b>REM</b>  | -34 (3)             | -45 (3)   | <b>0.04</b>  |


Upper part: Mean (SEM) time spent in wakefulness, NREM sleep, and REM sleep over 12- and 24h intervals (KO n=13, WT n=8). Lack of *Rev-erb $\alpha$*  caused animals to be significantly more awake in the light period (t-tests), an effect that was reversed in the dark, resulting in very similar 24h values (also see Figure 1). Lower part: Diurnal amplitude of sleep-wake states, quantified as the dark-light difference, was significantly smaller in KO mice. All values in minutes and based on two baseline days.

**Table S2. Sequences of primers and probes used for TaqMan qPCR analysis of selected genes and references genes.**

| <b>Gene Symbol</b> | <b>Direction</b>    | <b>Sequence 5' to 3'</b>   | <b>Accession #</b> |
|--------------------|---------------------|--|--------------------|
| <i>Homer1a</i>     | fwd<br>rev<br>probe | GCATTGCCATTCCACATAGG<br>ATGAACTTCCATATTTATCCACCTTACTT<br>ACACATTCAATTCAGCAATCATGA      | NM_011982          |
| <i>Bmal1</i>       | fwd<br>rev<br>probe | CAAGAAAGTATGGACACAGACAAA<br>GCATTCTTGATCCTTCCTTGGT<br>TGACCCTCATGGAAGGTTAGAATATGCAGAAC | NM_007489          |
| <i>Npas2</i>       | fwd<br>rev<br>probe | GGTCATCGGATTCTTGACAGAA<br>TCCAGTCTGCTGGATGTCA<br>CACAATGAAGTCTCAGCACAAACAGAAATC        | NM_008719          |
| <i>Per2</i>        | fwd<br>rev<br>probe | ATGCTCGCCATCCACAAGA<br>GCGGAATCGAATGGGAGAAT<br>ATCCTACAGGCCGGTGGACAGCC                 | NM_011066          |
| <i>Dbp</i>         | fwd<br>rev<br>probe | CGTGGAGGTGCTTAATGACCTTT<br>CATGGCCTGGAATGCTTGA<br>AACCTGATCCCCTGATCTCGCC               | NM_016974          |
| <i>Sgk1</i>        | fwd<br>rev<br>probe | ACGGTGGACTGGTGGTGTCT<br>GCCGTGTTCCGGCTATAAAA<br>TATGAGATGCTCTACGGCCTGCCCC              | NM_011361          |
| <i>GAPDH</i>       | fwd<br>rev<br>probe | CATGGCCTCCGTGTTCTTA<br>CCTGCTTCACAACTTTCTTGA<br>CCGCCTGGAGAAACCTGCCAAGTATG             | NM_008084          |
| <i>Tbp</i>         | fwd<br>rev<br>probe | TTGACCTAAAGACCATTGCACTTC<br>TTCTCATGATGACTGCAGCAAA<br>TGCAAGAAATGCTGAATATAATCCCAAGCG   | NM_013684          |
| <i>Rps9</i>        | fwd<br>rev<br>probe | GACCAGGAGCTAAAGTTGATTGGA<br>TCTTGGCCAGGGTAAACTTGA<br>AAACCTCACGTTTGTTCGGAGTCCATACT     | NM_029767          |
| <i>Eef1a</i>       | fwd<br>rev<br>probe | CCTGGCAAGCCCATGTGT<br>TCATGTCACGAACAGCAAAGC<br>TGAGAGCTTCTCTGACTACCCTCCACTTGGT         | NM_010106          |
| <i>Fabp7</i>       | fwd<br>rev<br>probe | TTTCTGCGCAACCTGGAAG<br>CACGTTTCCCACTTGCCTAG<br>AAGCTCTGGGCGTGGGCTTTGC                  | NM_021272          |
| <i>Clock</i>       | fwd<br>rev<br>probe | Mm00455950_m1<br>TaqMan Expression Assay, Life Technologies                            | NM_007715.5        |




**Figure S1. Number of brief awakenings during the 4 last hours of light period (ZT8-12).** The mean (+ 1 SEM; expressed per hour of sleep) number of brief awakenings did not differ between genotypes (t-test,  $p = 0.44$ ). The number of brief awakenings obtained at this time of day is used as a reference for the analysis presented in Figure 3C.



**Figure S2: Spectral composition of EEG activity during wakefulness, NREM sleep, and REM sleep in baseline. A)** Mean spectral profiles calculated over the two baseline days during wakefulness (upper, left), NREM sleep (upper, middle), and REM sleep (upper, right panel) together with their KO/WT spectral ratios (three lower panels). When averaged over the entire 24h day the NREM sleep EEG spectra do not differ between genotypes. In waking and REM sleep EEG power density between 1.5 - 5.5 Hz, 11 - 20 Hz, and, for wakefulness, between 33 - 42 Hz, was higher in KO mice (red horizontal lines at bottom; post-hoc t-test,  $P < 0.05$ ). The significant decrease in KO mice in the theta frequency range in wakefulness was not due to a lower theta activity but to a slowing of the dominant theta frequency. **B)** Theta peak frequency, but not theta peak power, was reduced in KO mice both in baseline (bsl, left panel; -0.9 Hz) and sleep deprivation (SD, right panel; -0.5 Hz; red stars; t-test  $P < 0.05$ ; mean  $\pm$  bi-directional SEM). Note that vertical scales differ between the two panels. Theta peak frequency during REM sleep was not affected by genotype (analysis not shown). **C)** Total EEG power calculated in baseline, which served as an individual reference to calculate relative EEG spectra, did not differ between genotypes. This reference was used for the analyses shown

Figures S2A, S3, 4, and 5. The genotype differences in the relative EEG spectra depicted in panel A were similarly observed for the absolute EEG spectra although frequency ranges that significantly differed were smaller due to a larger inter-individual variation (data not shown).



**Figure S3. Spectral profiles of the waking EEG obtained in the last 4h of the baseline light periods.** Upper panel: Mean spectral profiles of the baseline waking EEG between ZT8-12 used as reference for the analysis in Figure 4B (KO black, WT grey line). EEG power density was expressed as a percentage of individual total EEG power (see Methods and FigureS1C). Lower panel: Genotype differences (KO/WT) for the spectra in the upper panel were limited to the high delta frequencies (2.25 - 4.5 Hz; red squares; post-hoc t-tests,  $p < 0.05$ ).