



Original Research Article



Meta-analysis of chemotherapy in nasopharynx carcinoma (MAC-NPC): An update on 26 trials and 7080 patients

Pierre Blanchard^{a,b,*}, Anne W.M. Lee^c, Alexandra Carmel^{b,d}, Ng Wai Tong^c, Jun Ma^e, Anthony T.C. Chan^f, Ruey Long Hong^g, Ming-Yuan Chen^h, Lei Chen^h, Wen-Fei Li^h, Pei-Yu Huang^h, Dora L.W. Kwongⁱ, Sharon S.X. Poh^j, Roger Ngan^c, Hai-Qiang Mai^h, Camille Ollivier^{b,d}, George Fountzilas^k, Li Zhang^h, Jean Bourhis^l, Anne Aupérin^{b,d}, Benjamin Lacas^{b,d}, Jean-Pierre Pignon^{b,d}, on behalf of the MAC-NPC collaborative Group

^a Department of Radiation Oncology, Gustave Roussy Cancer Campus, Université Paris-Sud, Université Paris-Saclay, Villejuif, France Gustave-Roussy, Villejuif, France

^b Oncostat U1018 INSERM, labeled Ligue Contre le Cancer, Villejuif, France

^c University of Hong Kong – Shenzhen Hospital, University of Hong Kong, China

^d Service de Biostatistique et d'Epidémiologie, Gustave Roussy, Villejuif, France

^e Sun Yat-sen University Cancer Center, State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, Guangdong Key Laboratory of Nasopharyngeal Carcinoma Diagnosis and Therapy, Guangzhou 510060, China

^f State Key Laboratory of Translational Oncology, Hong Kong Cancer Institute, The Chinese University of Hong Kong, Hong Kong, China

^g National Taiwan University Hospital, Taipei, Taiwan

^h Sun Yat-sen University Cancer Center, Guangzhou, China

ⁱ Department of Clinical Oncology, University of Hong Kong, Hong Kong, China

^j National Cancer Center, Singapore

^k Aristotle University of Thessaloniki School of Medicine, Thessaloniki, and Hellenic Cooperative Oncology Group, Athens, Greece and German Oncology Center, Limassol, Cyprus

^l Department of Radiotherapy, Centre hospitalier universitaire vaudois, Lausanne, Switzerland

ARTICLE INFO

Keywords:

Individual patient data
Meta-analysis
Randomized trials
Chemotherapy
Nasopharynx carcinoma

ABSTRACT

Purpose: Chemotherapy, when added to radiotherapy, improves survival in locally advanced nasopharyngeal carcinoma (NPC). This article presents the second update of the Meta-Analysis of Chemotherapy in NPC.

Methods: Published or unpublished randomized trials assessing radiotherapy (\pm a second chemotherapy timing) with/without chemotherapy in non-metastatic NPC patients were identified. Updated data were sought for studies included in the previous rounds of the meta-analysis. The primary endpoint was overall survival. All trials were analyzed following the intent-to-treat principle using a fixed-effects model. Treatments were classified in five subsets according to chemotherapy timing. The statistical analysis plan was pre-specified.

Results: Eighteen new trials were identified. Individual patient data were available for seven. In total, the meta-analysis now included 26 trials and 7,080 patients. The addition of chemotherapy reduced the risk of death, with a hazard ratio (HR) of 0.79 (95% confidence interval (CI) [0.73; 0.85]), and an absolute survival increase at 5 and 10 years of 6.1% [+3.9; +8.3] and + 8.4% [+5.7; +11.1], respectively. The largest effect was observed for concomitant + adjuvant, induction (with concomitant in both arms) and concomitant chemotherapy, with respective HR [95%CI] of 0.68 [0.59; 0.79] (absolute survival increase at 5 years: 12.3% (7.0%;17.6%)), 0.73 [0.63; 0.86] (6.0% (2.5%;9.5%)) and 0.81 [0.70; 0.92] (5.2% (0.8%;9.6%)). The benefit of chemotherapy was also demonstrated by improvement in progression-free survival, cancer mortality, locoregional control and distant control. There was a significant interaction between patient age and chemotherapy effect.

Conclusion: This updated meta-analysis confirms the benefit of concomitant chemotherapy and concomitant + adjuvant chemotherapy, and suggests that addition of induction or adjuvant chemotherapy to concomitant chemotherapy improves tumor control and survival. The benefit of chemotherapy decreases with increasing patient age.

* Corresponding author at: Département de Radiothérapie Oncologique, GustaveRoussy, 114 rue Edouard Vaillant, 94805 Villejuif, France.

E-mail address: pierre.blanchard@gustaveroussy.fr (P. Blanchard).

<https://doi.org/10.1016/j.ctro.2021.11.007>

Received 8 November 2021; Received in revised form 18 November 2021; Accepted 18 November 2021

Available online 26 November 2021

2405-6308/© 2021 The Authors. Published by Elsevier B.V. on behalf of European Society for Radiotherapy and Oncology. This is an open access article under

the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

With >129 000 new cases globally in 2018, nasopharyngeal carcinoma (NPC) remains a major public health issue especially in endemic areas. In Asia, the age-standardized incidence rate is estimated to be 2.1 per 100 000 and up to 3.0 in China, compared to 0.44 in Europe or North America [1]. The majority of patients have non-keratinizing disease, which is related to Epstein Barr Virus (EBV) infection, and follows a different course compared to other head and neck cancers. Due to its anatomic location, in close vicinity to critical structures, its frequent lymphatic spread, risk for distant metastases and chemoradiosensitivity, the mainstay of treatment of locoregionally advanced NPC has long been a combination of chemotherapy (CT) and radiotherapy (RT) [2]. However, the best sequence remains to be found and there is a controversy over the benefit of adding induction or adjuvant chemotherapy when concurrent chemoradiotherapy is given.

Since the publication of the previous meta-analysis of chemotherapy in nasopharyngeal cancer MAC-NPC [3–4], multiple randomized trials have been conducted and novel induction regimens using taxane or gemcitabine have been studied [5–7]. In addition, quantitative plasma EBV DNA value at baseline and after chemoradiotherapy has a major prognostic role for recurrence and survival. Whether EBV DNA bears a predictive value for guiding personalized use of adjuvant chemotherapy remains to be demonstrated, as the first trial using EBV DNA after chemoradiotherapy to plan additional chemotherapy was reported negative [8].

The MAC-NPC collaborative group therefore decided in 2016 to update its analysis to focus on the role of induction chemotherapy as an adjunct to concomitant chemoradiotherapy.

Methods

This updated meta-analysis was performed according to a pre-specified protocol (available at <https://46.18.130.247/sites/default/files/mac-npc3-protocol.pdf>).

Selection criteria and search strategy

To be eligible, trials had to compare RT alone versus RT plus CT, or to compare a treatment strategy, i.e. RT plus concomitant CT (CRT) or RT plus induction CT (IC) or RT plus adjuvant CT (AC) with the same treatment strategy plus CT (other timing). They had to be properly randomized and include untreated non-metastatic NPC patients. Trials were eligible if at least 60 patients had been included (30 patients per arm for trials with more than two arms) and if all patients had undergone potentially curative loco-regional treatment. Accrual had to be completed before December 31, 2016.

Both published and unpublished trials meeting the criteria were eligible. Trials search combined search in electronic publication databases, trial registries and meeting proceedings (details in [web appendix 2](#)).

Individual patient data collection

Individual patient data were requested for each eligible trial and for all randomized patients. Data requested included characteristics of patients and tumours, date of randomization and treatment arm allocated, dates of failures and death, details on treatments received, acute and late toxicities. Toxicity was scored locally according to the scale used at the time of the trial and hence only graded according to NCI CTCAE v4 as severe (grade ≥ 3 for all except xerostomia where it was grade ≥ 2) vs not. Follow-up information was updated whenever possible.

All data were checked according to a standard procedure and compared with the trial protocol and published reports. Missing values and discrepancies were discussed with the trialists. Randomization validity was evaluated by checking patterns of treatment allocation over

time and balance of baseline characteristics between treatment arms. Follow-up of patients was also compared between treatment arms. [9] Each trial was reanalysed and the analyses were sent to the trialists for validation.

Endpoints

The primary endpoint was overall survival (OS), defined as the time from randomization until death from any cause. The secondary endpoints were progression-free survival (PFS), loco-regional failure (LRF), distant failure (DF), and cancer and non-cancer mortality. Progression-free survival was defined as the time from randomization to first progression (loco-regional or distant) or death from any cause. Non-cancer deaths were defined as deaths resulting from known causes other than nasopharynx cancer and without previous progression, and deaths from unknown cause occurring more than five years after randomization. Cancer deaths included deaths from nasopharynx cancer, deaths from any cause with previous progression and deaths from unknown cause within five years after randomization.

Statistical analysis

All analyses were performed on an intention-to-treat basis. Median follow-up was estimated with the reverse Kaplan-Meier method. [10] Analyses were stratified by trial. Individual and overall pooled hazard ratios (HR) with 95% confidence intervals (CI) were calculated through a fixed-effect model using the log-rank expected number of events and variance. [11] A similar model with chi-square instead of log-rank was used to estimate odds ratios (OR) for comparison of toxicity between arms. Rates of toxicity in the experimental arm were calculated using rate in the control arm and the OR. [12] Cumulative incidences of loco-regional failure and distant failure were analyzed with a competing risk method. [13] Only the first event was considered. If both a LRF and a DF occurred at the same time, the event was counted as DF only. For each endpoint, the studied type of failure was analyzed as the main event. The other type of failure and death without failure were analyzed as competing events. Subdistribution HR (sHR) for loco-regional failure and for distant failure were estimated in each trial using the Fine-Gray model and the global sHR were estimated with the Fine-Gray model stratified for trials. Chi-square heterogeneity tests and I^2 statistic were used to investigate the overall heterogeneity between trials. [14–15] In case of significant heterogeneity (defined by heterogeneity test p -value < 0.10), trials with 95% CI that did not cross the 95% CI of the pooled HR were excluded (i.e. outliers) as a sensitivity analysis. If heterogeneity remained significant, a random-effect model was used. With 7,000 patients, it would be possible to detect an absolute improvement in survival from 40 % to 45 % at 5-years with a power exceeding 95% (two-sided logrank test). Cancer mortality was obtained indirectly by subtracting the log-rank statistic for non-cancer mortality from the log-rank statistic for mortality from all causes. [14] Stratified survival curves were estimated for control and experimental groups using annual death rates and hazard ratios, and absolute benefit at five years with its 95% CI was calculated. [14]

Interaction between treatment effect on OS/PFS and patient subgroups (age, sex, performance status, and overall stage) was estimated directly in a single Cox model stratified on trial and containing treatment effect, covariate (for example age) effect and treatment-covariate interaction (“one-stage” model method), among the subset of trials using a “new” drug. [16] Bleomycin, epirubicin, floxuridine, hydroxyurea, oxaliplatin, mitomycin, methotrexate, vincristine or tegafur/uracil were considered as old drugs. Only trials including all subgroups could be included in a given subgroup analysis.

Subset analyses were performed to study the interaction between treatment effect on OS/PFS and trial level characteristics, using a test of heterogeneity between the different groups of trials (called interaction). Residual heterogeneity within trial subgroups was computed by

subtracting the χ^2 statistic of the heterogeneity test between groups from the χ^2 statistic of the overall heterogeneity test. [17] Predefined subsets were timing of randomized CT (adjuvant [after RT/CRT] (AC), induction [before RT] (IC), concomitant [during RT] (CRT), concomitant plus adjuvant (CRT-AC), induction [before concomitant chemoradiotherapy] (IC-(CRT))), CT drug (“old” vs “new” drug, with “new” drugs being the ones that are still used routinely nowadays), trial size, randomization method and radiotherapy technique.

Sensitivity analyses were performed after the exclusion of trials including <100 patients, trials including two different CT timings with only one being randomized, trials with a median follow-up shorter than five years, outliers, and patients with WHO type 1 cancer.

All p-values were two-sided. Analyses were performed using SAS software, version 9.4 (SAS Institute, Cary, NC), except for loco-regional failure and distant failure, which were analysed with the packages “cmprsk” and “crrSC” of the R software (version 3.6.3).

Role of the funding source

The funding sources had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The submission of the paper for publication was decided by the MAC-NPC collaborative group. PB, BL, JPP had access to the raw data. The corresponding author had full access to all of the data and the final responsibility to submit for publication.

Results

Description of the trials and patients

Eighteen new trials (3,746 patients) were identified. Individual patient data were not available for 10 trials (1,385 patients) because of data loss (n = 4), a change in activities (n = 2), the impossibility to obtain a final answer from the trial’s team (n = 3) or the impossibility to contact the trial’s authors (n = 1). Another trial was excluded post-checking (94 patients) because follow-up was inferior to two years and no deaths were recorded. [18] Therefore, seven new trials (2,274 patients) were included. All of them had CRT as control arm, six studied the addition of IC before CRT and one the addition of AC after CRT. The data of 29 patients (0.4%) excluded after randomization in their respective trials were retrieved for this analysis. The data of 13 patients (0.2%) included in two trials [19–20] could not be retrieved.

Overall, the meta-analysis included 26 trials (7,080 patients, see [web table 1](#) for the description of each trial, including patient population, chemotherapy regimen and timing, radiotherapy dose range and technique, sample size and follow-up). Among the two trials with 2x2 factorial plan, one [21] was counted as four comparisons and the other [22] as two comparisons, leading to 30 comparisons (7,302 patients). Fifteen trials (3,170 patients) compared CT vs. nil. In the 11 remaining trials (4,132 patients), all patients received one timing of CT and were randomized to an additional timing. The breakdown per CT subset is as follows:

- IC: 4 comparisons (830 pts),
- IC-(CRT): 8 comparisons (2,379 pts) – corresponds to trials in which all patients received CRT and only IC was randomized
- CRT: 7 comparisons (1,834 pts including 2 comparisons in which all patients received IC, VUMCA-95 (unpublished) and Guangzhou 2002–02 [23]),
- AC: 5 comparisons (992 pts including three comparisons with CRT in both arms (QMH-95 [21,24], Guangzhou 2006 [25], NPC 0502 [8]),
- CRT-AC: 6 comparisons (1,267 pts).

The median follow-up for all trials was 7.4 years (interquartile range: 5.5; 12.5), with 13 trials over 10 years of follow-up [21–23,26–36]. Follow-up varied widely according to the subset of trials, from 6.1 and

6.3 years for IC and IC-(CRT) trials, 6.8 years for AC trials, to 12.1 years for CRT trials and 15.1 for CRT-AC trials.

Patient characteristics, by trial subset and the breakdown by disease stage, can be found in [web tables 2, 3 and 4](#). Briefly, 74.4% of patients were male, with a median age of 45 years, and most had a non-keratinizing histology (97%). Stage III accounted for 44.9% and stage IV for 46.5%.

Overall survival

There were 2,879 (39.4%) deaths in 7,302 patients, which represents a 37% increase compared to the previous round of the meta-analysis. Causes of deaths are reported in [web table 5](#). The addition of chemotherapy reduced the risk of death, with a HR [95%CI] of 0.79 [0.73; 0.85] (p < 0.0001), and an absolute survival increase at 5 and 10 years of 6.1% [+3.9; +8.3] and + 8.4% [+5.7; +11.1] ([Figs. 1 and 2](#)). The heterogeneity (p = 0.08; I² = 28%) was reduced after exclusion of two outliers [26,34] without a change in the results. There was a significant interaction (p = 0.03) between chemotherapy timing and efficacy on overall survival. The largest effects were observed for CRT-AC, IC-(CRT) and CRT, with respective HR of 0.68 [0.59; 0.79], 0.73 [0.63; 0.86] and 0.81 [0.70; 0.92].

Secondary endpoints

The addition of chemotherapy to radiotherapy improved PFS ([Figs. 1 and 3](#)), cancer mortality, locoregional control and distant control, with respective HR/sHR of 0.75 [0.70; 0.80], 0.74 [0.68; 0.81], 0.79 [0.70; 0.89] and 0.70 [0.62; 0.78] (see forest plots and survival curves in [web Figs. 1-3](#), and patterns of events for PFS or cancer mortality in [web tables 6 and 7](#)). The use of chemotherapy did not significantly increase the rate of non-cancer death, with a HR of 1.16 [0.96; 1.40] ([Fig. 4](#)). Full results, on the entire population and by treatment timing, including absolute benefits at 5 and 10 years, are summarized in [Table 1](#).


Subgroup, subset and sensitivity analyses

There was a significant interaction between patient age and chemotherapy effect on survival, with the larger benefit observed in younger patients ([Table 2](#)). As an example, the absolute OS difference at 5 years decreased from 8.5% to 7.1%, 6.1% and 3.9% in patients younger than 40, aged between 40 and 49, aged between 50 and 59 and older than 60 respectively. When looking at specific mortality, there was a significant trend (p = 0.03) towards a decreased efficacy of chemotherapy in terms of cancer death with increasing age while there was no effect on non-cancer death, suggesting that increased treatment related death is not the major mechanism involved in the decreasing effect of treatment with increasing age ([web table 8](#)). There was no significant interaction for OS or PFS with the other patient level characteristics tested: gender, performance status, T Stage, N Stage, overall stage or imaging modality used. Due to missing data or small sample, analyses were not performed according to histology subtype or EBV DNA status.

Among subset of trials, there was a significant interaction between OS and PFS and the type of chemotherapy drug used. “New” drugs (i.e. those that remain used nowadays, representing 19 comparisons and 4,530 patients) yielded better outcomes than old drugs, with respective HR for death of 0.72 [0.65; 0.79] and 0.89 [0.79; 0.99] (interaction p = 0.0005). No interaction was seen between chemotherapy effect on OS or PFS and any of the other tested subset characteristics: trial size, randomization method and radiotherapy technique ([Web tables 9 and 10](#)). Sensitivity analyses showed similar results ([Web table 11](#)).

Toxicity

Among acute toxicity, neutropenia, thrombocytopenia, anemia, weight loss, dermatitis, mucositis, hearing loss and nausea/vomiting



Abbreviations: CI: Confidence Interval, E: Expected, HR: Hazard Ratio, O: Observed

Fig. 1. Forest plots for (A) overall survival and (B) progression-free survival with hazard ratios by timing of chemotherapy.

were increased with the use of chemotherapy (Table 3). For all of the above mentioned except dermatitis, there was also a significant interaction with chemotherapy timing (Web Table 12). In general, the addition of a second timing of chemotherapy, in the concomitant + adjuvant or induction (with concomitant) subsets, was responsible for more acute hematological toxicity. Mucositis was especially increased in arms with concomitant chemotherapy. Late toxicities were unfortunately poorly recorded, as seen in the overall small rate of toxicities observed compared with what would be expected in such a population. Only hearing deficit was increased with the use of chemotherapy (Odds Ratio (OR): 1.30 [1.08; 1.55]), and no interaction could be found between chemotherapy timing and late toxicity.

Discussion

Compared to the previous round of the meta-analysis, this update reconfirms that chemotherapy improves outcomes in patients with locally advanced nasopharyngeal cancer, especially when it is delivered as concomitant, concomitant + adjuvant, or induction in addition to concomitant. The benefit is maintained and even increased in the long term, with absolute survival benefits at 10 years that surpass those at five years. However, the benefit decreased in patients aged 60 or older. Acute toxicity was increased with the addition of chemotherapy, especially when two timings were used. This work supports NCCN and ASCO/ASCO guidelines which recommend IC + CRT or CRT + AC for locally advanced NPC [37–38].

The benefit risk ratio is key when prescribing a treatment. While the impact of age had not been demonstrated in previous MAC-NPC rounds, maybe due to a lack of power, it now seems clear that age is associated with treatment effect, the effect being larger in younger patients and not significant in patients older than 60 years. The cause of this age-effect is unclear, and competing causes of deaths are likely not involved as no increase in non-cancer death could be demonstrated with increasing patient age (Web table 8). Potential explanations could be related to

poor tolerance, increased risk of toxicity leading to lower chemotherapy dose intensity received. There is no clear pathophysiological explanation for such a decreased efficacy in older patients. As such a finding was replicated in other head and neck cancer meta-analyses evaluating chemotherapy or radiotherapy [39,40], the effect of chance is unlikely. Due to small patient numbers above 70, it was not reasonable to further stratify between 60 and 69 and 70+. No other disease or patient related factor, such as tumor stage or performance status, was associated with treatment effect. Hence it is not possible to define which patients could be the best candidates for the addition of chemotherapy, especially before or after concomitant. Physicians will need to consider both the patient's absolute benefit and the risk of additional toxicity before prescribing.

The strengths of the present work are the use of individual patient data, updated follow up, the use of standardized endpoints and a pre-planned analysis based on the intention to treat principle. However not all data could be retrieved, although it has been shown that trials with no individual patient data available are usually smaller, more often published in national journals, of lower quality and tend to show higher treatment effect than trial with IPD available [41]. For instance, among the 11 non-included trials, ten were reported only in the Chinese medical literature and median sample size was 121 patients, compared to 273 patients for the included trials. However not including those 11 trials could have partly affected the results of the meta-analysis.

There are many unanswered questions in the management of NPC that our analysis could unfortunately not address. Our analysis does not demonstrate a significant interaction between patient stage and treatment effect, although HR for progression or death for stage II patients was only 0.98 [0.57; 1.68] in our subgroup analysis. This analysis was greatly underpowered, with only 164 patients, as trials had to have all stages represented to be included. Hence one positive trial focusing mostly on stage II patients treated with 2D radiotherapy [34] was not included in this analysis due to the absence of all stage categories. Besides stage II is a highly heterogeneous category, with some patients




Fig. 2. Survival curves for overall survival by subsets of chemotherapy timing. A: Induction, B: Concomitant, C: Adjuvant, D: Concomitant + adjuvant, E: Induction (concomitant).

requiring CRT while others not. Another issue is the impact of EBV DNA before, during and after the course of radiotherapy and its influence on disease management. We attempted to collect EBV DNA levels pre-treatment or post-treatment, but those were missing for 3,948 patients

(87%) and 4,475 patients (99%) respectively. Therefore, EBV related parameters were not analyzed. Similarly, the management of keratinizing NPC usually follows the guidelines for endemic non-keratinizing NPC although there is no specific trial in this population. According to




Fig. 3. Survival curves for progression-free survival by subsets of chemotherapy timing. A: Induction, B: Concomitant, C: Adjuvant, D: Concomitant + adjuvant, E: Induction (concomitant).




Fig. 4. Survival curves for cancer and non-cancer deaths.

the protocol, histology type was to be analyzed as “WHO type I vs. II-III” due to its clinical relevance. But only three comparisons were eligible for such a subgroup analysis, which represented too few patients. In addition, staging and toxicity scoring evolved over time; hence, there is inherent heterogeneity between trials. Last, the included trials did not prospectively compare chemotherapy regimens used at the same timing,

so it was not possible to perform comparisons between different chemotherapy agents.

Looking at the absolute benefit of treatments at 5–10 years, a greater magnitude was achieved by Adjuvant + concomitant, followed by Induction (with concomitant in both arms). However, there are three reasons that can explain this difference. First, the induction (concomitant) group was more heterogeneous in terms of chemotherapy drugs. Second, the control arms were different: RT alone in the concomitant + adjuvant group and concomitant chemoradiation in the induction (concomitant) group, hence the control arm in the induction group did much better than in adjuvant + concomitant trials (5-year OS of 74% vs 58%). Last, the relative benefits of these two schedules appear similar, with respective HR for CRT-AC and IC-(CRT) of 0.68 [0.59; 0.79], 0.73 [0.63; 0.86] for death and 0.64 [0.56; 0.75] and 0.68 [0.60; 0.78] for progression or death.

Comparison of induction and adjuvant chemotherapy, both combined with concomitant chemoradiation, could not be performed in the setting of this meta-analysis, because we focused on the addition of a chemotherapy timing. While induction is easier to deliver than adjuvant, proponents of adjuvant claim that delivery of induction will impair the proper delivery of the concomitant phase, which is considered the cornerstone of treatment. The NPC-0501 trial has compared directly induction and adjuvant chemotherapy and suggests that induction could be associated with improved outcomes, especially when using conventionally fractionated radiotherapy [42]. Another way to perform such a comparison would be to conduct a network meta-analysis, which is planned as part of this update.

Even if outcomes are improved with the use of chemotherapy, a significant number of the patients will still relapse after treatment. In our report there are as many locoregional relapses as there are distant relapses. There is currently a lot of enthusiasm in the NPC community for checkpoint inhibitors, due to positive signals from two phase III

Table 1
Summary of the results, overall and by chemotherapy timing for all endpoints.

	Overall Survival	Progression-Free Survival	Cancer death	Loco-regional Failure	Distant Failure
All chemotherapy timings	HR: 0.79 [0.73; 0.85] AB5: +6.1% [+3.9; +8.3] AB10: +8.4% [+5.7; +11.1]	HR: 0.75 [0.70; 0.80] AB5: +8.5% [+6.2; +10.8] AB10: +9.3% [+6.7; +11.9]	HR: 0.74 [0.68; 0.81] AB5: +6.7% [+4.5; +8.9]	sHR: 0.79 [0.70; 0.89] AB5: -3.4% [-5.2; -1.6] AB10: -3.7% [-5.8; -1.6]	sHR: 0.70 [0.62; 0.78] AB5: -6.8% [-8.8; -4.8] AB10: -6.7% [-8.9; -4.5]
p-value	p < 0.0001	p < 0.0001	p < 0.0001	p = 0.0001	p < 0.0001
Interaction test	p = 0.03	p = 0.03	p = 0.255	p = 0.77	p = 0.55
Residual heterogeneity	p = 0.24	p = 0.27	NR	NR	NR
Induction	HR: 0.99 [0.80; 1.21] AB5: +1.7% [-5.8; +9.2] AB10: -1.4% [-9.9; +7.1]	HR: 0.82 [0.68; 0.97] AB5: +7.6% [+0.7; +14.5] AB10: +3.5% [-4.1; +11.1]	HR: 0.89 [0.71; 1.11]	sHR: 0.86 [0.67; 1.11]	sHR: 0.68 [0.52; 0.90]
Concomitant	HR: 0.81 [0.70; 0.92] AB5: +5.2% [+0.8; +9.6] AB10: +7.2% [+2.2; +12.2]	HR: 0.83 [0.73; 0.94] AB5: +6.3% [+1.7; +10.9] AB10: +6.1% [+1.2; +11.0]	HR: 0.72 [0.61; 0.87]	sHR: 0.82 [0.65; 1.03]	sHR: 0.76 [0.59; 0.97]
Adjuvant	HR: 0.90 [0.72; 1.11] AB5: +2.0% [-3.5; +7.5] AB10: +6.3% [-1.4; +14.0]	HR: 0.86 [0.71; 1.05] AB5: +3.4% [-2.6; +9.4] AB10: +7.7% [-0.6; +16.0]	HR: 0.82 [0.64; 1.04]	sHR: 0.73 [0.52; 1.04]	sHR: 0.84 [0.63; 1.11]
Concomitant and adjuvant	HR: 0.68 [0.59; 0.79] AB5: +12.3% [+7.0; +17.6] AB10: +13.9% [+8.1; +19.7]	HR: 0.64 [0.56; 0.75] AB5: +12.4% [+7.0; +17.8] AB10: +14.5% [+8.8; +20.2]	HR: 0.65 [0.54; 0.78]	sHR: 0.66 [0.49; 0.90]	sHR: 0.65 [0.51; 0.83]
Induction (concomitant)	HR: 0.73 [0.63; 0.86] AB5: +6.0% [+2.5; +9.5] AB10: +9.6% [+4.2; +15.0]	HR: 0.68 [0.60; 0.78] AB5: +10.2% [+6.3; +14.1] AB10: +11.6% [+6.6; +16.6]	HR: 0.72 [0.61; 0.85]	sHR: 0.81 [0.64; 1.02]	sHR: 0.63 [0.52; 0.77]

Abbreviations: AB5, absolute benefit at 5 years; AB10, absolute benefit at 10 years; HR, hazard ratio; NR, not relevant; sHR, subdistribution hazard ratio
Absolute benefits at 5 and 10 years were not calculated for each timing for cancer death, loco-regional failures or distant failures because interaction was not significant between treatment effect and chemotherapy timing.

Table 2
Efficacy of chemotherapy on overall and progression-free survival according to patient age.

	Overall Survival				Progression-Free Survival			
	No. deaths / No. patients		HR [95% CI]	AB5	No. events / No. patients		HR [95% CI]	AB5
	Control	Experimental			Control	Experimental		
<40	183/590	123/636	0.60 [0.48 ; 0.75]	+8.5%	231/590	159/636	0.58 [0.48 ; 0.71]	+12.3%
40–49	302/820	254/821	0.72 [0.61 ; 0.85]	+7.1%	366/820	298/821	0.70 [0.60 ; 0.81]	+10.0%
50–59	258/592	221/595	0.79 [0.66 ; 0.95]	+6.1%	288/592	255/595	0.80 [0.68 ; 0.95]	+6.9%
≥60	172/262	140/214	0.89 [0.71 ; 1.12]	+3.9%	178/262	137/214	0.89 [0.71 ; 1.12]	+4.1%
Interaction test	p = 0.08				p = 0.02			
Trend test	p = 0.01				p = 0.003			
Heterogeneity test (for interaction/trend)	p = 0.67/p = 0.23				p = 0.62/p = 0.19			

Subgroup analysis performed on the subset of trials using “modern” chemotherapy regimens (see Web table 9 for the definition of “old” drugs)
Abbreviations: AB5, Absolute benefit at 5 years; HR, Hazard Ratio

Table 3
Acute and late severe toxicities.

	Availability*		Incidence		OR [95% CI]	Efficacy	Heterogeneity	Interaction†
	No. comparisons	No. patients	CT‡	Control				
Acute								
Neutropenia	24	5,776	33.2%	7.3%	6.32 [5.50 ; 7.27]	p < 0.001	I ² = 82%, p < 0.001 ^a	p = 0.002
Febrile neutropenia	14	3,253	2.0%	1.8%	1.09 [0.68 ; 1.75]	p = 0.72	I ² = 7%, p = 0.38	p = 0.29
Thrombocytopenia	23	5,423	6.2%	1.6%	4.08 [3.16 ; 5.26]	p < 0.001	I ² = 62%, p < 0.001 ^b	p < 0.001
Anemia	24	5,776	5.8%	1.9%	3.17 [2.46 ; 4.08]	p < 0.001	I ² = 50%, p = 0.003 ^c	p = 0.047
Dermatitis	20	5,328	11.4%	9.3%	1.25 [1.04 ; 1.50]	p = 0.02	I ² = 35%, p = 0.06	p = 0.08
Weight loss	15	3,466	10.6%	6.1%	1.83 [1.43 ; 2.36]	p < 0.001	I ² = 45%, p = 0.03 ^d	p < 0.001
Mucositis	23	5,576	39.2%	33.5%	1.28 [1.15 ; 1.44]	p < 0.001	I ² = 61%, p < 0.001 ^e	p < 0.001
Hearing loss	19	4,633	1.9%	0.9%	2.16 [1.41 ; 4.14]	p < 0.001	I ² = 21%, p = 0.20	p < 0.001
Neurotoxicity	18	4,201	0.4%	0.3%	1.35 [0.67 ; 2.71]	p = 0.40	I ² = 0%, p > 0.99	p = 0.55
Nausea and vomiting	21	5,186	15.6%	8.6%	1.96 [1.65 ; 2.32]	p < 0.001	I ² = 63%, p < 0.001 ^f	p < 0.001
Late§								
Cutaneous fibrosis	15	4,064	2.2%	1.8%	1.24 [0.81 ; 1.90]	p = 0.32	I ² = 10%, p = 0.34	p = 0.18
Xerostomia	20	4,454	5.5%	4.5%	1.23 [0.94 ; 1.62]	p = 0.14	I ² = 0%, p = 0.86	p = 0.25
Bone necrosis	22	4,557	0.5%	0.6%	0.91 [0.50 ; 1.66]	p = 0.76	I ² = 0%, p > 0.99	p = 0.68
Hearing deficit	19	3,547	19.8%	16.0%	1.30 [1.08 ; 1.55]	p = 0.005	I ² = 0%, p = 0.28	p = 0.28
Cranial nerve palsy	18	3,483	4.6%	3.6%	1.28 [0.92 ; 1.79]	p = 0.14	I ² = 0%, p = 0.56	p = 0.78
Symptomatic temporal lobe necrosis	18	3,570	1.4%	1.5%	0.91 [0.55 ; 1.50]	p = 0.70	I ² = 0%, p = 0.97	p = 0.99
Brainstem or spinal cord damage	19	3,896	0.6%	0.5%	1.21 [0.64 ; 2.27]	p = 0.56	I ² = 0%, p = 0.98	p = 0.57
Trismus	21	4,341	4.2%	4.5%	0.93 [0.69 ; 1.25]	p = 0.61	I ² = 0%, p = 0.98	p = 0.31
Visual deficit	19	3,942	1.1%	1.1%	1.04 [0.60 ; 1.78]	p = 0.90	I ² = 0%, p = 0.99	p = 0.65
Massive bleeding	12	2,249	1.0%	1.2%	0.82 [0.40 ; 1.67]	p = 0.58	I ² = 0%, p = 0.98	p = 0.68

Toxicity was scored locally according to the scale used at the time of the trial and hence only graded according to NCI CTCAE v4 as severe (grade ≥ 3 for all except xerostomia where it was grade ≥ 2) vs not.

CI: Confidence Interval, CT: Chemotherapy, OR: Odds Ratio

* Only trials with available data for at least 60% of patients were included in the analyses

‡ Estimated with the Stewart et al method based on the toxicity rate in control arm and the odds ratio¹²

† Interaction between subsets of trials: induction, concomitant, adjuvant, concomitant + adjuvant, and induction (concomitant)

§ Only patients with a follow-up greater or equal to one year were included in the analyses

Residual heterogeneity:

a I² = 82%, p < 0.001; b I² = 9%, p = 0.34; c I² = 48%, p = 0.009; d I² = 0%, p = 0.77; e I² = 9%, p = 0.35; f I² = 0%, p = 0.58.

randomized trials of checkpoint inhibitors in combination with first line chemotherapy in recurrent or metastatic nasopharyngeal cancer [43–44]. Whether this will translate in an improvement for locally advanced disease is unknown. Indeed the recent trials of immunotherapy in head and neck squamous cell cancers have had disappointing results in the setting of chemoradiation [45].

In conclusion, this updated meta-analysis confirms the benefit of concomitant chemoradiation in locoregionally advanced NPC, especially with the addition of adjuvant or induction chemotherapy. The benefit of treatment decreases with increasing patient age.

Previous presentation

This work has not been presented at any meeting prior to this submission.

Funding

This research was funded by grants from Institut National du Cancer

(Programme Hospitalier de Recherche Clinique, grant PHRC-K15-189) and Ligue Nationale Contre le Cancer. The funding sources had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The submission of the paper for publication was decided by the MAC-NPC collaborative group. AA, PB, BL, and J-PP had access to the raw data. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication. All authors have seen and approved the final version and, after consultation with the collaborators, agreed to submit for publication.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary data

Supplementary data to this article and the protocol of the meta-analysis can be found online at <https://doi.org/10.1016/j.ctro.2021.11.007>.

References

- [1] Ferlay J, Ervik M, Lam F, et al: Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. [Internet][cited 2019 Sep 23] Available from: <https://gco.iarc.fr/today>, accessed.
- [2] Chen Y-P, Chan ATC, Le Q-T, Blanchard P, Sun Y, Ma J. Nasopharyngeal carcinoma. *Lancet* 2019;394(10192):64–80.
- [3] Blanchard P, Lee A, Marguet S, Leclercq J, Ng WT, Ma J, et al. Chemotherapy and radiotherapy in nasopharyngeal carcinoma: an update of the MAC-NPC meta-analysis. *Lancet Oncol* 2015;16(6):645–55.
- [4] Ribassin-Majed L, Marguet S, Lee AWM, Ng WT, Ma J, Chan ATC, et al. What Is the Best Treatment of Locally Advanced Nasopharyngeal Carcinoma? An Individual Patient Data Network Meta-Analysis. *J Clin Oncol* 2017;35(5):498–505.
- [5] Sun Y, Li W-F, Chen N-Y, et al. Induction chemotherapy plus concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in locoregionally advanced nasopharyngeal carcinoma: a phase 3, multicentre, randomised controlled trial. *Lancet Oncol* 2016;17:1509–20.
- [6] Frikha M, Auperin A, Tao Y, Elloumi F, Toumi N, Blanchard P, et al. A randomized trial of induction docetaxel–cisplatin–5FU followed by concomitant cisplatin-RT versus concomitant cisplatin-RT in nasopharyngeal carcinoma (GORTEC 2006–02). *Ann Oncol* 2018;29(3):731–6.
- [7] Zhang Y, Chen L, Hu G-Q, et al. Gemcitabine and Cisplatin Induction Chemotherapy in Nasopharyngeal Carcinoma. *N Engl J Med* 2019;381:1124–35.
- [8] Chan ATC, Hui EP, Ngan RKC, Tung SY, Cheng ACK, Ng WT, et al. Analysis of Plasma Epstein-Barr Virus DNA in Nasopharyngeal Cancer After Chemoradiation to Identify High-Risk Patients for Adjuvant Chemotherapy: A Randomized Controlled Trial. *JCO* 2018;36(31):3091–100.
- [9] Stewart LA. Practical methodology of meta-analyses (overviews) using updated individual patient data: Cochrane Working Group. *Stat Med* 1995;14(19):2057–79.
- [10] Schemper M, Smith TL. A note on quantifying follow-up in studies of failure time. *Control Clin Trials* 1996;17(4):343–6.
- [11] Yusuf S, Peto R, Lewis J, Collins R, Sleight P. Beta blockade during and after myocardial infarction: an overview of the randomized trials. *Prog Cardiovasc Dis* 1985;27(5):335–71.
- [12] Stewart LA, Parmar MKB. Meta-analysis of the literature or of individual patient data: is there a difference? *Lancet* 1993;341(8842):418–22.
- [13] Fine JP, Gray RJ. A Proportional Hazards Model for the Subdistribution of a Competing Risk. *J Am Stat Assoc* 1999;94(446):496–509.
- [14] Early Breast Cancer Trialists' Collaborative Group: Systemic treatment of early breast cancer by hormonal, cytotoxic, or immune therapy. 133 randomised trials involving 31,000 recurrences and 24,000 deaths among 75,000 women. *Lancet* 339:1–15, 1992.
- [15] Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21(11):1539–58.
- [16] Fisher DJ, Copas AJ, Tierney JF, Parmar MKB. A critical review of methods for the assessment of patient-level interactions in individual participant data meta-analysis of randomized trials, and guidance for practitioners. *J Clin Epidemiol* 2011;64(9):949–67.
- [17] Non-small Cell Lung Cancer Collaborative Group: Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials. *British medical journal* 311:899–909, 1995.
- [18] Chakraborty S, Chitapanarux I, Onchan W, Klunklin P, Supawongwattana B, Tharavichitkul E, et al. OP0005 Neoadjuvant chemotherapy plus chemoradiation (NACT-CCRT) versus chemoradiation plus adjuvant chemotherapy (CCRT-ACT) for locally advanced nasopharyngeal cancer: 3-Year results from a randomised trial. *Eur J Cancer* 2015;51:e2. <https://doi.org/10.1016/j.ejca.2015.06.009>.
- [19] Chan ATC, Teo PML, Leung TWT, Leung SF, Lee WY, Yeo W, et al. A prospective randomized study of chemotherapy adjunctive to definitive radiotherapy in advanced nasopharyngeal carcinoma. *Int J Radiat Oncol Biol Phys* 1995;33(3):569–77.
- [20] Tan T, Lim W-T, Fong K-W, Cheah S-L, Soong Y-L, Ang M-K, et al. Concurrent chemo-radiation with or without induction gemcitabine, Carboplatin, and Paclitaxel: a randomized, phase 2/3 trial in locally advanced nasopharyngeal carcinoma. *Int J Radiat Oncol Biol Phys* 2015;91(5):952–60.
- [21] Kwong DLW, Sham JST, Au GKH, Chua DTT, Kwong PWK, Cheng ACK, et al. Concurrent and adjuvant chemotherapy for nasopharyngeal carcinoma: A factorial study. *J Clin Oncol* 2004;22(13):2643–53.
- [22] Lee AWM, Tung SY, Chan ATC, Chappell R, Fu Y-T, Lu T-X, et al. A randomized trial on addition of concurrent-adjuvant chemotherapy and/or accelerated fractionation for locally-advanced nasopharyngeal carcinoma. *Radiother Oncol* 2011;98(1):15–22.
- [23] Huang P-Y, Zeng Qi, Cao K-J, Guo X, Guo L, Mo H-Y, et al. Ten-year outcomes of a randomised trial for locoregionally advanced nasopharyngeal carcinoma: A single-institution experience from an endemic area. *Eur J Cancer* 2015;51(13):1760–70.
- [24] Kwong DL, Sham JS, Au GK, Choy DT. Long-term results of concurrent and adjuvant chemotherapy for advanced nasopharyngeal carcinoma. *JCO* 2008;26(15 suppl):6056. 6056.
- [25] Chen L, Hu C-S, Chen X-Z, Hu G-Q, Cheng Z-B, Sun Y, et al. Adjuvant chemotherapy in patients with locoregionally advanced nasopharyngeal carcinoma: Long-term results of a phase 3 multicentre randomised controlled trial. *Eur J Cancer* 2017;75:150–8.
- [26] Al-Sarraf M, LeBlanc M, Giri PG, Fu KK, Cooper J, Vuong T, et al. Chemoradiotherapy versus radiotherapy in patients with advanced nasopharyngeal cancer: phase III randomized Intergroup study 0099. *J Clin Oncol* 1998;16(4):1310–7.
- [27] Chi K-H, Chang Y-C, Guo W-Y, Leung M-J, Shiau C-Y, Chen S-Y, et al. A phase III study of adjuvant chemotherapy in advanced nasopharyngeal carcinoma patients. *Int J Radiat Oncol Biol Phys* 2002;52(5):1238–44.
- [28] Chan ATC, Leung SF, Ngan RKC, Teo PML, Lau WH, Kwan WH, et al. Overall survival after concurrent cisplatin-radiotherapy compared with radiotherapy alone in locoregionally advanced nasopharyngeal carcinoma. *J Natl Cancer Inst* 2005;97(7):536–9.
- [29] Wee J, Tan EH, Tai BC, Wong HB, Leong SS, Tan T, et al. Randomized trial of radiotherapy versus concurrent chemoradiotherapy followed by adjuvant chemotherapy in patients with American Joint Committee on Cancer/International Union against cancer stage III and IV nasopharyngeal cancer of the endemic variety. *J Clin Oncol* 2005;23(27):6730–8.
- [30] Lee AWM, Tung SY, Chua DTT, Ngan RKC, Chappell R, Tung R, et al. Randomized trial of radiotherapy plus concurrent-adjuvant chemotherapy vs radiotherapy alone for regionally advanced nasopharyngeal carcinoma. *J Natl Cancer Inst* 2010;102(15):1188–98.
- [31] Fountzilas G, Ciuleanu E, Bobos M, Kologera-Fountzila A, Eleftheraki AG, Karayannopoulou G, et al. Induction chemotherapy followed by concomitant radiotherapy and weekly cisplatin versus the same concomitant chemoradiotherapy in patients with nasopharyngeal carcinoma: a randomized phase II study conducted by the Hellenic Cooperative Oncology Group (HeCOG) with biomarker evaluation. *Ann Oncol* 2012;23(2):427–35.
- [32] Wu X, Huang PY, Peng PJ, Lu LX, Han F, Wu SX, et al. Long-term follow-up of a phase III study comparing radiotherapy with or without weekly oxaliplatin for locoregionally advanced nasopharyngeal carcinoma. *Ann Oncol* 2013;24(8):2131–6.
- [33] Huang P-Y, Cao K-J, Guo X, Mo H-Y, Guo L, Xiang Y-Q, et al. A randomized trial of induction chemotherapy plus concurrent chemoradiotherapy versus induction chemotherapy plus radiotherapy for locoregionally advanced nasopharyngeal carcinoma. *Oral Oncol* 2012;48(10):1038–44.
- [34] Chen Q-Y, Wen Y-F, Guo L, Liu H, Huang P-Y, Mo H-Y, et al. Concurrent chemoradiotherapy vs radiotherapy alone in stage II nasopharyngeal carcinoma: phase III randomized trial. *J Natl Cancer Inst* 2011;103(23):1761–70.
- [35] Hong RL, Hsiao CF, Ting LL, Ko JY, Wang CW, Chang JTC, et al. Final results of a randomized phase III trial of induction chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in patients with stage IVA and IVB nasopharyngeal carcinoma-Taiwan Cooperative Oncology Group (TCOG) 1303 Study. *Ann Oncol* 2018;29(9):1972–9.
- [36] International Nasopharynx Cancer Study Group, VUMCA I Trial: Preliminary results of a randomized trial comparing neoadjuvant chemotherapy (cisplatin, epirubicin, bleomycin) plus radiotherapy vs. radiotherapy alone in stage IV(> or = N2, M0) undifferentiated nasopharyngeal carcinoma: a positive effect on progression-free survival. *Int J Radiat Oncol Biol Phys* 35:463–469, 1996.
- [37] NCCN Guidelines - Head and Neck Cancers - 2021 [Internet][cited 2021 Jul 21] Available from: https://www.nccn.org/professionals/physician_gls/pdf/head-and-neck.pdf.
- [38] Chen Y-P, Ismaila N, Chua MLK, Colevas AD, Haddad R, Huang SH, et al. Chemotherapy in Combination With Radiotherapy for Definitive-Intent Treatment of Stage II-IVA Nasopharyngeal Carcinoma: CSCO and ASCO Guideline. *JCO* 2021;39(7):840–59.
- [39] Lacas B, Bourhis J, Overgaard J, et al. Role of radiotherapy fractionation in head and neck cancers (MARCH): an updated meta-analysis. *Lancet Oncol* 2017;18:1221–37.
- [40] Lacas B, Carmel A, Landais C, et al. Meta-analysis of chemotherapy in head and neck cancer (MACH-NC): An update on 107 randomized trials and 19,805 patients, on behalf of MACH-NC Group. *Radiother Oncol* 2021;156:281–93.
- [41] Fayard F, Petit C, Lacas B, Pignon JP. Impact of missing individual patient data on 18 meta-analyses of randomised trials in oncology: Gustave Roussy experience. *BMJ Open* 2018;8(8):e020499.
- [42] Lee AWM, Ngan RKC, Ng W-T, Tung SY, Cheng AAC, Kwong DLW, et al. NPC-0501 trial on the value of changing chemoradiotherapy sequence, replacing 5-fluorouracil with capecitabine, and altering fractionation for patients with advanced nasopharyngeal carcinoma. *Cancer* 2020;126(16):3674–88.
- [43] Yang Y, Qu S, Li J, Hu C, Xu M, Li W, et al. Camrelizumab versus placebo in combination with gemcitabine and cisplatin as first-line treatment for recurrent or metastatic nasopharyngeal carcinoma (CAPTAIN-1st): a multicentre, randomised, double-blind, phase 3 trial. *Lancet Oncol* 2021;22(8):1162–74.
- [44] Rui-hua Xu H-QM, Sun Yat-sen University Cancer Center SKL of O in SC, Department of Nasopharyngeal Carcinoma SYUCC, et al. JUPITER-02: Randomized, double-blind, phase III study of toripalimab or placebo plus gemcitabine and cisplatin as first-line treatment for recurrent or metastatic nasopharyngeal carcinoma (NPC). [Internet][cited 2021 Jun 28] Available from: <https://meetinglibrary.asco.org/record/196620/abstract>.
- [45] Lee NY, Ferris RL, Psyrri A, Haddad RI, Tahara M, Bourhis J, et al. Avelumab plus standard-of-care chemoradiotherapy versus chemoradiotherapy alone in patients with locally advanced squamous cell carcinoma of the head and neck: a randomised, double-blind, placebo-controlled, multicentre, phase 3 trial. *Lancet Oncol* 2021;22(4):450–62.

Further reading

- [46] Yuan C, Xu X-H, Luo S-W, et al: Which neoadjuvant chemotherapy regimen should be recommended for patients with advanced nasopharyngeal carcinoma?: A network meta-analysis. *Medicine (Baltimore)* 97:e11978, 2018.
- [47] Shen J, Sun C, Zhou M, et al. Combination treatment with cetuximab in advanced nasopharyngeal carcinoma patients: a meta-analysis. *ONCOTARGETS AND THERAPY* 2019;12:2477–94.
- [48] Tan TH, Soon YY, Cheo T, Ho F, Wong LC, Tey J, et al. Induction chemotherapy for locally advanced nasopharyngeal carcinoma treated with concurrent chemoradiation: A systematic review and meta-analysis. *Radiother Oncol* 2018; 129(1):10–7.
- [49] Zhang B, Li MM, Chen WH, Zhao JF, Chen WQ, Dong YH, et al. Association of Chemoradiotherapy Regimens and Survival Among Patients With Nasopharyngeal Carcinoma: A Systematic Review and Meta-analysis. *JAMA Netw Open* 2019;2(10): e1913619. <https://doi.org/10.1001/jamanetworkopen.2019.13619>.
- [50] Chua DTT, Sham JST, Choy D, Kwong DLW, Au GKH, Kwong PWK, et al. Patterns of failure after induction chemotherapy and radiotherapy for locoregionally advanced nasopharyngeal carcinoma: the Queen Mary Hospital experience. *Int J Radiat Oncol Biol Phys* 2001;49(5):1219–28.
- [51] Cvitkovic E, Eschwege F, Rahal M, et al. Preliminary results of a randomized trial comparing neoadjuvant chemotherapy (cisplatin, epirubicin, bleomycin) plus radiotherapy vs. radiotherapy alone in stage IV (\geq N2, M0) undifferentiated nasopharyngeal carcinoma: A positive effect on progression-free survival. *Int J Radiat Oncol Biol Phys* 1996;35:463–9.
- [52] Hareyama M, Sakata K-I, Shirato H, Nishioka T, Nishio M, Suzuki K, et al. A prospective, randomized trial comparing neoadjuvant chemotherapy with radiotherapy alone in patients with advanced nasopharyngeal carcinoma. *Cancer* 2002;94(8):2217–23.
- [53] Low WK, Toh ST, Wee J, Fook-Chong SMC, Wang DY. Sensorineural Hearing Loss After Radiotherapy and Chemoradiotherapy: A Single, Blinded, Randomized Study. *JCO* 2006;24(12):1904–9.
- [54] Lee AWM, Tung SY, Ng WT, Lee V, Ngan RKC, Choi HCW, et al. A multicenter, phase 3, randomized trial of concurrent chemoradiotherapy plus adjuvant chemotherapy versus radiotherapy alone in patients with regionally advanced nasopharyngeal carcinoma: 10-year outcomes for efficacy and toxicity: 10-Year Outcomes of CRT for Advanced NPC. *Cancer* 2017;123(21):4147–57.
- [55] Hui EP, Ma BB, Leung SF, King AD, Mo F, Kam MK, et al. Randomized phase II trial of concurrent cisplatin-radiotherapy with or without neoadjuvant docetaxel and cisplatin in advanced nasopharyngeal carcinoma. *J Clin Oncol* 2009;27(2):242–9.
- [56] Chen Y, Sun Y, Liang S-B, Zong J-F, Li W-F, Chen Mo, et al. Progress report of a randomized trial comparing long-term survival and late toxicity of concurrent chemoradiotherapy with adjuvant chemotherapy versus radiotherapy alone in patients with stage III to IVB nasopharyngeal carcinoma from endemic regions of China. *Cancer* 2013;119(12):2230–8.
- [57] Li X-Y, Chen Q-Y, Sun X-S, Liu S-L, Yan J-J, Guo S-S, et al. Ten-year outcomes of survival and toxicity for a phase III randomised trial of concurrent chemoradiotherapy versus radiotherapy alone in stage II nasopharyngeal carcinoma. *Eur J Cancer* 2019;110:24–31.
- [58] Yang Q, Cao S-M, Guo L, et al. Induction chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in locoregionally advanced nasopharyngeal carcinoma: long-term results of a phase III multicentre randomised controlled trial. *Eur J Cancer* 2019;119:87–96.
- [59] Li W, Chen N, Zhang N, et al. Concurrent chemoradiotherapy with/without induction chemotherapy in locoregionally advanced nasopharyngeal carcinoma: Long-term results of phase 3 randomized controlled trial. *Int J Cancer* 2019;145: 295–305.