

# Trends in overweight and obesity levels in Geneva: time for optimism?

**Pedro Marques-Vidal<sup>1</sup>, Verónica Gomez<sup>2</sup> and Idris Guessous<sup>3</sup>**

<sup>1</sup> Institute of Social and Preventive Medicine (IUMSP), Lausanne University Hospital, Lausanne, Switzerland ;<sup>2</sup> National Institute of Health Ricardo Jorge, Lisbon, Portugal and <sup>3</sup> Geneva University Hospital, Geneva, Switzerland

## Aim

To assess trends in overweight and obesity levels using objectively measured data.

## Methods

Data collected between 2005 and 2011 by the Bus Santé study on representative samples of the Geneva population.

## Results

**Table 1:** Trends in anthropometric data, by study year

	2005	2006	2007	2008	2009	2010	2011	p-value
Sample size	210	254	257	457	1023	1010	930	
BMI (Kg/m <sup>2</sup> )	24.90 ± 4.26	25.14 ± 3.84	24.97 ± 4.34	24.87 ± 4.08	24.97 ± 4.05	25.46 ± 4.48	25.38 ± 4.30	0.02
BMI groups (%)								
Normal	55.7	52.0	56.4	57.1	53.7	51.3	52.7	
Overweight	34.3	37.8	33.9	30.9	35.5	34.0	33.6	0.14
Obese	10.0	10.2	9.7	12.0	10.9	14.8	13.8	
Waist (cm)	88.0 ± 14.2	89.0 ± 13.4	87.1 ± 12.9	84.9 ± 13.0	88.6 ± 42.9	88.6 ± 19.7	88.0 ± 19.2	0.58
Abd obesity (%)	24.5	28.1	21.6	18.2	23.1	27.6	22.3	0.003
WHR	0.87 ± 0.10	0.80 ± 0.10	0.87 ± 0.09	0.86 ± 0.10	0.88 ± 0.28	0.88 ± 0.17	0.90 ± 0.18	0.005
High WHR (%)	47.1	48.6	43.5	42.1	43.8	49.3	59.4	<0.001

Results are expressed as percentage or as mean ± Standard deviation. BMI, body mass index; Abd obesity, abdominal obesity; WHR; waist/hip ratio. Statistical analysis by chi-square or linear regression using calendar year as an independent variable.

**Table 2:** Trends in anthropometric data, by study year, stratified by gender

	2005	2006	2007	2008	2009	2010	2011	p-value
<b>Men (N)</b>								
BMI (Kg/m <sup>2</sup> )	26.05 ± 3.87	26.13 ± 2.60	25.60 ± 3.91	25.84 ± 3.62	26.06 ± 3.65	26.14 ± 3.90	26.12 ± 3.67	0.39
BMI groups (%)								
Normal	40.2	38.2	45.4	45.8	40.8	43.7	42.9	
Overweight	48.5	49.6	46.9	41.0	46.0	40.2	42.4	0.37
Obese	11.3	12.2	7.7	13.2	13.2	16.1	14.7	
Waist (cm)	95.6 ± 13.1	95.3 ± 12.1	92.1 ± 11.2	91.2 ± 10.9	93.5 ± 11.7	94.8 ± 23.6	93.3 ± 10.6	0.83
Abd obesity (%)	25.5	26.7	16.4	15.6	20.7	23.0	19.2	0.09
WHR	0.94 ± 0.07	0.94 ± 0.07	0.92 ± 0.07	0.93 ± 0.09	0.93 ± 0.08	0.95 ± 0.21	0.95 ± 0.08	0.08
High WHR (%)	72.3	68.7	59.4	62.4	65.1	67.3	75.7	0.001
<b>Women (N)</b>								
BMI (Kg/m <sup>2</sup> )	23.91 ± 4.35	24.09 ± 3.82	24.32 ± 4.66	23.92 ± 4.28	23.99 ± 4.14	24.80 ± 4.89	24.66 ± 4.71	0.008
BMI groups (%)								
Normal	69.0	66.7	67.7	68.3	65.2	58.7	62.1	
Overweight	22.1	25.2	20.5	20.9	26.0	27.9	25.1	0.15
Obese	8.9	8.1	11.8	10.9	8.7	13.5	12.8	
Waist (cm)	81.4 ± 11.7	82.3 ± 11.3	82.0 ± 12.6	78.5 ± 11.8	84.2 ± 57.8	82.6 ± 12.2	82.9 ± 23.6	0.44
Abd obesity (%)	23.6	29.5	26.8	20.9	25.3	32.1	25.3	0.05
WHR	0.80 ± 0.07	0.81 ± 0.07	0.81 ± 0.07	0.80 ± 0.07	0.82 ± 0.37	0.82 ± 0.08	0.85 ± 0.23	0.02
High WHR (%)	25.5	27.1	27.6	21.4	24.8	31.7	43.8	<0.001

Results are expressed as percentage or as mean ± Standard deviation. BMI, body mass index; Abd obesity, abdominal obesity; WHR; waist/hip ratio. Statistical analysis by chi-square or linear regression using calendar year as an independent variable.

## Conclusion

Overweight and obesity levels appear to have levelled in Geneva, with a possible decrease in overweight in men.

Still, the decrease in overweight appears to be partly compensated by an increase in obesity,

These favourable findings should be confirmed in other locations.

**Table 3:** Trends in anthropometric data, by study year, stratified by education

	2005	2006	2007	2008	2009	2010	2011	p-value
<b>Non-university (N)</b>								
BMI (Kg/m <sup>2</sup> )	25.63 ± 4.10	24.42 ± 4.02	25.74 ± 5.04	25.39 ± 4.30	25.47 ± 4.11	25.85 ± 4.53	25.89 ± 4.67	0.15
BMI groups (%)								
Normal	46.4	49.1	53.5	49.6	47.5	46.8	48.4	
Overweight	41.6	37.9	31.0	35.3	39.5	36.8	34.1	0.49
Obese	12.0	13.0	15.5	15.1	13.1	16.4	17.5	
Waist (cm)	90.8 ± 14.0	89.3 ± 13.7	88.7 ± 14.2	85.7 ± 13.1	91.4 ± 54.1	89.8 ± 23.3	89.5 ± 23.8	0.79
Abd obesity (%)	30.1	32.0	27.7	20.6	27.3	31.7	27.2	0.06
WHR	0.89 ± 0.10	0.88 ± 0.10	0.88 ± 0.10	0.87 ± 0.10	0.89 ± 0.35	0.89 ± 0.21	0.91 ± 0.23	0.16
High WHR (%)	58.5	50.3	48.2	45.9	51.2	55.0	61.0	0.001
<b>University (N)</b>								
BMI (Kg/m <sup>2</sup> )	23.81 ± 4.29	24.59 ± 3.40	24.01 ± 3.03	24.21 ± 3.68	24.23 ± 3.84	24.93 ± 4.37	24.78 ± 3.73	0.003
BMI groups (%)								
Normal	69.4	57.7	60.0	66.8	62.9	57.3	57.7	
Overweight	23.5	37.7	37.4	25.1	29.5	30.1	32.9	0.009
Obese	7.1	4.7	2.6	8.0	7.6	12.6	9.4	
Waist (cm)	83.7 ± 13.5	88.5 ± 12.8	85.2 ± 10.9	83.9 ± 12.8	84.3 ± 13.8	87.0 ± 13.1	86.3 ± 11.5	0.103
Abd obesity (%)	16.1	20.2	14.0	15.0	16.9	22.1	16.6	0.21
WHR	0.83 ± 0.09	0.87 ± 0.09	0.85 ± 0.08	0.85 ± 0.11	0.85 ± 0.11	0.87 ± 0.09	0.88 ± 0.09	<0.001
High WHR (%)	29.6	45.2	37.7	37.0	32.7	41.6	57.5	<0.001

Results are expressed as percentage or as mean ± Standard deviation. BMI, body mass index; Abd obesity, abdominal obesity; WHR; waist/hip ratio. Statistical analysis by chi-square or linear regression using calendar year as an independent variable.

**Table 4:** multivariate-adjusted anthropometric data, by study year

	2005	2006	2007	2008	2009	2010	2011	P-value *
BMI (Kg/m <sup>2</sup> )	24.93 ± 0.28	24.82 ± 0.37	24.95 ± 0.25	24.89 ± 0.19	24.96 ± 0.13	25.37 ± 0.13	25.37 ± 0.13	0.005
BMI groups								
Overweight	1 (ref.)	0.94 (0.56 - 1.59)	0.94 (0.62 - 1.42)	0.86 (0.59 - 1.25)	1.06 (0.75 - 1.48)	1.04 (0.74 - 1.46)	1.02 (0.73 - 1.44)	0.39
Obese	1 (ref.)	0.86 (0.38 - 1.96)	0.95 (0.50 - 1.81)	1.19 (0.68 - 2.08)	1.12 (0.67 - 1.88)	1.56 (0.94 - 2.61)	1.48 (0.88 - 2.48)	0.003
Waist (cm)	88.1 ± 1.8	87.1 ± 2.3	86.9 ± 1.6	84.7 ± 1.3	88.6 ± 0.8	88.2 ± 0.8	88.0 ± 0.8	0.33
Abd. obesity	1 (ref.)	1.07 (0.63 - 1.81)	0.84 (0.53 - 1.33)	0.68 (0.45 - 1.05)	0.92 (0.64 - 1.33)	1.15 (0.80 - 1.65)	0.90 (0.62 - 1.31)	0.43
WHR	0.87 ± 0.01	0.87 ± 0.02	0.86 ± 0.01	0.86 ± 0.01	0.88 ± 0.01	0.88 ± 0.01	0.90 ± 0.01	<0.005
High WHR	1 (ref.)	0.82 (0.48 - 1.40)	0.76 (0.50 - 1.18)	0.71 (0.48 - 1.05)	0.80 (0.56 - 1.14)	1.00 (0.71 - 1.43)	1.86 (1.30 - 2.67)	<0.001

Results are expressed as relative risk ratio (95% confidence interval) or as multivariate-adjusted mean ± standard error. Statistical analysis by multinomial logistic regression or multivariate analysis of variance. Adjusting for age, gender, education (2 groups), smoking (3 groups), nationality (Swiss and non-Swiss).