

## RESEARCH ARTICLE

# Association of IBD specific treatment and prevalence of pain in the Swiss IBD cohort study

Lorenz Bon<sup>1</sup>, Sylvie Scharl<sup>1</sup>, Stephan Vavricka<sup>1</sup>, Gerhard Rogler<sup>1,2</sup>, Nicolas Fournier<sup>3</sup>, Valerie Pittet<sup>3</sup>, Michael Scharl<sup>1,2</sup>, Thomas Greuter<sup>1</sup>, Philipp Schreiner<sup>1</sup>, Pascal Frei<sup>4</sup>, Benjamin Misselwitz<sup>1</sup>, Luc Biedermann<sup>1</sup>, Jonas Zeitz<sup>1,5</sup>\*, for the Swiss IBD Cohort Study Group<sup>†</sup>

**1** Division of Gastroenterology and Hepatology, University Hospital Zurich, University of Zurich, Zurich, Switzerland, **2** Zurich Center for Integrative Human Physiology, University of Zurich, Zurich, Switzerland, **3** Institute of Social and Preventive Medicine, University of Lausanne, Lausanne, Switzerland, **4** Gastroenterology Bethanien, Zurich, Switzerland, **5** Center of Gastroenterology, Clinic Hirslanden, Zurich, Switzerland

\* These authors contributed equally to this work.

† Membership of the SIBDCS study group is provided in the Acknowledgments.

\* [jonas.zeitz@gastrozentrum.ch](mailto:jonas.zeitz@gastrozentrum.ch)



## OPEN ACCESS

**Citation:** Bon L, Scharl S, Vavricka S, Rogler G, Fournier N, Pittet V, et al. (2019) Association of IBD specific treatment and prevalence of pain in the Swiss IBD cohort study. PLoS ONE 14(4): e0215738. <https://doi.org/10.1371/journal.pone.0215738>

**Editor:** Stefanos Bonovas, Humanitas University, ITALY

**Received:** September 17, 2018

**Accepted:** April 8, 2019

**Published:** April 25, 2019

**Copyright:** © 2019 Bon et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data may be made available upon request due to ethical restrictions imposed by the Swiss IBD cohort study group which collected this data. Interested researchers may request access to the collected data from this study in the same manner as the authors did. Data is available upon ethical approval and a request to the Head of the cohort, Dr. Phil Gerhard Rogler, at [gerhard.rogler@usz.ch](mailto:gerhard.rogler@usz.ch). Data access requests may also be made to the Swiss IBD Cohort at [sibdcs-submission@chuv.ch](mailto:sibdcs-submission@chuv.ch) or <http://ibdcohort.ch/index.php/informationen-fuer-forscher.html>.

## Abstract

### Background

Extraintestinal manifestations (EIM) contribute significantly to the burden of disease in inflammatory bowel disease (IBD). Pain is a leading symptom in IBD and could be seen as an EIM itself. Treatment of IBD associated pain is challenging and insufficiently studied. A better knowledge on the association of pain and IBD specific treatment is warranted to improve the management of IBD patients.

### Methods

All patients of the Swiss IBD Cohort Study (SIBDCS) (n = 2152) received a questionnaire regarding pain localization, pain character, and the use of IBD specific medication.

### Results

1263 completed questionnaires were received. Twenty-one out of 184 patients (10%) receiving anti-TNF treatment compared to 142 out of 678 patients (21%) not receiving anti-TNF medication reported elbow pain (p = 0.002) while 28 out of 198 patients (14%) receiving steroid treatment significantly more often reported elbow pain compared to 59 from 696 patients (8%) not receiving steroids (p = 0.021). Furthermore, we found significantly more female patients under anti-TNF treatment to report knee/ lower leg pain and ankle/ foot pain compared to their male counterparts (36% vs. 20% and 22% vs. 10%, respectively, p = 0.015 for both comparisons). The frequency of knee, lower leg, ankle and foot pain was especially low in male patients under anti-TNF treatment, indicating a high benefit of male patients from anti-TNF therapy regarding EIM.

**Funding:** This research was supported by a research grant from the Swiss National Science Foundation (<http://www.snf.ch/de/Seiten/default.aspx>) to GR for the Swiss IBD Cohort (Grant No. 3347CO-108792). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Conclusions

The frequency of elbow pain was lower in IBD patients treated with anti-TNF but higher in patients treated with steroids.

## Introduction

Pain is a common symptom in patients with inflammatory bowel disease (IBD) [1, 2]. In a recent study we showed that the vast majority of patients (71%) within the Swiss IBD Cohort Study experienced pain during their disease course and that for 52% of the patients pain was a longstanding problem [3]. Abdominal pain can be a direct or indirect consequence of intestinal inflammation; however, extraintestinal manifestations (EIM) of IBD can also cause pain and pain in itself can be seen as an EIM [4–6]. The most common EIM of IBD are arthropathies [5, 7–17]. Also, in our former study we could show that pain has a substantial impact on health-related quality of life (HRQOL) of IBD patients, as the general quality of life was significantly lower in patients suffering from pain compared to those without pain [3]. Such a relationship has also been described in other chronic diseases [18–21].

Treatment of both, IBD and IBD associated pain is challenging. The mainstay of IBD treatment includes systemic immunosuppressive medications, such as corticosteroids, anti-tumor necrosis factor (TNF) antibodies or immunomodulators. Furthermore, the management of an acute flare differs from the strategies for maintenance of remission [22, 23]. Moreover, presence of EIM will also influence the choice of a treatment regime. For instance, anti-TNF therapy is known to be very effective regarding gut inflammation as well as arthropathies/ arthritis.

Furthermore, non-steroidal anti-inflammatory drugs (NSAIDs) can very effectively mediate pain relieve due to their analgetic and also anti-inflammatory effects. However, due to the risk of disease exacerbation and induction of flares their use in IBD is limited [24–30].

Here, we used the well-characterized patient collective of the Swiss IBD Cohort Study (SIBDCS) to study the association of pain and IBD treatment with a focus on anti-TNF treatment.

## Methods

### Ethics consideration

Ethics approval was obtained from the regional Swiss Ethics Committees in which cohort participants were enrolled (Commission d'éthique du Canton de Vaud, Lausanne, Switzerland/ Protocol no. 33/06). Written, informed consent was obtained from each patient included in the study. The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in a priori approval by the institution's human research committee.

### Study design

Patients of the nationwide SIBDCS have been prospectively included since 2006 with a yearly follow-up. The cohort goals and methodology of SIBDC have been described elsewhere [31].

A questionnaire addressing various aspects of pain including pain duration localization and frequency was mailed to 2152 SIBDC patients, representing the entire cohort. The questionnaire also inquired about the use of pain specific medication in detail. Our questionnaire contained several questions from a validated German pain questionnaire [32]. The questionnaire was used in a German and in a French version. Further details of the questionnaires including

the fully originally used French and German versions are described elsewhere [3]. Basic epidemiological and clinical data including the use of IBD specific therapy was retrieved from the SIBDCS databases. All data are stored in Microsoft Access (Microsoft Corporation) databases.

## Statistical analysis

Descriptive statistical analyses were performed: Categorical variables were summarized as frequencies and percentages, whereas quantitative variables as median and range. To assess differences in categorical data distribution between groups of different sizes, Fisher's exact test was used.

The statistical analysis was performed using GraphPad Prism 7 for MacOS. A p-value of <0.05 was considered statistically significant.

## Results

### Patient's characteristics

The patients' characteristics shown in Table 1 have been described previously [3]. In brief: 1263 out of 2152 patients completed the questionnaires (response rate 59%). 599 out of 1263 patients were male (47%) and 664 female (53%). The median age was 47 years. Extraintestinal manifestations (EIM) of IBD were present in 699 patients (55%). The median IBD disease duration was 15 years (mean: 15 years, range: 0–57 years). The vast majority of patients (894/1263, 71%) reported the experience of pain in general during the course of the disease. Table 2 shows the frequency of IBD specific treatment.

### Association between IBD specific treatment and pain localization

When comparing the use of IBD specific medication and ten different pain localizations, we found several significant differences. Regarding elbow pain, only 21 patients (10%) receiving anti-TNF treatment compared to 142 patients (21%) not receiving anti-TNF were affected ( $p = 0.002$ ). Other pain localizations did not reveal significant differences regarding anti-TNF treatment (Table 3).

Comparing other IBD specific therapy and the different pain localization, we found patients not receiving steroid treatment significantly less often to be suffering from elbow pain compared to patients receiving steroids (8% vs. 14%;  $p = 0.021$ ). For the evaluation of other pain

**Table 1. Patient characteristics.**

Patient characteristics		Number of patients (%)
Gender	Female	664 (53)
	Male	599 (47)
Diagnosis	CD	679 (54)
	UC	556 (44)
	IC	28 (2)
	Sum	1263 (100)
Pain	Yes	894 (71)
	No	369 (29)
EIM	Yes	699 (55)
	No	564 (45)
Disease duration (Years)	Average	15
	Min-Max	0–57

<https://doi.org/10.1371/journal.pone.0215738.t001>

Table 2. IBD specific treatment.

IBD treatment	Number of patients (%)	
	with pain	without pain
Anti-TNF	216 (24.2)	100 (27.1)
Steroids	198 (22.1)	73 (19.8)
5-aminosalicylic acid (5-ASA)	334 (37.4)	136 (36.9)
Antibiotics	11 (1.2)	6 (1.6)
Calcineurin-Inhibitors	12 (1.3)	5 (1.4)
Immunomodulators	316 (35.3)	125(33.9)

<https://doi.org/10.1371/journal.pone.0215738.t002>

localizations and other IBD specific therapy (5-ASA, calcineurin-inhibitors, immunomodulators) no significant differences were observed (S1–S5 Tables).

### Association between IBD specific treatment and duration of pain

Duration of pain did not differ between patients on anti-TNF treatment versus those not on anti-TNF treatment (Table 4). The duration of pain was also not influenced by other IBD specific medications (Steroids, 5-ASA, Antibiotics, Calcineurin-inhibitors, Immunomodulators) also, no significant differences were observed (S6–S10 Tables).

### Association between IBD specific treatment and frequency of pain

The frequency of pain in patients with and without anti-TNF treatment did not significantly differ (Table 5). When comparing the pain frequencies of patients taking other IBD specific medications (steroids, 5-ASA, antibiotics, calcineurin-inhibitors, immunomodulators), also no significant differences were observed (S11–S15 Tables).

### Association between IBD specific treatment and pain character

Further, there was no association between the pain character and the use of IBD specific medication. 36 patients (20%) on anti-TNF treatment described their pain to be constant with slight fluctuations compared to 115 patients (19%) without anti-TNF treatment ( $p > 0.999$ ). Constant pain with strong fluctuations was reported by 15 patients (8%) using anti-TNF treatment and by 64 patients (11%) not receiving anti-TNF treatment ( $p = 0.331$ ). 115 patients (61%) with TNF treatment and 349 (58%) without anti-TNF treatment experienced pain attacks with pain

Table 3. Pain localization.

Pain localization	Anti-TNF	No anti-TNF	p-value
	N (%)	N (%)	
Head	56 (26)	147 (21.7)	0.193
Neck	28 (13)	95 (14)	0.735
Finger/hand	53 (24.5)	142 (20.9)	0.297
Elbow	21 (9.7)	142 (20.9)	<b>0.002</b>
Shoulder	44 (20.4)	138 (20.4)	>0.999
Back	77 (35.6)	236 (34.8)	0.869
Hip/thigh	52 (24)	162 (23.9)	>0.999
Knee/lower leg	61 (28.2)	181 (26.7)	0.660
Ankle/foot	35 (16.2)	109 (16.1)	>0.999
Abdomen	105 (48.6)	375 (55.3)	0.099

<https://doi.org/10.1371/journal.pone.0215738.t003>

Table 4. Duration of pain.

	Anti-TNF	No anti-TNF	
Pain peroid	N (%)	N (%)	p-value
<1 month	6 (2.7)	9 (1.3)	0.218
1 month-½ year	20 (9.3)	37 (5.5)	0.054
½ year-1 year	13 (6)	46 (6.8)	0.608
1–2 years	18 (8.3)	61 (9)	0.890
2–5 years	51 (23.6)	164 (24.2)	0.927
>5 years	108 (50)	361 (53.2)	0.434

<https://doi.org/10.1371/journal.pone.0215738.t004>

free intervals ( $p = 0.554$ ). Pain attacks with constant pain were reported by 30 patients (16%) receiving anti-TNF treatment compared to 75 (12%) not receiving anti-TNF ( $p = 0.268$ ). When comparing the pain character of patients receiving other IBD specific medication (steroids, 5-ASA, antibiotics, calcineurin-inhibitors, immunomodulators), no significant differences across treatment groups were seen (S16–S21 Tables).

### Association between IBD specific treatment and duration of pain attacks

Moreover, the duration of pain attacks was not influenced by IBD specific medication (Table 6), neither with regards to anti-TNF nor other agents to treat IBD, including steroids, 5-ASA, Antibiotics, Calcineurin-Inhibitors, Immunomodulators (S22–S26 Tables).

### Comparison of pain localization of male and female patients with and without anti-TNF therapy

From a total of 894 patients, a similar fraction of male and female patients (24% for both) received anti-TNF therapy (Fig 1).

When comparing the pain localizations of male and female patients receiving anti-TNF treatment significantly fewer male patients with anti-TNF treatment suffered from knee/ lower leg pain compared to female patients receiving anti-TNF therapy (20% vs. 36%;  $p = 0.015$ ). Also, significantly fewer male patients receiving anti-TNF treatment reported ankle/ foot pain compared to female patients with anti-TNF treatment (10% vs. 22%;  $p = 0.015$ ). For the other pain localizations, no differences regarding gender were seen (Table 7).

We did not observe any differences in pain localizations in patients with versus without anti-TNF-therapy, neither in male nor female patients (Tables 8 and 9).

Table 5. Frequency of pain.

	Anti-TNF	No anti-TNF	
Pain Frequency	N (%)	N (%)	p-value
Several times daily	49 (28.2)	115 (22)	0.099
1x/day	11 (6.3)	34 (6.5)	>0.999
Several times per week	34 (19.5)	100 (19.1)	0.911
1x/week	11 (6.3)	26 (5)	0.557
Several times per month	28 (16.1)	102 (19.5)	0.369
1x/month	12 (6.9)	55 (10.5)	0.182
<1x/month	29 (16.7)	91 (17.4)	0.907

<https://doi.org/10.1371/journal.pone.0215738.t005>

Table 6. Duration of pain attacks.

	Anti-TNF	No anti-TNF	
Duration of pain attacks	N (%)	N (%)	p-value
Seconds	26 (14.9)	61 (11.9)	0.295
Minutes	56 (32)	158 (30.8)	0.777
Hours	55 (31.4)	175 (34)	0.577
<3 days	21 (12)	67 (13)	0.793
>5 days	17 (9.7)	53 (10.3)	0.885

<https://doi.org/10.1371/journal.pone.0215738.t006>

## Discussion

In our study population 5-aminosalicylic acid (5-ASA) (37%) was the most frequently used IBD specific medication, followed by immunomodulators (35%) and anti-TNF antibodies (24%). As for anti-TNF, Vavricka et al. showed that in more than 40% of the cases, this therapy regime is initiated to treat EIM rather than bowel inflammation and over 70% showed a clinical response of EIM to anti-TNF therapy [6]. Our study supports these findings: we could show that significantly less patients on anti-TNF reported elbow pain compared to patients not on anti-TNF. Of note, significantly more patients on steroid treatment reported elbow pain. References to support these findings are lacking.

Regarding gender specific differences in treatment of EIM/ pain in IBD patients, data is not consistent. Concerning IBD treatment, Lopetusa et al. found no general influence of the gender on the therapy of ulcerative colitis (UC) with anti-TNF (infliximab) [33]. However, female patients with steroid-refractory UC and successive anti-TNF treatment showed an increased 1-year remission rate and a cumulative non-colectomy rate. In contradiction, Lopetusa et al. found a lower rate of response to treatment and of disease remission in female patients under TNF inhibitors with axial spondyloarthritis [34]. As for possible explanations, Nguyen et al. showed that the three biomarkers praealbumin, platelet factor 4 and S100A12 accurately predict the response of patients with rheumatoid arthritis to TNF inhibitors [35]. Further studies about a gender-specific correlation of these marker could reveal useful findings. In our study, we found that statistically significant less male patients with anti-TNF treatment reported knee/ lower leg and foot/ ankle pain compared to female patients with anti-TNF. This data may indicate that there is a gender difference regarding the effect of anti-TNF therapy for EIM.

One strength of our study is the size of the cohort with 1263 completed questionnaires. Together with our former study evaluating pain in the SIBDCS [3] it is, to the best of our knowledge, the largest evaluation of pain and the use of IBD specific therapy in IBD up to date.

However, our study also has limitations. Due to the study design and the lack of control regarding unreturned questionnaires, a reporting bias cannot be excluded. Patients who actually suffer from pain due to IBD therefore might be overrepresented compared to patients without pain, since the former might be more motivated to return the questionnaire. The patients not responding to the survey might have represented a different phenotype regarding our topic of interest. The existing data of the SIBDCS doesn't include any information about pain, preventing us from comparing pain specific parameters between responders and non-responders. Furthermore, regarding the use of IBD specific therapy and pain localizations, we do not have information on the reason to initiate medical therapy (i.e. EIM vs. intestinal activity of IBD or both) and how high the prevalence of pain has been before treatment initiation. Our statistical evaluation of the data represents another limitation. We have performed a mostly descriptive analysis of the dataset. To remain a high response rate and not no




Fig 1. Male and female patients with and without Anti-TNF.

<https://doi.org/10.1371/journal.pone.0215738.g001>

overstrain the goodwill of the patients, we intended to keep our questionnaire on a simplistic level. We further see the sizes of our subgroups as a potential limitation. We are aware that small subgroups may be linked to random positive findings. In comparison to other studies about pain in IBD, the subgroups examined here are not considerably small. Furthermore, we aimed to include as many pain localizations as possible to thoroughly analyse the distribution of pain. Additionally, our findings, particularly regarding Anti-TNF, match the clinical observations, depicting a genuine outcome.

In summary, we could show that the frequency of elbow pain was lower in patients treated with anti-TNF but higher under steroid treatment. There were no significant differences regarding the use IBD specific therapy and the character, duration and frequency of pain. Furthermore, our data point towards a higher treatment benefit of anti-TNF with regards to EIM in male patients which should be followed up in future studies.

Table 7. Pain localization of male vs. female patients with anti-TNF.

Pain localization	Anti-TNF/ male	Anti-TNF/ female	p-value
	N (%)	N (%)	
Back	32 (30.8)	45 (40.2)	0.158
Knee/lower leg	21 (20.2)	40 (35.7)	<b>0.015</b>
Elbow	11 (10.6)	10 (8.9)	0.819
Hip/thigh	22 (21.2)	30 (26.8)	0.344
Finger/hand	21 (20.2)	32 (28.6)	0.158
Ankle/foot	10 (9.6)	25 (22.3)	<b>0.015</b>

<https://doi.org/10.1371/journal.pone.0215738.t007>

Table 8. Pain localization of male patients with vs. without anti-TNF.

Pain localization	Anti-TNF/ male	No anti-TNF/ male	p-value
	N (%)	N (%)	
Back	32 (30.8)	122 (38)	0.198
Knee/lower leg	21 (20.2)	89 (27.7)	0.156
Elbow	11 (10.6)	28 (8.7)	0.561
Hip/thigh	22 (21.2)	73 (22.7)	0.787
Finger/hand	21 (20.2)	69 (21.5)	0.890
Ankle/foot	10 (9.6)	51 (15.9)	0.146

<https://doi.org/10.1371/journal.pone.0215738.t008>

Table 9. Pain localization of female patients with vs. without anti-TNF.

Pain Localisation	anti-TNF/female	No anti-TNF/female	p-value
	N (%)	N (%)	
Back	45 (40.2)	114 (31.9)	0.110
Knee/lower leg	40 (35.7)	92 (25.8)	0.053
Elbow	10 (8.9)	38 (10.6)	0.721
Hip/thigh	30 (26.8)	89 (24.9)	0.709
Finger/hand	32 (28.6)	73 (20.4)	0.090
Ankle/foot	25 (22.3)	58 (16.2)	0.156

<https://doi.org/10.1371/journal.pone.0215738.t009>

## Supporting information

### S1 Table. Pain localization (Steroids).

(PDF)

### S2 Table. Pain localization (5-aminosalicylic acid).

(PDF)

### S3 Table. Pain localization (Immunomodulators).

(PDF)

### S4 Table. Pain localization (Antibiotics).

(PDF)

### S5 Table. Pain localization (Calcineurin-Inhibitors).

(PDF)

### S6 Table. Duration of pain (Steroids).

(PDF)

### S7 Table. Duration of pain (5-aminosalicylic acid).

(PDF)

### S8 Table. Duration of pain (Antibiotics).

(PDF)

### S9 Table. Duration of pain (Calcineurin-Inhibitors).

(PDF)

### S10 Table. Duration of pain (Immunomodulators).

(PDF)

- S11 Table. Frequency of pain (Steroids).**  
(PDF)
- S12 Table. Frequency of pain (5-aminosalicylic acid).**  
(PDF)
- S13 Table. Frequency of pain (Antibiotics).**  
(PDF)
- S14 Table. Frequency of pain (Calcineurin-Inhibitors).**  
(PDF)
- S15 Table. Frequency of pain (Immunomodulators).**  
(PDF)
- S16 Table. Pain character (Anti-TNF).**  
(PDF)
- S17 Table. Pain character (Steroids).**  
(PDF)
- S18 Table. Pain character (5-aminosalicylic acid).**  
(PDF)
- S19 Table. Pain character (Antibiotics).**  
(PDF)
- S20 Table. Pain character (Calcineurin-Inhibitors).**  
(PDF)
- S21 Table. Pain character (Immunomodulators).**  
(PDF)
- S22 Table. Duration of pain attacks (Steroids).**  
(PDF)
- S23 Table. Duration of pain attacks (5-aminosalicylic acid).**  
(PDF)
- S24 Table. Duration of pain attacks (Antibiotics).**  
(PDF)
- S25 Table. Duration of pain attacks (Calcineurin-Inhibitors).**  
(PDF)
- S26 Table. Duration of pain attacks (Immunomodulators).**  
(PDF)

## Acknowledgments

The authors thank all the patients for their collaboration and the members of the Swiss Inflammatory Bowel Disease Cohort Study for their contribution.

Members of the SIBDCS study group: Claudia Anderegg; Peter Bauerfeind; Christoph Beglinger; Stefan Bègré; Dominique Belli; José M. Bengoa; Luc Biedermann; Beat Bigler; Janek Binek; Mirjam Blattmann; Stephan Boehm; Jan Borovicka; Christian P. Braegger; Nora Brunner; Patrick Bühr; Bernard Burnand; Emanuel Burri; Sophie Buyse; Matthias Cremer; Dominique H. Criblez; Philippe de Saussure; Lukas Degen; Joakim Delarive; Christopher Doerig;

Barbara Dora; Gian Dorta; Mara Egger; Tobias Ehmann; Ali El-Wafa; Matthias Engelmann; Jessica Ezri; Christian Felley; Markus Fliegner; Nicolas Fournier; Montserrat Fraga; Pascal Frei; Remus Frei; Michael Fried; Florian Froehlich; Christian Funk; Raoul Ivano Furlano; Suzanne Gallot-Lavallée; Martin Geyer; Marc Girardin; Delphine Golay; Tanja Grandinetti; Beat Gysi; Horst Haack; Johannes Haarer; Beat Helbling; Peter Hengstler; Denise Herzog; Cyrill Hess; Klaas Heyland; Thomas Hinterleitner; Philippe Hiroz; Claudia Hirschi; Petr Hruz; Rika Iwata; Res Jost; Pascal Juillerat; Vera Kessler Brondolo; Christina Knellwolf; Christoph Knoblauch; Henrik Köhler; Rebekka Koller; Claudia Krieger-Grübel; Gerd Kullak-Ublick; Patrizia Künzler; Markus Landolt; Rupprecht Lange; Frank Serge Lehmann; Andrew Macpherson; Philippe Maerten; Michel H. Maillard; Christine Manser; Michael Manz; Urs Marbet; George Marx; Christoph Matter; Valérie McLin; Rémy Meier; Martina Mendanova; Christa Meyenberger; Pierre Michetti; Benjamin Misselwitz; Darius Moradpour; Bernhard Morell; Patrick Mosler; Christian Mottet; Christoph Müller; Pascal Müller; Beat Müllhaupt; Claudia Münger-Beyeler; Leilla Musso; Andreas Nagy; Michaela Neagu; Cristina Nichita; Jan Niess; Natacha Noël; Andreas Nydegger; Nicole Obialo; Carl Oneta; Cassandra Oropesa; Ueli Peter; Daniel Peternac; Laetitia Marie Petit; Franziska Piccoli-Gfeller; Julia Beatrice Pilz; Valérie Pittet; Nadia Raschle; Ronald Rentsch; Sophie Restellini; Jean-Pierre Richterich; Sylvia Rihs; Marc Alain Ritz; Jocelyn Roudit; Daniela Rogler; Gerhard Rogler; Jean-Benoît Rossel; Markus Sagmeister; Gaby Saner; Bernhard Sauter; Mikael Sawatzki; Michela Schäppi; Michael Scharl; Martin Schelling; Susanne Schibli; Hugo Schlauri; Sybille Schmid Uebelhart; Jean-François Schnegg; Alain Schoepfer; Frank Seibold; Mariam Seirafi; Gian-Marco Semadeni; David Semela; Arne Senning; Marc Sidler; Christiane Sokollik; Johannes Spalinger; Holger Spangenberg; Philippe Stadler; Michael Steuerwald; Alex Straumann; Bigna Straumann-Funk; Michael Sulz; Joël Thorens; Sarah Tiedemann; Radu Tutuian; Stephan Vavricka; Francesco Viani; Jürg Vöglin; Roland Von Känel; Alain Vonlaufen; Dominique Vouillamoz; Rachel Vulliamy; Jürg Wermuth; Helene Werner; Paul Wiesel; Reiner Wiest; Tina Wylie; Jonas Zeitz; Dorothee Zimmermann.

## Author Contributions

**Conceptualization:** Luc Biedermann, Jonas Zeitz.

**Data curation:** Nicolas Fournier, Valerie Pittet.

**Formal analysis:** Lorenz Bon, Luc Biedermann, Jonas Zeitz.

**Investigation:** Lorenz Bon.

**Methodology:** Lorenz Bon, Luc Biedermann, Jonas Zeitz.

**Project administration:** Stephan Vavricka, Gerhard Rogler, Philipp Schreiner, Pascal Frei, Luc Biedermann, Jonas Zeitz.

**Supervision:** Luc Biedermann, Jonas Zeitz.

**Writing – original draft:** Lorenz Bon, Jonas Zeitz.

**Writing – review & editing:** Sylvie Scharl, Valerie Pittet, Michael Scharl, Thomas Greuter, Benjamin Misselwitz, Luc Biedermann, Jonas Zeitz.

## References

1. Wagtmans MJ, Verspaget HW, Lamers CB, van Hogezaand RA. Crohn's disease in the elderly: a comparison with young adults. *J Clin Gastroenterol.* 1998; 27(2):129–33. PMID: [9754773](https://pubmed.ncbi.nlm.nih.gov/9754773/).

2. Aghazadeh R, Zali MR, Bahari A, Amin K, Ghahghaie F, Firouzi F. Inflammatory bowel disease in Iran: a review of 457 cases. *J Gastroenterol Hepatol*. 2005; 20(11):1691–5. <https://doi.org/10.1111/j.1440-1746.2005.03905.x> PMID: 16246187.
3. Zeitz J, Ak M, Muller-Mottet S, Scharl S, Biedermann L, Fournier N, et al. Pain in IBD Patients: Very Frequent and Frequently Insufficiently Taken into Account. *PLoS One*. 2016; 11(6):e0156666. Epub 2016/06/23. <https://doi.org/10.1371/journal.pone.0156666> PMID: 27332879; PubMed Central PMCID: PMC4917102.
4. Veloso FT, Carvalho J, Magro F. Immune-related systemic manifestations of inflammatory bowel disease. A prospective study of 792 patients. *J Clin Gastroenterol*. 1996; 23(1):29–34. PMID: 8835896.
5. Lakatos L, Pandur T, David G, Balogh Z, Kuronya P, Tollas A, et al. Association of extraintestinal manifestations of inflammatory bowel disease in a province of western Hungary with disease phenotype: results of a 25-year follow-up study. *World journal of gastroenterology: WJG*. 2003; 9(10):2300–7. <https://doi.org/10.3748/wjg.v9.i10.2300> PMID: 14562397; PubMed Central PMCID: PMC4656482.
6. Vavricka SR, Schoepfer A, Scharl M, Lakatos PL, Navarini A, Rogler G. Extraintestinal Manifestations of Inflammatory Bowel Disease. *Inflamm Bowel Dis*. 2015; 21(8):1982–92. Epub 2015/07/15. <https://doi.org/10.1097/MIB.0000000000000392> PMID: 26154136; PubMed Central PMCID: PMC4511685.
7. Salvarani C, Vlachonikolis IG, van der Heijde DM, Fornaciari G, Macchioni P, Beltrami M, et al. Musculoskeletal manifestations in a population-based cohort of inflammatory bowel disease patients. *Scandinavian journal of gastroenterology*. 2001; 36(12):1307–13. PMID: 11761022.
8. Vavricka SR, Brun L, Ballabeni P, Pittet V, Prinz Vavricka BM, Zeitz J, et al. Frequency and risk factors for extraintestinal manifestations in the Swiss inflammatory bowel disease cohort. *Am J Gastroenterol*. 2011; 106(1):110–9. <https://doi.org/10.1038/ajg.2010.343> PMID: 20808297.
9. Palm O, Bernklev T, Moum B, Gran JT. Non-inflammatory joint pain in patients with inflammatory bowel disease is prevalent and has a significant impact on health related quality of life. *The Journal of rheumatology*. 2005; 32(9):1755–9. PMID: 16142874.
10. D'Inca R, Podswiadek M, Ferronato A, Punzi L, Salvagnini M, Sturniolo GC. Articular manifestations in inflammatory bowel disease patients: a prospective study. *Dig Liver Dis*. 2009; 41(8):565–9. <https://doi.org/10.1016/j.dld.2009.01.013> PMID: 19278908.
11. Bernstein CN, Blanchard JF, Rawsthorne P, Yu N. The prevalence of extraintestinal diseases in inflammatory bowel disease: a population-based study. *Am J Gastroenterol*. 2001; 96(4):1116–22. <https://doi.org/10.1111/j.1572-0241.2001.03756.x> PMID: 11316157.
12. Lanna CC, Ferrari Mde L, Rocha SL, Nascimento E, de Carvalho MA, da Cunha AS. A cross-sectional study of 130 Brazilian patients with Crohn's disease and ulcerative colitis: analysis of articular and ophthalmologic manifestations. *Clin Rheumatol*. 2008; 27(4):503–9. <https://doi.org/10.1007/s10067-007-0797-5> PMID: 18097711.
13. Turkcapar N, Toruner M, Soykan I, Aydinoglu OT, Cetinkaya H, Duzgun N, et al. The prevalence of extraintestinal manifestations and HLA association in patients with inflammatory bowel disease. *Rheumatology international*. 2006; 26(7):663–8. <https://doi.org/10.1007/s00296-005-0044-9> PMID: 16136311.
14. de Vlam K, Mielants H, Cuvelier C, De Keyser F, Veys EM, De Vos M. Spondyloarthritis is underestimated in inflammatory bowel disease: prevalence and HLA association. *J Rheumatol*. 2000; 27(12):2860–5. Epub 2000/12/29. PMID: 11128677.
15. Orchard TR, Wordsworth BP, Jewell DP. Peripheral arthropathies in inflammatory bowel disease: their articular distribution and natural history. *Gut*. 1998; 42(3):387–91. PMID: 9577346; PubMed Central PMCID: PMC49172027.
16. Brynskov J, Binder V. Arthritis and the gut. *Eur J Gastroenterol Hepatol*. 1999; 11(9):997–9. PMID: 10503836.
17. van Erp SJ, Brakenhoff LK, van Gaalen FA, van den Berg R, Fidder HH, Verspaget HW, et al. Classifying Back Pain and Peripheral Joint Complaints in Inflammatory Bowel Disease Patients: A Prospective Longitudinal Follow-up Study. *Journal of Crohn's & colitis*. 2015. <https://doi.org/10.1093/ecco-jcc/jjv195> PMID: 26512134.
18. Kukreja M, Bryant AS, Cleveland DC, Dabal R, Hingorani N, Kirklín JK. Health-Related Quality of Life in Adult Survivors After the Fontan Operation. *Semin Thorac Cardiovasc Surg*. 2015; 27(3):299–306. <https://doi.org/10.1053/j.semtcvs.2015.08.007> PMID: 26708372.
19. Hussain KB, Fontana RJ, Moyer CA, Su GL, Sneed-Pee N, Lok AS. Comorbid illness is an important determinant of health-related quality of life in patients with chronic hepatitis C. *Am J Gastroenterol*. 2001; 96(9):2737–44. <https://doi.org/10.1111/j.1572-0241.2001.04133.x> PMID: 11569704.
20. Naliboff BD, Kim SE, Bolus R, Bernstein CN, Mayer EA, Chang L. Gastrointestinal and psychological mediators of health-related quality of life in IBS and IBD: a structural equation modeling analysis. *Am J*

- Gastroenterol. 2012; 107(3):451–9. <https://doi.org/10.1038/ajg.2011.377> PMID: 22085819; PubMed Central PMCID: PMC3855477.
21. Luo J, Hendryx M, Safford MM, Wallace R, Rossom R, Eaton C, et al. Newly Developed Chronic Conditions and Changes in Health-Related Quality of Life in Postmenopausal Women. *J Am Geriatr Soc*. 2015; 63(11):2349–57. <https://doi.org/10.1111/jgs.13796> PMID: 26503351.
  22. Dignass A, Lindsay JO, Sturm A, Windsor A, Colombel JF, Allez M, et al. Second European evidence-based consensus on the diagnosis and management of ulcerative colitis part 2: current management. *J Crohns Colitis*. 2012; 6(10):991–1030. <https://doi.org/10.1016/j.crohns.2012.09.002> PMID: 23040451.
  23. Dignass A, Van Assche G, Lindsay JO, Lemann M, Soderholm J, Colombel JF, et al. The second European evidence-based Consensus on the diagnosis and management of Crohn's disease: Current management. *J Crohns Colitis*. 2010; 4(1):28–62. <https://doi.org/10.1016/j.crohns.2009.12.002> PMID: 21122489.
  24. Ananthakrishnan AN, Higuchi LM, Huang ES, Khalili H, Richter JM, Fuchs CS, et al. Aspirin, nonsteroidal anti-inflammatory drug use, and risk for Crohn disease and ulcerative colitis: a cohort study. *Ann Intern Med*. 2012; 156(5):350–9. <https://doi.org/10.7326/0003-4819-156-5-201203060-00007> PMID: 22393130; PubMed Central PMCID: PMC3369539.
  25. Long MD, Kappelman MD, Martin CF, Chen W, Anton K, Sandler RS. Role of Nonsteroidal Anti-Inflammatory Drugs in Exacerbations of Inflammatory Bowel Disease. *J Clin Gastroenterol*. 2015. <https://doi.org/10.1097/MCG.0000000000000421> PMID: 26485106.
  26. Takeuchi K, Smale S, Premchand P, Maiden L, Sherwood R, Thjodleifsson B, et al. Prevalence and mechanism of nonsteroidal anti-inflammatory drug-induced clinical relapse in patients with inflammatory bowel disease. *Clinical gastroenterology and hepatology: the official clinical practice journal of the American Gastroenterological Association*. 2006; 4(2):196–202. PMID: 16469680.
  27. Bjarnason I, Hayllar J, MacPherson AJ, Russell AS. Side effects of nonsteroidal anti-inflammatory drugs on the small and large intestine in humans. *Gastroenterology*. 1993; 104(6):1832–47. PMID: 8500743.
  28. Kaufmann HJ, Taubin HL. Nonsteroidal anti-inflammatory drugs activate quiescent inflammatory bowel disease. *Ann Intern Med*. 1987; 107(4):513–6. PMID: 3498419.
  29. Miner PB Jr. Factors influencing the relapse of patients with inflammatory bowel disease. *Am J Gastroenterol*. 1997; 92(12 Suppl):1S–4S. PMID: 9395345.
  30. Kefalakes H, Stylianides TJ, Amanakis G, Kolios G. Exacerbation of inflammatory bowel diseases associated with the use of nonsteroidal anti-inflammatory drugs: myth or reality? *European journal of clinical pharmacology*. 2009; 65(10):963–70. <https://doi.org/10.1007/s00228-009-0719-3> PMID: 19711064.
  31. Pittet V, Juillerat P, Mottet C, Felley C, Ballabeni P, Burnand B, et al. Cohort profile: the Swiss Inflammatory Bowel Disease Cohort Study (SIBDCS). *Int J Epidemiol*. 2009; 38(4):922–31. Epub 2008/09/11. <https://doi.org/10.1093/ije/dyn180> PMID: 18782896.
  32. Nagel B, Gerbershagen HU, Lindena G, Pflingsten M. [Development and evaluation of the multidimensional German pain questionnaire]. *Schmerz*. 2002; 16(4):263–70. Epub 2002/08/23. <https://doi.org/10.1007/s00482-002-0162-1> PMID: 12192435.
  33. Lopetuso LR, Gerardi V, Papa V, Scaldaferrri F, Rapaccini GL, Gasbarrini A, et al. Can We Predict the Efficacy of Anti-TNF-alpha Agents? *Int J Mol Sci*. 2017; 18(9). Epub 2017/09/15. <https://doi.org/10.3390/ijms18091973> PMID: 28906475; PubMed Central PMCID: PMC5618622.
  34. Lubrano E, Perrotta FM, Manara M, D'Angelo S, Addimanda O, Ramonda R, et al. The Sex Influence on Response to Tumor Necrosis Factor-alpha Inhibitors and Remission in Axial Spondyloarthritis. *J Rheumatol*. 2018; 45(2):195–201. Epub 2018/02/09. <https://doi.org/10.3899/jrheum.170666> PMID: 29419448.
  35. Nguyen MVC, Baillet A, Romand X, Trocme C, Courtier A, Marotte H, et al. Prealbumin, platelet factor 4 and S100A12 combination at baseline predicts good response to TNF alpha inhibitors in rheumatoid arthritis. *Joint Bone Spine*. 2018. Epub 2018/06/10. <https://doi.org/10.1016/j.jbspin.2018.05.006> PMID: 29885551.