



UNIL | Université de Lausanne

Unicentre

CH-1015 Lausanne

<http://serval.unil.ch>

Year : 2025

Associations Between Physical Activity Patterns and
Cardiovascular Events and Risk Factors. Cross-Sectional
and Prospective Studies
Fabrizio Cominetti

Cominetti Fabrizio, 2025, Associations Between Physical Activity Patterns and
Cardiovascular Events and Risk Factors. Cross-Sectional and Prospective Studies

Originally published at : Thesis, University of Lausanne

Posted at the University of Lausanne Open Archive <http://serval.unil.ch>
Document urn:nbn:ch:serval-BIB_369FCEC9C5A25

Droits d'auteur

L'Université de Lausanne attire expressément l'attention des utilisateurs sur le fait que tous les documents publiés dans l'Archive SERVAL sont protégés par le droit d'auteur, conformément à la loi fédérale sur le droit d'auteur et les droits voisins (LDA). A ce titre, il est indispensable d'obtenir le consentement préalable de l'auteur et/ou de l'éditeur avant toute utilisation d'une oeuvre ou d'une partie d'une oeuvre ne relevant pas d'une utilisation à des fins personnelles au sens de la LDA (art. 19, al. 1 lettre a). A défaut, tout contrevenant s'expose aux sanctions prévues par cette loi. Nous déclinons toute responsabilité en la matière.

Copyright

The University of Lausanne expressly draws the attention of users to the fact that all documents published in the SERVAL Archive are protected by copyright in accordance with federal law on copyright and similar rights (LDA). Accordingly it is indispensable to obtain prior consent from the author and/or publisher before any use of a work or part of a work for purposes other than personal use within the meaning of LDA (art. 19, para. 1 letter a). Failure to do so will expose offenders to the sanctions laid down by this law. We accept no liability in this respect.



UNIL | Université de Lausanne

Faculté de biologie
et de médecine

UNIVERSITE DE LAUSANNE - FACULTE DE BIOLOGIE ET DE MEDECINE

Département de Médecine

Service de Médecine Interne

**Associations Between Physical Activity Patterns and
Cardiovascular Events and Risk Factors.
Cross-Sectional and Prospective Studies**

THESE

préparée sous la direction du Professeur Pedro MARQUES-VIDAL
avec la co-direction de la Docteure Vanessa KRAEGE
et avec la collaboration du Professeur Julien VAUCHER

et présentée à la Faculté de biologie et de médecine de
l'Université de Lausanne pour l'obtention du grade de

DOCTEUR EN MEDECINE

par

Fabrizio COMINETTI

Médecin diplômé de la Confédération Suisse
Originaire de Lausanne (Vaud)

Lausanne
2025

IMPRIMATUR

La Faculté de biologie et médecine de l'Université de Lausanne, sur proposition du jury, autorise le dépôt de la thèse de doctorat rédigée par

Fabrizio COMINETTI

intitulée

**Associations Between Physical Activity Patterns
and Cardiovascular Events and Risk Factors.
Cross-Sectional and Prospective Studies**

sans se prononcer sur les opinions exprimées dans cette thèse.

Directeur	Professeur Pedro Manuel Marques-Vidal
Co-directrice	Docteure Vanessa Kraege
Experte interne	Professeure Patrizia D'Amelio
Vice-directeur de l'Ecole doctorale	Professeur John Prior

Lausanne, le 04.02.2025



pour Le Doyen
de la Faculté de Biologie et de Médecine

Monsieur le Professeur John Prior
Vice-Directeur de l'Ecole doctorale

ASSOCIATIONS BETWEEN PHYSICAL ACTIVITY PATTERNS AND
CARDIOVASCULAR EVENTS AND RISK FACTORS.
CROSS-SECTIONAL AND PROSPECTIVE STUDIES

Doctorant : Fabrizio Cominetti
Directeur de thèse : Prof Pedro Marques-Vidal
Co-directrice de thèse : Dre Vanessa Kraege

Contexte

L'activité physique (AP) d'intensité modérée est recommandée pour ses effets bénéfiques sur la santé, mais le moment optimal pour celle-ci, en ce qui concerne les maladies cardiovasculaires (MCV), reste débattu.

Objectifs

Nous avons évalué l'impact de différents schémas de répartition journalière de l'activité physique sur les facteurs de risque cardiovasculaires, ainsi que sur l'incidence des maladies cardiovasculaires.

Méthodes

Données de deux enquêtes (suivis S1 et S2) de l'étude de cohorte CoLaus|PsyCoLaus (2'465 et 1'692 participants, respectivement ; 55.3% [54.3%] femmes ; moyenne \pm déviation standard 61.2 \pm 9.7 ans [64.4 \pm 9.5]), menées à Lausanne, Suisse. L'AP a été évaluée à l'aide d'un accéléromètre porté au poignet. Les participants ont été répartis en quatre groupes selon la similitude de leurs schémas de répartition journalière de l'AP par la méthode des *K-means*.

Résultats

L'AP du matin était positivement associée à l'hypertension (odds ratio ajusté, multivariable, [intervalle de confiance à 95 %] : 1,36 [1,00-1,84]) lors du S1, avec une tendance similaire pour S2. Aucune association significative n'a été trouvée entre les groupes d'AP et le cholestérol total, le cholestérol HDL, le cholestérol LDL ou les triglycérides. L'AP prédominant le matin (S2) était positivement associée à un traitement médicamenteux hypolipémiant : 1,88 (1,07-3,30). L'AP uniformément répartie sur la journée (S2) était positivement associée au diabète : 1,82 (1,06-3,12), avec une tendance similaire en S1. Dans l'étude prospective, le groupe d'AP prédominant tôt le matin (7 à 12 heures) et le groupe d'AP uniformément répartie sur la journée étaient associés à un risque plus élevé d'événements cardiovasculaires (rapport de risque [IC 95 %] : 3,33 [1,08-10,3] et 3,16 [1,04-9,57], respectivement).

Conclusion

Dans cette étude populationnelle, nous avons observé un risque plus élevé d'événements cardiovasculaires chez les participants dont l'AP quotidienne se déroulait principalement tôt le matin (de 7 à 12 heures) ou était répartie uniformément tout au long de la journée. Aucune répartition journalière d'AP n'a été associée de manière consistante à l'hypertension, aux lipides sanguins ou aux marqueurs du diabète.

ORIGINAL RESEARCH

Associations Between Physical Activity Patterns and Cardiovascular Events and Risk Factors



Cross-Sectional and Prospective Studies

Fabrizio Cominetti, MD CANDIDATE,^a Julien Vaucher, MD,^{a,b} Pedro Marques-Vidal, MD, PhD,^{a,*} Vanessa Kraege, MD, MBA^{c,*}

ABSTRACT

BACKGROUND Moderate-intensity physical activity (PA) is recommended for health benefits, but optimal PA timing regarding cardiovascular disease (CVD) is debated.

OBJECTIVES The authors assessed the impact of differing PA patterns on CVD risk factors and outcomes.

METHODS Data from 2 surveys (S1 and S2) of the CoLaus-PsyCoLaus study (2,465 and 1,692 participants, respectively; 55.3% [54.3%] females; mean age 61.2 ± 9.7 years [64.4 ± 9.5]), conducted in Lausanne, Switzerland. PA was assessed using a wrist-worn accelerometer, and PA patterns were assessed using K-means clustering.

RESULTS Morning PA was positively associated with hypertension (multivariable-adjusted OR: 1.36 [95% CI: 1.00-1.84]) in S1, similar trend in S2. No significant association was found between PA clusters and total, HDL-, and LDL-cholesterol or triglycerides. Morning PA was positively associated with hypolipidemic drug treatment: 1.88 (1.07-3.30) in S2. Evenly distributed daily PA was positively associated with diabetes: 1.82 (95% CI: 1.06-3.12) in S2, with a similar trend in S1. In the outcome analysis, the early morning PA cluster (7 AM-12 AM) and the evenly distributed daily PA cluster led to a higher risk of CVD events (HR: 3.33 [95% CI: 1.08-10.3] and 3.16 [95% CI: 1.04-9.57], respectively).

CONCLUSIONS In a population-based study, we observed a higher risk for cardiovascular events in participants whose daily PA occurred predominantly in the early morning (7 AM-12 AM) or was evenly distributed throughout the day. No PA pattern was consistently associated with hypertension, blood lipids, or diabetes markers. (JACC Adv. 2024;3:101324) © 2024 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

International guidelines recommend 150 to 300 minutes of moderate-intensity physical activity (PA) per week to maintain good health.¹ Research has long focused on frequency, intensity,

and total amount of PA, but the optimal timing of PA remains unclear regarding general health benefits.²

A few studies have focused on health and cardiometabolic effects of PA timing. A systematic

From the ^aInternal Medicine Division, Department of Medicine, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland; ^bDivision of Internal Medicine, Department of Internal Medicine and Specialties, Hospital and University of Fribourg, Fribourg, Switzerland; and the ^cMedical and Innovation and Clinical Research Directorates, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland *Drs Marques-Vidal and Kraege are joint last co-authors. The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

Manuscript received June 16, 2024; revised manuscript received August 30, 2024, accepted September 4, 2024.

**ABBREVIATIONS
AND ACRONYMS****BP** = blood pressure**CVD** = cardiovascular disease**DBP** = diastolic blood pressure**MVPA** = moderate to vigorous
physical activity**PA** = physical activity**S1/S2** = survey 1/survey 2**SBP** = systolic blood pressure

review in 2020 showed no conclusive evidence favoring PA at a certain time of day for effect on adiposity markers and cardiometabolic biomarkers.² Still, a recent study using accelerometer-based PA evaluation in the UK Biobank showed an association between morning PA and lower risk of cardiovascular disease (CVD), irrespective of daily total PA.³ Another study on the same cohort assessed the effects of the timing of moderate to vigorous PA (MVPA) on all-cause

and specific mortality; the authors concluded that CVD mortality was lower in participants whose MVPA was higher at midday and in the afternoon.⁴ Indeed, timing of PA and its concordance with chronotype appear to be an important factor in CVD mortality risk, with early morning and nighttime PA being less favorable than midday activity.⁵

In a post-hoc analysis of a randomized controlled trial, Albalak et al showed an association between increased morning PA and several metabolic parameters, including body mass index (BMI).⁶ A cross-sectional analysis of a diabetic population showed sex differences, morning MVPA being associated with better cardiorespiratory fitness in males, whereas evening MVPA was associated with better cardiorespiratory fitness in females.⁷ In a Dutch cohort, insulin resistance was reduced in groups with higher afternoon and evening MVPA compared to a group with evenly distributed MVPA throughout the day.⁸ Similarly, evening PA was associated with better blood pressure outcomes in Japanese male workers,⁹ and with better health-related fitness in Finnish workers.¹⁰ Still, the analysis of existing data is complicated as most authors used different methods to quantify PA and its timing.

Hence, our study aimed to assess the impact of the time of day of accelerometry-assessed PA on cardiometabolic parameters (BMI, waist circumference, blood pressure, blood glucose levels, lipid levels) using data from a population-based study.

METHODS

PARTICIPANTS. The study used data from the CoLaus-PsyCoLaus study, a prospective study conducted in the population of Lausanne, Switzerland. Details of the recruitment procedure have been reported previously.¹¹ Three follow-ups were performed in 2009 to 2012, 2014 to 2017, and 2018 to 2021. They included 5,064 (75.2%), 4,881 (72.5%), and 3,751 (55.7%), respectively, of the 6,733 initial participants. Objectively assessed PA was conducted in the second and third follow-ups; hence, data from those 2

follow-ups was used in this study. For simplicity, the second and third follow-ups will be named survey 1 (S1) and survey 2 (S2), respectively. Each survey was considered and analyzed independently in both cross-sectional analyses. In other words, the cohort's population was cross-sectionally analyzed at 2 distinct moments, in the first survey and then 3.8 years (median) later in the second. For the outcome analysis, S1 participants were studied, as they had 2 consecutive PA assessments (in second and third follow-up).

PHYSICAL ACTIVITY. PA was objectively assessed using a wrist-worn triaxial accelerometer (GENEActiv, Activinsights Ltd, www.activinsights.com). These devices are the same as those used in the UK biobank study,¹² weigh 16 g and allow continuous monitoring of PA for a maximum of 45 days. The devices were preprogrammed with a 50 Hz sampling frequency and subsequently attached to the participants' right wrist. Participants were requested to wear the device continuously for 14 days in their free-living conditions. Raw accelerometry data were downloaded using the GENEActiv software version 2.9 (GENEActiv, Activinsights Ltd) and transformed into 1-minute epoch files using the GENEActiv MACRO,^{6,13} which also computes the vector magnitude (vectogram).

Data from days 2 to 8 of recording was used, as data from day 1 is not representative while wearers get used to the accelerometer. For each participant and day of recording, the average vectogram for each hour was computed to assess PA. Then, for each individual, the daily hourly values of PA were standardized by the 7-day mean of total daily PA as follows:

$$PAstd_i = \frac{\overline{PA}_i}{\overline{PA}_{24}}, \text{ where } i = 1 \text{ to } 24$$

The standardized PA for hour *i* is the ratio of the average PA for time *i* divided by the average of the 24-hour PA. For example, hour *i* = 1 corresponds to the time between 0 and 59 minutes AM. The 24 standardized hourly values were then grouped into clusters using K-means clustering analysis as performed by Albalak et al³ to obtain standardized relative PA profiles. We determined the number of 4 clusters with proportional reduction in error testing. As each cluster has a definite relative PA pattern, they will be referred to as "PA timing cluster." This number also allowed comparison with previously published results.

CARDIOMETABOLIC RISK FACTORS. Body weight and height were measured with participants barefoot and in light indoor clothes. Body weight was measured in kilograms to the nearest 100 g using a Seca scale. Height was measured to the nearest 5 mm

using a Seca height gauge. BMI was categorized as normal ($<25 \text{ kg/m}^2$), overweight (≥ 25 - $<30 \text{ kg/m}^2$), and obesity ($\geq 30 \text{ kg/m}^2$).

Waist circumference was measured midway between the lowest rib and the iliac crest using a non-stretchable tape, and the average of 2 measurements was taken. Abdominal obesity was defined as a waist circumference $>102 \text{ cm}$ (males) or $>88 \text{ cm}$ (females).

Blood pressure (BP) was measured using an Omron HEM-907 automated oscillometric sphygmomanometer after at least a 10-minute rest in a seated position, and the average of the last 2 measurements was used. Hypertension was defined by a systolic blood pressure (SBP) $\geq 140 \text{ mm Hg}$ or a diastolic blood pressure (DBP) $\geq 90 \text{ mm Hg}$ or presence of antihypertensive drug treatment.

Biological assays were performed by the Centre Hospitalier Universitaire Vaudois Clinical Laboratory on fresh blood samples within 2 hours of blood collection. Fasting plasma glucose was assessed by glucose hexokinase. Diabetes was defined as the presence of antidiabetic treatment and/or a fasting blood glucose $\geq 7 \text{ mmol/L}$ or glycated hemoglobin level (HbA_{1c}) above 48 mmol/mol .

Total cholesterol was assessed by cholesterol oxidase phenol 4-aminoantipyrine peroxidase; high-density lipoprotein-cholesterol was assessed by cholesterol oxidase phenol 4-aminoantipyrine peroxidase + polyethylene glycol + cyclodextrin. Triglycerides were assessed by glycerol phosphate oxidase phenol 4-aminoantipyrine peroxidase; low-density lipoprotein-cholesterol levels were computed using the Friedewald formula.

CARDIOVASCULAR OUTCOMES. During the follow-up period, first incident CVD events and deaths were prospectively collected and independently adjudicated according to established recommendations and similar definitions detailed elsewhere.¹¹ Details of the adjudication procedure are provided in the [Supplemental Appendix](#).

COVARIATES. Socio-economic and lifestyle data were collected by questionnaire. Education was categorized as mandatory education, apprenticeship, high school, and university education. Smoking was categorized as never, former, and current. Medications were collected and classified according to the World Health Organization ATC criteria. Alcohol consumption was categorized as nondrinker vs number of units per week (1-13, 14-27, 28 or more). Morning/evening chronotype was assessed by questionnaire.

INCLUSION AND EXCLUSION CRITERIA. For the cross-sectional analyses, participants were excluded if they had: 1) no or <1 week of valid PA data; 2)


presented with previous CVD; or 3) missing data for outcomes or covariates. For the outcome analysis, participants were also excluded if they; 4) already presented with CVD; or 5) had no follow-up after the first survey.

STATISTICAL ANALYSIS. Statistical analyses were conducted using Stata version 18.0 for Windows (Stata Corp). Participant characteristics were expressed as number (percentage) for categorical variables or as mean \pm SD for continuous variables. Between-group comparisons were performed using chi-square or Fisher's exact test for categorical variables and a Student's *t*-test, analysis of variance, or Kruskal-Wallis test for continuous variables.

Cross-sectional multivariable analyses were conducted using logistic regression for categorical outcomes (hypertension, hypolipidemic drug treatment, diabetes [yes, no]), and results were expressed as OR (95% CI). For continuous outcomes, multivariable analyses were conducted using analysis of variance, and results were expressed as adjusted mean \pm SEM. Multivariable analyses were adjusted on sex, age, education, smoking, and alcohol consumption for all outcomes. For hypertension, diabetes and hypolipidemic drug treatment, further adjustment on BMI categories was performed. For blood pressure, glycemia, and lipids, further adjustment on antihypertensive, antidiabetic, and hypolipidemic drug treatment was performed.

For the outcome analysis, the incidence of CVD events, CVD-specific, and overall mortality were computed for each PA timing cluster. Unadjusted and adjusted analyses were performed using Cox regression for overall mortality and incident fatal and nonfatal CVD events. Proportional hazards assumption was tested using the Stata command `estat phtest` and no deviations were found. Results were expressed as HR (95% CI). Multivariable analyses were performed adjusting for sex, age, education, smoking, alcohol consumption, and BMI categories in a first adjusted model. In the second model, we conducted supplementary adjustments for hypertension, diabetes, and hypolipidemic drug treatment. A Fine-Gray model using non-CVD death as competing risk was also applied, adjusting for the same variables as before (third model).

Furthermore, analysis of each PA timing cluster was also stratified by age (<65 or >65 years of age) and PA level (less or more active groups: a participant is categorized in the more active group if its mean PA vector over the entire study period is above the sample's median). A second analysis investigated the association between mean hourly physical activity of



total study population and cardiovascular disease events using Cox-proportional hazard model, adjusted for age and sex. Finally, we performed inverse probability weighting in the outcome analysis. Statistical significance was considered for a 2-sided test with $P < 0.05$.

ETHICAL STATEMENT. The institutional Ethics Committee of the University of Lausanne, which afterward became the Ethics Commission of Canton Vaud, approved the baseline CoLaus study (reference 16/03). The approval was renewed for the first (reference 33/09), the second (reference 26/14), and the third (reference PB_2018-00040) follow-ups. The approval for the entire CoLaus|PsyCoLaus study was confirmed in 2021 (reference PB_2018-00038, 239/09). The full decisions of the Ethics Committee of Canton Vaud can be obtained from the authors upon request. The study was performed in agreement with the Helsinki Declaration and its former amendments and in accordance with the applicable Swiss legislation. All participants gave their signed informed consent before entering the study.

RESULTS


STUDY SAMPLES. Of the initial 4,881 and 3,751 participants in the first and second surveys, 2,465

(50.5%) and 1,692 (45.1%), respectively, were retained for analysis. The reasons for exclusion are indicated in [Figure 1](#) and the comparison between included and excluded participants is provided in [Supplemental Table 1](#). Excluded participants were older, less well educated, with a different chronotype, more frequently teetotalers, had higher levels of BMI, waist circumference, fasting glucose and insulin, lipids, and systolic blood pressure, and were more likely to have an antidiabetic, antihypertensive, or cholesterol-lowering medication.

CHARACTERISTICS OF PHYSICAL ACTIVITY CLUSTERS.

Within each follow-up, each participant was classified into a single cluster. The number of clusters was selected according to the proportional reduction in error. For the first survey, the number of clusters was 4, and for the second survey, it could be either 4 or 5 ([Supplemental Figures 1 and 2](#)). We selected 4 clusters as it agreed with the results and also allowed comparison with the literature. The PA clusters were quite similar between surveys ([Figure 2](#)). In cluster 1, PA peaked between 10 AM and 2 PM; in cluster 2, PA peaked between 7 AM and 12 AM; in cluster 3, PA peaked between 8 AM and 1 PM and in cluster 4, no clear peak of PA was observed.

FIGURE 2 Physical Activity Timing Clusters for Survey 1 and 2, CoLaus-PsyCoLaus Study, Lausanne, Switzerland



Patterns of standardized vectogram showing mean relative PA of each cluster in the first and second surveys. Cluster 1: PA peaks between 10 AM and 2 PM. Cluster 2: PA peaks between 7 AM and 12 AM. Cluster 3: PA peaks between 8 AM and 1 PM. Cluster 4 demonstrates no clear peak of PA. FU2 = second follow-up; FU3 = third follow-up; PA = physical activity.

TABLE 1 Cardiometabolic Risk Factors by Physical Activity Timing Clusters, Survey 1 (Second Follow-Up, 2014-2017), CoLaus-PsyCoLaus Study, Lausanne, Switzerland

	Cluster Number				P Value
	1 (n = 481)	2 (n = 453)	3 (n = 790)	4 (n = 741)	
Blood pressure markers					
SBP (mm Hg)					
Unadjusted	124 ± 17	128 ± 18	127 ± 18	123 ± 17	<0.001
Adjusted	124 ± 1	126 ± 1	126 ± 1	125 ± 1	0.188
DBP (mm Hg)					
Unadjusted	77 ± 11	77 ± 10	78 ± 11	78 ± 11	0.735
Adjusted	77 ± 1	78 ± 1	78 ± 1	77 ± 1	0.448
Hypertension					
Unadjusted	171 (35.6)	223 (49.2)	338 (42.8)	260 (35.1)	<0.001
Adjusted	1.00 (reference)	1.36 (1.00-1.84)	1.14 (0.87-1.49)	1.14 (0.87-1.50)	
Lipid markers					
Total cholesterol (mmol/L)					
Unadjusted	5.4 ± 0.9	5.4 ± 1.0	5.5 ± 1.0	5.3 ± 0.9	<0.001
Adjusted	5.34 ± 0.04	5.36 ± 0.05	5.50 ± 0.04	5.37 ± 0.04	0.018
HDL cholesterol (mmol/L)					
Unadjusted	1.6 ± 0.5	1.7 ± 0.5	1.6 ± 0.4	1.5 ± 0.5	<0.001
Adjusted	1.62 ± 0.02	1.65 ± 0.02	1.60 ± 0.01	1.59 ± 0.02	0.126
LDL cholesterol (mmol/L)					
Unadjusted	3.2 ± 0.9	3.2 ± 0.9	3.3 ± 0.9	3.2 ± 0.8	0.007
Adjusted	3.14 ± 0.04	3.17 ± 0.04	3.30 ± 0.03 ^a	3.21 ± 0.03	0.008
Triglycerides (mmol/L)					
Unadjusted	1.3 ± 1.0	1.2 ± 0.8	1.3 ± 1.1	1.3 ± 1.1	0.597 ^a
Adjusted	1.30 ± 0.05	1.22 ± 0.05	1.33 ± 0.04	1.27 ± 0.04	0.057 ^a
Hypolipidemic drug treatment					
Unadjusted	53 (11.3)	78 (17.9)	124 (16.2)	84 (11.7)	0.002
Adjusted	1.00 (reference)	1.13 (0.73-1.73)	1.26 (0.86-1.86)	1.44 (0.96-2.15)	
Diabetes markers					
Glucose (mmol/L)					
Unadjusted	5.3 ± 0.9	5.4 ± 1.1	5.4 ± 0.9	5.4 ± 1.0	0.573
Adjusted	5.36 ± 0.04	5.41 ± 0.04	5.37 ± 0.03	5.38 ± 0.03	0.865
HbA _{1c} (mmol/mol)					
Unadjusted	37.4 (6.3)	38.0 (6.7)	37.9 (5.7)	37.1 (6.0)	0.036
Adjusted	37.6 ± 0.2	37.6 ± 0.3	37.5 ± 0.2	37.7 ± 0.2	0.923
Insulin (μIU/mL)					
Unadjusted	8.8 ± 5.3	9.3 ± 6.5	9.1 ± 5.9	9.0 ± 6.8	0.618 ^a
Adjusted	8.9 ± 0.3	9.2 ± 0.3	9.0 ± 0.2	9.3 ± 0.2	0.821 ^a
Diabetes ^b					
Unadjusted	33 (6.9)	37 (8.2)	66 (8.4)	54 (7.3)	0.737
Adjusted	1.00 (reference)	0.84 (0.48-1.46)	1.04 (0.64-1.71)	1.19 (0.72-1.96)	

Values are mean ± SD, n (%), or OR (95% CI). ^aBased on log-transformed data. ^bDefined as a fasting plasma glucose ≥7 mmol/L and/or presence of antidiabetic drug treatment. Between-group comparisons were performed using Pearson chi-squared test for categorical variables and analysis of variance for continuous variables. Statistically significant ($P < 0.05$) ORs are indicated in **bold**. Multivariable analysis is performed using logistic regression for categorical variables and ANOVA for continuous variables. For continuous variables, post-hoc bivariate comparisons were performed using Scheffe's method: ^acomparison of clusters 2, 3, and 4 to cluster 1, ^bcomparison of clusters 3 and 4 to cluster 2, and ^ccomparison of cluster 4 to cluster 3; presence of the subscript indicates that clusters differ at $P < 0.05$. All multivariable models adjusted for sex (male, female), age (continuous), educational level (basic, apprenticeship, secondary, high school, and university), alcohol consumption (none, 1-13, 14-27, and 27+ per week), smoking categories (never, former, and current), and BMI categories (normal, overweight, and obese). For blood pressure levels, a further adjustment on antihypertensive drug treatment (yes, no) was performed; for lipid levels, a further adjustment on hypolipidemic drug treatment (yes, no) was performed; and for diabetes markers, a further adjustment on antidiabetic drug treatment (yes, no) was performed.

ANOVA = analysis of variance; BMI = body mass index; DBP = diastolic blood pressure; HDL cholesterol = high density lipoprotein cholesterol; LDL cholesterol = low-density lipoprotein cholesterol; SBP = systolic blood pressure.

The sociodemographic characteristics of the PA clusters according to survey are summarized in [Supplemental Table 2](#). In both surveys, clusters differed regarding age, sex, educational level, and smoking status distribution. In S2 (third follow-up),

clusters also differed significantly regarding BMI, waist circumference, and sleep chronotype. Cluster 4 was the youngest, with less females, more educated participants, and a higher prevalence of current smokers.

TABLE 2 Cardiometabolic Risk Factors by Physical Activity Timing Clusters, Survey 2 (Third Follow-Up, 2018-2021), CoLaus-PsyCoLaus Study, Lausanne, Switzerland

	Cluster Number				P Value
	1 (n = 458)	2 (n = 413)	3 (n = 480)	4 (n = 341)	
Blood pressure markers					
SBP (mm Hg)					
Unadjusted	128 ± 17	128 ± 18	129 ± 18	126 ± 17	0.047
Adjusted	128 ± 1	128 ± 1	127 ± 1	126 ± 1	0.245
DBP (mm Hg)					
Unadjusted	78 ± 10	78 ± 11	77 ± 10	78 ± 11	0.769
Adjusted	78 ± 1	77 ± 1	78 ± 1	77 ± 1	0.268
Hypertension					
Unadjusted	199 (43.5)	185 (44.8)	239 (49.8)	146 (42.8)	0.144
Adjusted	1.00 (reference)	1.14 (0.84-1.54)	0.99 (0.74-1.33)	1.01 (0.73-1.39)	
Lipid markers					
Total cholesterol (mmol/L)					
Unadjusted	5.4 ± 0.9	5.2 ± 0.9	5.3 ± 1.0	5.2 ± 1.0	0.049
Adjusted	5.46 ± 0.08	5.32 ± 0.08	5.34 ± 0.07	5.30 ± 0.09	0.496
HDL cholesterol (mmol/L)					
Unadjusted	1.6 ± 0.4	1.6 ± 0.5	1.6 ± 0.4	1.5 ± 0.4	<0.001
Adjusted	1.47 ± 0.03	1.53 ± 0.03	1.51 ± 0.03	1.47 ± 0.04	0.450
LDL cholesterol (mmol/L)					
Unadjusted	3.2 ± 0.8	3.1 ± 0.8	3.1 ± 0.9	3.1 ± 0.9	0.234
Adjusted	3.29 ± 0.07	3.17 ± 0.08	3.17 ± 0.07	3.14 ± 0.08	0.431
Triglycerides (mmol/L)					
Unadjusted	1.3 ± 0.7	1.4 ± 1.2	1.3 ± 0.9	1.4 ± 1.0	0.558 ^a
Adjusted	1.53 ± 0.08	1.50 ± 0.09	1.44 ± 0.07	1.54 ± 0.09	0.387 ^a
Hypolipidemic drug treatment					
Unadjusted	49 (38.3)	56 (47.9)	86 (54.1)	48 (49.0)	0.065
Adjusted	1.00 (reference)	1.88 (1.07-3.30)	1.33 (0.78-2.25)	1.39 (0.77-2.53)	
Diabetes markers					
Glucose (mmol/L)					
Unadjusted	5.4 ± 0.7	5.5 ± 1.1	5.5 ± 1.1	5.6 ± 1.2	0.107
Adjusted	5.50 ± 0.04	5.51 ± 0.04	5.48 ± 0.04	5.57 ± 0.05	0.577
HbA _{1c} (mmol/mol)					
Unadjusted	36.6 (5.0)	36.7 (5.8)	37.4 (6.3)	37.7 (7.6)	0.021
Adjusted	36.9 ± 0.2	37.0 ± 0.3	37.0 ± 0.2	37.7 ± 0.3	0.143
Insulin (μIU/mL)					
Unadjusted	9.9 ± 8.6	9.6 ± 5.7	9.8 ± 5.6	11.3 ± 9.2	0.007 ^a
Adjusted	10.3 ± 0.3	9.9 ± 0.3	9.7 ± 0.3	11.1 ± 0.4	0.054 ^a
Diabetes ^b					
Unadjusted	28 (6.1)	32 (7.8)	41 (8.5)	38 (11.1)	0.080
Adjusted	1.00 (reference)	1.26 (0.73-2.19)	1.19 (0.70-2.02)	1.82 (1.06-3.12)	

Values are mean ± SD, n (%), or OR (95% CI). ^aBased on log-transformed data. ^bDefined as a fasting plasma glucose ≥7 mmol/L and/or presence of antidiabetic drug treatment. Between-group comparisons were performed using Pearson chi-squared test for categorical variables and analysis of variance for continuous variables. Statistically significant ($P < 0.05$) ORs are indicated in **bold**. Multivariable analysis is performed using logistic regression for categorical variables and ANOVA for continuous variables. For continuous variables, post-hoc bivariate comparisons were performed using Scheffe's method: ^ccomparison of clusters 2, 3, and 4 to cluster 1, ^dcomparison of clusters 3 and 4 to cluster 2, and ^ecomparison of cluster 4 to cluster 3; presence of the subscript indicates that clusters differ at $P < 0.05$. All multivariable models adjusted for sex (male, female), age (continuous), educational level (basic, apprenticeship, secondary, high school, and university), alcohol consumption (none, 1-13, 14-27, and 27+ per week), smoking categories (never, former, and current), and BMI categories (normal, overweight, and obese). For blood pressure levels, a further adjustment on antihypertensive drug treatment (yes, no) was performed; for lipid levels, a further adjustment on hypolipidemic drug treatment (yes, no) was performed; and for diabetes markers, a further adjustment on antidiabetic drug treatment (yes, no) was performed.

Abbreviations as in Table 1.

ASSOCIATION OF PHYSICAL ACTIVITY TIMING CLUSTERS WITH CARDIOMETABOLIC RISK FACTORS. The unadjusted and adjusted analysis of cardiometabolic markers according to PA timing clusters is provided in

Tables 1 and 2 for S1 and S2, respectively. After multivariable adjustment, no association was found between clusters and SBP or DBP. Cluster 2 was positively associated with hypertension in S1, with a

TABLE 3 Associations Between Physical Activity Timing Clusters and Cardiovascular Events, CoLaus-PsyCoLaus Study, Lausanne, Switzerland

	Cluster Number			
	1 (n = 386)	2 (n = 364)	3 (n = 670)	4 (n = 639)
Number of events	4	19	30	17
Incidence	0.2 (0.1-0.6)	1.2 (0.7-1.8)	1.0 (0.7-1.4)	0.6 (0.4-1.0)
Unadjusted	1.00 (reference)	5.16 (1.76-15.2)	4.40 (1.55-12.5)	2.56 (0.86-7.62)
Adjusted model 1	1.00 (reference)	3.90 (1.29-11.8)	2.93 (1.00-8.59)	3.36 (1.12-10.1)
Adjusted model 2	1.00 (reference)	3.33 (1.08-10.3)	2.83 (0.96-8.34)	3.16 (1.04-9.57)
Adjusted model 3	1.00 (reference)	3.34 (1.11-10.1)	2.84 (0.99-8.17)	3.16 (1.06-9.43)

Incidence rates are expressed as rate per 1,000 person-years (and 95% CIs). For unadjusted and adjusted analysis, results are expressed as HR (95% CI). Analyses conducted using Cox regression for CVD events. Adjusted model 1: adjusted for sex (male, female), age (continuous), educational level (basic, apprenticeship, secondary, high school, and university), alcohol consumption (none, 1-13, 14-27, and 27+ per week), smoking categories (never, former, current), and BMI categories (normal, overweight, and obese). Adjusted model 2: as model 1, plus hypertension (yes, no), diabetes (yes, no), and hypolipidemic drug treatment (yes, no). Significant ($P < 0.05$) coefficients are indicated in **bold**. Adjusted model 3: Fine-Gray model, using non-CVD death as a competing risk, and adjusting as for model 2.

BMI = body mass index; CVD = cardiovascular disease.

nonsignificant trend in S2 (Tables 1 and 2). Regarding lipids, no association was found between clusters and HDL cholesterol or triglycerides. Cluster 3 had the highest total and LDL cholesterol levels in S1, but this was not confirmed in S2. Cluster 2 was positively associated with hypolipidemic drug treatment in S2, but this was not confirmed in S1 (Tables 1 and 2). Regarding diabetes, no association was found between clusters and glucose, HbA_{1c} or insulin levels. Cluster 4 was positively associated with diabetes in S2, but this was not confirmed in S1 (Tables 1 and 2).

ASSOCIATION OF PHYSICAL ACTIVITY TIMING CLUSTERS WITH INCIDENCE OF CARDIOVASCULAR EVENTS. For the outcome analysis, 2,059 participants from the first survey were included and followed for a median period of 46 months (IQR: 43-48 months). The incidence rates and association between PA timing clusters and incidence of CVD events are provided in Table 3 and the incidence curve in Figure 3. Cluster 1 had the lowest incidence rate for CVD and was selected as reference. On unadjusted analysis, participants in clusters 2 and 3 had a higher risk of developing CVD, but after multivariable adjustment (models 1, 2, and 3), clusters 2 and 4 presented with a higher risk (Table 3 and Central Illustration). The increased CVD risk in Cluster 2 and Cluster 4 persisted in the inverse probability weighted analyses, where Cluster 3 also had a higher CVD risk in the model 2 (with adjustments for hypertension, diabetes, and hypercholesterolemia) (Supplemental Table 5). Moreover, stratified analysis by age demonstrated a higher CVD risk in participants <65 years of age in Cluster 2 and Cluster 3, whereas the stratification by PA group showed that CVD risk was not increased in the more active

participants of Cluster 2 and Cluster 3 (Supplemental Table 6). The second analysis of CVD risk according to mean physical activity per hour for the total study population demonstrated a nonsignificant trend for an increased risk for participants more active between 22 PM and 2 AM (Supplemental Figure 3).

DISCUSSION


MAIN FINDINGS. Distinct clusters of daily PA patterns were established with K-means clustering. In the cross-sectional analysis, no consistent association was found between the 4 PA clusters and cardiovascular risk factor levels. In the outcome analysis, we found a higher risk of cardiovascular events in PA clusters with an early morning peak or no clear peak.

PHYSICAL ACTIVITY TIMING CLUSTERS. The patterns obtained in our study are comparable to those described by Albalak et al, who used accelerometer-based PA evaluation in the UK Biobank study.³ To some extent, they are also comparable with those obtained by Li et al on the same Biobank, although in our study, Cluster 4 (no clear PA peak) demonstrated a higher variability in daily PA.¹⁴ Cluster cross-over between S1 and S2 was observed for some participants and was more pronounced between Cluster 4 and Cluster 2, which may explain the different shape of PA patterns between follow-ups for these clusters. Nevertheless, the fact that the same clusters were obtained in different populations and settings is encouraging, as it facilitates comparisons between studies. Further, the reproducibility of those clusters suggests that they might correspond to global PA behaviors, at least in European countries.

ASSOCIATION OF PHYSICAL ACTIVITY TIMING CLUSTERS WITH CARDIOMETABOLIC RISK FACTORS. No consistent differences were found between PA clusters regarding cardiometabolic risk factors.

Regarding BP and hypertension, results are conflicting regarding the best PA timing to reduce BP or prevent hypertension. Imamura et al demonstrated that evening PA was associated with better BP outcomes in Japanese male workers⁹; Li et al¹⁴ observed a benefit in early morning PA and early morning plus evening PA on hypertension incidence, while the Liu et al¹⁵ systematic review of intervention studies concluded to a benefit of morning PA on BP. The differences in SBP or DBP observed between PA groups in the UK biobank were relatively small (maximum of 6.6 mm Hg in SBP), and similar findings were reported in a systematic review.^{2,3}

Regarding lipids, no conclusive trend was found in our study. This result is in keeping with published systematic reviews.^{2,15} In a review of 35 studies,



Janssen et al concluded that there was no consistent evidence that PA at 1 specific period of the day provided more favorable health benefits.² Similarly, in a review of 4 studies, Liu et al¹⁵ regrouped reported mixed results and no clear favorable moment of PA on lipid metabolism.


Regarding diabetes markers, conflicting results have been published. A post-hoc study of a randomized controlled trial showed a positive association between increased morning PA and fasting glucose, HbA_{1c}, and insulin,⁶ while a Dutch cohort showed a decreased insulin resistance in groups with higher afternoon and evening PA, compared to a group with evenly distributed PA throughout the day.⁸ In Tian et al's prospective study on type 2 diabetes risk, morning and afternoon PA were beneficial, whereas evening PA had a neutral effect.¹⁶ Finally, the systematic review by Janssen et al² included 2 studies favoring evening PA, 1 favoring morning PA, and 4 remaining neutral regarding diabetes. Studies among diabetics also show conflicting results. A study in a Hispanic/Latino diabetic population demonstrated better HbA_{1c} levels in the evening PA group.¹⁷ Another reported difference in the association according to sex was morning PA being more favorable for males with diabetes, while evening PA was more favorable for females with the disease.⁷

Overall, our results show no consistent associations between PA timing clusters and cardiometabolic risk factors, a finding somewhat in line with the inconclusive or conflicting results of the literature.

ASSOCIATION OF PHYSICAL ACTIVITY TIMING CLUSTERS WITH INCIDENCE OF CARDIOVASCULAR EVENTS.

However, a difference in cardiovascular events was observed between our clusters, with early morning PA (7 AM-12 AM) and no clear PA peak (or evenly distributed daily PA) being associated with an increased CVD risk. The association was maintained in the sensitivity analysis according to different adjustment models. This result is part of the unequivocal body of literature on the best time of day for PA regarding CVD risk. Several studies assessed the association between timing of PA and CVD events using data from the UK Biobank. Feng et al⁴ found a decreased CVD mortality risk in participants whose PA was higher at mid-day and during the afternoon, independently of the total PA volume. Ma et al⁵ reported that mid-day PA had the lowest CVD mortality risk in individuals without pre-existing CVD. Finally, Albalak et al³ reported that early and late morning peaks of PA were associated with a lower CVD risk. Interestingly, different conclusions were obtained using data from the same study, which suggests that the results of the studies on the effects of timing of PA

CENTRAL ILLUSTRATION Associations Between Physical Activity Patterns and Cardiovascular Events and Risk Factors



Cominetti F, et al. *JACC Adv.* 2024;3(11):101324.

Participants had their physical activity objectively assessed for 7 days. They were regrouped into 4 clusters according to their relative daily physical activity patterns. Multivariable analysis after a median of 46 months of follow-up found an association of early morning PA and CVD events, also for participants with no PA peak during the day (results according to adjusted model 2 outcome analysis). CVD = cardiovascular disease; PA = physical activity.

and CVD events are likely dependent on the inclusion criteria and/or methods used to identify clusters. Noteworthy, in our study, cluster cross-over between S1 and S2 was observed, being more pronounced between Cluster 4 and Cluster 2. This suggests that PA behavior changes with time, which might partly explain the conflicting results between studies. Eventually, the magnitude of the OR between clusters in our study should be cautiously interpreted, since the large CIs may indicate that our relatively small study sample could have led to a decreased precision.

STRENGTHS AND LIMITATIONS. Our study has many strengths. First, we replicated a previously used method to define clusters of PA activity, allowing an easier comparison with the literature. Secondly, our study population is drawn from a well-characterized population-based cohort. Moreover, we used objectively assessed PA data to minimize recall bias, and a potential Hawthorne effect was mitigated by excluding the first day of recording for the analyses. Finally, self-reported and clinically measured

outcomes were used to study patient characteristics of each PA cluster in detail.

Still, our study also has limitations. As almost one-half of the initial sample did not have objectively assessed PA, we can hypothesize that less active participants could have been more reluctant to collect this data, leading to potential selection bias. Moreover, our study population was in a single Swiss city and mainly constituted of Caucasian subjects. Whether our results also apply in other settings should be verified. Finally, as in all studies on objectively assessed PA, we relied on PA data processed by a software, which can lead to variations in PA level estimations,¹³ and on specific methods to determine clusters of PA patterns. Nevertheless, to mitigate this effect, we used common software and replicated the clustering-methods, as stated above.^{3,5}

CONCLUSIONS

In a population-based study, we observed a higher risk for cardiovascular events in participants whose

daily PA occurred predominantly in the early morning (7-12 AM) or was evenly distributed throughout the day. No association was found between PA timing and hypertension, blood lipids, or diabetes markers.

ACKNOWLEDGMENTS The authors would like to thank all the people who participated in the recruitment of the participants, data collection and validation, particularly Nicole Bonvin, Yolande Barreau, Mathieu Firmann, François Bastardot, Panagiotis Antiochos, Cédric Gubelmann, Marylène Bay and Benoît Delabays. They would also like to thank Céline Maglieri for double-checking analysis data.

FUNDING SUPPORT AND AUTHOR DISCLOSURES

The CoLaus-PsyCoLaus study was supported by unrestricted research grants from GlaxoSmithKline, the Faculty of Biology and Medicine of Lausanne, the Swiss National Science Foundation (grants 3200B0-105993, 3200B0-118308, 33CS0-122661, 33CS30-139468, 33CS30-148401, 33CS30-177535, and 3247730_204523), and the Swiss Personalized Health Network (grant 2018DR101). The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

ADDRESS FOR CORRESPONDENCE: Dr Fabrizio Cominetti, Lausanne University Hospital, Office 10-672, Rue du Bugnon 46, CH-1011, Lausanne, Switzerland. E-mail: fabrizio.cominetti@chuv.ch.

PERSPECTIVES

COMPETENCY IN MEDICAL KNOWLEDGE: Physical activity counseling is part of daily medical routine, but it is unclear if a certain time of day is most beneficial in terms of cardiovascular health, as results are conflicting. In this study, no specific time of day seems unequivocally beneficial for cardiovascular risk-factor reduction. Further, early morning physical activity is associated with higher cardiovascular disease events.

TRANSLATIONAL OUTLOOK: Additional research is needed to explore if direct causality of early morning physical activity on cardiovascular disease events can be confirmed. Still, physical activity should be promoted in a way that fits a person's preferences and capacities.


REFERENCES

- Piercy KL, Troiano RP, Ballard RM, et al. The physical activity guidelines for Americans. *JAMA*. 2018;320(19):2020-2028. <https://doi.org/10.1001/jama.2018.14854>
- Janssen I, Campbell JE, Zahran S, Saunders TJ, Tomasone JR, Chaput JP. Timing of physical activity within the 24-hour day and its influence on health: a systematic review. *Health Promot Chronic Dis Prev Can Res Policy Pract*. 2022;42(4):129-138. <https://doi.org/10.24095/hpcdp.42.4.02>
- Albalak G, Stijntjes M, van Bodegom D, et al. Setting your clock: associations between timing of objective physical activity and cardiovascular disease risk in the general population. *Eur J Prev Cardiol*. 2023;30(3):232-240. <https://doi.org/10.1093/eurjpc/zwac239>
- Feng H, Yang L, Liang YY, et al. Associations of timing of physical activity with all-cause and cause-specific mortality in a prospective cohort study. *Nat Commun*. 2023;14(1):930. <https://doi.org/10.1038/s41467-023-36546-5>
- Ma T, Jennings L, Sirard JR, Xie YJ, Lee CD. Association of the time of day of peak physical activity with cardiovascular mortality: findings from the UK Biobank study. *Chronobiol Int*. 2023;24:1-11. <https://doi.org/10.1080/07420528.2023.2170240>
- Albalak G, Stijntjes M, Wijsman CA, et al. Timing of objectively-collected physical activity in relation to body weight and metabolic health in sedentary older people: a cross-sectional and prospective analysis. *Int J Obes*. 2022;46(3):515-522. <https://doi.org/10.1038/s41366-021-01018-7>
- Qian J, Walkup MP, Chen SH, et al. Association of objectively measured timing of physical activity bouts with cardiovascular health in type 2 diabetes. *Diabetes Care*. 2021;44(4):1046-1054. <https://doi.org/10.2337/dc20-2178>
- van der Velde JHPM, Boone SC, Winters-van Eekelen E, et al. Timing of physical activity in relation to liver fat content and insulin resistance. *Diabetologia*. 2023;66(3):461-471. <https://doi.org/10.1007/s00125-022-05813-3>
- Imamura M, Tahara Y, Suiko T, Nagamori Y, Shibata S. Association between blood pressure and circadian timing of physical activity of Japanese workers. *Front Physiol*. 2022;13:992945. <https://doi.org/10.3389/fphys.2022.992945>
- Stenholm S, Pulakka A, Leskinen T, et al. Daily physical activity patterns and their association with health-related physical fitness among aging workers—the Finnish retirement and aging study. *J Gerontol A Biol Sci Med Sci*. 2021;76(7):1242-1250. <https://doi.org/10.1093/gerona/glaa193>
- Beuret H, Hausler N, Nanchen D, Méan M, Marques-Vidal P, Vaucher J. Comparison of Swiss and European risk algorithms for cardiovascular prevention in Switzerland. *Eur J Prev Cardiol*. 2021;28(2):204-210. <https://doi.org/10.1177/2047487320906305>
- Cassidy S, Fuller H, Chau J, Catt M, Bauman A, Trenell MI. Accelerometer-derived physical activity in those with cardio-metabolic disease compared to healthy adults: a UK Biobank study of 52,556 participants. *Acta Diabetol*. 2018;55(9):975-979. <https://doi.org/10.1007/s00592-018-1161-8>
- Verhoog S, Gubelmann C, Bano A, Muka T, Franco OH, Marques-Vidal P. Comparison of different software for processing physical activity measurements with accelerometry. *Sci Rep*. 2023;13:2879. <https://doi.org/10.1038/s41598-023-29872-7>
- Li X, Zeng J, Chen B, et al. Effects of the timing of intense physical activity on hypertension risk in a general population: a UK-biobank study. *Curr Hypertens Rep*. 2023;26(2):81-90. <https://doi.org/10.1007/s11906-023-01278-w>
- Liu H, Liu S, Wang K, et al. Time-dependent effects of physical activity on cardiovascular risk factors in adults: a systematic review. *Int J Environ Res Publ Health*. 2022;19(21):14194. <https://doi.org/10.3390/ijerph192114194>
- Tian C, Bürki C, Westerman KE, Patel CJ. Association between timing and consistency of physical activity and type 2 diabetes: a cohort study on participants of the UK Biobank. *Diabetologia*. 2023;66(12):2275-2282. <https://doi.org/10.1007/s00125-023-06001-7>
- Kerr D, Abbasi M, Bevier W, Glantz N, Larez A, Sabharwal A. Patterns of timing and intensity of physical activity and HbA1c levels in hispanic/latino adults with or at risk of type 2 diabetes. *J Diabetes Sci Technol*. 2024;18(1):106-112. <https://doi.org/10.1177/19322968221105531>

KEY WORDS cardiovascular, chronotype, chronoactivity, epidemiology, physical activity


APPENDIX For supplemental tables, figures, and the adjudication procedure for cardiovascular disease events, please see the online version of this paper.

Supplemental Figure 1 – K-means clustering analysis, sum of squares plot, selection of clusters for Survey 1 (Follow-up 2, 2014-17)




WSS, within groups sum of squares. PRE, proportional reduction of error. η squared measures the proportional reduction of the WSS for each cluster solution k compared with the total sum of squares. PRE indicates the proportional reduction of the WSS for solution with i clusters compared with the previous solution with $i - 1$ clusters.

Supplemental Figure 2 – K-means clustering analysis, sum of squares plot, selection of clusters for Survey 2 (Follow-up 3, 2018-21)



WSS, within groups sum of squares. PRE, proportional reduction of error. η squared measures the proportional reduction of the WSS for each solution with k clusters compared with the total sum of squares. PRE indicates the proportional reduction of the WSS for solution with i clusters compared with the previous solution with $i - 1$ clusters.

Supplemental Figure 3 – Cardiovascular disease risk according to time of day mean physical activity, CoLausPsyCoLaus study, Lausanne, Switzerland.



CI, confidence interval. Mean physical activity per hour (for each hour of the day), for the total study population and its association with cardiovascular disease risk, using Cox-proportional hazard model, adjusted for age and sex.

Supplemental Table 1 – Characteristics of included and excluded participants, survey 1 (Follow-up 2, 2014-17) and 2 (Follow-up 3, 2018-21), CoLaus|PsyCoLaus study, Lausanne, Switzerland

	Survey 1			Survey 2		
	Included	Excluded	P-value	Included	Excluded	P-value
N	2465	2416		1692	2059	
Female Sex, N (%)	1362 (55.3)	1327 (54.9)	0.818	920 (54.4)	1164 (56.5)	0.185
Age, mean (years)	61.2 ± 9.7	64.7 ± 10.9	<0.001	64.4 ± 9.5	65.9 ± 10.2	<0.001
Educational level, N (%)			<0.001			<0.001
Compulsory education	353 (14.3)	486 (20.1)		214 (12.7)	352 (17.1)	
Apprenticeship	883 (35.8)	866 (35.9)		586 (34.6)	726 (35.3)	
Secondary school	282 (11.4)	264 (10.9)		197 (11.6)	254 (12.4)	
High school degree	378 (15.3)	334 (13.8)		297 (17.6)	277 (13.5)	
University degree	568 (23.1)	463 (19.2)		398 (23.5)	448 (21.8)	
Sleep chronotype, N (%)			<0.001			<0.001
Morning	871 (35.3)	579 (29.2)		568 (33.6)	478 (30.0)	
Evening	916 (37.2)	577 (29.1)		597 (35.3)	481 (30.1)	
Other	678 (27.5)	824 (41.6)		527 (31.2)	637 (39.9)	
Alcohol consumption, N (%)			<0.001			<0.001
None	551 (23.7)	624 (32.6)		550 (32.5)	1142 (55.5)	
1-13/week	1450 (62.3)	1038 (54.3)		935 (55.3)	730 (35.5)	
14-27/week	270 (11.6)	203 (10.6)		172 (10.2)	148 (7.2)	
28+/week	56 (2.4)	48 (2.5)		35 (2.1)	39 (1.9)	
Smoking status, N (%)			0.086			0.050
Never	1052 (42.7)	834 (41.1)		761 (45.0)	734 (43.1)	
Former	973 (39.5)	781 (38.5)		686 (40.5)	670 (39.3)	
Current	440 (17.9)	415 (20.4)		245 (14.5)	299 (17.6)	
Body mass index (kg/m ²)	26.3 ± 4.6	26.6 ± 4.8	0.036	26.1 ± 4.5	26.5 ± 4.9	0.015
Waist circumference (cm)	91.2 ± 13.2	92.7 ± 13.9	<0.001	91.3 ± 13.5	92.5 ± 14.0	0.013
Diabetes, N (%)	190 (7.7)	308 (14.8)	<0.001	139 (8.2)	244 (13.1)	<0.001

Antidiabetic treatment, N (%)	139 (5.6)	258 (10.7)	<0.001	107 (6.3)	189 (9.2)	0.001
Fasting glucose (mmol/l)	5.4 ± 1.0	5.6 ± 1.2	<0.001	5.5 ± 1.0	5.6 ± 1.2	0.013
Fasting insulin (µU/mL)	7.5 [5.1 - 11.2]	8.1 [5.5 - 12.8]	§ <0.001	8.2 [5.7 - 12.4]	8.8 [5.9 - 13.4]	§ 0.001
Lipid levels (mmol/L)						
Total cholesterol	5.4 ± 1.0	5.2 ± 1.1	<0.001	5.3 ± 1.0	5.1 ± 1.1	<0.001
HDL cholesterol	1.6 ± 0.5	1.6 ± 0.5	0.210	1.6 ± 0.4	1.6 ± 0.4	0.295
LDL cholesterol	3.2 ± 0.9	3.0 ± 1.0	<0.001	3.1 ± 0.8	3.0 ± 0.9	<0.001
Triglycerides	1.1 [0.8 - 1.5]	1.1 [0.8 - 1.5]	§ 0.013	1.1 [0.9 - 1.6]	1.1 [0.8 - 1.6]	§ 0.759
Lipid-lowering medication, N (%)	339 (14.2)	600 (25.8)	<0.001	239 (47.6)	396 (59.2)	<0.001
Blood pressure (mmHg)						
Systolic (mean, SD)	126 ± 17	129 ± 19	<0.001	128 ± 18	129 ± 19	0.008
Diastolic (mean, SD)	77 ± 11	77 ± 11	0.788	78 ± 10	78 ± 11	0.642
Antihypertensive treatment, N (%)	677 (27.5)	1020 (42.2)	<0.001	522 (30.9)	826 (40.1)	<0.001

Results are expressed as number of participants (column percentage) for categorical variables and as average ± standard deviation or median [interquartile range] for continuous variables. Between group comparisons performed using Pearson chi-square test for categorical variables and student's t-test or Kruskal-Wallis test (§) for continuous variables. For excluded participants, the total number of participants may vary due to missing data.

Supplemental Table 2 – Participant characteristics by PA timing clusters for the survey 1 and 2 (Follow-ups 2 and 3), CoLaus|PsyColaus study, Lausanne, Switzerland

Cluster number	Survey 1				P-value	Survey 2				P-value
	1	2	3	4		1	2	3	4	
N	481	453	790	741		458	413	480	341	
Female Sex, N (%)	261 (54.3)	260 (57.4)	500 (63.3)	341 (46.0)	<0.001	262 (57.2)	193 (46.7)	307 (64.0)	158 (46.3)	<0.001
Age, mean (years)	59.8 ± 9.4	64.2 ± 10.2	64.1 ± 9.3	57.1 ± 8.4	<0.001	64.5 ± 8.6	62.6 ± 9.4	67.6 ± 9.2	62.0 ± 9.9	<0.001
Educational level, N (%)					<0.001					<0.001
Compulsory education	66 (13.7)	98 (21.6)	129 (16.4)	60 (8.1)		54 (11.8)	47 (11.4)	82 (17.1)	31 (9.1)	
Apprenticeship	142 (29.5)	199 (43.9)	292 (37.0)	250 (33.7)		131 (28.6)	150 (36.3)	211 (44.0)	94 (27.6)	
Secondary school	71 (14.8)	42 (9.3)	77 (9.8)	92 (12.4)		67 (14.6)	39 (9.4)	43 (9.0)	48 (14.1)	
High-school degree	74 (15.4)	55 (12.1)	138 (17.5)	111 (15.0)		90 (19.7)	78 (18.9)	67 (14.0)	62 (18.2)	

University degree	128 (26.6)	59 (13.0)	153 (19.4)	228 (30.8)		116 (25.3)	99 (24.0)	77 (16.0)	106 (31.1)	
Body mass index (kg/m ²)	26.2 ± 4.9	26.4 ± 4.5	26.1 ± 4.5	26.5 ± 4.7	0.513	25.7 ± 4.0	25.9 ± 4.4	26.3 ± 4.6	26.9 ± 4.9	0.001
Waist circumference (cm)	90.9 ± 13.5	91.3 ± 13.1	90.7 ± 12.9	92 ± 13.3	0.293	89.6 ± 12.2	91.0 ± 13.6	91.3 ± 13.2	94.0 ± 15.0	<0.001
Sleep chronotype, N (%)					0.293					<0.001
Morning person	53 (11.0)	259 (57.2)	266 (33.7)	293 (39.5)		91 (19.9)	217 (52.5)	198 (41.3)	62 (18.2)	
Evening person	308 (64.0)	82 (18.1)	305 (38.6)	221 (29.8)		210 (45.9)	89 (21.6)	113 (23.5)	185 (54.3)	
Other	120 (25.0)	112 (24.7)	219 (27.7)	227 (30.6)		157 (34.3)	107 (25.9)	169 (35.2)	94 (27.6)	
Alcohol consumption, N (%)					0.062					0.464
None	121 (26.5)	122 (29.3)	165 (22.0)	143 (20.3)		151 (33.0)	124 (30.0)	164 (34.2)	111 (32.6)	
1-13/week	270 (59.1)	240 (57.7)	484 (64.5)	456 (64.9)		248 (54.2)	244 (59.1)	262 (54.6)	181 (53.1)	
14-27/week	55 (12.0)	43 (10.3)	84 (11.2)	88 (12.5)		53 (11.6)	38 (9.2)	42 (8.8)	39 (11.4)	
28+/week	11 (2.4)	11 (2.6)	18 (2.4)	16 (2.3)		6 (1.3)	7 (1.7)	12 (2.5)	10 (2.9)	
Smoking status, N (%)					<0.001					0.018
Never	182 (37.8)	192 (42.4)	355 (44.9)	323 (43.6)		191 (41.7)	183 (44.3)	233 (48.5)	154 (45.2)	
Former	182 (37.8)	192 (42.4)	323 (40.9)	276 (37.3)		206 (45.0)	166 (40.2)	192 (40.0)	122 (35.8)	
Current	117 (24.3)	69 (15.2)	112 (14.2)	142 (19.2)		61 (13.3)	64 (15.5)	55 (11.5)	65 (19.1)	

Results are expressed as number of participants (column percentage) for categorical variables and as average ± standard deviation for continuous variables. Between-group comparisons performed using chi-square for categorical variables and analysis of variance for continuous variables.

Supplemental Table 3 – Participant characteristics by physical activity timing clusters based on METs, multivariable analysis, survey 1 (Follow-up 2, 2014-17), CoLaus|PsyCoLaus study, Lausanne, Switzerland

Cluster number	1	2	3	4	P-value §
Hypertension	1 (reference)	1.20 (0.92 - 1.57)	0.97 (0.75 - 1.26)	1.13 (0.87 - 1.46)	
Hypolipidemic drug treatment	1 (reference)	1.15 (0.81 - 1.63)	0.91 (0.63 - 1.33)	0.97 (0.68 - 1.38)	
Diabetes	1 (reference)	1.48 (0.95 - 2.29)	1.22 (0.74 - 1.99)	0.90 (0.55 - 1.47)	
Blood pressure (mm Hg)					
Systolic	126 ± 1	125 ± 1	124 ± 1	126 ± 1	0.281
Diastolic	77 ± 1	78 ± 1	77 ± 1	78 ± 1	0.561
Lipid levels (mmol/L)					
Total cholesterol	5.51 ± 0.04	5.32 ± 0.04 ^a	5.37 ± 0.04	5.40 ± 0.04	0.007
HDL cholesterol	1.61 ± 0.02	1.58 ± 0.02	1.59 ± 0.02	1.66 ± 0.02 ^b	0.012
LDL cholesterol	3.31 ± 0.03	3.16 ± 0.04 ^a	3.18 ± 0.04	3.19 ± 0.04	0.015
Triglycerides	1.31 ± 0.04	1.28 ± 0.05	1.31 ± 0.05	1.24 ± 0.05 ^c	0.014 ‡
Diabetes markers					
Glucose (mmol/L)	5.37 ± 0.03	5.34 ± 0.04	5.39 ± 0.04	5.43 ± 0.04	0.305
Insulin (µIU/mL)	9.1 ± 0.2	9.4 ± 0.2	9.0 ± 0.2	9.1 ± 0.2	0.373 ‡
HbA _{1c} (mmol/mol)	37.5 ± 0.2	37.5 ± 0.2	37.7 ± 0.2	37.9 ± 0.2	0.536

§ for ANOVA; ‡ based on log-transformed data. Results are expressed as multivariable-adjusted odds ratio and (95% confidence interval) for hypertension, hypolipidemic drug treatment and diabetes, and as multivariable-adjusted marginal mean ± standard error for the other variables. Statistically significant ($p < 0.05$) odds ratios are indicated in bold characters. Multivariable analysis performed using logistic regression for categorical variables and by ANOVA for continuous variables. For continuous variables, post-hoc bivariate comparisons were performed using Scheffe's method: subscript a refers to comparison of clusters 2, 3 and 4 to cluster 1, subscript b to comparison of clusters 3 and 4 to cluster 2, and subscript c to comparison of cluster 4 to cluster 3; presence of the subscript indicates that clusters differ at $p < 0.05$. All multivariable models adjusted for sex (male, female), age (continuous), educational level (basic, apprenticeship, secondary, high school, and university), alcohol consumption (none, 1-13, 14-27 and 27+ per week), smoking categories (never, former, current) and BMI categories (normal, overweight, obese). For blood pressure levels, a further adjustment on antihypertensive drug treatment (yes, no) was performed; for lipid levels, a further adjustment on hypolipidemic drug treatment (yes, no) was performed, and for diabetes markers, a further adjustment on antidiabetic drug treatment (yes, no) was performed.

Supplemental Table 4 – Participant characteristics by physical activity timing clusters based on METs, multivariable analysis, survey 2 (Follow-up 3, 2018-21), CoLaus|PsyCoLaus study, Lausanne, Switzerland

Cluster number	1	2	3	4	P-value §
Hypertension	1 (reference)	1.06 (0.79 - 1.42)	1.34 (0.98 - 1.84)	1.38 (1.01 - 1.91)	
Hypolipidemic drug treatment	1 (reference)	0.87 (0.51 - 1.49)	1.30 (0.72 - 2.36)	0.97 (0.53 - 1.75)	
Diabetes	1 (reference)	0.90 (0.52 - 1.56)	1.16 (0.67 - 2.01)	1.51 (0.88 - 2.57)	
Blood pressure (mm Hg)					
Systolic	127 ± 1	127 ± 1	129 ± 1	127 ± 1	0.310
Diastolic	77 ± 1	78 ± 1	78 ± 1	77 ± 1	0.991
Lipid levels (mmol/L)					
Total cholesterol	5.43 ± 0.09	5.40 ± 0.07	5.27 ± 0.09	5.32 ± 0.09	0.507
HDL cholesterol	1.46 ± 0.03	1.54 ± 0.03	1.52 ± 0.03	1.44 ± 0.03	0.127
LDL cholesterol	3.23 ± 0.08	3.22 ± 0.06	3.13 ± 0.08	3.17 ± 0.08	0.743
Triglycerides	1.59 ± 0.09	1.40 ± 0.07	1.49 ± 0.09	1.56 ± 0.09	0.058‡
Diabetes markers					
Glucose (mmol/L)	5.54 ± 0.04	5.46 ± 0.04	5.51 ± 0.04	5.55 ± 0.05	0.353
Insulin (µIU/mL)	10.3 ± 0.3	9.6 ± 0.3	9.7 ± 0.4	11.4 ± 0.4 ^b	0.004‡
HbA _{1c} (mmol/mol)	37.1 ± 0.3	37.0 ± 0.2	36.9 ± 0.3	37.4 ± 0.3	0.483

§ for ANOVA; ‡ based on log-transformed data. Results are expressed as multivariable-adjusted odds ratio and (95% confidence interval) for hypertension, hypolipidemic drug treatment and diabetes, and as multivariable-adjusted marginal mean ± standard error for the other variables. Statistically significant ($p < 0.05$) odds ratios are indicated in bold characters. Multivariable analysis performed using logistic regression for categorical variables and by ANOVA for continuous variables. For continuous variables, post-hoc bivariate comparisons were performed using Scheffe's method: subscript a refers to comparison of clusters 2, 3 and 4 to cluster 1, subscript b to comparison of clusters 3 and 4 to cluster 2, and subscript c to comparison of cluster 4 to cluster 3; presence of the subscript indicates that clusters differ at $p < 0.05$. All multivariable models adjusted for sex (male, female), age (continuous), educational level (basic, apprenticeship, secondary, high school, and university), alcohol consumption (none, 1-13, 14-27 and 27+ per week), smoking categories (never, former, current) and BMI categories (normal, overweight, obese). For blood pressure levels, a further adjustment on antihypertensive drug treatment (yes, no) was performed; for lipid levels, a further adjustment on hypolipidemic drug treatment (yes, no) was performed, and for diabetes markers, a further adjustment on antidiabetic drug treatment (yes, no) was performed.

Supplemental Table 5 – Analysis of the associations between physical activity timing clusters and cardiovascular events, CoLaus|PsyCoLaus study, Lausanne, Switzerland, using inverse probability weighting to account for excluded participants.

Cluster number	1	2	3	4
Bivariate	1 (reference)	4.13 (1.36 - 12.6)	3.52 (1.20 - 10.3)	2.57 (0.84 - 7.81)
Multivariable 1	1 (reference)	3.39 (1.09 - 10.5)	2.89 (0.99 - 8.47)	3.21 (1.04 - 9.94)
Multivariable 2	1 (reference)	3.51 (1.12 - 11.0)	2.99 (1.02 - 8.76)	3.21 (1.04 - 9.96)

For bivariate and multivariable analysis, results are expressed as hazard ratio and (95% confidence interval). Analyses conducted using Cox regression. Multivariable model 1: adjusted for sex (male, female), age (continuous), educational level (basic, apprenticeship, secondary, high school, and university), alcohol consumption (none, 1-13, 14-27 and 27+ per week), smoking categories (never, former, current), and BMI categories (normal, overweight, obese). Multivariable model 2: as model 1, plus hypertension (yes, no), diabetes (yes, no) and hypolipidemic drug treatment (yes, no). Significant ($p < 0.05$) coefficients are indicated in bold.

Supplemental Table 6 – Analysis of the associations between physical activity timing clusters and cardiovascular events, CoLaus|PsyCoLaus study, Lausanne, Switzerland, stratified by age or physical activity group.

Cluster	Age¹	Cases	Non-cases	HR (95% CI)	P-value	Interaction
Cluster 1	<65 years	2	283	1		
	≥65 years	2	99	1		
Cluster 2	<65 years	8	185	6.37 (1.35 - 30.0)	0.019	
	≥65 years	11	160	3.35 (0.74 - 15.1)	0.116	0.564
Cluster 3	<65 years	11	350	4.82 (1.06 - 21.8)	0.041	
	≥65 years	19	290	3.20 (0.74 - 13.7)	0.118	0.748
Cluster 4	<65 years	10	538	2.46 (0.54 - 11.3)	0.245	
	≥65 years	7	84	4.49 (0.92 - 21.8)	0.063	0.715
Physical activity²						
Cluster 1	Less active	2	196	1		
	More active	2	186	1		
Cluster 2	Less active	14	140	7.20 (1.61 - 32.2)	0.010	
	More active	5	205	1.62 (0.31 - 8.37)	0.567	0.258
Cluster 3	Less active	15	301	3.97 (0.90 - 17.5)	0.068	
	More active	15	339	2.78 (0.63 - 12.3)	0.176	0.866
Cluster 4	Less active	11	281	4.36 (0.95 - 19.9)	0.057	
	More active	6	341	1.88 (0.38 - 9.36)	0.443	0.426

Results are expressed as hazard ratio and (95% confidence interval). Analyses conducted using Cox regression. ¹ adjusted on sex; ² adjusted on age and sex. Interaction : effect interaction between the less active / more active groups if p-value for interaction is <0.05.

Annex 1- Adjudication procedure for cardiovascular disease events

Vital status of the participants was assessed at each follow-up exam. If the follow-up information letter was returned by the post office, or the participant did not respond to the successive phone call to schedule the follow-up exam, the civil register of the community in which the subject was living at the time of the last exam was contacted to determine the participant's new address, or the occurrence and date of death. If the participant moved within Switzerland, the new address was systematically recorded, and a new contact was established. If no contact could be obtained, vital status was controlled through the civil register of their new community. If the participant moved outside Switzerland, no information regarding vital status could be obtained and the last contact date was used.

Cause of death was established using all available sources of information: general practitioners, any hospital in Switzerland (if death occurred in this setting), the pre-hospital emergency care unit of the City of Lausanne, the forensic medicine department of the University of Lausanne, and the Swiss Statistics of the Swiss Confederation. If medical records were available, cause of death was adjudicated by two internal medicine specialists. Official death certificates were used only if medical reports were unavailable.¹

1. Beuret H, Hausler N, Nanchen D, Méan M, Marques-Vidal P, Vaucher J. Comparison of Swiss and European risk algorithms for cardiovascular prevention in Switzerland. *Eur J Prev Cardiol.* 2021;28(2):204-210. doi:10.1177/2047487320906305