

# Sperm of more colourful males are better adapted to ovarian fluids in lake char (*Salmonidae*)

David Nusbaumer | Laura Garaud | Christian de Guttry | Laurie Ançay |  
Claus Wedekind 

Department of Ecology & Evolution,  
University of Lausanne, Lausanne,  
Switzerland

## Correspondence

Claus Wedekind, Department of Ecology  
& Evolution, University of Lausanne,  
Lausanne, Switzerland.  
Email: [claus.wedekind@unil.ch](mailto:claus.wedekind@unil.ch)

## Funding information

Schweizerischer Nationalfonds zur  
Förderung der Wissenschaftlichen  
Forschung, Grant/Award Number:  
31003A\_159579 and 31003A\_182265

Handling Editor: Michael M. Hansen

## Abstract

Fish often spawn eggs with ovarian fluids that have been hypothesized to support the sperm of some males over others (cryptic female choice). Alternatively, sperm reactions to ovarian fluids could reveal male strategies. We used wild-caught lake char (*Salvelinus umbla*) to experimentally test whether sperm react differently to the presence of ovarian fluid, and whether any differential sperm reaction could be predicted by male breeding coloration, male inbreeding coefficients (based of 4150 SNPs) or the kinship coefficients between males and females. Male coloration was positively linked to body size and current health (based on lymphocytosis and thrombocytosis) but was a poor predictor of inbreeding or kinship coefficients. We found that sperm of more colourful males were faster in diluted ovarian fluids than in water only, while sperm of paler males were faster in water than in ovarian fluids. We then let equal numbers of sperm compete for fertilizations in the presence or absence of ovarian fluids and genetically assigned 1464 embryos (from 70 experimental trials) to their fathers. The presence of ovarian fluids significantly increased the success of the more colourful competitors. Sperm of less inbred competitors were more successful when tested in water only than in diluted ovarian fluids. The kinship coefficients had no significant effects on sperm traits or fertilization success in the presence of ovarian fluids, although parallel stress tests on embryos had revealed that females would profit more from mating with least related males rather than most coloured ones. We conclude that sperm of more colourful males are best adapted to ovarian fluids, and that the observed reaction norms suggest male strategies rather than cryptic female choice.

## KEYWORDS

cryptic female choice, inbreeding coefficient, kinship coefficient, loaded raffle, male reproductive strategy, sperm competition, sperm reaction norm, sperm velocity

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Molecular Ecology* published by John Wiley & Sons Ltd.

## 1 | INTRODUCTION

Ejaculate economics predicts that males invest strategically into sperm according to the expected microecology they will experience after ejaculation (Parker & Pizzari, 2010). This microecology is strongly defined by competitors and the level of sperm competition. Males are predicted to increase sperm number and/or sperm velocity with increasing risk of sperm competition (Parker & Pizzari, 2010). This prediction is meanwhile well supported in many taxa (Magris, 2021). Another aspect of the microecology that sperm have to deal with is the biochemical environment that females create around their eggs with their reproductive fluids. Such female reproductive fluids occur in internal and external fertilizers and include, for example, ovarian, follicular, oviductal or coelomic fluids (Gasparini et al., 2020). The functional significance of such female reproductive fluids is not well understood yet (Gasparini et al., 2020).

In externally spawning fishes, the fluids with which females expel their eggs are called 'ovarian fluids' and consist of various inorganic and organic compounds (Lahnsteiner et al., 1995). Ovarian fluids have repeatedly been found to support sperm velocity, motility and longevity (Zadmajid et al., 2019), that is, key targets of sexual selection (Evans et al., 2013). They have also been discussed as allowing for certain forms of cryptic female choice (Firman et al., 2017). Their release with eggs can, for example, increase sperm competition by slowing down the fertilization process (Bakker et al., 2006) or specifically promote paternity of dominant males over parasitic spawners ('sneaker males') (Alonzo et al., 2016; Egeland et al., 2016; Lehnert et al., 2017; Makiguchi et al., 2016). The support of sperm function can be even more specific: Ovarian fluids of guppies (*Poecilia reticulata*) were found to slow down sperm of full-sibs as compared to unrelated males, potentially to reduce inbreeding (Gasparini & Pilastro, 2011). The presence of ovarian fluids during spawning has also been found to reduce inbreeding by differentially supporting some sperm over others in Chinook salmon (*Oncorhynchus tshawytscha*; Geßner et al., 2017; Lehnert et al., 2017; Rosengrave et al., 2016), while ovarian fluids of lake trout (*Salvelinus namaycush*) were found to promote sperm velocity of more related males (Butts et al., 2012), potentially to reduce outbreeding depression in this salmonid. Indeed, Yeates et al. (2013) found that ovarian fluids specifically support sperm of the same *Salmo* species over sperm of another *Salmo* species, suggesting a species-specific chemoattraction promoting reproductive isolation.

So far, likely effects of ovarian fluids on sperm have mainly been discussed in the light of possible adaptive female reproductive strategies. However, sperm characteristics can also reveal male reproductive strategies, as predicted by the theory of ejaculate economics (Parker & Pizzari, 2010) and experimentally demonstrated in different taxa (Jeannerat et al., 2018; Rudolfen et al., 2006). It is therefore possible that many of the observed effects of ovarian fluids on sperm function reveal variation in male reproductive strategies. If so,

males would be expected to tailor sperm production and/or ejaculate characteristics to the likely exposure of their sperm to ovarian fluids, that is, to the synchrony with, and the position to, the female during egg release (Egeland et al., 2016).

Salmonids are externally spawning fish whose ovarian fluids vary in amount and composition among species and even within populations (Lahnsteiner et al., 1995; Zadmajid et al., 2019). Viscosity of undiluted ovarian fluids is about two to three times higher than that of water (Zadmajid et al., 2019), but ovarian fluids will quickly be diluted when expelled with eggs into water. During spawning, freshly spawned eggs and ovarian fluids are swirled around by male and female movements. Therefore, some eggs may no more be surrounded by significant amounts of ovarian fluids at the time of fertilizations. A likely predictor of how much sperm will be exposed to ovarian fluids is the position of the male during spawning. Arguably the closer a male is with its urogenital opening to a female's vent during egg release, the more likely its sperm will be exposed to ovarian fluids.

Salmonid males can play various roles during spawning (Esteve, 2005), and many of these roles have been found to affect milt and sperm characteristics (Magris, 2021). In migratory populations, for example, there is usually a fraction of males that avoid the costs of migration but then remain small and subdominant to their migratory rivals at spawning. They, therefore, reproduce mainly surreptitiously and invest disproportionately into sperm number and sperm velocity in order to increase the competitiveness of their sperm (Young et al., 2013). Evidence for strategic male investments has also been found within non-migratory populations. Male dominance at the spawning place is usually size dependent (Esteve, 2005), and older males and larger male whitefish (*Coregonus muelleri*) (Selz & Seehausen, 2023) have indeed been found to invest less into gonad weight and to have slower sperm than young and small males who may typically spawn further away from the female's vent (Rudolfen et al., 2008). However, the links between male size and sperm traits can vary among populations (Perroud et al., 2021) and be plastic in response to the perceived social environment (Bartlett et al., 2017). When Rudolfen et al. (2006), for example, led size-matched male Arctic char (*Salvelinus alpinus*) develop a dominance hierarchy, sperm characteristics changed quickly: as predicted from ejaculate theory, dominance led to reduced sperm velocity. Egeland et al. (2016) then used an analogous set-up to show that sperm of dominant males swim better in ovarian fluids. One possible explanation for this increased performance is that ovarian fluids interact with milt of dominant males and specifically promote their sperm (cryptic female choice). An alternative hypothesis is that dominant males can expect to be close to the females during gamete release and may therefore produce sperm that do well when exposed to ovarian fluids. If so, any other male trait that is likely to affect the distance between male and female during spawning would be predicted to affect sperm reaction to ovarian fluids. Here, we use wild-caught lake char (*Salvelinus umbla*) from Lake Geneva (Switzerland) to test the reaction of sperm of different kind of males to the presence of ovarian fluids.

The lake char is a non-migratory salmonid endemic to Alpine lakes. It spawns in winter in lek-like mating system where male competition is typically intense and male mating success can be expected to be positively correlated with their sexual ornamentation (Fiske et al., 1998). The lake char is closely related to the Arctic char (Kottelat & Freyhof, 2007) that has often been used to study effects of ovarian fluids on sperm motility (e.g. Egeland et al., 2016; Turner & Montgomerie, 2002; Urbach et al., 2005). Both species develop spawning colorations. Arctic char often develop strong red colorations in both sexes (Janhunen et al., 2011). Male lake char of our study population are mostly yellow during spawning season while females hardly develop any spawning coloration. This sex difference and the colour differences within males are, to the best of our knowledge, not sufficiently understood yet.

In other fish families, sexual ornaments have sometimes been found to be positively linked to aspects of sperm performance, for example, the carotenoid-based skin colours in three-spined sticklebacks (*Gasterosteus aculeatus*; Mehliis & Bakker, 2013), European minnows (*Phoxinus phoxinus*; Kekäläinen et al., 2014) or cherry barbs (*Puntius titteya*; Fukuda & Karino, 2014; Kortet et al., 2004). A common argument here is that both types of traits can react to induced stress, suggesting that high-quality sperm in well-ornamented males is an indicator of health and vigour rather than of alternative mating strategies. The pattern in char seems less clear. It has been suggested that skin colours in Arctic char reveal health and vigour and hence dominance (Skarstein & Folstad, 1996; but see Backström et al., 2015). Rudolfson et al. (2006) found in wild Arctic char that dominant males have reduced sperm velocity. However, Janhunen et al. (2009) found in a captive population of Arctic char that the intensity of male coloration was positively correlated with sperm velocity when tested in water only.

Here, we use haematological analyses (Seibel et al., 2021) to first test whether male coloration or their inbreeding coefficient ( $F_{\text{beta}}$ ) are indicators of current health and vigour in wild-caught lake char. High leucocyte counts and especially relative lymphocytosis (percentage of lymphocytes among leucocytes) have been associated with acute infections (Haenen et al., 2010) or other forms of stress (Meuthen et al., 2020; Oluah et al., 2020). Thrombocytosis (elevated platelet counts) is another possible indicator of an acute infection (Rose et al., 2012; Yan et al., 2013). We also test whether male coloration is correlated to inbreeding coefficients ( $F_{\text{beta}}$ ).

In a first series of experiments, we then test whether (i) male size, skin coloration or  $F_{\text{beta}}$  predicts sperm performance in water, (ii) sperm of different types of males react differently to the presence of ovarian fluids and (iii) ovarian fluids vary in their effects on sperm. In a second series of experiments, we use sperm competition trials and genotyping of the resulting offspring to test whether (iv) the competitive success can be predicted by male colour, male  $F_{\text{beta}}$  or their kinship coefficient to the female that would determine the average inbreeding coefficient of the offspring, and (v) the presence of ovarian fluids change the outcome of sperm competition.

## 2 | METHODS

### 2.1 | Sampling and handling of fish, gametes and ovarian fluids

Wild char were caught in December 2017 and 2018 from three spawning sites in Lake Geneva (Ripaille, 46.3957N–6.4718E; Locum, 46.4049N–6.7588E; Meillerie, 46.4113N–6.7216E) by local fishermen using gill nets (48 mm mesh size, set over night) and transported to a research facility (INRA) where they were kept in a 1000L circular tank with free-flowing lake water before further processing. In 2017, 10 males and 4 females were haphazardly chosen for the experiment. Males were processed on the first day after catching and females on the second day. Each fish was anaesthetized in 10L of water containing 3 mL of Eugenol (clove oil in ethanol at a 1:9 ratio), immediately photographed in a custom-made photo box under standardized conditions (17 mm, f/5.0, 1/200s, ISO 400, WB 4000K, JPG 24 Mpx) together with a standard colour scale (Tiffen), and stripped to collect gametes. Tissue samples were taken from the anal fin and stored in 70% ethanol at  $-20^{\circ}\text{C}$  for further analysis. In 2018, further 24 males were sampled and handled as in 2017 to test again for correlations between skin colorations and sperm characteristics.

Milt was directly stripped into 50-mL conical tubes (Falcon, BD Biosciences), carefully avoiding contamination by urine, faeces or water. Between 2 and 5 mL of milt were obtained per male. These samples were stored at  $4^{\circ}\text{C}$  (<2 h) until further use. When all males had been stripped, 1.5 mL of each milt sample was diluted 1:9 in Storfish (IMV Technologies; an inactivating medium) in 15-mL tubes (Falcon, BD Biosciences). These tubes were then placed on ice and are referred to as 'diluted milt.'

Eggs and ovarian fluids were stripped into individual plastic containers. About 20 mL of ovarian fluid each was separated from the eggs with a syringe and stored in 50 mL tubes at  $4^{\circ}\text{C}$ . Eggs were kept (<1 h) in the dark at  $4^{\circ}\text{C}$  in about 20-mL Ovafish (IMV Technologies) to prevent drying.

### 2.2 | Colour measurement

Custom-made macros in Fiji (Schindelin et al., 2012) were used to analyse male skin colorations. First, the white balance of each image was standardized with the help of the white and black values of the colour scale. Skin grey value was then determined as the overall body grey value in the RGB colour space (range from 0 = black, to 255 = white; using a mean of all pixels in each of the RGB channels). Redness and yellowness were measured from three squares (about 1% of the total body area each) in the pectoral region, the ventral region and the anal region, as described by Parolini et al. (2018) (Figure S1). Redness and yellowness were measured as the  $a^*$  and  $b^*$  components of the CIE- $L^*a^*b^*$  colour space respectively (the  $a^*$  axis for greenness–redness and the  $b^*$  axis for blueness–yellowness both range from  $-100$  to  $100$ ). All

measures of yellowness were significantly correlated (Pearson's  $r > .92$ ,  $p$  always  $< .001$ ). Similarly, all measures of redness were significantly correlated (Pearson's  $r > .75$ ,  $p < .013$ ). Therefore, the means of the three measurements per fish were used for each colour. Grey values were correlated with yellowness (the yellower the lighter) and with redness (the redder the darker; [Figure S1](#)) and were therefore ignored in further analyses.

## 2.3 | Sperm characteristics

Sperm characteristics of all males caught in 2017 and 2018 were measured on the day of sampling with Qualisperm® (AKYmed). Briefly, 20  $\mu$ L of the diluted milt was activated in a microtube with 980- $\mu$ L ice-cooled standardized water (OECD, 1992) and vortexed for 5 s. Then, 2  $\mu$ L of this activated solution was pipetted into the well of a 4-well chamber slide (Leja, AKYmed), on a cooling station set at 4.5°C (the expected water temperature at the spawning ground). Sperm concentration, sperm velocity (average path velocity) and percentage of immotile sperm (sperm motility) were recorded under a phase-contrast microscope at 20 $\times$  magnification 20 s after activation in 2017 and 25 s after activation in 2018. Sperm longevity was recorded with a stopwatch from the time of activation until no progressive sperm motion could be observed in the frame of capture. Two replicated measures were taken for each sperm sample to test for consistency and means of these measurements were used for statistics. The concentration of active spermatozoa was calculated for each male by multiplying the total sperm concentration with the rate of motile sperm (e.g. 2000 Mio/mL times 80% motile sperm = 1600 Mio/mL active spermatozoa).

## 2.4 | Haematological analyses

In order to determine leucocyte counts, relative lymphocytosis and thrombocytosis, a blood sample (2–3 mL) was drawn from the caudal vasculature of the 10 males sampled in 2017 with a 3-mL syringe mounted with a 23-G needle heparinized with heparin 1 IU/ $\mu$ L. The blood was emptied in 2-mL microtubes, centrifuged 10 min at 10,000g and 4°C. The plasma was then carefully collected, disposed in 1.5-mL microtubes and frozen in the vapour of liquid nitrogen for transportation and stored at -80°C in the lab for further analysis. A drop of blood from the syringe was used to make two blood smears per individual, on microscope frosted slides. Blood smears were dried in the field and fixed for 3 min in absolute ethanol back in the lab. Slides were stained for 20 min in Giemsa stain and rinsed twice in PBS. From each slide, 10 photographs (i.e. 20 per individual) were taken on an EVOS XL Core inverted light transmission microscope (ThermoFisher) at 40 $\times$  magnification. Counts of thrombocytes, lymphocytes and granulocytes were done manually by a naïve observer ([Figure S2](#)). Counts of total number of cells were done automatically with a custom-made macro in Fiji ([Figure S2](#)). They were used to determine 'leucocyte counts' (% of all blood cells), 'relative

lymphocytosis' (% of lymphocytes among leucocytes) and 'thrombocytosis' (thrombocytes counts per 100 blood cells).

## 2.5 | Experiment 1: Sperm characteristics in response to ovarian fluids

Experiments 1 and 2 were done with the 2017 sample. Sperm characteristics were determined again 1 day after the fish had been sampled from the wild, to determine each male's sperm characteristics in each female's ovarian fluid (40 combinations of milt samples and ovarian fluids) and in water (controls, 2 replicates each; [Table S1](#)). To do so, 10  $\mu$ L of the diluted milt was activated either with 490- $\mu$ L standardized water or with 490- $\mu$ L ovarian fluid solution. The concentration of ovarian fluid in this solution was 50% as this would reflect a possible concentration of ovarian fluid in the water when sperm swim towards the egg (Rosengrave et al., 2016).

## 2.6 | Experiment 2: Competitive fertilization trials and embryo monitoring

The 10 males were haphazardly assigned to 5 dyads (pairs of competitors) whose sperm then competed for fertilization of 24 eggs/female, either activated in water only or in water with ovarian fluid. Each possible combination of dyad  $\times$  female was tested in both environments twice, resulting in 80 sperm competition trials (5 dyads  $\times$  4 females  $\times$  2 treatments  $\times$  2 replicates; [Table S2](#)) that involved in total 1920 eggs. Due to an accident during handling, 10 of these 80 experimental cells were lost (all of the same female, prepared to be exposed to sperm activated in water only, see [Table S2](#)), reducing the number of eggs that could be monitored to  $N = 1680$ .

In preparation of these sperm competition trials, the eggs of each female were first washed twice with 200-mL Ovafish® (IMV Technologies) to remove ovarian fluids. Then, 20 batches of 24 eggs each were placed in wells of six-well plates (Falcon, BD Biosciences). Ten millilitres of diluted milt of two males each was prepared such that each male was represented with the same concentration of active sperm (25 Mio/mL) in the mix. One millilitre of each mix was then used to fertilize the two batches of 24 eggs/female. This 1 mL of mix was activated in a separated tube with either 4 mL of standardized water or 4 mL of standardized water with ovarian fluid (ratio ovarian fluid to water = 1:2) and vortexed for 5 s. The solution was poured in a well with eggs. Two minutes later, that is, after fertilization could be expected to have happened, standardized water was added to fill the wells (16.8 mL) and the eggs were left undisturbed in a dark environment for 2 h to allow hardening.

Each batch of 24 supposedly fertilized eggs was transferred to a 50-mL tube filled with standardized water and transported on ice to a climate chamber where they were rinsed for 30 s under running tap water (4 L/min) in a sterilized tea strainer. The eggs were then distributed singly to wells of 24-well plates (Falcon, BD Biosciences) filled with 1.8 mL of autoclaved standardized water (OECD, 1992).

Embryos were raised at 4.5°C in a 12h:12h light-dark cycle. Eggs were classified as non-fertilized if no embryo could be found at the neurula stage. Embryo mortality, the timing of hatching, hatchling size and hatchling growth were monitored from then on in the course of two parallel studies, one that combines these data with further stress experiments (by a pathogen and a chemical pollutant) on embryos produced in a regular full-factorial breeding design to test for parental effects on embryo performance (Garaud et al., 2023), and another one on sex-specific stress tolerance (Nusbaumer et al., 2021). Embryos and larvae that died during the observational period were transferred to individual microtubes and stored in 96% ethanol at -20°C for further analysis. Larvae were euthanized 14 days after hatching with a spike of 100 µL of Koi Med® Sleep 4.85% (Koi & Bonsai Zimmermann) in the well before being transferred to individual microtubes.

## 2.7 | Genotyping and paternity assignment based on microsatellite markers

In the case of larvae and adults, DNA was extracted from up to 25 mg tissue with an extraction robot and the DNAeasy Blood & Tissue kit (Qiagen) following manufacturer's instructions. Unhatched embryos were homogenized and DNA was extracted using the Qiaamp Fast DNA Stool mini kit protocol (Qiagen) as in Wilkins et al. (2015). A first multiplex PCR of three polymorphic microsatellite markers (Savary et al., 2017) turned out not sufficient to assign all embryos and larvae to their fathers (Bernatchez & Duchesne, 2000). A second multiplex PCR of three further microsatellite markers (Savary et al., 2017) was therefore used to assign the remaining offspring (Table S3). In the first multiplex PCR, the *sdY* sex marker (Yano et al., 2013) was included to sex offspring (Table S3) for a parallel study on sex-specific stress tolerance (Nusbaumer et al., 2021). Multiplex PCR were run on a Biometra thermocycler in 20-µL reaction volume using HotStarTaq DNA Polymerase (Qiagen) reagents (i.e. 2-µL PCR buffer 10×, 4-µL QSolution 5×, 0.4-µL HotStar Taq DNA Polymerase, 0.4-µL of dNTPs 10mM and 0.4 µL of each primer). PCR cycles were run as follow: initial heat activation of 15 min at 95°C; 35 cycles of 30s denaturation at 94°C, 90s annealing at 57°C and 60s extension at 72°C; followed by 30min of final extension at 72°C and 10min at 4°C. DNA extracted from unhatched embryo was amplified using the same protocol, but in 10-µL reaction volume and with 40 cycles. PCR products were diluted 2X and run on an ABI3100 sequencer (Applied Biosystems). Genotypes were read using Genemapper v4.0 (Applied Biosystems). Genotyping of three dead embryos was not successful possibly due to poor DNA quality, bringing the final number of genotyped offspring to  $N = 1475$ .

Adults were genotyped twice (no discrepancies observed). Paternity was assigned using CERVUS v3.0 (Kalinowski et al., 2007) as recommended for categorical assignments in controlled experiments (Jones et al., 2010). Input simulation parameters were set with six loci, 10,000 offspring, two candidate fathers with a proportion of sampling of one and a proportion of typing of one, a minimum of

two typed loci and an allelic dropout/mistyping rate of 0.05 as recommended by Wang (2004), and an assignment confidence of 95%.

## 2.8 | Double-digested RAD sequencing and SNPs calling

DNA from all males and females of experiment 2 was used to also determine their multilocus heterozygosity (MLH) and the genomic similarity ('kinship coefficient') between males and females of all possible combinations ( $N = 40$ ). DNA concentration was measured using Qubit 2.0 (Thermo Fisher Scientific) while its integrity was verified on a 1% agarose gel. Each individual's DNA concentration was then standardized to 20 ng/µL. Two sequencing libraries were produced using all 10 males and four females each following the Brelford et al. (2016) protocol adapted from Parchman et al. (2012). Briefly, DNA was digested using the enzymes EcoRI-HF and MspI (New England Biolab, Ipswich, Massachusetts, United States) and a unique EcoRI barcode was ligated to each individual. After library purification, PCR amplification was performed and fragments size-selected in between 400 and 550 bp. Libraries were then single-end genotyped on two lanes of Illumina HiSeq 2500 with fragments of 125 bp length at Lausanne Genomic Technologies Facility (University of Lausanne).

After the quality control on the resulting fastq files done with FASTQC v0.11.7 (Andrews, 2010), reads were trimmed to 110 bp given the low per-base quality of the sequences (<20 Phred score). The resulting RAD data were analysed using Stacks 1.48 (Catchen et al., 2013) using the default parameters unless otherwise specified. Individual sequences were demultiplexed using *process\_radtags* (Stacks 1.48) and reads mapped to the *Salvelinus* spp. reference genome (NCBI accession number SRP101753) using BWA (Li & Durbin, 2010). *Pstacks* (Stacks 1.48) was done selecting a minimum stack depth of 5 ( $-m$  5) and using the bounded SNP model. The catalogue of loci was created using *Cstacks* (Stacks 1.48) allowing two mismatches ( $-n$  2) between loci. *Populations* (Stacks 1.48) was used to generate the VCF file considering only loci present in 100% of the individuals and markers heterozygosity of 0.5 ( $--max\_obs\_het$  0.50). None of the individuals was excluded based on genotyping rate. Further filtering was done using *vcftools*, v0.1.15 (Danecek et al., 2011). To reduce incorrect heterozygosity call and remove paralogs, loci were filtered for a minimum coverage of 10× ( $--min-meanDP$  10) and a maximum coverage of 50× ( $--max-meanDP$  50; two times the mean coverage). Only loci under Hardy-Weinberg equilibrium were retained with a  $p$ -value threshold of .05 ( $--hwe$  0.05). Finally, all the loci that were not shared between all individuals were discarded ( $--max-missing$  1). No filtering was made based on minor allele frequencies. A total of 4150 SNPs was retained after filtering with a mean coverage per individual of 29×.

Individual MLH, that is, the number of heterozygous loci on the genotyped portion of the genome, was calculated using *vcftools* ( $--het$ ) (Danecek et al., 2011). The kinship coefficient, that is, the expected inbreeding coefficient for the progeny of one family, was estimated



using the *beta.dosage* function of the package *Hierfstat* (Goudet, 2005). This function generates a genomic matrix based on allele dosage to estimate kinship coefficients considering the proportion of alleles shared between individuals (Goudet et al., 2018). The inbreeding coefficients for females and males,  $F_{\text{beta}}$ , were obtained by extracting the diagonal of the genomic matrix obtained with *beta.dosage*. Figure S3 shows how MLH (range in males: 0.242–0.259) relates to  $F_{\text{beta}}$  (range in males: 0.029–0.092). Further analyses were focused on  $F_{\text{beta}}$ .

## 2.9 | Statistical analysis

Analyses were done in R (R Development Core Team, 2015) and in JMP® 15.2.1. Pearson correlation coefficients ( $r$ ) were used to test for correlations when visual inspection of the plots suggested that the conditions for parametric statistics were not significantly violated. Linear and logistic regression models were run on continuous and categorical variables respectively. In experiment 2, the likely effects of ovarian fluids were tested on one data set but for three different types of focal males within the dyads. Therefore, the


Holm–Bonferroni method was used to adjust the critical  $p$ -values when testing for effects of ovarian fluids.

## 3 | RESULTS

### 3.1 | Male phenotypes, $F_{\text{beta}}$ , leucocyte counts and milt characteristics

Colourful males were mostly yellow (Figure 1) and in few cases also slightly reddish. However, 19 of 34 males had negative  $a^*$  values and the mean ( $\pm 95\%$  CI)  $a^*$  value for the males was  $0.95 \pm 1.78$ . Because redness was also positively correlated with yellowness ( $r = .45$ ,  $p < .005$ ), it was ignored in further analyses.

Yellowness increased with male size in both years (Figure 1a; linear regression, effect of yellowness:  $F = 5.6$ ,  $p = .02$ , year:  $F = .3$ ,  $p = .59$ , yellowness  $\times$  year:  $F = .1$ ,  $p = .74$ ) but was not significantly correlated with  $F_{\text{beta}}$  (Figure 1b;  $r = .16$ ,  $p = .65$ ). While the correlation between leucocyte counts and skin coloration was not significant ( $r = -.47$ ,  $p = .17$ ), skin coloration turned out to be a good predictor



**FIGURE 1** Male yellowness versus size, inbreeding coefficient, immune parameters and characteristics of sperm when activated in water only. The two fish illustrate the two extremes of observed yellowness in 2017. Yellowness versus (a) body size (cm), (b) inbreeding coefficients ( $F_{\text{beta}}$ ), the immune parameters (c) relative lymphocytosis (% of lymphocytes among leucocytes) and (d) thrombocytosis (thrombocytes counts per 100 blood cells), and the sperm characteristics (e) velocity ( $\mu\text{m/s}$ ), (f) rate of immotile sperm (%) and (g) longevity (sec). Non-hatched and hatched regression lines illustrate significant and non-significant relationships, respectively, after potentially confounding effects of sampling year were taken into account. Symbols and regressions in light grey indicate samples collected in 2017, darker symbols and regression lines those from 2018. See text for statistics.

of relative lymphocytosis (Figure 1c;  $r = -.73$ ,  $p = .02$ ) and of thrombocytosis (Figure 1d;  $r = -.74$ ,  $p = .01$ ).  $F_{\text{beta}}$  was not corrected to any immune parameters (Figure S4; leucocyte count:  $r = .03$ , lymphocytosis:  $r = .18$ , thrombocytosis:  $r = -.12$ ,  $p$  always  $> .61$ ).


Yellowness did not correlate with sperm concentration (linear regression, effect of yellowness:  $F = .8$ ,  $p = .38$ , year:  $F = 1.0$ ,  $p = .32$ , yellowness  $\times$  year:  $F = .1$ ,  $p = .72$ ). When their milt was activated in water only, yellowness did also not significantly correlate with sperm velocity (Figure 1e; effect of yellowness:  $F = 3.8$ ,  $p = .06$ , year:  $F = 1.7$ ,  $p = .20$ , yellowness  $\times$  year:  $F = .1$ ,  $p = .74$ ), the rate of immotile sperm when controlling for significant year effects (Figure 1f; effect of yellowness:  $F = 2.1$ ,  $p = .16$ , year:  $F = 6.4$ ,  $p = .02$ , yellowness  $\times$  year:  $F = .3$ ,  $p = .61$ ), nor sperm longevity (Figure 1g;  $r = -.22$ ,  $p = .30$ ).

Mean male size differed between sampling years (Figure 1a;  $F = 4.9$ ,  $p = .03$ ). When controlling for this year effect, none of the sperm traits could be predicted by male size (linear regressions, effect of male size:

$F$  always  $< 1.8$ ,  $p$  always  $> .19$ ).  $F_{\text{beta}}$  was no significant predictor of sperm velocity (Figure S4;  $r = -.25$ ,  $p = .48$ ), but males with higher  $F_{\text{beta}}$  had higher counts of immotile sperm (Figure S4;  $r = .74$ ,  $p = .01$ ).

### 3.2 | Experiment 1: Sperm performance in ovarian fluid

Activating sperm with or without ovarian fluids did not significantly affect average sperm velocity, but sperm velocity differed between males, and sperm of different males reacted differently to the presence or absence of ovarian fluids (Figure 2a; linear regression, effect of activation medium:  $F = 1.2$ ,  $p = .28$ , effect of male identity:  $F = 10.5$ ,  $p < .001$ , activation medium  $\times$  male:  $F = 2.6$ ,  $p = .01$ ). Activating sperm with ovarian fluids generally reduced the rate of immotile sperm (Figure 2b; effect of activation medium:  $F = 5.6$ ,



**FIGURE 2** The effects of ovarian fluids on sperm of different types of males (experiment 1). Panels (a–c) show average sperm characteristics (Turkey box plots with quartiles and whiskers) for each of the 10 males after their sperm were activated in water only or in diluted ovarian fluids: (a) sperm velocity ( $\mu\text{m}/\text{sec}$ ), (b) rate of immotile sperm (%) and (c) sperm longevity (sec). Panels (d–f) give the mean ( $\pm$ SD) differences between of sperm characteristics in ovarian fluid and in water (8 measurements per male in ovarian fluids compared to the respective mean in water each) relative to male yellowness: (d) differences in sperm velocity, (e) differences in rates of immotile sperm, (f) differences in longevity. The regression line illustrates the significant correlation, the dotted line in (d) and (e) marks zero difference. The asterisks indicate a significant effect of the activation medium, either as main effect (b, c) or in interaction with male identity (a). See text for statistics.

$p=.02$ , male:  $F=21.7$ ,  $p<.001$ , activation medium $\times$ male:  $F=1.6$ ,  $p=.14$ ) and increased sperm longevity (Figure 2c; effect of activation medium:  $F=263.8$ ,  $p<.0001$ , male:  $F=1.7$ ,  $p=.11$ , activation medium $\times$ male:  $F=.9$ ,  $p=.55$ ).

Table 1 focuses on sperm traits after activation with ovarian fluids. Males significantly varied in all sperm traits that were monitored here (main effects of male identity in Table 1). Females significantly varied in how their ovarian fluids affected average sperm velocity and longevity but not the percentage of immotile sperm (see main effects of female identity in Table 1). Sperm of different males responded differently to the ovarian fluids of different females: average sperm velocity and longevity but not the percentage of immotile sperm depended on the combination of male and female identities (male $\times$ female interaction terms in Table 1; see also Figure S5).

The male-specific responses to ovarian fluids could partly be predicted by male skin coloration. With increasing yellowness, sperm

TABLE 1 Experiment 1: Sperm characteristics of different males after activation with diluted ovarian fluid of different females.

Factor	df	F	p
(A) Sperm velocity			
Male	9	7.1	<b>&lt;.001</b>
Female	3	6.2	<b>.002</b>
Male $\times$ female	27	1.9	<b>.03</b>
(B) Sperm motility			
Male	9	27.9	<b>&lt;.001</b>
Female	3	.6	.64
Male $\times$ female	27	.8	.74
(C) Longevity			
Male	9	446.5	<b>&lt;.001</b>
Female	3	1185.3	<b>&lt;.001</b>
Male $\times$ female	27	145.8	<b>&lt;.001</b>

Note: Significant  $p$ -values are highlighted in bold.

velocity increased more in the presence of ovarian fluids (Figure 2d; regression on means per male:  $r=.65$ ,  $n=10$ ,  $p=.04$ ). Male yellowness did, however, not predict the reduced rate of immotile sperm that ovarian fluids caused (Figure 2e;  $r=.42$ ,  $p=.23$ ), nor did it predict some of the increase in sperm longevity linked to the presence of ovarian fluids (Figure 2f;  $r=-.36$ ,  $p=.31$ ). None of the analogous correlations to male  $F_{\text{beta}}$  nor to the average kinship between males and females were significant ( $r$  always  $<.55$ ,  $n=10$ ,  $p$  always  $>.10$ ), that is, the effects that ovarian fluids had on sperm traits were not linked to male  $F_{\text{beta}}$  or the average kinship to the donors of ovarian fluids.

### 3.3 | Experiment 2: Males success in sperm competition

In total, 1475 offspring were genotyped of which 1464 could be assigned to one of the two competitors within the dyads (99.3% success). In the following, three types of focal males within the dyads are considered: the most yellow, the least inbred and the least related to the respective female. Male yellowness did not correlate with their  $F_{\text{beta}}$  (Figure 1b; see above) nor with the kinship with the females (Figure S6a;  $r=.13$ ,  $n=40$ ,  $p=.42$ ), and kinship could not be predicted by male  $F_{\text{beta}}$  (Figure S6b;  $r=.12$ ,  $n=40$ ,  $p=.47$ ).

Figure 3 shows the outcome of the individual sperm competition trials when run in water only or with ovarian fluids of the different females, plotted each for the three types of focal males within the dyads (see Table S4 for the identities of the focal males in the different scenarios). Overall, the average (SD) success rates of focal males were 0.534 (0.180), 0.562 (0.169) and 0.532 (0.179) if the focal male was the most yellow, the least inbred or the least related to the female respectively (Figure 3). Dyad identity always had a strong effect on the focal male's success rate (Table 2).

If the focal male was the yellower of the two competitors, its fertilization success was significantly enhanced in the presence of




FIGURE 3 Outcome of the sperm competition trials (experiment 2). The mean fertilization rates per dyad, female and whether or not ovarian fluids were present of (a) the yellower male of the two competitors each, (b) the less inbred male each and (c) the male each that is least related to the female whose ovarian fluids the sperm was exposed to. Turkey box plots with quartiles and whiskers. The asterisks indicate a significant difference linked to the activation medium, n.s., not significant. See Table 2 for statistics.



**TABLE 2** Experiment 2: Correlates to fertilization success of the focal male in the sperm competition trials, if the focal male is (a) the more yellow one of the two competitors, (b) the one with the lower inbreeding coefficient  $F_{\text{beta}}$  or (c) the least related to the female.

Factor	df	$\chi^2$	<i>p</i>
<b>(A) Focal male: more yellow</b>			
Female	3	6.3	.10
Dyad	4	156.4	<.001
Ovarian fluid	1	7.8	<b>.005</b>
Difference in sperm velocity <sup>a</sup>	2	7.8	.02
<b>(B) Focal male: lower <math>F_{\text{beta}}</math></b>			
Female	3	6.0	.11
Dyad	4	122.1	<.001
Ovarian fluid	1	5.8	<b>.02</b>
Difference in sperm velocity <sup>a</sup>	2	7.9	.02
<b>(C) Focal male: lower kinship</b>			
Female	3	20.4	<.001
Dyad	4	26.7	<.001
Ovarian fluid	1	1.8	.18
Difference in sperm velocity <sup>a</sup>	2	6.2	.04
Difference in sperm longevity <sup>a</sup>	2	39.0	<.001

Note: Logistic regressions with female (egg donor), dyad (pair of competing males), ovarian fluid (present or absent) and the difference in mean sperm velocity, rates of immotile sperm and sperm longevity between rivals (focus–opponent). These differences in sperm traits had been determined in experiment 1 both in water and in ovarian fluid and were therefore nested in the factor ‘ovarian fluid.’ The table shows the outcome after stepwise removal of non-significant effects of sperm traits. Significant effects of ovarian fluids (after Holm–Bonferroni correction for multiple comparison) are highlighted in bold.

<sup>a</sup>Nested in ‘ovarian fluid.’

ovarian fluid as compared to water only (Table 2a; Figure 3a) while female identity played no significant role in this scenario (Table 2a). If the focal male was the one with the lower  $F_{\text{beta}}$ , its fertilization success was higher when sperm were activated with water only instead of diluted ovarian fluids (Table 2b; Figure 3b). As before, female identity played no significant role (Table 2b). In both previous scenarios, the different effects that ovarian fluids had on sperm velocity contributed significantly to the outcome (Table 2a,b). No effects of ovarian fluids could be observed if the focal male was the least related to the egg donor (Table 2c; Figure 3c). In the latter scenario, male success was best explained by female identity and the difference in sperm velocity and longevity between the rivals (Table 2c).

## 4 | DISCUSSION

The mating systems of salmonid fish typically lead to intense male–male competition (Esteve, 2005; Fiske et al., 1998) that is likely to affect male reproductive strategies (Magris, 2021; Parker & Pizzari, 2010). Previous studies on male reproductive strategies have often focused on traits that may only partly determine fertilization success under

ecologically relevant conditions (Dougherty et al., 2022). In fish, for example, it is still largely unclear whether and to what degree sperm reaction to ovarian fluids may reveal male adaptations to the microecology that their sperm are likely exposed to during fertilization. Here, we asked whether male skin coloration or their inbreeding coefficient  $F_{\text{beta}}$  reveals factors that may matter in male–male competition, and whether sperm traits and especially the reaction of sperm to ovarian fluids can be predicted from male coloration or  $F_{\text{beta}}$ .

Secondary sexual traits like conspicuous skin colours are expected to reveal general health and vigour (Andersson, 1994), and they have repeatedly been observed to do so in other species (Johnson & Fuller, 2015; Milinski & Bakker, 1990). In our study, paler males suffered from increased relative lymphocytosis and thrombocytosis, two potential indicators of acute infections or other physiological stress (Haenen et al., 2010; Johansen et al., 2019; Rose et al., 2012; Yan et al., 2013), than yellower males. We therefore conclude that male yellowness is a good predictor of current male health condition. We also found a positive correlation between male yellowness and body size that often determines male dominance in fish (Jacob et al., 2007, 2009). However, neither male yellowness nor body size correlated significantly with any milt or sperm trait when sperm were activated with water only. Our study therefore provides no support of the phenotype-linked fertility hypothesis of mate selection (e.g. Mehlis et al., 2013; Monteiro et al., 2017).

Theory also predicts a correlation between an individual's inbreeding coefficient and general health and vigour (Allendorf & Luikard, 2007), and such links have indeed been found in other studies (Fox & Reed, 2011). We did not find any significant correlations between male  $F_{\text{beta}}$  and immune parameters, but this may need to be studied again in larger samples that provide more statistical power. There was also no significant correlation between male  $F_{\text{beta}}$  and milt or sperm traits when tested in water only, except that the rates of immotile sperm increased with increasing male  $F_{\text{beta}}$ . This latter observation supports the hypothesis that sperm motility can still be an indicator of certain aspects of health and vigour (Kowalski & Cejko, 2019).

$F_{\text{beta}}$  and skin colorations were not significantly correlated in our sample. In previous studies, such a correlation was sometimes found (e.g. Herdegen et al., 2014; Zajitschek & Brooks, 2010) and sometimes not (e.g. Frommen et al., 2008; Marsh et al., 2017), including a study on Arctic char (Janhunen et al., 2011). One possibility is that skin coloration is a plastic trait that quickly reacts to acute stress (Johnson & Fuller, 2015) while  $F_{\text{beta}}$  affects more the overall tolerance to environmental stress. If so, the link between  $F_{\text{beta}}$  and skin colorations is expected to change with changing environmental conditions.

Our first experiment was designed to test whether and how ovarian fluids affect sperm traits, and whether such effects differ for different types of males. We found that ovarian fluids reduced the percentage of immotile sperm and nearly doubled mean sperm longevity. There was no significant link between these supportive effects of ovarian fluids and male yellowness, their  $F_{\text{beta}}$ , nor to the kinship between males and females. With regard to these sperm traits, all types of males profited similarly from the presence of ovarian fluids.

Sperm velocity was not generally increased by ovarian fluids, apparently contrary to some previous findings in other fishes (Zadmajid et al., 2019). However, when comparing the effects of ovarian fluids on sperm characteristics of different types of males, it turned out that the yellower a male, the more does the velocity of its sperm increase when in contact with ovarian fluids. Interestingly, sperm of paler males were faster in water only than in diluted ovarian fluids. Such a link between the effects of ovarian fluids and male coloration suggests that sperm of yellower males are better prepared to ovarian fluids while sperm of paler males are better prepared to the absence of ovarian fluids. Our findings suggest that yellower males are typically closer to the female vent during spawning than paler males. This could be due to male–male dominance, female preference, or both.

Females varied in how much their ovarian fluids generally increased sperm velocity and sperm longevity, but not in how much their ovarian fluids reduced the rate of immotile sperm. Females also varied in how much their ovarian fluids specifically increased velocity and longevity of sperm of certain males over others, that is, we found the corresponding females  $\times$  male interactions to be significant (but again not in the case of sperm motility). However, effects of ovarian fluids on sperm traits were not significantly linked to male  $F_{\text{beta}}$  or the average kinship between males and females.

Our second experiment was to see whose sperm benefit most from exposure or non-exposure to ovarian fluids when in sperm competition, and whether the most successful males are best characterized by their skin coloration, their  $F_{\text{beta}}$  or their kinship to a given female. We found that the presence of ovarian fluids led to enhanced fertilization success of the more yellow of the two competitors. This result further supports the conclusion from the first experiment that sperm of more yellow males seem better adapted to the presence of ovarian fluid than sperm of pale males. If the focal male was the one with the lower  $F_{\text{beta}}$ , its fertilization success turned out to be higher when sperm were activated with water only than with diluted ovarian fluids. No significant effects of ovarian fluids were observed if the focal male was the least related to the female.

Competition trials typically reveal a limited range of possible situations. In our case, the overall sperm to egg ratios were so high that variation in sperm longevity may not have affected the outcome of the sperm competition, while late fertilizations may often be relevant under natural conditions. However, no significant correlations between sperm longevity and male yellowness,  $F_{\text{beta}}$  or kinship to the donor of the ovarian fluid was observed in the first experiment. We therefore conclude that the competitive advantage that sperm of more yellow males have over sperm of pale males depends on the presence of ovarian fluids.

In a parallel study, Garaud et al. (2023) used an extended sample of our study population to test whether females would profit from mating with more yellow, less inbred or less related males. They used full-factorial in vitro fertilizations (crossing 10 male and 6 females in all possible combinations) and single rearing of large

numbers of offspring to study embryo development under different environmental conditions. Contrary to expectations from 'good-genes' hypotheses of sexual selection (Andersson, 1994), offspring of yellower males were on average smaller and less tolerant to pathogen stress than offspring of paler males. These results confirm previous findings on Arctic char (Janhunen et al., 2011) and suggest that females would not profit from giving yellower males a selective advantage during sperm competition. Interestingly, Garaud et al. (2023) found significant effects of the genetic relatedness between males and females: Embryos grew smaller with increasing kinship of their parents.

In conclusion, male yellowness correlated positively with health and vigour and body size, that is, with traits that are likely to determine the outcome of male–male competition. Yellower males may therefore typically be closer to the females during egg release than pale males. If so, yellow males could expect their sperm to be more likely exposed to ovarian fluids than pale males. We conclude from two different types of experiments that sperm of yellow males profit more from the presence of ovarian fluid than sperm of pale males. No such beneficial effects of ovarian fluid could be seen with regard to male  $F_{\text{beta}}$  or the kinship between males and females, while parallel breeding experiments with the same study population concluded that females would profit most from supporting sperm of least related males. Combined, these results suggest that the increased performance of sperm of yellower males in ovarian fluids is an adaptation to the microenvironment that the sperm are likely to be exposed to, rather than cryptic female choice by ovarian fluids. Studies on sexual selection at the gamete level (e.g. testing for cryptic female choice) should therefore always distinguish between likely effects of male strategies and likely effects of female strategies.

## AUTHOR CONTRIBUTIONS

David Nusbaumer and Claus Wedekind designed the experiment. David Nusbaumer and Laura Garaud organized and conducted the fieldwork (with the help of Christian de Guttry) and determined sperm characteristics. David Nusbaumer performed the two experiments, determined male colours and genotyped the offspring for parental assignments. Christian de Guttry did the ddRADseq library preparation and genomic analysis. Laurie Ançay stained the blood smears and classified the blood cells. David Nusbaumer & Claus Wedekind did the statistics and wrote the manuscript that was revised by all authors.

## ACKNOWLEDGEMENTS

We thank G. Levray from the Pistolet fishery for catching the fish, the APERA association and the *pisciculture des Rives* for access to the fish, L. Adhia Eya, A. Atherton, C. Berney, J. Bideaux, F. Dolivo, L. Espinat, S. Kreuter, E. Longange, L. Marques da Cunha and N. Sironi for assistance in the field or in the lab, F. Schütz for discussion on statistics, E. Lasne for discussion and access to the INRA facilities, and handling editor M. M. Hansen and four reviewers for helpful comments. Open access funding provided by Universite de Lausanne.

## FUNDING INFORMATION

This project was supported by the Swiss National Science Foundation (31003A\_159579 and 31003A\_182265).

## DATA AVAILABILITY STATEMENT

The data used for this study have been deposited in the Dryad depository (doi: [10.5061/dryad.rn8pk0pct](https://doi.org/10.5061/dryad.rn8pk0pct) and [10.5061/dryad.gqnk98sp3](https://doi.org/10.5061/dryad.gqnk98sp3)) (de Guttry et al., 2022; Nusbaumer et al., 2022).

## CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

## ETHICS STATEMENT

This work complied with the national, cantonal and university regulations where it was carried out. The handling and transport of adults and transport of embryos was approved by the French authorities (INTRA.FR.2017.0109258).

## ORCID

Claus Wedekind  <https://orcid.org/0000-0001-6143-4716>

## REFERENCES

- Allendorf, F. W., & Luikard, G. (2007). *Conservation and the genetics of populations*. Oxford University Press.
- Alonzo, S. H., Stiver, K. A., & Marsh-Rollo, S. E. (2016). Ovarian fluid allows directional cryptic female choice despite external fertilization. *Nature Communications*, 7, 12452. <https://doi.org/10.1038/ncomm12452>
- Andersson, M. B. (1994). *Sexual selection*. Princeton University Press.
- Andrews, S. (2010). *FastQC: A quality control tool for high throughput sequence data*. Babraham Bioinformatics. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>
- Backström, T., Heynen, M., Brannas, E., Nilsson, J., & Magnhagen, C. (2015). Dominance and stress signalling of carotenoid pigmentation in Arctic charr (*Salvelinus alpinus*): Lateralization effects? *Physiology & Behavior*, 138, 52–57. <https://doi.org/10.1016/j.physbeh.2014.10.003>
- Bakker, T. C., Zbinden, M., Frommen, J. G., Weiss, A., & Largiadèr, C. R. (2006). Slow fertilization of stickleback eggs: The result of sexual conflict? *BMC Ecology*, 6, 7. <https://doi.org/10.1186/1472-6785-6-7>
- Bartlett, M. J., Steeves, T. E., Gemmill, N. J., & Rosengrave, P. C. (2017). Sperm competition risk drives rapid ejaculate adjustments mediated by seminal fluid. *eLife*, 6, e28811. <https://doi.org/10.7554/eLife.28811>
- Bernatchez, L., & Duchesne, P. (2000). Individual-based genotype analysis in studies of parentage and population assignment: How many loci, how many alleles? *Canadian Journal of Fisheries and Aquatic Sciences*, 57, 1–12.
- Breltsford, A., Dufresnes, C., & Perrin, N. (2016). High-density sex-specific linkage maps of a European tree frog (*Hyla arborea*) identify the sex chromosome without information on offspring sex. *Heredity*, 116, 177–181. <https://doi.org/10.1038/hdy.2015.83>
- Butts, I. A. E., Johnson, K., Wilson, C. C., & Pitcher, T. E. (2012). Ovarian fluid enhances sperm velocity based on relatedness in lake trout, *Salvelinus namaycush*. *Theriogenology*, 78, 2105–2109. <https://doi.org/10.1016/j.theriogenology.2012.06.031>
- Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: An analysis tool set for population genomics. *Molecular Ecology*, 22, 3124–3140. <https://doi.org/10.1111/mec.12354>
- Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000 Genomes Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*, 27, 2156–2158. <https://doi.org/10.1093/bioinformatics/btr330>
- de Guttry, C., Nusbaumer, D., Garaud, L., Ançay, L., & Wedekind, C. (2022). Inbreeding and kinship coefficients of parents of sperm competition experiment with lake char (Lake Geneva 2017/18). *Dryad Digital Repository*. <https://doi.org/10.5061/dryad.rn8pk0pct>
- Dougherty, L. R., Skirrow, M. J. A., Jennions, M. D., & Simmons, L. W. (2022). Male alternative reproductive tactics and sperm competition: A meta-analysis. *Biological Reviews of the Cambridge Philosophical Society*, 97, 1365–1388. <https://doi.org/10.1111/brv.12846>
- Egeland, T. B., Rudolfson, G., Nordeide, J. T., & Folstad, I. (2016). Status specific tailoring of sperm behavior in an external fertilizer. *Frontiers in Ecology and Evolution*, 4, 135–143. <https://doi.org/10.3389/fevo.2016.00135>
- Esteve, M. (2005). Observations of spawning behaviour in salmoninae: *Salmo*, *Oncorhynchus* and *Salvelinus*. *Reviews in Fish Biology and Fisheries*, 15, 1–21. <https://doi.org/10.1007/s11160-005-7434-7>
- Evans, J. P., Rosengrave, P., Gasparini, C., & Gemmill, N. J. (2013). Delineating the roles of males and females in sperm competition. *Proceedings of the Royal Society B-Biological Sciences*, 280, 20132047.
- Firman, R. C., Gasparini, C., Manier, M. K., & Pizzari, T. (2017). Postmating female control: 20 years of cryptic female choice. *Trends in Ecology & Evolution*, 32, 368–382. <https://doi.org/10.1016/j.tree.2017.02.010>
- Fiske, P., Rintamäki, P. T., & Karvonen, E. (1998). Mating success in lekking males: A meta-analysis. *Behavioral Ecology*, 9, 328–338. <https://doi.org/10.1093/beheco/9.4.328>
- Fox, C. W., & Reed, D. H. (2011). Inbreeding depression increases with environmental stress: An experimental study and meta-analysis. *Evolution*, 65, 246–258. <https://doi.org/10.1111/j.1558-5646.2010.01108.x>
- Frommen, J. G., Luz, C., Mazzi, D., & Bakker, T. C. M. (2008). Inbreeding depression affects fertilization success and survival but not breeding coloration in threespine sticklebacks. *Behaviour*, 145, 425–441. <https://doi.org/10.1163/156853908792451458>
- Fukuda, S., & Karino, K. (2014). Male red coloration, female mate preference, and sperm longevity in the cyprinid fish *Puntius titteya*. *Environmental Biology of Fishes*, 97, 1197–1205. <https://doi.org/10.1007/s10641-013-0207-6>
- Garaud, L., Nusbaumer, D., de Guttry, C., Marques da Cunha, L., Ançay, L., Atherton, A., Lasne, E., & Wedekind, C. (2023). Genetic compatibility exceeds possible good genes effects of sexual selection in lake char. *bioRxiv*. <https://doi.org/10.1101/2022.03.03.482834>
- Gasparini, C., & Pilastro, A. (2011). Cryptic female preference for genetically unrelated males is mediated by ovarian fluid in the guppy. *Proceedings of the Royal Society B-Biological Sciences*, 278, 2495–2501. <https://doi.org/10.1098/rspb.2010.2369>
- Gasparini, C., Pilastro, A., & Evans, J. P. (2020). The role of female reproductive fluid in sperm competition. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 375, 20200077. <https://doi.org/10.1098/rstb.2020.0077>
- Geßner, C., Johnson, S. L., Fisher, P., Clarke, S., Rutherford, K., Symonds, J., & Gemmill, N. J. (2017). Male-female relatedness at specific SNP-linkage groups influences cryptic female choice in Chinook salmon (*Oncorhynchus tshawytscha*). *Proceedings of the Royal Society B-Biological Sciences*, 284, 20170853. <https://doi.org/10.1098/rspb.2017.0853>
- Goudet, J. (2005). HIERFSTAT, a package for R to compute and test hierarchical *F*-statistics. *Molecular Ecology Notes*, 5, 184–186. <https://doi.org/10.1111/j.1471-8286.2004.00828.x>
- Goudet, J., Kay, T., & Weir, B. S. (2018). How to estimate kinship. *Molecular Ecology*, 27, 4121–4135. <https://doi.org/10.1111/mec.14833>
- Haenen, O. L. M., Lehmann, J., Engelsma, M. Y., Stürenberg, F. J., Roozenburg, I., Kerkhoff, S., & Breteler, J. K. (2010). The health

- status of European silver eels, *Anguilla Anguilla*, in the Dutch River Rhine watershed and Lake IJsselmeer. *Aquaculture*, 309, 15–24. <https://doi.org/10.1016/j.aquaculture.2010.08.026>
- Herdegen, M., Dudka, K., & Radwan, J. (2014). Heterozygosity and orange coloration are associated in the guppy (*Poecilia reticulata*). *Journal of Evolutionary Biology*, 27, 220–225. <https://doi.org/10.1111/jeb.12290>
- Jacob, A., Evanno, G., Renai, E., Sermier, R., & Wedekind, C. (2009). Male body size and breeding tubercles are both linked to intrasexual dominance and reproductive success in the minnow. *Animal Behaviour*, 77, 823–829. <https://doi.org/10.1016/j.anbehav.2008.12.006>
- Jacob, A., Nusslé, S., Britschgi, A., Evanno, G., Müller, R., & Wedekind, C. (2007). Male dominance linked to size and age, but not to 'good genes' in brown trout (*Salmo trutta*). *BMC Evolutionary Biology*, 7, 207. <https://doi.org/10.1186/1471-2148-7-207>
- Janhunen, M., Peuhkuri, N., Primmer, C. R., Kolari, I., & Piironen, J. (2011). Does breeding ornamentation signal genetic quality in Arctic charr, *Salvelinus alpinus*? *Evolutionary Biology*, 38, 68–78. <https://doi.org/10.1007/s11692-010-9100-9>
- Janhunen, M., Rudolfsen, G., Kekäläinen, J., Figenschou, L., Peuhkuri, N., & Kortet, R. (2009). Spawning coloration and sperm quality in a large lake population of Arctic charr (Salmonidae: *Salvelinus alpinus* L.). *Biological Journal of the Linnean Society*, 98, 794–802. <https://doi.org/10.1111/j.1095-8312.2009.01317.x>
- Jeannerat, E., Marti, E., Berney, C., Janett, F., Bollwein, H., Sieme, H., Burger, D., & Wedekind, C. (2018). Stallion semen quality depends on major histocompatibility complex matching to teaser mare. *Molecular Ecology*, 27, 1025–1035. <https://doi.org/10.1111/mec.14490>
- Johansen, I. B., Henriksen, E. H., Shaw, J. C., Mayer, I., Amundsen, P. A., & Øverli, Ø. (2019). Contrasting associations between breeding coloration and parasitism of male Arctic charr relate to parasite species and life cycle stage. *Scientific Reports*, 9, 1–9. <https://doi.org/10.1038/s41598-019-47083-x>
- Johnson, A. M., & Fuller, R. C. (2015). The meaning of melanin, carotenoid, and pterin pigments in the bluefin killifish, *Lucania goodei*. *Behavioral Ecology*, 26, 158–167. <https://doi.org/10.1093/beheco/aru164>
- Jones, A. G., Small, C. M., Paczolt, K. A., & Ratterman, N. L. (2010). A practical guide to methods of parentage analysis. *Molecular Ecology Resources*, 10, 6–30. <https://doi.org/10.1111/j.1755-0998.2009.02778.x>
- Kalinowski, S. T., Taper, M. L., & Marshall, T. C. (2007). Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology*, 16, 1099–1106. <https://doi.org/10.1111/j.1365-294X.2007.03089.x>
- Kekäläinen, J., Pirhonen, J., & Taskinen, J. (2014). Do highly ornamented and less parasitized males have high quality sperm?—An experimental test for parasite-induced reproductive trade-offs in European minnow (*Phoxinus phoxinus*). *Ecology and Evolution*, 4, 4237–4246. <https://doi.org/10.1002/ece3.1267>
- Kortet, R., Vainikka, A., Rantala, M. J., & Taskinen, J. (2004). Sperm quality, secondary sexual characters and parasitism in roach (*Rutilus rutilus* L.). *Biological Journal of the Linnean Society*, 81, 111–117.
- Kottelat, M., & Freyhof, J. (2007). *Handbook of European freshwater fishes*. Kottelat and Freyhof.
- Kowalski, R. K., & Cejko, B. I. (2019). Sperm quality in fish: Determinants and affecting factors. *Theriogenology*, 135, 94–108. <https://doi.org/10.1016/j.theriogenology.2019.06.009>
- Lahnsteiner, F., Weismann, T., & Patzner, R. A. (1995). Composition of the ovarian fluid in 4 salmonid species: *Oncorhynchus mykiss*, *Salmo trutta f lacustris*, *Salvelinus alpinus* and *Hucho hucho*. *Reproduction Nutrition Development*, 35, 465–474.
- Lehnert, S. J., Butts, I. A. E., Flannery, E. W., Peters, K. M., Heath, D. D., & Pitcher, T. E. (2017). Effects of ovarian fluid and genetic differences on sperm performance and fertilization success of alternative reproductive tactics in Chinook salmon. *Journal of Evolutionary Biology*, 30, 1236–1245. <https://doi.org/10.1111/jeb.13088>
- Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with burrows-wheeler transform. *Bioinformatics*, 26, 589–595. <https://doi.org/10.1093/bioinformatics/btp698>
- Magris, M. (2021). Strategic adjustment of ejaculate quality in response to variation of the socio-sexual environment. *Behavioral Ecology and Sociobiology*, 75. <https://doi.org/10.1007/s00265-021-03032-1>
- Makiguchi, Y., Torao, M., Kojima, T., & Pitcher, T. E. (2016). Reproductive investment patterns and comparison of sperm quality in the presence and absence of ovarian fluid in alternative reproductive tactics of masu salmon, *Oncorhynchus masou*. *Theriogenology*, 86, 2189–2193. <https://doi.org/10.1016/j.theriogenology.2016.07.009>
- Marsh, J. N., Vega-Trejo, R., Jennions, M. D., & Head, M. L. (2017). Why does inbreeding reduce male paternity? Effects on sexually selected traits. *Evolution*, 71, 2728–2737. <https://doi.org/10.1111/evo.13339>
- Mehlis, M., & Bakker, T. C. M. (2013). Male reproductive traits of full-sibs of different age classes in three-spined sticklebacks (*Gasterosteus aculeatus*). *Springerplus*, 2. <https://doi.org/10.1186/2193-1801-2-175>
- Mehlis, M., Hilke, L. K., & Bakker, T. C. M. (2013). Attractive males have faster sperm in three-spined sticklebacks *Gasterosteus aculeatus*. *Current Zoology*, 59, 761–768. <https://doi.org/10.1093/czoolo/59.6.761>
- Meuthen, D., Meuthen, I., Bakker, T. C. M., & Thunken, T. (2020). Anticipatory plastic response of the cellular immune system in the face of future injury: Chronic high perceived predation risk induces lymphocytosis in a cichlid fish. *Oecologia*, 194, 597–607. <https://doi.org/10.1007/s00442-020-04781-y>
- Milinski, M., & Bakker, T. C. M. (1990). Female sticklebacks use male coloration in mate choice and hence avoid parasitized males. *Nature*, 344, 330–333.
- Monteiro, N. M., Carneiro, D., Antunes, A., Queiroz, N., Vieira, M. N., & Jones, A. G. (2017). The lek mating system of the worm pipefish (*Nerophis lumbriciformis*): A molecular maternity analysis and test of the phenotype-linked fertility hypothesis. *Molecular Ecology*, 26, 1371–1385. <https://doi.org/10.1111/mec.13931>
- Nusbaumer, D., Ançay, L., Garaud, L., de Guttry, C., & Wedekind, C. (2022). Sperm competition experiment with lake char (Lake Geneva 2017/18). *Dryad Digital Repository*. <https://doi.org/10.5061/dryad.gqnk98sp3>
- Nusbaumer, D., Garaud, L., Ançay, L., & Wedekind, C. (2021). Sex-specific stress tolerance in embryos of lake char (*Salvelinus umbla*). *Frontiers in Ecology and Evolution*, 9, 768263. <https://doi.org/10.3389/fevo.2021.768263>
- OECD. (1992). *OECD guideline for the testing of chemicals 203 (fish, acute toxicity test)*. OECD Publishing.
- Oluah, N. S., Aguzie, I. O., Ekechukwu, N. E., Madu, J. C., Ngene, C. I., & Oluah, C. (2020). Hematological and immunological responses in the African catfish *Clarias gairepinus* exposed to sub-lethal concentrations of herbicide Ronstar (R). *Ecotoxicology and Environmental Safety*, 201, 110824. <https://doi.org/10.1016/j.ecoenv.2020.110824>
- Parchman, T. L., Gompert, Z., Mudge, J., Schilkey, F. D., Benkman, C. W., & Buerkle, C. A. (2012). Genome-wide association genetics of an adaptive trait in lodgepole pine. *Molecular Ecology*, 21, 2991–3005. <https://doi.org/10.1111/j.1365-294X.2012.05513.x>
- Parker, G. A., & Pizzari, T. (2010). Sperm competition and ejaculate economics. *Biological Reviews*, 85, 897–934. <https://doi.org/10.1111/j.1469-185X.2010.00140.x>
- Parolini, M., Iacobuzio, R., Possenti, C. D., Bassano, B., Pennati, R., & Saino, N. (2018). Carotenoid-based skin coloration signals antioxidant defenses in the brown trout (*Salmo trutta*). *Hydrobiologia*, 815, 267–280. <https://doi.org/10.1007/s10750-018-3571-6>
- Perroud, G., Nusbaumer, D., de Guttry, C., & Wedekind, C. (2021). Male sexual signaling and expected effects of hatchery-induced sperm



- competition vary with water depth at which whitefish are caught. *Current Zoology*, 67, 337–340. <https://doi.org/10.1093/cz/zoab007>
- R Development Core Team. (2015). R: A language and environment for statistical computing. R Foundation for Statistical Computing.
- Rose, S. R., Petersen, N. J., Gardner, T. J., Hamill, R. J., & Trautner, B. W. (2012). Etiology of thrombocytosis in a general medicine population: Analysis of 801 cases with emphasis on infectious causes. *Journal of Clinical Medicine Research*, 4, 415–423. <https://doi.org/10.4021/jocmr1125w>
- Rosengrave, P., Montgomerie, R., & Gemmell, N. (2016). Cryptic female choice enhances fertilization success and embryo survival in Chinook salmon. *Proceedings of the Royal Society B-Biological Sciences*, 283, 20160001. <https://doi.org/10.1098/rspb.2016.0001>
- Rudolfson, G., Figenschou, L., Folstad, I., Tveiten, H., & Figenschou, M. (2006). Rapid adjustments of sperm characteristics in relation to social status. *Proceedings of the Royal Society B-Biological Sciences*, 273, 325–332. <https://doi.org/10.1098/rspb.2005.3305>
- Rudolfson, G., Müller, R., Urbach, D., & Wedekind, C. (2008). Predicting the mating system from phenotypic correlations between life-history and sperm quality traits in the alpine whitefish *Coregonus zugensis*. *Behavioral Ecology and Sociobiology*, 62, 561–567. <https://doi.org/10.1007/s00265-007-0480-5>
- Savary, R., Dufresnes, C., Champigneulle, A., Caudron, A., Dubey, S., Perrin, N., & Fumagalli, L. (2017). Stocking activities for the Arctic charr in Lake Geneva: Genetic effects in space and time. *Ecology and Evolution*, 7, 5201–5211. <https://doi.org/10.1002/ece3.3073>
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, 9, 676–682. <https://doi.org/10.1038/nmeth.2019>
- Seibel, H., Bassmann, B., & Rebl, A. (2021). Blood will tell: What hematological analyses can reveal about fish welfare. *Frontiers in Veterinary Science*, 8, 616955. <https://doi.org/10.3389/fvets.2021.616955>
- Selz, O. M., & Seehausen, O. (2023). A taxonomic revision of ten whitefish species from the lakes Lucerne, Sarnen, Sempach and Zug, Switzerland, with descriptions of seven new species (Teleostei, Coregonidae). *Zookeys*, 1144, 95–169. <https://doi.org/10.3897/zookeys.1144.67747>
- Skarstein, F., & Folstad, I. (1996). Sexual dichromatism and the immunocompetence handicap: An observational approach using Arctic charr. *Oikos*, 76, 359–367.
- Turner, E., & Montgomerie, R. (2002). Ovarian fluid enhances sperm movement in Arctic charr. *Journal of Fish Biology*, 60, 1570–1579.
- Urbach, D., Folstad, I., & Rudolfson, G. (2005). Effects of ovarian fluid on sperm velocity in Arctic charr (*Salvelinus alpinus*). *Behavioral Ecology and Sociobiology*, 57, 438–444.
- Wang, J. (2004). Sibship reconstruction from genetic data with typing errors. *Genetics*, 166, 1963–1979. <https://doi.org/10.1534/genetics.166.4.1963>
- Wilkins, L. G. E., Rogivue, A., Fumagalli, L., & Wedekind, C. (2015). Declining diversity of egg-associated bacteria during development of naturally spawned whitefish embryos (*Coregonus* spp.). *Aquatic Sciences*, 77, 481–497. <https://doi.org/10.1007/s00027-015-0392-9>
- Yan, S. L., Russell, J., Harris, N. R., Senchenkova, E. Y., Yildirim, A., & Granger, D. N. (2013). Platelet abnormalities during colonic inflammation. *Inflammatory Bowel Disease*, 19, 1245–1253. <https://doi.org/10.1097/MIB.0b013e318281f3df>
- Yano, A., Nicol, B., Jouanno, E., Quillet, E., Fostier, A., Guyomard, R., & Guiguen, Y. (2013). The sexually dimorphic on the Y-chromosome gene (sdY) is a conserved male-specific Y-chromosome sequence in many salmonids. *Evolutionary Applications*, 6, 486–496. <https://doi.org/10.1111/eva.12032>
- Yeates, S. E., Diamond, S. E., Einum, S., Emerson, B. C., Holt, W. V., & Gage, M. J. G. (2013). Cryptic choice of conspecific sperm controlled by the impact of ovarian fluid on sperm swimming behavior. *Evolution*, 67, 3523–3536. <https://doi.org/10.1111/evo.12208>
- Young, B., Conti, D. V., & Dean, M. D. (2013). Sneaker “jack” males out-compete dominant “hooknose” males under sperm competition in Chinook salmon (*Oncorhynchus tshawytscha*). *Ecology and Evolution*, 3, 4987–4997. <https://doi.org/10.1002/ece3.869>
- Zadmajid, V., Myers, J. N., Sorensen, S. R., & Ernest Butts, I. A. (2019). Ovarian fluid and its impacts on spermatozoa performance in fish: A review. *Theriogenology*, 132, 144–152. <https://doi.org/10.1016/j.theriogenology.2019.03.021>
- Zajitschek, S. R. K., & Brooks, R. C. (2010). Inbreeding depression in male traits and preference for outbred males in *Poecilia reticulata*. *Behavioral Ecology*, 21, 884–891. <https://doi.org/10.1093/beheco/arq077>

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Nusbaumer, D., Garaud, L., de Guttry, C., Ançay, L., & Wedekind, C. (2023). Sperm of more colourful males are better adapted to ovarian fluids in lake char (Salmonidae). *Molecular Ecology*, 00, 1–13. <https://doi.org/10.1111/mec.17103>