

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Redirecting NK cells mediated tumor cell lysis by a new recombinant bifunctional protein.

Authors: Germain C, Campigna E, Salhi I, Morisseau S, Navarro-Teulon I, Mach JP, Pèlegri A, Robert B

Journal: Protein engineering, design & selection : PEDS

Year: 2008 Nov

Volume: 21

Issue: 11

Pages: 665-72

DOI: [10.1093/protein/gzn047](https://doi.org/10.1093/protein/gzn047)

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.



Protein Eng Des Sel. Author manuscript; available in PMC 2009 Oct 9.

PMCID: PMC2759461

Published in final edited form as:

HALMS: HALMS325786

[Protein Eng Des Sel. 2008 Nov; 21\(11\): 665–672.](#)

Published online 2008 Sep 11. doi: [10.1093/protein/gzn047](https://doi.org/10.1093/protein/gzn047)

Inserm subrepository

Redirecting NK cells mediated tumor cell lysis by a new recombinant bifunctional protein

[Claire Germain](#),¹ [Emmanuelle Campigna](#),¹ [Imed Salhi](#),¹ [Sébastien Morisseau](#),¹ [Isabelle Navarro-Teulon](#),¹ [Jean-Pierre Mach](#),² [André Pèlerin](#),¹ and [Bruno Robert](#)^{1,*}

¹IRCM, Institut de recherche en cancérologie de Montpellier INSERM : U896, Université Montpellier I, FR

²Department of Biochemistry Université de Lausanne, BIL Biomedical Research Center University of Lausanne CH-1066 Epalinges, CH

* Correspondence should be addressed to: Bruno Robert ; Email: bruno.robert@valdorel.fnclcc.fr

[Copyright notice](#) and [Disclaimer](#)

Abstract

Natural killer (NK) cells are at the crossroad between innate and adaptive immunity and play a major role in cancer immunosurveillance. NK cell stimulation depends on a balance between inhibitory and activating receptors, such as the stimulatory lectinlike receptor NKG2D. To redirect NK cells against tumor cells we designed bifunctional proteins able to specifically bind tumor cells and to induce their lysis by NK cells, after NKG2D engagement. To this aim, we used the “knob into hole” heterodimerization strategy, in which “knob” and “hole” variants were generated by directed mutagenesis within the CH3 domain of human IgG1 Fc fragments fused to an anti-CEA or anti-HER2 scFv or to the H60 murine ligand of NKG2D, respectively. We demonstrated the capacity of the bifunctional proteins produced to specifically coat tumor cells surface with H60 ligand. Most importantly, we demonstrated that these bifunctional proteins were able to induce an NKG2D-dependent and antibody-specific tumor cell lysis by murine NK cells. Overall, the results show the possibility to redirect NK cytotoxicity to tumor cells by a new format of recombinant bispecific antibody, opening the way of potential NK cell-based cancer immunotherapies by specific activation of the NKG2D receptor at the tumor site.

Keywords: Animals, Blotting, Western, Cell Death, immunology, Cell Line, Tumor, Cytotoxicity, Immunologic, Electrophoresis, Gel, Two-Dimensional, Humans, Intercellular Signaling Peptides and Proteins, immunology, Killer Cells, Lymphokine-Activated, immunology, Killer Cells, Natural, immunology, metabolism, Mice, Minor Histocompatibility Antigens, immunology, Recombinant Proteins, immunology, Tumor Cells, Cultured, metabolism

Keywords: Bifunctional proteins, « Knob into hole » strategy, Natural Killer cells, NKG2D, Tumor targeting

Introduction

Natural Killer (NK) cells are cytotoxic cells, belonging to innate immunity, which play a major role in host response against different pathogens, such as viruses ([Bukowski et al., 1985](#)), bacteria or parasites ([Tay et al., 1998](#)). It is also now well established that NK cells are involved in cancer immunosurveillance ([Kim et al., 2000](#); [Smyth et al., 2002](#)). Despite no expression of antigen-specific receptors, such as T- or B-cell receptors (TcR and BcR), NK cells are able to recognize infected or malignant cells by several inhibitory or activating receptors, which act in balance either to induce NK cell tolerance towards normal cells or to promote NK cell activation and lysis of infected or tumor cells ([Raulet and Vance, 2006](#)). While inhibitory receptors, such as the killer immunoglobulin-like inhibitory receptors (KIR), are able to recognize the classical major histocompatibility class I (MHC I) proteins, whose expression is often lost or downregulated on infected ([Lodoen and Lanier, 2005](#)) or malignant cells ([Garrido et al., 1997](#)), activating receptors, such as the natural cytotoxicity receptors (NCR) NKp30, NKp44 and NKp46 or the natural killer group 2 D

(NKG2D) receptor, are able to recognize different ligands over-expressed on transformed cells ([Moretta et al., 2001](#)).

The activating NKG2D receptor has been particularly studied, both in humans and in mice. It is homogeneously expressed on all human and mouse NK cells ([Bauer et al., 1999](#); [Jamieson et al., 2002](#)), but also on all human and mouse activated CD8⁺ T cells, where it acts as a TcR co-activator ([Groh et al., 2001](#); [Jamieson et al., 2002](#)). Several NKG2D ligands are now well described. The human NKG2D ligands comprise the MHC class I Chain-related proteins A and B (MICA and MICB) ([Bahram et al., 1994](#)) and the retinoic acid early transcripts (RAET1E, G, H, I, L, N) (IMGT Repertoire (RPI); <http://imgt.cines.fr>) ([Cosman et al., 2001](#); [Jan Chalupny et al., 2003](#)). In mice, NKG2D ligands include the minor histocompatibility antigen H60 ([Malarkannan et al., 1998](#)), the retinoic acid early transcript 1 molecules α , β , γ , δ and ϵ (Rae1 α - ϵ) ([Cerwenka et al., 2000](#)) and the murine UL16-binding protein-like transcript 1 molecule (Mult1) ([Carayannopoulos et al., 2002](#)). All these ligands are stress-induced proteins and have been identified on various tumor cell types, such as human epithelial carcinomas, melanomas, gliomas and leukemias ([Pende et al., 2002](#); [Vetter et al., 2002](#); [Friese et al., 2003](#); [Salih et al., 2003](#)) or mouse epithelial carcinomas and lymphomas ([Cerwenka et al., 2000](#); [Diefenbach et al., 2000](#); [Carayannopoulos et al., 2002](#)). However, during tumor progression, malignant cells lose frequently NKG2D ligand expression, as a consequence of tumor variant selection ([Vetter et al., 2004](#); [Carbone et al., 2005](#)), contact inhibition ([Zou et al., 2006](#)) or proteolytic shedding ([Salih et al., 2002](#); [Salih et al., 2003](#)), and thus escape to NK cell recognition.

Several studies have established that transfection of human tumor cells with genes encoding NKG2D ligands sensitizes these cells to NK cell-mediated lysis *in vitro*, in an NKG2D-dependent pathway ([Friese et al., 2003](#); [Busche et al., 2006](#)). It has also been demonstrated *in vivo* that transfected murine tumor cells expressing Rae1 or H60 are rejected by NK cells, while parental tumor cells critically grow, and that rejection is associated with T cell response priming ([Diefenbach et al., 2001](#)).

To bypass the need of a gene therapy, we designed here bifunctional proteins able to bind tumor cells through specific antibody structures and to coat tumor cell surface with NKG2D ligands. These new bifunctional proteins were produced according to the “knob into hole” strategy, which allows the preferential heterodimerization of two heavy chains of distinct origin by changing two amino acids in their CH3 domains, and thus the production of recombinant bispecific antibodies ([Ridgway et al., 1996](#); [Merchant et al., 1998](#)). In the present study, we assembled through this strategy two different anti-tumor associated antigen (TAA) scFv with the mouse NKG2D ligand H60. After showing the specificity of these new bifunctional proteins for two targeted TAAs, we demonstrated their capacity to stimulate NKG2D dependent tumor cell lysis by murine NK cells.

Materials and methods

Tumor cell lines

Human acute monocytic leukemia THP-1 [expressing Fc receptors (FcR) Fc γ RI and Fc γ RII] and human breast carcinoma SK-BR-3 (expressing HER2) were obtained from the American Type Culture Collection (Rockville, MD) and maintained as described by the manufacturer. The murine colon carcinoma MC38 expressing CEA (C15.4.3.AP) cell line (kindly provided by Dr. J. Primus, Department of Pathology, Vanderbilt University Medical Center, Nashville, TN ([Clarke et al., 1998](#))), the murine melanoma B16 expressing HER2 cell line (kindly provided by Dr A. Donda, Department of Biochemistry, Faculty of Biology and Medicine, University of Lausanne, Switzerland ([Stirnemann et al., 2008](#))) and the murine T lymphoma RMA expressing CEA cell line, obtained by electro-transfection (250 V, 960 μ F, 5 μ g of DNA) of RMA cells with plasmid containing human CEA cDNA ([Pelegri et al., 1992](#)), were maintained in DMEM/F12 (1:1) supplemented with 2 mmol/L Lglutamine, 10% fetal calf serum (FCS), penicillin (100 units/ml)/streptomycin (100 μ g/ml), and geneticin (0.5 mg/ml; Life Technologies, Grand Island, NY).

Vectors construction

The sequences of the different primers used for the construction of the bifunctional proteins are given in Table 1. First, the genes encoding anti-human CEA scFv 511 (in pFLAG-1, kindly provided by Dr A.V. Terskikh, The Burnham Institute of Medical Research, La Jolla, CA, derived from the mouse anti-human

CEA mAb 511 hybridoma) and anti-human HER2 scFv 4D5 (in Bluescript® II KS +/-, kindly provided by Dr Alena Donda, Department of Biochemistry, Faculty of Biology and Medicine, University of Lausanne, Switzerland, derived from the humanized anti-human HER2 mAb 4D5 ([Deyev et al., 2003](#))) were amplified by PCR using primers P1/P2 and P3/P4. The amplified fragments were next digested and inserted in frame in plasmid ps878 (kindly provided by Dr P. Schneider, Department of Biochemistry, Faculty of Biology and Medicine, University of Lausanne, Switzerland ([Schneider, 2000](#))) encoding human IGHG1 Fc fragment (hinge, CH2 and CH3; GenBank/EMBL/IMG/CLUSTAL-DB accession No. [X70421](#)), with the signal sequence of the influenza virus hemagglutinin (HA) to allow protein secretion by mammalian cells. Knob variants of each construction were obtained by site-directed mutagenesis (PROMEGA Kit) in the IGHG1 CH3 sequence using primers P5/P6 ([Xie et al., 2005](#)). Plasmid ps521-rH60-Fc (kindly provided by Dr W. Held, Ludwig Institute for Cancer Research, Lausanne branch, Switzerland ([Coudert et al., 2005](#))) contains the gene encoding the extracellular portion including the endogenous leader peptide of H60 (amino acids 1–213), amplified by polymerase chain reaction (PCR) from A20 B-cell cDNA, using the following primers (restriction sites are underlined): H60 5' sense, actg AAGCTT tgaggaagacc ATG GCAAAG G; H60 3' antisense, actg GTCGAC CTG GTT GTC AGA ATT ATG TCG GAA G. The PCR product was directionally cloned into modified PCR-3 vectors (Invitrogen, San Diego, CA) to add a COOH-terminal FLAG epitope or to generate an Fc fusion protein (kindly provided by P. Schneider, Department of Biochemistry, University of Lausanne, Switzerland). A hole variant of this construction was obtained by site-directed mutagenesis using primers P7/P8 ([Xie et al., 2005](#)). Site-directed mutagenesis efficiency was verified by sequencing.

Bifunctional proteins expression and purification

HEK 293 cells (ATCC CRL-1573) were grown in DMEM/F12 (1:1) supplemented with 2 mmol/L L-glutamine, 10% FCS and penicillin (100 units/ml)/streptomycin (100 µg/ml) and transfected with each mutant vector using FuGene 6 reagent (Roche Applied Science, Indianapolis, IN), as recommended by the manufacturer. Clones were selected by limit dilution with medium containing geneticin (0.8 mg/ml). Culture supernatant of each clone was assayed for bifunctional proteins production by western blotting, as below. As control, HEK 293 cells were also transfected with the original ps521-rH60-Fc vector encoding rH60-Fc, or with the original ps878 vector, encoding (scFv 511)-Fc or (scFv 4D5)-Fc, or untransfected. For production of homodimeric and bifunctional proteins, selected clones were cultured for 12 days in medium without geneticin and FCS. Supernatant of these production batches were purified by affinity chromatography using a protein G column (Amersham Pharmacia Biotech AB, Uppsala, Sweden) according to the manufacturer's instructions, and by gel filtration on a 16/60 Superdex 200 L column (Amersham Pharmacia Biotech AB). Average yield of each of the recombinant proteins produced were: 3, 5.5 and 1.5 µg/ml for H60-Fc, 4D5-Fc and 511-Fc respectively and 2 and 4 µg/ml for (scFv 511/H60)-Fc and (scFv 4D5/H60)-Fc respectively. A schematic representation of the different homodimeric and bifunctional proteins produced is proposed in [Figure 1](#).

SDS-Page and Western Blot analysis of the produced bifunctional proteins

Purified bifunctional proteins, namely (scFv 511/rH60)-Fc and (scFv 4D5/rH60)-Fc, were analysed on a 6% SDS-PAGE gel, in comparison with homodimeric control proteins, rH60-Fc and (scFv 511)-Fc or (scFv 4D5)-Fc, under non-reducing and reducing conditions. For western blot analysis, reduced proteins were transferred to a nitrocellulose membrane. The membrane was saturated for two hours in PBS-0.1% Tween 20 containing 5 % nonfat dried milk at room temperature under agitation, incubated after washing with a rat anti-H60 mAb (R&D Systems Inc., Minneapolis, MN) and then with an HRP-conjugated goat anti-rat IgG whole molecule mAb (Sigma Aldrich, St. Louis, MO), or directly incubated with an HRP-conjugated goat anti-human IgG Fc specific mAb (Sigma-Aldrich). Finally, membrane was revealed after washing using the ECL Western Blotting System (Amersham Pharmacia Biotech AB).

Flow cytometry analyses

TAA and H60 expression on tumor cell lines was assessed by flow cytometry with mouse anti-CEA 511 mAb, humanized anti-human HER2 trastuzumab (Herceptin®, Genentech, Inc., San Francisco, CA) and a rat anti-H60 mAb, revealed by a FITC-conjugated goat anti-mouse IgG Fc specific mAb (Sigma-Aldrich), a FITC-conjugated goat anti-human IgG Fc specific mAb (Sigma-Aldrich) or a FITC-conjugated mouse anti-

rat IgG2a Fc specific mAb (BD Pharmingen, San Diego, CA), respectively. NKG2D expression on murine NK cells was assessed with a rat anti-mouse NKG2D mAb (R&D Systems, Minneapolis, MN), revealed by a FITC-conjugated mouse anti-rat IgG whole molecule mAb (Sigma-Aldrich). Specificity of the homodimeric proteins (scFv 511)-Fc and (scFv 4D5)-Fc for CEA and HER2, respectively, was assessed as follows: tumor cells (5×10^5) were incubated with each of the two proteins for 1 h at 4°C, washed, and then incubated with a FITC-conjugated goat anti-human IgG Fc specific mAb for 45 minutes at 4°C. Washed cells were finally analyzed on a FACScan (Becton Dickinson, San Jose, CA). Capacity of the bifunctional proteins (scFv 511/rH60)-Fc and (scFv 4D5/rH60)-Fc to coat tumor cells with H60 was assessed as follows: tumor cells were first incubated with each of the two proteins for 1 h at 4°C, washed, then incubated with a rat anti-H60 mAb for 1 h at 4°C, washed, and finally detected with a FITC-conjugated mouse anti-rat IgG2a Fc specific mAb. As controls, cells were first incubated with the parental homodimeric proteins, i.e. rH60-Fc and (scFv 511)-Fc or (scFv 4D5)-Fc, washed, then incubated with a rat anti-H60 mAb for 1 h at 4°C, washed, and finally incubated with a FITC-conjugated mouse anti-rat IgG2a Fc specific mAb for 45 minutes at 4°C, or the cells were incubated with the different secondary antibodies. Washed cells were similarly analyzed on a FACScan.

Murine Natural Killer cells preparation

Murine NK cells used as effectors in chromium release assays were prepared from a C57BL/6 spleen. Briefly, spleen was dilacerated, aggregates were eliminated on a 30 µm filter (Miltenyi Biotec, Auburn, CA), and splenic cells were cultured for 5 days in DMEM/F12 (1/1) supplemented with 2 mmol/L L-glutamine, 10% FCS, penicillin (100 units/ml)/streptomycin (100 µg/ml), non-essential amino acids, 1 µmol/L sodium pyruvate (Life Technologies), 50 µmol/L β-mercaptoethanol, and recombinant human interleukin-2 (5000 units/ml) (Proleukin, Chiron, Emeryville, CA). After 5 days, adherent cells, corresponding to lymphokine activated killer (LAK) cells were recovered and used as effectors in chromium release assays.

Chromium-release assays

Natural Killer cell-mediated cytotoxicity was measured by the chromium release assay. Tumor target cells (2×10^5 in 100 µl) were co-incubated for 1 h at 37°C with, at the same time, 100 µCi of ^{51}Cr and 5 µg, or less if specified, of the studied bifunctional protein or antibody. After washing, the labeled tumor target cells were plated in a 96-well plate (2000 cells per well) and incubated with LAK cells at different effector-to-target cell ratios (E:T ratios). For the comparison of trastuzumab antibody with our anti-HER2 bifunctional protein an E:T ratio of 30:1 was used. After incubation for 4 h at 37°C, 100 µl of supernatant was taken to measure the amount of ^{51}Cr released, reflecting the number of target cells killed by LAK cells. Spontaneous release of ^{51}Cr was determined by incubating the target cells with medium alone. Maximum release was determined by adding 0.1 M HCl. The percentage of specific lysis was calculated as: $100 \times [(\text{experimental} - \text{spontaneous release}) / (\text{total} - \text{spontaneous release})]$ (Ogg *et al.*, 2000). In control experiments, ^{51}Cr -labeled tumor target cells were preincubated without any bifunctional protein, with the irrelevant bifunctional protein, or with the parental homodimeric protein scFv-Fc, and tested with LAK cells preincubated or not for 30 minutes at room temperature with a rat anti-mouse NKG2D mAb used at 20 µg/ml. In the experiment comparing trastuzumab antibody and anti-HER2 bifunctional protein, the spontaneous lysis of SK-BR-3 cells alone was subtracted for each value. In a reverse antibody-dependent cell-mediated cytotoxicity lysis assay, THP-1 cells were incubated with 100 µCi of ^{51}Cr and rH60-Fc, after or without preincubation with an FcR blocking reagent which is composed of intact rat IgG2 antibodies (Miltenyi Biotec), to prevent rH60-Fc binding to the Fc receptors expressed on THP-1 cells. They were then incubated with LAK cells.

Results

Production of bifunctional proteins

Supernatants of several G418-resistant HEK 293 clones, expressing the soluble homodimeric and bifunctional proteins designed (Figure 1), were analysed by western blot to identify the ones producing the attempted proteins in terms of molecular weight (MW). Production supernatant batches for five different selected clones were purified on protein G columns, followed by a gel filtration chromatography.

When loaded on a SDS-PAGE gel under non-reducing conditions, purified (scFv 4D5/rH60)-Fc bifunctional protein appeared as a single band of approximately 160 kDa, i. e. at an intermediate molecular weight between the homodimeric rH60-Fc and (scFv 4D5)-Fc proteins, with MW of approximately 180 kDa and 120 kDa, respectively (Figure 2A, left part of the gel). Under reducing conditions, bifunctional protein appeared as two bands of 90 kDa and 60 kDa, corresponding to the respective MW of the reduced parental homodimeric proteins, rH60-Fc and (scFv 4D5)-Fc (Figure 2A, right part of the gel). Western blot analysis (Figure 2B), using an HRP-conjugated goat anti-human IgG Fc specific mAb (left membrane) and a rat anti-H60 mAb (right membrane), confirmed that purified bifunctional (scFv 4D5/rH60)-Fc protein comprises the mouse NKG2D ligand H60 associated by an Fc part to the considered scFv. The same results were obtained for the purified bifunctional (scFv 511/rH60)-Fc protein (data not shown).

Functional evaluation of the recombinant homodimeric proteins

Homodimeric anti-CEA or anti-HER2 scFv-Fc proteins were tested for their capacity to bind specifically CEA or HER2 on tumor cells by flow cytometry. Using a FITC-conjugated anti-human IgG Fc specific mAb, an efficient and specific binding of the two homodimeric proteins (scFv 511)-Fc and (scFv 4D5)-Fc to MC38 expressing CEA and B16 expressing HER2 cells, respectively, was obtained (Figure 3A, right panels). Flow cytometry signals of similar magnitudes to those obtained with specific monoclonal antibodies, 511 mAb and trastuzumab, were observed (Figure 3A, left panels).

Recombinant H60-Fc protein was tested in a reverse antibody-dependent cell-mediated cytotoxicity (ADCC) lysis assay for its capacity to induce NK-mediated tumor cell lysis. In a chromium release assay using NKG2D-expressing IL-2 activated murine NK cells (LAK cells) as effectors (Figure 3B), pre-incubation of the THP-1 cells, which are known to express FcRs, with rH60-Fc induced a specific increase of THP-1 cells lysis (Figure 3C, black bars). This lysis was reduced to background level (Figure 3C, white bars) when THP-1 cells were pre-incubated with an FcR blocking reagent (Figure 3C, hatched bars), confirming that this lysis was exclusively triggered by the coating of these cells with H60.

TAA binding capacity of the bifunctional proteins

Binding capacity of the two purified bifunctional proteins to tumor cells expressing relevant antigens was assessed by flow cytometry. MC38 cells expressing CEA and B16 cells expressing HER2 (Figure 4, first column) did not express endogenous H60, as assayed with a rat anti-H60 mAb (Figure 4, second column). These two cell lines clearly became positive for H60 detection when incubated with the relevant bifunctional protein, (scFv 511/rH60)-Fc for MC38-CEA⁺ cells and (scFv 4D5/rH60)-Fc for B16-HER2⁺ cells (Figure 4, fifth column). By contrast, no detection of this ligand could be observed when tumor cells were first incubated with the different homodimeric proteins (Figure 4, third and fourth column), neither when tumor cells were first incubated with the irrelevant bifunctional protein (data not shown). These results thus clearly demonstrate the capacity of these two bifunctional proteins to specifically bind and coat tumor cell lines expressing CEA or HER2 with H60.

Cytotoxicity assays

Among all the cell lines negative for endogenous H60 expression tested (the human, LS174T, LoVo and HT29 colon carcinoma; SK-BR-3 breast carcinoma; SK-OV-3 ovarian carcinoma and the murine, MC38 colon carcinoma; RMA T lymphoma both transfected with CEA), SK-BR-3 expressing HER2 and CEA-transfected RMA cells (Figure 5A, black-filled peaks) were chosen for their high resistance to lysis mediated by NK cells. Lysis of these two tumor cell lines was evaluated in presence of the different bifunctional and control proteins (Figure 5B–D). In presence of the (scFv 4D5/rH60)-Fc bifunctional protein, SK-BR-3 cells were much more lysed by murine activated NK cells (Figure 5B, black bars). In contrast, no increase of SK-BR-3 cells lysis was observed when these cells, which do not express CEA (Figure 5A, left panel, black-dotted peak), were incubated with the irrelevant bifunctional protein (scFv 511/rH60)-Fc (Figure 5B, dark grey bars), neither in presence of the homodimeric protein (scFv 4D5)-Fc (Figure 5B, light grey bars), excluding any Fc receptor-engagement by the Fc part of the protein and so any antibody dependent cell cytotoxicity (ADCC). Similarly, (scFv 511/rH60)-Fc bifunctional protein significantly sensitized RMA cells expressing CEA antigen to lysis by activated murine NK cells (Figure 5D, black bars). Finally, specific lysis induced by the two bifunctional proteins studied was abrogated by pre-incubation of murine NK cells with

an anti-mouse NKG2D antibody ([Figure 5B and D](#), hatched bars). This clearly demonstrated that the increase in NK cell-mediated lysis was only the result of H60 engagement, brought exogenously thanks to these two bifunctional proteins, by the activating receptor NKG2D present on NK cells. Moreover, stimulation of NKG2D by our anti-HER2 (scFv 4D5/rH60)-Fc bifunctional protein seems to be more efficient than activation of Fc receptors by anti-HER2 trastuzumab antibody to induce NK-mediated SK-BR-3 tumor cell lysis, whatever the dose tested ([Figure 5C](#)).

Discussion

It has been previously described the *in vivo* rejection of several types of tumor cells when these ones are genetically manipulated to express at the cell surface ligands of NKG2D, an activating receptor expressed on NK and T cells ([Cerwenka et al., 2001](#); [Diefenbach et al., 2001](#); [Friese et al., 2003](#); [Oppenheim et al., 2005](#); [Busche et al., 2006](#)). In this study, we designed and purified new bifunctional proteins, associating a ligand of NKG2D to anti-tumor antibody fragments, and demonstrated their capacity to specifically coat this ligand on different tumor cells and, more importantly, to induce their lysis by NK cells. Here, actually, we demonstrate that the exogenous coating of tumor cells with NKG2D ligands by our bifunctional proteins can induce an efficient and specific NK-mediated lysis of these cells, in an NKG2D-dependent pathway.

Bispecific antibodies and bifunctional molecules have been largely exploited in cancer immunotherapy ([Koelemij et al., 1999](#); [Peipp and Valerius, 2002](#); [Kontermann, 2005](#)), notably in order to favour the recruitment and/or the activation of effector cells to the tumoral site, by immuno-targeted cytokines ([Gillies et al., 1991](#); [Hoogenboom et al., 1991](#)) or agonist antibodies against activating receptors expressed on cytotoxic cells ([Van Dijk et al., 1989](#)), thanks to antibody structures specific for TAAs. Many strategies have been developed to this aim, such as hybrid hybridomas (quadromas ([Milstein and Cuello, 1983](#))) or chemical conjugation ([Karpovsky et al., 1984](#); [Robert et al., 1996](#)). Nevertheless, production of this type of molecules is frequently limited in terms of quality and/or quantity, two major criteria for their use in humans as therapeutic agents. While quadroma method leads to the formation of many different contaminant forms (up to nine ([Carter, 2001](#))) via uncertain rearrangement of the two heavy chains and the two light chains in presence, chemical conjugation, using for example Fab' fragments of monoclonal antibodies ([Robert et al., 1999](#)), is often associated with poor production outputs. Genetic engineering allowed the development of other formats such as diabody ([Arndt et al., 1999](#)), single-chain bispecific antibody ([Loffler et al., 2000](#)), or single-chain diabody, but unfortunately frequently associated with short half-lives *in vivo* ([Kontermann, 2005](#)). To bypass this last point, however, an original study recently proposed the fusion of these small molecules with human serum albumin, strongly improving their pharmacokinetic properties ([Muller et al., 2007](#)).

In a previous study ([Germain et al., 2005](#)), we demonstrated the ability of chemical conjugates associating recombinant MICA molecules with Fab' fragments of different monoclonal anti-tumor antibodies to specifically bind TAAs and to stimulate NKG2D-dependent lysis of resistant tumor cells by human NK cells. In the same way, an antitumor activity against human myeloma has been described with a recombinant ULBP2-scFv anti CD138 molecule ([von Strandmann et al., 2006](#)). Here, in the perspective to produce higher amounts of *in vivo* usable bifunctional proteins we chose to use the “knob into hole” strategy, which was developed to produce bispecific antibodies easily purified from culture supernatant ([Ridgway et al., 1996](#); [Merchant et al., 1998](#)), based on the preferential heterodimerization of two distinct heavy chains of human immunoglobulin. Recently, Xie et al. applied similarly this technology for the production of a bispecific antibody associating an anti-HER2 scFv to an anti-CD16 scFv to stimulate *in vitro* CD16-dependent tumor cell lysis by PBMC ([Xie et al., 2005](#)). In contrast to this report, the (scFv anti-TAA)/rH60-Fc bifunctional proteins produced here associate an anti-tumor scFv to a recombinant form of one of the natural ligands of the mouse NKG2D receptor, H60, able as shown for MICA in humans to induce mouse NK cells cytotoxicity when expressed on tumor cells ([Cerwenka et al., 2000](#); [Diefenbach et al., 2000](#)). We postulated that the intermediate affinity of H60 for mouse NKG2D, compared with other mouse NKG2D ligands ([O'Callaghan et al., 2001](#); [Carayannopoulos et al., 2002](#)), is high enough to bind and activate NK cells when coated on tumor cells, in correlation with TAA expression degree, and not sufficiently high to induce a dramatic, or ineffective, NK cell activation in the whole blood circulation. This hypothesis is supported by results previously obtained during the evaluation of the ability of Fab-HLA-A2/Flu conjugates to activate T-lymphocytes, in soluble form or coated on tumor cells ([Robert et al., 2001](#)). Using Ca²⁺ mobilization assays, we demonstrated that these conjugates, associating monomeric HLA-A2/Flu complexes

to anti-TAA Fab' fragments, can efficiently activate CTL only when oligomerized on the surface of tumor cells.

In the present study, we demonstrated that the two produced (scFv anti-TAA)/rH60- Fc bifunctional proteins are able to specifically coat CEA- or HER2-expressing tumor cells with H60, and to induce their lysis by mouse activated NK cells in an NKG2D-dependent manner.

Acknowledgments

Funding: Comité de l'Hérault Ligue Nationale Contre le Cancer (to G.C.); Association pour la Recherche sur le Cancer (to G.C.); Caisse d'Assurance Maladie des Professions Libérales Provinces; Fondation Gustave Prévot.

Authors thank Marie-Paule Lefranc, Thierry Chardes and Eric Vives for the detailed reading of the manuscript and valuable suggestions, and Geneviève Heintz, Sabine Bousquié, Imade Ait-Arsa and Michel Brissac for precious technical assistance.

Footnotes

Conflict of interest: The authors declare no conflict of interest.

References

- Arndt MA, Krauss J, Kipriyanov SM, Pfreundschuh M, Little M. *Blood*. 1999;94:2562–2568. [PubMed: 10515858]
- Bahram S, Bresnahan M, Geraghty DE, Spies T. *Proc Natl Acad Sci U S A*. 1994;91:6259–6263. [PMCID: PMC44180] [PubMed: 8022771]
- Bauer S, Groh V, Wu J, Steinle A, Phillips JH, Lanier LL, Spies T. *Science*. 1999;285:727–729. [PubMed: 10426993]
- Bukowski JF, Warner JF, Dennert G, Welsh RM. *J Exp Med*. 1985;161:40–52. [PMCID: PMC2187554] [PubMed: 2981954]
- Busche A, Goldmann T, Naumann U, Steinle A, Brandau S. *Hum Gene Ther*. 2006;17:135–146. [PubMed: 16454647]
- Carayannopoulos LN, Naidenko OV, Fremont DH, Yokoyama WM. *J Immunol*. 2002;169:4079–4083. [PubMed: 12370332]
- Carbone E, Neri P, Mesuraca M, Fulciniti MT, Otsuki T, Pende D, Groh V, Spies T, Pollio G, Cosman D, et al. *Blood*. 2005;105:251–258. [PubMed: 15328155]
- Carter P. *Nat Rev Cancer*. 2001;1:118–129. [PubMed: 11905803]
- Cerwenka A, Bakker AB, McClanahan T, Wagner J, Wu J, Phillips JH, Lanier LL. *Immunity*. 2000;12:721–727. [PubMed: 10894171]
- Cerwenka A, Baron JL, Lanier LL. *Proc Natl Acad Sci U S A*. 2001;98:11521–11526. [PMCID: PMC58762] [PubMed: 11562472]
- Clarke P, Mann J, Simpson JF, Rickard-Dickson K, Primus FJ. *Cancer Res*. 1998;58:1469–1477. [PubMed: 9537250]
- Cosman D, Mullberg J, Sutherland CL, Chin W, Armitage R, Fanslow W, Kubin M, Chalupny NJ. *Immunity*. 2001;14:123–133. [PubMed: 11239445]
- Coudert JD, Zimmer J, Tomasello E, Cebecauer M, Colonna M, Vivier E, Held W. *Blood*. 2005;106:1711–1717. [PubMed: 15886320]
- Deyev SM, Waibel R, Lebedenko EN, Schubiger AP, Pluckthun A. *Nat Biotechnol*. 2003;21:1486–1492. [PubMed: 14634668]

Diefenbach A, Jamieson AM, Liu SD, Shastri N, Raulet DH. *Nat Immunol.* 2000;1:119–126. [PubMed: 11248803]

Diefenbach A, Jensen R, Jamieson AM, Raulet DH. *Nature.* 2001;413:165–171. [PMCID: PMC3900321] [PubMed: 11557981]

Friese MA, Platten M, Lutz SZ, Naumann U, Aulwurm S, Bischof F, Buhning HJ, Dichgans J, Rammensee HG, Steinle A, et al. *Cancer Res.* 2003;63:8996–9006. [PubMed: 14695218]

Garrido F, Ruiz-Cabello F, Cabrera T, Perez-Villar JJ, Lopez-Botet M, Duggan-Keen M, Stern PL. *Immunol Today.* 1997;18:89–95. [PubMed: 9057360]

Germain C, Larbouret C, Cesson V, Donda A, Held W, Mach JP, Pelegrin A, Robert B. *Clin Cancer Res.* 2005;11:7516–7522. [PubMed: 16243826]

Gillies SD, Young D, Lo KM, Foley SF, Reisfeld RA. *Hybridoma.* 1991;10:347–356. [PubMed: 1916847]

Groh V, Rhinehart R, Randolph-Habecker J, Topp MS, Riddell SR, Spies T. *Nat Immunol.* 2001;2:255–260. [PubMed: 11224526]

Hoogenboom HR, Volckaert G, Raus JC. *Mol Immunol.* 1991;28:1027–1037. [PubMed: 1922108]

Jamieson AM, Diefenbach A, McMahon CW, Xiong N, Carlyle JR, Raulet DH. *Immunity.* 2002;17:19–29. [PubMed: 12150888]

Jan Chalupny N, Sutherland CL, Lawrence WA, Rein-Weston A, Cosman D. *Biochem Biophys Res Commun.* 2003;305:129–135. [PubMed: 12732206]

Karpovsky B, Titus JA, Stephany DA, Segal DM. *J Exp Med.* 1984;160:1686–1701. [PMCID: PMC2187539] [PubMed: 6239899]

Kim S, Iizuka K, Aguila HL, Weissman IL, Yokoyama WM. *Proc Natl Acad Sci U S A.* 2000;97:2731–2736. [PMCID: PMC15998] [PubMed: 10694580]

Koelmeij R, Kuppen PJ, van de Velde CJ, Fleuren GJ, Hagens M, Eggermont AM. *J Immunother (1997)* 1999;22:514–524.

Kontermann RE. *Acta Pharmacol Sin.* 2005;26:1–9. [PubMed: 15659107]

Lodoen MB, Lanier LL. *Nat Rev Microbiol.* 2005;3:59–69. [PubMed: 15608700]

Loffler A, Kufer P, Lutterbuse R, Zettl F, Daniel PT, Schwenkenbecher JM, Riethmuller G, Dorken B, Bargou RC. *Blood.* 2000;95:2098–2103. [PubMed: 10706880]

Malarkannan S, Shih PP, Eden PA, Horng T, Zuberi AR, Christianson G, Roopenian D, Shastri N. *J Immunol.* 1998;161:3501–3509. [PubMed: 9759870]

Merchant AM, Zhu Z, Yuan JQ, Goddard A, Adams CW, Presta LG, Carter P. *Nat Biotechnol.* 1998;16:677–681. [PubMed: 9661204]

Milstein C, Cuello AC. *Nature.* 1983;305:537–540. [PubMed: 6137772]

Moretta A, Bottino C, Vitale M, Pende D, Cantoni C, Mingari MC, Biassoni R, Moretta L. *Annu Rev Immunol.* 2001;19:197–223. [PubMed: 11244035]

Muller D, Karle A, Meissburger B, Hofig I, Stork R, Kontermann RE. *J Biol Chem.* 2007;282:12650–12660. [PubMed: 17347147]


O’Callaghan CA, Cerwenka A, Willcox BE, Lanier LL, Bjorkman PJ. *Immunity.* 2001;15:201–211. [PubMed: 11520456]

Ogg GS, Dunbar PR, Cerundolo V, McMichael AJ, Lemoine NR, Savage P. *Br J Cancer.* 2000;82:1058–1062. [PMCID: PMC2374430] [PubMed: 10737389]

- Oppenheim DE, Roberts SJ, Clarke SL, Filler R, Lewis JM, Tigelaar RE, Girardi M, Hayday AC. *Nat Immunol*. 2005;6:928–937. [PubMed: 16116470]
- Peipp M, Valerius T. *Biochem Soc Trans*. 2002;30:507–511. [PubMed: 12196124]
- Pelegrin A, Terskikh A, Hayoz D, Chalandon Y, Olsson NO, Folli S, Buchegger F, Kromer B, Schwarz K, Martin M, et al. *Int J Cancer*. 1992;52:110–119. [PubMed: 1500216]
- Pende D, Rivera P, Marcenaro S, Chang CC, Biassoni R, Conte R, Kubin M, Cosman D, Ferrone S, Moretta L, et al. *Cancer Res*. 2002;62:6178–6186. [PubMed: 12414645]
- Raulet DH, Vance RE. *Nat Rev Immunol*. 2006;6:520–531. [PubMed: 16799471]
- Ridgway JB, Presta LG, Carter P. *Protein Eng*. 1996;9:617–621. [PubMed: 8844834]
- Robert B, Dorvillius M, Buchegger F, Garambois V, Mani JC, Pugnieres M, Mach JP, Pelegrin A. *Int J Cancer*. 1999;81:285–291. [PubMed: 10188732]
- Robert B, Guillaume P, Luescher T, Doucey MA, Cerottini JC, Romero P, Mach JP. *Cancer Immun*. 2001;1:2. [PubMed: 12747763]
- Robert B, Mach JP, Mani JC, Ychou M, Folli S, Artus JC, Pelegrin A. *Cancer Res*. 1996;56:4758–4765. [PubMed: 8840995]
- Salih HR, Antropius H, Gieseke F, Lutz SZ, Kanz L, Rammensee HG, Steinle A. *Blood*. 2003;102:1389–1396. [PubMed: 12714493]
- Salih HR, Rammensee HG, Steinle A. *J Immunol*. 2002;169:4098–4102. [PubMed: 12370336]
- Schneider P. *Methods Enzymol*. 2000;322:325–345. [PubMed: 10914028]
- Smyth MJ, Hayakawa Y, Takeda K, Yagita H. *Nat Rev Cancer*. 2002;2:850–861. [PubMed: 12415255]
- Stirnemann K, Romero JF, Baldi L, Robert B, Cesson V, Besra GS, Zauderer M, Wurm F, Corradin G, Mach JP, et al. *J Clin Invest*. 2008;118:994–1005. [PMCID: PMC2230658] [PubMed: 18259610]
- Tay CH, Szomolanyi-Tsuda E, Welsh RM. *Curr Top Microbiol Immunol*. 1998;230:193–220. [PubMed: 9586357]
- Van Dijk J, Tsuruo T, Segal DM, Bolhuis RL, Colognola R, van de Griend RJ, Fleuren GJ, Warnaar SO. *Int J Cancer*. 1989;44:738–743. [PubMed: 2793246]
- Vetter CS, Groh V, Straten P, Spies T, Brocker EB, Becker JC. *J Invest Dermatol*. 2002;118:600–605. [PubMed: 11918705]
- Vetter CS, Lieb W, Brocker EB, Becker JC. *Br J Cancer*. 2004;91:1495–1499. [PMCID: PMC2409941] [PubMed: 15381927]
- von Strandmann EP, Hansen HP, Reiners KS, Schnell R, Borchmann P, Merkert S, Simhadri VR, Draube A, Reiser M, Purr I, et al. *Blood*. 2006;107:1955–1962. [PubMed: 16210338]
- Xie Z, Guo N, Yu M, Hu M, Shen B. *J Immunol Methods*. 2005;296:95–101. [PubMed: 15680154]
- Zou Y, Mirbaha F, Stastny P. *Hum Immunol*. 2006;67:183–187. [PubMed: 16698440]



Figures and Tables

Fig. 1




Schematic representation of the different homodimeric and bifunctional proteins produced. Upper illustrations represent the parental homodimeric proteins (scFv 511)-Fc, rH60-Fc and (scFv 4D5)-Fc. Lower illustrations represent the two bifunctional proteins obtained after knob and hole variants generation, namely (scFv 511/rH60)-Fc and (scFv 4D5/rH60)-Fc.

Fig. 2




SDS-PAGE and western blot analysis of the bifunctional protein (scFv 4D5/rH60)-Fc in comparison with the two parental homodimeric proteins rH60-Fc and (scFv 4D5)-Fc. (A) SDS-PAGE analysis. Under non-reducing conditions (left part of the gel) (scFv 4D5/rH60)-Fc appeared as a single band of intermediate molecular weight compared with the bands obtained for rH60-Fc and (scFv 4D5)-Fc. Under reducing conditions (right part of the gel) (scFv 4D5/rH60)-Fc appeared as two bands, each one of MW similar to the single bands obtained for the two related homodimeric parental proteins. (B) Western blot analysis. Left membrane, revealed by an HRP-conjugated goat anti-human IgG Fc specific mAb, confirmed that these two different chains contain a human Fc part, and that only one (the heavier) contains H60, as demonstrated in right membrane, first incubated with a rat anti-H60 mAb and then revealed by an HRP-conjugated goat anti-rat IgG whole molecule mAb. Similar profiles were obtained for (scFv 511/rH60)-Fc SDS-PAGE and western blot analysis.

Fig. 3




Functional evaluation of the three homodimeric parental molecules. (A) Flow cytometry evaluation of the control molecules (scFv 511)-Fc and (scFv 4D5)-Fc on CEA expressing MC38 cells and HER2 expressing B16 cells. Cells were first incubated with scFv-Fc proteins, then with a FITC-conjugated goat anti-human IgG Fc specific mAb, and finally analysed on a FACScan. Each scFv-Fc protein efficiently bound the TAA considered (black-filled peaks, right panels). As control, CEA expression on MC38-CEA cells and HER2 expression on B16-HER2 cells (black filled peaks, left panels) was evaluated using the specific monoclonal antibodies 511 mAb and trastuzumab, respectively. Cells were first incubated with monoclonal antibodies, then with a FITC-conjugated goat anti-mouse IgG Fc specific mAb or with a FITC-conjugated goat anti-human IgG Fc specific mAb, respectively, and finally analysed on a FACScan. (B) NKG2D expression on murine LAK cells. Cells were first incubated with a rat anti-mouse NKG2D mAb, then with a FITC-conjugated mouse anti-rat IgG whole molecule mAb, and finally analysed on a FACScan. LAK cells, obtained after incubation of murine splenocytes for 5 days in presence of rhIL-2 (5000 units/ml), significantly expressed NKG2D (black-filled peak). (C) Chromium release evaluation of the control homodimeric rH60-Fc. Preincubation of THP-1 cells with rH60-Fc (black bars) significantly increased targets specific lysis by murine LAK cells in comparison with control THP-1 cells (white bars). Addition of an FcR blocking reagent reduced targets specific lysis to control (hatched bars).

Fig. 4



Flow cytometry evaluation of the two bifunctional proteins produced. CEA, HER2 and H60 expression on tumor cells was evaluated using anti-CEA 511 mAb, anti-HER2 trastuzumab or a rat anti-H60 mAb, revealed by a FITC-conjugated goat anti-mouse IgG Fc specific mAb, a FITC conjugated goat anti-human IgG Fc specific mAb or a FITC conjugated mouse anti-rat IgG2a Fc specific mAb, respectively. Neither CEA expressing MC38 cells nor HER2 expressing B16 cells (black-filled peaks, first column) did express H60 (black-filled peaks, second column). In a second time, tumor cells were first incubated with the homodimeric parental proteins, rH60-Fc, (scFv 511)-Fc or (scFv 4D5)-Fc, or with the relevant bifunctional molecule, (scFv 511/rH60)-Fc or (scFv 4D5/rH60)-Fc, then with a rat anti-H60 mAb, and finally with a FITC-conjugated mouse anti-rat IgG2a Fc specific mAb. As opposed to control molecules (black-filled peaks, third and fourth columns), each bifunctional molecule specifically coated tumor cells with H60 (black-filled peaks, fifth column).

Fig. 5



Antigenic profile and susceptibility to LAK cells lysis, without and with recombinant bifunctional proteins (scFv anti-TAA/rH60)-Fc, of the different tumor cell lines tested. (A) CEA, HER2 and H60 expression on SK-BR-3 and CEA transfected RMA cell lines. Neither HER2 expressing SK-BR-3 cells nor CEA expressing RMA cells (black-filled peaks) did express H60 (black-thick peaks). SK-BR-3 cells neither expressed CEA (black-dotted peak). (B–D) Functional evaluation in chromium release assay of the bifunctional proteins (scFv 4D5/rH60)-Fc and (scFv 511/rH60)-Fc. (B) Preincubation of SK-BR-3 cells with (scFv 4D5/rH60)-Fc (black bars) significantly increased targets lysis by murine LAK cells in comparison with SK-BR-3 alone (white bars), as opposed to SK-BR-3 cells preincubated with the homodimeric parental protein (scFv 4D5)-Fc (light grey bars) or with the irrelevant bifunctional protein (scFv 511/rH60)-Fc (dark grey bars). (C) Higher cytotoxicity is obtained when SK-BR-3 cells are preincubated with (scFv 4D5/rH60)-Fc (black bars) than with trastuzumab (white bars). (D) RMA-CEA cells: Preincubation with (scFv 511/rH60)-Fc (black bars) significantly increased targets lysis by murine LAK cells compared to RMA-CEA alone (white bars), as opposed to RMA-CEA cells preincubated with the irrelevant bifunctional protein (scFv 4D5/rH60)-Fc (dark grey bars) or with anti-mNKG2D (hatched bars).

with Herceptin (trastuzumab; white bars) at every doses tested in presence of murine LAK cells. (D) Preincubation of RMA-CEA⁺ cells with the bifunctional protein (ScFv 511/rH60)-Fc (black bars) significantly sensitized targets to lysis by murine LAK cells in comparison with control (white bars). Preincubation of murine LAK cells with a rat anti-mouse NKG2D mAb completely prevented tumor cell lysis increase induced by the bifunctional protein considered (B, D hatched bars).