










Swiss consensus on the management of acute diverticulitis

Timothée Girardin¹ , David Martin¹, Enrique Lázaro-Fontanet¹ , Daniel Clerc¹ , Martin Hübner¹, Lukas Brügger², Matthias Turina³, Walter Brunner⁴ , Dimitri Christoforidis⁵, Frederic Ris⁶, Michel Adamina⁷ , Marco von Strauss⁸ , Dieter Hahnloser^{1,*}  and the Swiss Colorectal Working Group

¹Department of Visceral Surgery, Lausanne University Hospital (CHUV), University of Lausanne (UNIL), Lausanne, Switzerland

²Department of Visceral Surgery, Bern University Hospital (Inselspital), Bern, Switzerland

³Department of Visceral Surgery, Zurich University Hospital, Zurich, Switzerland

⁴Department of Visceral Surgery, St-Gallen Kantonsspital, St-Gallen, Switzerland

⁵Department of Visceral Surgery, Ospedale Regionale di Lugano, Lugano, Switzerland

⁶Department of Visceral Surgery, Hôpitaux Universitaires de Genève (HUG), Genève, Switzerland

⁷Department of Visceral Surgery, Winterthur Kantonsspital, Winterthur, Switzerland

⁸Department of Visceral Surgery, St Claraspital und Universitätsspital Basel, Basel, Switzerland

*Correspondence to: Dieter Hahnloser, Department of Visceral Surgery, Lausanne University Hospital (CHUV), Rue du Bugnon 46, Lausanne, 1011, Switzerland (e-mail: dieter.hahnloser@chuv.ch)

Members of the Swiss Colorectal Working Group are co-authors of this study and are listed under the heading Collaborators.

Presented as an original abstract at the Swiss Surgical Annual Meeting (SGC-SCC Congress), Bern, Switzerland, June 2022 and at the Annual Swiss Gastroenterology and Visceral Surgery Meeting, Interlaken, Switzerland, September 2022.

Introduction

Acute diverticulitis (AD) is one of the most common clinical conditions encountered in the emergency setting¹. Possible complications include abscess, perforation, fistulas, obstruction, and bleeding. Fistulas are often the consequence of repeated acute episodes, with localized perforation into adjacent viscera². Multiple AD episodes can also increase the risk of stricture formation. The risk of developing an AD in a patient with diverticulosis is estimated at 4% throughout a patient's lifetime³.

Management of this condition has recently changed, with guidelines published by the World Society of Emergency Surgery (WSES) and the European Society of Coloproctology (ESCP)^{4,5}. However, the levels of evidence are inconsistent and recommendations may vary according to the health systems and the resources available. There can be a lag of years from publication to implementation⁶.

Considering these variations and the latest published management recommendations, the aim of the present study was to establish a consensus among colorectal surgeons on the diagnosis and management of AD based on Swiss national practice.

Methods

All Swiss hospitals that performed, according to the national statistics (2015–2017), at least 50 colorectal resections annually, over a 3-year interval, were contacted (52 hospitals). Each centre could include up to three board-certified surgeons at a consultant level, regularly performing colorectal resection for benign and malignant indications. No minimal individual surgeon volume was

required. A three-round Delphi consensus was conducted, which is detailed in the [Supplementary Methods](#). Consensus was defined as greater than or equal to 70% agreement, either obtained within one single answer or with multiple answers pointing in the same direction (either positive or negative), for questions on a Likert scale (1–2 versus 4–5).

Results

Initial evaluation, abdominal imaging, complicated diverticulitis definition, and outpatient management (Fig. 1)

Abdominal CT reached consensus as the recommended imaging modality (100%) for diagnosis of AD. Complicated diverticulitis was defined as the presence of an abscess (88%), fistula (84%), pericolic (86%) or distant (83%) free air, and rectal contrast extravasation (95%). Stenosis (62%) and free fluid in the pelvis (40%) or in the upper abdomen (48%) were not deemed to be complicated diverticulitis.

Fever (58%), findings of localized peritonitis (47%), antibiotic requirement (14%), and leucocytes greater than $20 \times 10^9/l$ (49%) were not considered as contraindications for outpatient management if taken individually. Additional results are available in the [Supplementary material](#).

Non-operative/operative treatment and elective surgery (Fig. 2)

Antibiotic treatment and percutaneous drainage are described in [Fig. 2](#). In the case of drainage, no consensus was reached on the necessity to perform control abdominal imaging (51%).

Received: August 17, 2023. Revised: November 23, 2023. Accepted: November 27, 2023

© The Author(s) 2024. Published by Oxford University Press on behalf of BJS Foundation Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com




Fig. 1 Initial evaluation, abdominal imaging, complicated diverticulitis definition and contraindications to outpatient management

Likert scale: 1, strongly disagree; 2, partially agree; 3, neutral; 4, agree; and 5, strongly agree. CRP, C-reactive protein; GP, general practitioner; AD, acute diverticulitis.

No procedure reached consensus for surgical management in the emergency setting (Fig. 2). Primary anastomosis was the preferred option when the patient is stable (77%) (stapled end to end, 73%). From a technical point of view, no consensus was reached on the type of dissection (28% for total mesorectal excision *versus* 42% for close-to-bowel dissection). Insertion of a pelvic drain (90%) reached consensus for emergency procedures.

Elective surgical resection was recommended in patients with persistent symptoms (95%) or immunosuppression (89%). Age (14%) and number of episodes of recurrent AD (27%) were not considered criteria for elective surgery. Standard planned elective sigmoidectomy after a complicated case of AD (42%) or after a percutaneously drained abscess (65%) did not gain consensus. Laparoscopic resection was the favoured approach for elective sigmoidectomy (98%). Additional results are available in the [Supplementary material](#).

Discussion

Swiss colorectal surgeons reached consensus on the management of AD with regard to diagnosis, outpatient management, and indication for elective surgery. Emergency surgical management did not reach consensus.

Many radiological classifications have described the different stages of diverticular disease; however, none is currently universally accepted⁷. This was also apparent in this study. The definitive treatment for each stage also depends on the clinical condition and patient co-morbidity. More recent classifications (Sallinen and WSES) that combine clinical, radiological, and

physiological parameters were not considered in this study, limiting the generalization of the results in clinical practice^{8,9}.

Non-operative management of AD has changed over the past few decades. Two randomized controlled studies (AVOD and DIABOLO trials) confirmed that this approach was not associated with a higher recurrence or rate of sigmoidectomy^{10,11}. However, controversies and habits tend to persist, as highlighted in the present study, where a consensus was reached in favour of antibiotic treatment¹⁰⁻¹³. In ESCP guidelines, antibiotics are recommended for immunocompromised patients or sepsis⁵. Percutaneous drainage of a pericolic abscess is another unresolved question. A systematic review revealed comparable treatment failure rates for antibiotics (19.9%) and percutaneous drainage (20.8%)¹⁴. Based on these results, it was suggested that abscesses greater than 4 cm should be drained, confirming the results of the present study.

A systematic review demonstrated that outpatient management of AD was safe, but no criteria of failure were identified, which makes patient selection difficult¹⁵. The present study identified subjective contraindications for outpatient management, such as generalized peritonitis, immunosuppression, or complicated diverticulitis. These criteria were assessed individually, not in combination. Management decisions should be made based on the general and physiological condition of the patient.

The present study reached a consensus on the necessity for emergency surgery in patients with sepsis. However, the levels of evidence are inconsistent in patients with perforation without haemodynamic instability⁵. WSES guidelines suggest a Hartmann procedure in patients with faecal peritonitis or sepsis⁴. Conversely,




Fig. 2 Non-operative/operative treatment and elective surgery

Likert scale: 1, strongly disagree; 2, partially agree; 3, neutral; 4, agree; and 5, strongly agree. AD, acute diverticulitis; PA, primary anastomosis.

ESCP guidelines state that the choice of the procedure should be left to the surgeon, according to their experience, but resection is recommended⁵. The present study has not established a consensus on the most appropriate surgical technique. The experience and the logistics according to the hospital (ICU, theatres, and out-of-hours access) probably play an important role in decision-making. Damage-control surgery (DCS) is an option in unstable patients. A retrospective study of 203 consecutive patients treated with DCS by applying a decision algorithm for severe intra-abdominal sepsis confirmed that it was feasible and 65% of patients were stoma free at hospital discharge¹⁶. However, none of the guidelines recommends DCS over a Hartmann procedure^{4,5}. Indications, timing, and techniques of the DCS approach need to be further assessed (for example primary anastomosis with a diverting loop ileostomy or Hartmann procedure).

The use of laparoscopic lavage for purulent peritonitis is controversial; however, due to the lack of proof supporting its inferiority, this technique may be used according to the ESCP guidelines⁵. The present study demonstrated no consensus for the role of laparoscopic lavage. The SCANDIV trial results suggested that the use of laparoscopic lavage *versus* primary resection did not reduce severe postoperative complications and led to a higher reoperation rate^{17,18}.

The target surgeon group was chosen to include colorectal experts, aiming to provide a well-founded basis for the diagnosis and management of AD. This study addressed the question in a methodologically robust way. The Delphi technique is a widely used and validated method to transform opinion into group consensus among experts and specific questions or issues¹⁹. The

aim of the study was not to generate new guidelines for AD, but to confirm/challenge them within a group of national experts, as published guidelines are not systematically accepted or applied in practice today.

Collaborators

Swiss Colorectal Working Group

Aamr Orban (Seespital, Horgen, Switzerland); Adrienne Imhof (Praxis Imhof, Schaffhausen, Switzerland); Alex Ochsner (Spitallimmattal, Schlieren, Switzerland); Alfred Kuhrmeier (Clinica Luganese Moncucco, Lugano, Switzerland); Andrea Donadini (Clinica Luganese Moncucco, Lugano, Switzerland); Andreas Keerl (Kantonsspital Baden, Baden, Switzerland); Andreas Kohler (Inselspital, Bern, Switzerland); Andreas Scheiwiller (Luzerner Kantonsspital, Luzern, Switzerland); Andreas Zerz (eSwiss Medical & Surgical Center, St Gallen, Switzerland); Anita Scheiwiller (Klinik Beau-Site, Bern, Switzerland); Annette Ringger (Solethurner Spitaler, Solothurn, Switzerland); Antje Lechleiter (Inselspital, Bern, Switzerland); Antonio Nocito (Kantonsspital Baden, Baden, Switzerland); Beat Muggli (Praxis Beat Muggli, Bern, Switzerland); Beat Schnuriger (Inselspital, Bern, Switzerland); Beatrice Kern (Clarunis, Basel, Switzerland); Bernhard Egger (HFR, Fribourg, Switzerland); Boumediene Mustapha Guendil (Hopital du Valais, Sion, Switzerland); Carsten Thomas Viehl (Spitalzentrum Biel, Biel/Bienne, Switzerland); Cedric Vallet (Clinique de la Source, Lausanne, Switzerland); Christian Nebiker (Kantonsspital Aarau, Aarau, Switzerland); Christine Maurus (Hopital du Valais, Sion,

Switzerland); Claudio Soravia (Clinique générale Baulieu, Genève, Switzerland); Cornelia Frei (Spital Zollikerberg, Zollikerberg, Switzerland); Daniel Steinemann (Clarunis, Basel, Switzerland); David Petermann (EHC, Morges, Switzerland); Duri Gianom-Campell (Clinica Luganese Moncucco, Lugano, Switzerland); Emanuel Gmuer (GZO Spital Wetzikon, Wetzikon, Switzerland); Eric Grossen (Seespital, Horgen, Switzerland); Fabian Hauswirth (Spital Thurgau, Frauenfeld, Switzerland); Franc Hetzer (Viszeralchirurgie Bellaria, Zürich, Switzerland); François Pugin (HFR, Fribourg, Switzerland); Frank-Rainer Müller (Stadtspital, Zürich, Switzerland); Gaël Simoné (Clinique de Genolier, Genolier, Switzerland); Georg Linke (Spital Thun, Thun, Switzerland); Giacinto Basilicata (Spital Bülach, Bülach, Switzerland); Ida Füglistaler (Claraspital, Basel, Switzerland); Ivo Fischer (Spital Bülach, Bülach, Switzerland); Jan Schmidt (Praxis Swiss Surgery, Zürich, Switzerland); Jens Forberger (Spital Männedorf, Männedorf, Switzerland); Karoline Horisberger (USZ, Zurich, Switzerland); Luca Di Mare (EHC, Morges, Switzerland); Marc Daniel Worreth (RHNe, Neuchâtel, Switzerland); Marc-Olivier Guenin (Clarunis, Basel, Switzerland); Marc-Olivier Sauvain (RHNe, Neuchâtel, Switzerland); Mark Hartel (Kantonsspital Aarau, Aarau, Switzerland); Markus Gass (Luzerner Kantonsspital, Luzern, Switzerland); Markus Müller (Spital Thurgau, Frauenfeld, Switzerland); Michel Erne (eHnv, Yverdon-les-Bains, Switzerland); Nicolas Buchs (Hôpital de la Tour, Genève, Switzerland); Olivier Gié (Clinica Luganese Moncucco, Lugano, Switzerland); Olivier Pittet (Hôpital Riviera-Chablais, Rennaz, Switzerland); Othmar Schöb (Chirurgisches Zentrum, Zürich, Switzerland); Peter Sandera (Spitäler Schaffhausen, Schaffhausen, Switzerland); Peter Studer (VISCERA Bauchmedizin, Bern, Switzerland); Raffaele Galli (Kantonsspital Baselland, Liestal, Switzerland); Rebecca Kraus (Kantonsspital Graubünden, Chur, Switzerland); Riadh Ksontini (Hôpital Riviera-Chablais, Rennaz, Switzerland); Robert Rosenberg (Kantonsspital Baselland, Liestal, Switzerland); Roland Chautems (RHNe, Neuchâtel, Switzerland); Sebastian Lamm (Kantonsspital Baselland, Liestal, Switzerland); Stefan Wildi (Stadtspital, Zürich, Switzerland); Stephan Sager (Seespital, Horgen, Switzerland); Stephan Vorburger (Spital Emmental, Burgdorf, Switzerland); Teresa Cereser (Kantonsspital Winterthur, Winterthur, Switzerland); Thomas Kinsbergen (SRO Klinik, Langenthal, Switzerland); Thomas Simon (Hôpital du Valais, Visp, Switzerland); Urs Zingg (Spital Limattal, Schlieren, Switzerland).

Funding

The authors have no funding to declare.

Acknowledgements

T.G. and D.M. contributed equally and share first authorship.

Disclosure

The online voting system was made available by Johnson&Johnson, Switzerland. The authors declare no other conflict of interest.

Supplementary material

[Supplementary material](#) is available at *BJS Open* online.

Data availability

Data are available on request.

References

- Wilkins T, Embry K, George R. Diagnosis and management of acute diverticulitis. *Am Fam Physician* 2013;**87**:612–620
- Bahadursingh AM, Virgo KS, Kaminski DL, Longo WE. Spectrum of disease and outcome of complicated diverticular disease. *Am J Surg* 2003;**186**:696–701
- Shahedi K, Fuller G, Bolus R, Cohen E, Vu M, Shah R et al. Long-term risk of acute diverticulitis among patients with incidental diverticulosis found during colonoscopy. *Clin Gastroenterol Hepatol* 2013;**11**:1609–1613
- Sartelli M, Weber DG, Kluger Y, Ansaloni L, Coccolini F, Abu-Zidan F et al. 2020 update of the WSES guidelines for the management of acute colonic diverticulitis in the emergency setting. *World J Emerg Surg* 2020;**15**:32
- Schultz JK, Azhar N, Binda GA, Barbara G, Biondo S, Boermeester MA et al. European Society of Coloproctology: guidelines for the management of diverticular disease of the colon. *Colorectal Dis* 2020;**22**:5–28
- Faes S, Hübner M, Demartines N, Hahnloser D, Martin D; Swiss Snapshot Diverticulitis Group. Elective surgery for diverticulitis in Swiss hospitals. *Front Surg* 2021;**8**:717228
- Klarenbeek BR, de Korte N, van der Peet DL, Cuesta MA. Review of current classifications for diverticular disease and a translation into clinical practice. *Int J Colorectal Dis* 2012;**27**:207–214
- Sallinen VJ, Leppäniemi AK, Mentula PJ. Staging of acute diverticulitis based on clinical, radiologic, and physiologic parameters. *J Trauma Acute Care Surg* 2015;**78**:543–551
- Sartelli M, Moore FA, Ansaloni L, Di Saverio S, Coccolini F, Griffiths EA et al. A proposal for a CT driven classification of left colon acute diverticulitis. *World J Emerg Surg* 2015;**10**:3
- Chabok A, Pählman L, Hjern F, Haapaniemi S, Smedh K; AVOD Study Group. Randomized clinical trial of antibiotics in acute uncomplicated diverticulitis. *Br J Surg* 2012;**99**:532–539
- van Dijk ST, Daniels L, Ünlü Ç, de Korte N, van Dieren S, Stockmann HB et al. Long-term effects of omitting antibiotics in uncomplicated acute diverticulitis. *Am J Gastroenterol* 2018;**113**:1045–1052
- Isacson D, Smedh K, Nikberg M, Chabok A. Long-term follow-up of the AVOD randomized trial of antibiotic avoidance in uncomplicated diverticulitis. *Br J Surg* 2019;**106**:1542–1548
- Daniels L, Ünlü Ç, de Korte N, van Dieren S, Stockmann HB, Vrouwenraets BC et al. Randomized clinical trial of observational versus antibiotic treatment for a first episode of CT-proven uncomplicated acute diverticulitis. *Br J Surg* 2017;**104**:52–61
- Gregersen R, Mortensen LQ, Burcharth J, Pommergaard HC, Rosenberg J. Treatment of patients with acute colonic diverticulitis complicated by abscess formation: a systematic review. *Int J Surg Lond Engl* 2016;**35**:201–208
- Cirocchi R, Randolph JJ, Binda GA, Gioia S, Henry BM, Tomaszewski KA et al. Is the outpatient management of acute diverticulitis safe and effective? A systematic review and meta-analysis. *Tech Coloproctol* 2019;**23**:87–100
- Faes S, Hübner M, Girardin T, Demartines N, Hahnloser D. Rate of stoma formation following damage-control surgery for severe

- intra-abdominal sepsis: a single-centre consecutive case series. *BJS Open* 2021;**5**:zrab106
17. Schultz JK, Wallon C, Blecic L, Forsmo HM, Folkesson J, Buchwald P et al. One-year results of the SCANDIV randomized clinical trial of laparoscopic lavage versus primary resection for acute perforated diverticulitis. *Br J Surg* 2017;**104**:1382–1392
 18. Schultz JK, Yaqub S, Wallon C, Blecic L, Forsmo HM, Folkesson J et al. Laparoscopic lavage vs primary resection for acute perforated diverticulitis: the SCANDIV randomized clinical trial. *JAMA* 2015;**314**:1364–1375
 19. Jones J, Hunter D. Consensus methods for medical and health services research. *BMJ* 1995;**311**:376–380