P. Pascale B. Quartenoud

"Thrombolysis" by a neuromuscular blocking agent

A 72-year-old man was brought to our intensive care unit for a community-acquired pneumonia complicated by septic shock with multiple organ failure. Mechanical ventilation was initiated upon admission. Three days later, the patient developed striking ST segment elevation on electrocardiogram raising concern of an ongoing cardiac event (Fig. 1). However, no hemodynamic repercussion was present. Serum electrolytes were normal.

A further electrocardiogram provided evidence of a noncardiac electrical activity with a baseline undulation of the electrocardiogram asynchronous with


Fig. 1

Received: 16 November 2005 Accepted: 25 November 2005 Published online: 27 February 2006


Patrizio Pascale, MD · B. Quartenoud, MD Division of Cardiology
Department of Internal Medicine
University Hospital Zurich
8091 Zurich, Switzerland
Tel.: +41-442551111
Fax: +41-442554251
E-Mail: patrizio.pascale@usz.ch

Authors' institutions: Intensive Care Unit Cantonal Hospital Fribourg, Switzerland the regular superimposed QRS complex (Fig. 2). The electrocardiogram normalized after administration of a neuromuscular blocking agent for the ventilatory management of the acute respiratory distress syndrome (Fig. 3). The echocardiogram performed subsequently showed diaphragmatic myoclonus at a frequency similar to the heart rate [1–3]. On the first electrocardiogram performed, the superimposed electromyographic activity of the diaphragmatic flutter was incidentally in phase with the QRS complex and had therefore been misinterpreted as cardiac electrical activity.

Fig. 2







References

- Rigatto M, De Medeiros NP (1962)
 Diaphragmatic flutter: report of a case and review of the literature. Am J Med 32:103–109
- Cvietusa PJ, Nimmagadda SR, Wood R, Liu AH (1995) Diaphragmatic flutter presenting as inspiratory stridor. Chest 107:872–875
- Hoffman R, Yahr W, Krieger B (1990)
 Diaphragmatic flutter resulting on failure to wean from mechanical ventilator support after coronary artery bypass surgery. Crit Care Med 18:499–501