



New Reference Genome Sequences for 17 Bacterial Strains of the Honey Bee Gut Microbiota

Kirsten M. Ellegaard,^a Philipp Engel^a

^aDepartment of Fundamental Microbiology, University of Lausanne, Lausanne, Switzerland

ABSTRACT We sequenced the genomes of 17 strains isolated from the gut of honey bees, including strains representing the genera *Lactobacillus*, *Bifidobacterium*, *Gilliamella*, *Snodgrassella*, *Frischella*, and *Commensalibacter*. These genome sequences represent an important step forward in the development of a comprehensive reference database to aid future analysis of this emerging gut microbiota model.

The honey bee gut is colonized by a remarkably simple community dominated by only 8 to 10 bee-specific phylotypes (1). However, genome-level analyses have shown that several of the phylotypes comprise highly divergent strains (2–4). As such, the honey bee is a promising future model for studying strain-level evolution and function in gut-associated bacterial communities (5). Here, we present 17 new genome sequences of strains isolated from the gut of honey bees, which were generated to facilitate the development of a reference genome database for this community. All strains were isolated from honey bees collected from our apiary in Lausanne, Switzerland, by culturing gut homogenates on agar plates (6) under microaerophilic or anaerobic conditions (7).

Four strains of the genus *Lactobacillus* (Table 1) were selected for sequencing with PacBio 20K (Pacific Biosciences) single-molecule real-time (SMRT) technology. The strains were grown overnight in MRS broth supplemented with fructose and cysteine (8) at 35°C under anaerobic conditions, and total genomic DNA was extracted using a cetyltrimethylammonium bromide-based extraction protocol (7). *De novo* genome assembly was done using the Hierarchical Genome Assembly Process (HGAP) version 2.3. Another 13 strains representing the genera *Bifidobacterium*, *Gilliamella*, *Snodgrassella*, *Frischella*, and *Commensalibacter* (Table 1) were selected for sequencing with Illumina technology. The strains were cultured as described previously (7), and total genomic DNA was extracted with the GenElute bacterial genomic DNA kit according to the manufacturer's instructions (Table 1). Sequencing libraries were prepared with the TruSeq DNA kit and sequenced on the MiSeq platform (Illumina) using the paired-end 2 × 250-bp protocol. All 13 genomes were sequenced to a minimum depth of 50× (Table 1). The resulting FASTQ files were trimmed with Trimmomatic (9) to remove eventual adapter sequences and low-quality reads using the following parameters: LEADING, 20; TRAILING, 20; SLIDINGWINDOW, 4:15; and MINLEN, 50. The reads were assembled with SPAdes version 3.7.1 (10) using the “-careful” flag and multiple *k*-mer sizes (21, 33, 55, 77, 99, 127). Small contigs (less than 500 bp) and contigs with low *k*-mer coverage (less than 5×) were removed from the assemblies, resulting in 6 to 40 contigs per assembly, with a median N_{50} of 529,190 bp. For strains with related complete genome sequences or scaffolds available, the contigs were reordered with Mauve (11).

Assembly qualities were checked by remapping reads to assemblies with the Burrows-Wheeler Aligner (12) and by GC-skew visualization with DNAPlotter (13). For strain ESL0184, the main chromosome was cut into three contigs due to assembly

Received 14 June 2018 Accepted 22 June 2018 Published 26 July 2018

Citation Ellegaard KM, Engel P. 2018. New reference genome sequences for 17 bacterial strains of the honey bee gut microbiota. *Microbiol Resour Announc* 7:e00834-18. <https://doi.org/10.1128/MRA.00834-18>.

Editor J. Cameron Thrash, Louisiana State University

Copyright © 2018 Ellegaard and Engel. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Philipp Engel, philipp.engel@unil.ch.

TABLE 1 Genome assembly statistics and strain information

Genus	Species	Phylotype ^a	Sublineage	Strain	Extracted DNA (μg) ^b	No. of contigs	N_{50} (bp)	Assembly size (bp)	Coverage (x)	GC content (%)	No. of genes ^c
<i>Lactobacillus</i>	<i>L. apis</i>	Firm5	Firm5-1	ESL0185	17.9	1	1,683,102	1,683,102	420	37	1,578
<i>Lactobacillus</i>	<i>L. helsingborgensis</i>	Firm5	Firm5-2	ESL0183	19.3	2	1,856,015	1,867,232	300	37	1,780
<i>Lactobacillus</i>	<i>L. melliventris</i>	Firm5	Firm5-3	ESL0184	19.8	4	1,505,590	2,036,181	320	36	2,015
<i>Lactobacillus</i>	<i>L. kullabergensis</i>	Firm5	Firm5-4	ESL0186	16.1	1	2,018,944	2,018,944	290	36	1,915
<i>Bifidobacterium</i>	<i>B. asteroides</i>	Bifido	Bifido-1	ESL0170	1.1	7	1,162,986	2,175,262	200	60	1,771
<i>Bifidobacterium</i>	<i>B. asteroides</i>	Bifido	Bifido-1	ESL0198	1.3	12	618,428	2,235,610	280	60	1,820
<i>Bifidobacterium</i>	<i>B. asteroides</i>	Bifido	Bifido-1	ESL0199	5.3	7	558,059	2,167,340	50	59	1,741
<i>Bifidobacterium</i>	<i>B. asteroides</i>	Bifido	Bifido-1	ESL0200	4.7	16	500,320	1,933,421	300	60	1,621
<i>Bifidobacterium</i>	<i>B. indicum</i> / <i>B. coryneforme</i>	Bifido	Bifido-2	ESL0197	1.1	6	1,389,647	1,715,238	300	61	1,408
<i>Gilliamella</i>	<i>G. apicola</i>	Gilliamella	Gilli-1	ESL0178	0.3	18	364,598	2,885,657	200	34	2,602
<i>Gilliamella</i>	<i>G. apis</i>	Gilliamella	Gilli-2	ESL0169	3.9	13	481,163	2,430,778	270	35	2,227
<i>Gilliamella</i>	<i>G. apis</i>	Gilliamella	Gilli-2	ESL0172	1.8	17	374,672	2,685,772	200	34	2,468
<i>Gilliamella</i>	NA ^d	Gilliamella	Gilli-3	ESL0177	2.0	19	953,736	3,086,198	50	35	2,868
<i>Gilliamella</i>	NA	Gilliamella	Gilli-3	ESL0182	1.2	31	255,373	3,537,173	160	35	3,257
<i>Snodgrassella</i>	<i>S. alvi</i>	Snodgrassella	NA	ESL0196	3.7	15	1,281,809	2,446,304	130	41	2,224
<i>Frischella</i>	<i>F. perrara</i>	Frischella	NA	ESL0167	3.1	40	277,847	2,558,525	200	34	2,313
<i>Commensalibacter</i>	<i>Commensalibacter</i> sp.	Commensalibacter	NA	ESL0284	1.3	13	471,180	1,948,862	50	38	1,767

^aBased on 16S rRNA amplicon sequencing.

^bTotal amount of extracted DNA.

^cGene count based on the JGI Microbial Genome Annotation Pipeline.

^dNA, not applicable.

uncertainty generated by a duplicated prophage sequence. Strains ESL0183, ESL0185, and ESL0186 were submitted as complete genomes, with strain ESL0183 having a small plasmid contig of 11.3 kb.

Core phylogenies were generated for the *Lactobacillus*, *Bifidobacterium*, and *Gilliamella* strains, including previously published isolates derived from honey bees, using OrthoFinder (14) for ortholog prediction and RAxML (15) for phylogenetic inference. Based on the phylogenies, the *Lactobacillus* and *Bifidobacterium* strains represent members of previously reported sublineages, whereas two strains of the genus *Gilliamella* (ESL0177 and ESL0182) represent a new sublineage, with strain ESL182 having the largest genome size reported for this genus to date (3.5 Mbp) (Table 1).

Data availability. The complete genome sequences for the strains reported here have been deposited in GenBank under the accession numbers CP029476, CP029544/CP029545, and CP029477, and the whole-genome shotgun projects have been deposited under the accession numbers QGLH00000000, QGLJ00000000, QGLK00000000, QGLL00000000, QGLI00000000, QGLG00000000, QGLQ00000000, QGLN00000000, QGLO00000000, QGLP00000000, QGLR00000000, QGLS00000000, QGLM00000000, and QGLT00000000. Additionally, the genomes were annotated using the JGI Microbial Genome Annotation Pipeline, where they have been deposited under the genome identification numbers 2684622912, 2684622914, 2684622911, 2684622916, 2684622918, 2684622919, 2684622920, 2684622917, 2684622913, 2684622925, 2684622922, 2684622923, 2684622924, 2684622926, 2684622927, 2684622921, and 2756170209.

ACKNOWLEDGMENTS

This work was funded by Human Frontier Science Program (HFSP) Young Investigator grant RGY0077/2016, the European Research Council Starting Grant (ERC-StG) "MicroBeeOme," Swiss National Science Foundation grant 31003A_160345, and the Fondation Herbettes at the University of Lausanne.

REFERENCES

- Kwong WK, Moran NA. 2016. Gut microbial communities of social bees. *Nat Rev Microbiol* 14:374–384. <https://doi.org/10.1038/nrmicro.2016.43>.
- Ellegaard KM, Tamarit D, Javelind E, Olofsson TC, Andersson SGE, Vasquez A. 2015. Extensive intra-phylogroup diversity in lactobacilli and bifidobacteria from the honeybee gut. *BMC Genomics* 16:284. <https://doi.org/10.1186/s12864-015-1476-6>.
- Engel P, Stepanauskas R, Moran NA. 2014. Hidden diversity in honey bee gut symbionts detected by single-cell genomics. *PLoS Genet* 10:e1004596. <https://doi.org/10.1371/journal.pgen.1004596>.
- Engel P, Martinson VG, Moran NA. 2012. Functional diversity within the simple gut microbiota of the honey bee. *Proc Natl Acad Sci U S A* 109:11002–11007. <https://doi.org/10.1073/pnas.1202970109>.

5. Ellegaard KM, Engel P. 2016. Beyond 16S rRNA community profiling: intra-species diversity in the gut microbiota. *Front Microbiol* 7:1475.
6. Engel P, James RR, Koga R, Kwong WK, McFrederick QS, Moran NA. 2013. Standard methods for research on *Apis mellifera* gut symbionts. *J Apiculi Res* 52:1–24. <https://doi.org/10.3896/IBRA.1.52.4.07>.
7. Kesnerova L, Mars RAT, Ellegaard KM, Troilo M, Sauer U, Engel P. 2017. Disentangling metabolic functions of bacteria in the honey bee gut. *PLoS Biol* 15:e2003467. <https://doi.org/10.1371/journal.pbio.2003467>.
8. Olofsson TC, Alsterfjord M, Nilson B, Butler E, Vasquez A. 2014. *Lactobacillus apinorum* sp. nov., *Lactobacillus mellifer* sp. nov., *Lactobacillus mellis* sp. nov., *Lactobacillus melliventris* sp. nov., *Lactobacillus kimbladii* sp. nov., *Lactobacillus helsingborgensis* sp. nov. and *Lactobacillus kullaber-gensis* sp. nov., isolated from the honey stomach of the honeybee *Apis mellifera*. *Int J Syst Evol Microbiol* 64:3109–3119. <https://doi.org/10.1099/ijs.0.059600-0>.
9. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
10. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
11. Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* 5:e11147. <https://doi.org/10.1371/journal.pone.0011147>.
12. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.
13. Carver T, Thomson N, Bleasby A, Berriman M, Parkhill J. 2009. DNAPlotter: circular and linear interactive genome visualization. *Bioinformatics* 25: 119–120. <https://doi.org/10.1093/bioinformatics/btn578>.
14. Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. *Genome Biol* 16:157. <https://doi.org/10.1186/s13059-015-0721-2>.
15. Stamatakis A. 2006. RAXML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22:2688–2690. <https://doi.org/10.1093/bioinformatics/btl446>.