

Prevalence and diversity of *Chlamydiales* and other amoeba-resisting bacteria in domestic drinking water systems

J. Lienard¹, A. Croxatto¹, A. Gervais³, Y. Lévi⁴, J.-F. Loret⁵, K. M. Posfay-Barbe³ and G. Greub^{1,2}

1) Center for Research on Intracellular Bacteria, Institute of Microbiology, Lausanne University Hospital and University of Lausanne, 2) Infectious Diseases Service, Lausanne University Hospital, Lausanne, Switzerland, 3) Children's Hospital of Geneva, University Hospitals of Geneva and Medical School of the University of Geneva, Geneva, Switzerland, 4) University of Paris-Sud XI, Faculty of Pharmacy, Paris and 5) Suez Environnement CIRSEE, Le Pecq, France

Abstract

A growing number of human infections incriminate environmental bacteria that have evolved virulent mechanisms to resist amoebae and use them as a replicative niche. These bacteria are designated amoeba-resisting bacteria (ARB). Despite the isolation of these ARB in various human clinical samples, the possible source of infection remains undetermined in most cases. However, it is known that the ARB *Legionella pneumophila*, for instance, causes a respiratory infection in susceptible hosts after inhalation of contaminated water aerosols from various sources. The *Chlamydiales* order contains many ARB, such as *Parachlamydia acanthamoebae* or *Simkania negevensis*, previously implicated in human respiratory infections with no identified contamination sources. We thus investigated whether domestic water systems are a potential source of transmission of these *Chlamydiales* to humans by using amoebal culture and molecular methods. Other important ARB such as mycobacteria and *Legionella* were also investigated, as were their possible amoebal hosts. This work reports for the first time a very high prevalence and diversity of *Chlamydiales* in drinking water, being detected in 35 (72.9%) of 48 investigated domestic water systems, with members of the *Parachlamydiaceae* family being dominantly detected. Furthermore, various *Legionella* and mycobacteria species were also recovered, some species of which are known to be causal agents of human infections.

© 2016 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

Keywords: Amoebal co-culture, amoebal enrichment, biofilm, *Criblamydiaceae*, *Parachlamydiaceae*

Original Submission: 8 June 2016; **Revised Submission:** 10 October 2016; **Accepted:** 14 October 2016

Article published online: 14 November 2016

Corresponding author: G. Greub, MD, PhD, Center for Research on Intracellular Bacteria (CRIB), Institute of Microbiology, Lausanne University Hospital and University of Lausanne, 1011 Lausanne, Switzerland

E-mail: gilbert.greub@chuv.ch

Introduction

Free-living amoebae are ubiquitous in the environment, especially water. In case of unfavourable growth conditions, such as starvation or desiccation, these protists can exhibit a resistant form, termed cysts. The cyst structure helps the amoebae to survive various disinfection treatments [1–3]. Thus, amoebae may bypass all the barriers present in drinking water treatment

plants [4] and may reach the water distribution system, where they may colonize biofilms and sediments.

Amoebae have been shown to be natural hosts of different bacteria that can resist intracellular killing through several mechanisms [5]. Some of these amoeba-resisting bacteria (ARB) have been shown to reside in the amoebal cyst, where they are protected from biocides and disinfection treatments [6–8]. The evolution of traits that result in bacterial resistance to amoebae may explain the ability of some ARB to also resist other phagocytic cells, such as macrophages [9–12]. The observation that some ARB are able to infect both amoebae and macrophages supports this hypothesis [13,14].

Humans may be exposed to these ARB through various water systems such as cooling towers, humidifier aerosols, drinking water, spas or swimming pools, all of which have previously been shown to be reservoirs of ARB. For instance, the

ARB *Legionella pneumophila* was discovered after an outbreak of pneumonia in 1976 in Philadelphia in which dozens of people were infected by a contaminated air-conditioning system [15]. Breiman *et al.* [16] later showed a correlation between Legionnaire's disease due to *Legionella pneumophila* and the use of showers. Newly discovered ARB are emerging as potential respiratory pathogens, such as *Parachlamydia acanthamoebae* [17] and *Simkania negevensis* [18], both able to replicate in amoebae [7,19,20]. However, the mode of transmission of these *Chlamydia*-related bacteria remains to be determined.

Recently a *Chlamydiales*-specific quantitative PCR was developed and was applied to 422 nasopharyngeal swabs from patients [21]. This study showed that 48 patients were positive for a member of the *Chlamydiales* order, among which 38 corresponded to *Chlamydia*-related bacteria, demonstrating that these bacteria can reach the human respiratory tract.

Thus, in the present work, domestic drinking waters and biofilms from plumbing systems were investigated for the presence of *Chlamydiales* by PCR and culture methods. These samples were also screened for other ARB belonging to the families *Legionellaceae* and *Mycobacteriaceae*, from which several members are established as human pathogens. Finally, the screening of potential amoebal hosts was also performed.

Materials and Methods

Sample

Water ($n = 48$) and biofilm ($n = 48$) samples were collected from 48 different domestic water systems in the regions of Geneva ($n = 37$), Lausanne ($n = 7$) and Sion ($n = 4$), Switzerland. Sampling was performed from September 2010 to August 2011. One litre of first-flow water was first sampled from the shower, filtered through a 0.22 μm membrane, which was then resuspended in 10 mL of filtrated water. The mean temperature of the water was $20.6 \pm 3.8^\circ\text{C}$. Then, using a sterile swab, biofilms were collected from the flexible pipe connected to the shower head (after unscrewing the shower head) and was then resuspended on site in about 3 mL of shower water. Aliquots of 100 μL of concentrated water and 100 μL of resuspended biofilm were kept at -20°C for DNA extraction (Fig. 3) while the samples were processed immediately for analyses.

Screening of ARB with amoebal co-culture

Acanthamoeba castellanii ATCC 30010 was used to cultivate ARB. *A. castellanii* was grown in the rich peptone yeast-extract glucose (PYG) medium [22,23], at 28°C without CO_2 , in 75 cm^2 surface cell culture flasks (Becton Dickinson, Allschwil, Switzerland). Amoebae were collected by centrifugation

($1500 \times g$, 10 minutes) and washed with phosphate-buffered saline and finally resuspended in poor medium Page amoeba saline (PAS) [22,23] to avoid extracellular overgrowth of bacteria. Amoebae were seeded in a 24-well culture microplate (Milian, Wohlen, Switzerland) at 5×10^5 amoebal cells/mL. An aliquot (100 μL) of biofilm or concentrated water sample was then inoculated, and tenfold serial dilutions were performed. The microplates were immediately centrifuged at $1790 \times g$ for 15 minutes, and the cells were incubated for 1 hour at 28°C . Cells were gently washed once with PAS and incubated at 32°C in a humidified atmosphere without CO_2 . Amoebae were observed daily for amoebal lysis, and the co-cultures were reseeded on fresh confluent amoebae in PAS after 7 and 14 days [24]. At day 7 and day 14, 100 μL of each amoebae-containing well was collected and stored at -20°C until DNA extraction.

Screening of amoebae with amoebal enrichment

Nonnutrient agar plates were covered with a solution of live *Escherichia coli* ATCC 25922. About 20 μL of concentrated water or biofilm samples was seeded onto the agar and incubated at 28°C in a humidified atmosphere. Plates were observed daily under an optical microscope for the presence of amoebae. When positive, subcultures were performed [24], and amoebae were collected and frozen at -20°C until DNA extraction.

PCR on water samples and biofilms

DNAs were automatically extracted by the LC automated system (Roche, Rotkreuz, Switzerland) and the MagNA Pure LC DNA isolation kit I (Roche) using 100 μL of water and 100 μL of biofilm sample. For each run of extraction, a negative extraction control was included. Water samples ($n = 48$) and biofilm samples ($n = 48$) were analysed by 16S rRNA gene-directed PCR for the presence of DNA from *Legionella* spp. (Leg225/Leg858 primers [25]), *Mycobacterium* spp. (TB285F/TB264R primers [26]) and *Chlamydiales* (panCh16F2/panCh16R2 primers and panCh16S probe [21]). Finally, amoebae were identified by sequencing a part of the 18S rRNA gene, amplified using the Ami6F1/Ami9R primers [43]. The *Chlamydiales*-specific real-time PCR targeting the 16S rRNA gene was performed as previously described [21]. Briefly, using the primers panCh16F2, panCh16R2 and a probe panCh16S, 5 μL of DNA was analysed in duplicate with 50 cycles consisting of denaturing for 15 seconds at 95°C , annealing for 15 seconds at 67°C and amplification for 15 seconds at 72°C .

When the PCR or quantitative real-time PCR was positive, the PCR product was purified with the MSB Spin PCRapace kit and sequenced with the same primers. In the case of positive samples for mycobacteria with the 16S rRNA PCR, a second PCR targeting the *rpoB* gene and using the primers MycoF/

Mycor was used for precise identification by sequencing [28]. Concerning the PCR products obtained with the *Chlamydiales*-specific real-time PCR, they were purified using the GenElute PCR Clean-Up Kit (Sigma, Buchs, Switzerland), and sequencing was performed with inner primers as described elsewhere [21]. All newly generated nucleotide sequences were submitted to GenBank; the accession numbers may be found in the Supplementary Tables.

PCR on amoebal culture and amoebal enrichment

Amoebal co-culture wells were screened by PCR for the presence of *Legionella* spp., *Mycobacterium* spp. and *Chlamydia*-related bacteria after 1 and 2 weeks of incubation. DNA was extracted from 100 µL of the culture using the Wizard genomic DNA purification kit (Promega, Duedendorf, Switzerland) in the presence of proteinase K (20 mg/mL) following the manufacturer’s protocol for animal tissues. For each run of extraction, a negative extraction control was included. Detection by PCR and sequencing of mycobacteria, *Legionella* and amoebae was performed as described above. For the *Chlamydiales*, the 16SigF/Rp2Chlam primers were used, as described elsewhere [29].

Results

ARB documented in water and biofilm samples

The number of bacteria and amoebae detected in this study are represented in Fig. 1. In addition, for each domestic water system, all bacterial and amoebal species identified by sequencing are presented in Table 1.

***Chlamydiales* species**

Among the 48 domestic water systems investigated, 35 (72.9%) were positive for *Chlamydiales* detected by specific real-time PCR (rtPCR) in the water, the biofilm or both samples (Fig. 1 and Table 1). Sequencing of the rtPCR products gave a sequence of about 200 bp that was used to classify the bacteria at the family level following the criteria of Everett et al. [29]. A total of 55 *Chlamydiales* sequences could be obtained for 33 of these 35 positive households. The classification could be achieved for 51 DNA sequences (Table 1 and Supplementary Table S1), and four remained unclassified. Among these 55 sequences, 28 (50.9%) may correspond to new species-level




FIG. 1. Distribution of type of samples and detection methods for each bacterial groups and amoebae detected. (A) Parts of whole representing number of positive households for *Chlamydiales*, *Legionella*, *Mycobacterium* or amoeba, detected in water, biofilm or both in water and biofilm samples. Corresponding number of positive samples is also indicated. (B) Distribution of detection methods among positive samples (water and biofilm) by PCR only, culture only or both PCR and culture.

TABLE 1. Summary of all *Chlamydiales* spp, *Legionella* spp., *Mycobacterium* spp. and amoebae detected in each water system of 48 households investigated

Household ID	Water	Detection	Biofilm	Detection
	Species		Species	
GE10016	<i>Criblamydiaceae</i> putative species 1 [Chlam]	qP	<i>Criblamydiaceae</i> putative species 2 [Chlam]	qP
GE10027	100% <i>L. waltersii</i> [Legio]	P		
	<i>Criblamydiaceae</i> [Chlam]	qP		
GE10028	<i>Parachlamydiaceae</i> [Chlam]	qP		
	<i>Parachlamydiaceae</i> [Chlam]	qP	<i>Parachlamydiaceae</i> [Chlam]	qP
GE10032	99% uncultured bacterium clone ncd843d07c1 [Legio]	C	95% <i>M. moriokaense</i> or <i>M. barrassiae</i> ^a [Myco]	P
GE10037	Unclassified <i>Chlamydiales</i> [Chlam]	qP		
GE10044	100% <i>Mycobacterium iranicum</i> strain CCUG 52297 ^a [Myco]	C	100% <i>M. gilvum</i> [Myco]	C
	99% <i>H. vermiformis</i> [Amoeba]	C	100% <i>M. phocaicum</i> strain MBWY-1 ^b [Myco]	C
GE10049	<i>Criblamydiaceae</i> [Chlam]	qP		
	100% <i>L. anisa</i> [Legio]	C		
GE10056	<i>Chlamydiales</i> (failed sequencing)	qP	<i>Parachlamydiaceae</i> [Chlam]	qP
GE10061	<i>Criblamydiaceae</i> [Chlam]	qP	<i>Parachlamydiaceae</i> [Chlam]	P
	100% <i>Mycobacterium</i> sp. FI-10135 ^a [Myco]	C	97% <i>M. tusciae</i> ^a [Myco]	P+C
GE10062	97% <i>M. tusciae</i> ^a [Myco]	P		
GE10064	<i>Parachlamydiaceae</i> [Chlam]	qP	<i>Parachlamydiaceae</i> [Chlam]	qP
	Unclassified <i>Chlamydiales</i> [Chlam]	C	Unclassified <i>Chlamydiales</i> [Chlam]	C
GE10068	100% <i>L. taurinensis</i> [Legio]	P+C	100% <i>L. taurinensis</i> [Legio]	P
	99% <i>H. vermiformis</i> [Amoeba]	qP	99% <i>H. vermiformis</i> [Amoeba]	qP
GE10088	<i>Criblamydiaceae</i> putative species 1 [Chlam]	P	<i>Criblamydiaceae</i> putative species 2 [Chlam]	qP
GE10096	98% uncultured bacterium clone F20 [Legio]	qP		
GE10143	<i>Criblamydiaceae</i> [Chlam]	qP	Unclassified <i>Chlamydiales</i> [Chlam]	qP
	<i>Parachlamydiaceae</i> [Chlam]	qP	<i>Parachlamydiaceae</i> [Chlam]	qP
GE10148	98% <i>L. longbeachae</i> [Legio]	C	99% <i>Mycobacterium</i> sp. [Myco]	C
GE10150	<i>Criblamydiaceae</i> putative species 1 [Chlam]	qP	<i>Criblamydiaceae</i> putative species 2 [Chlam]	qP
	<i>Parachlamydiaceae</i> putative species 2 [Chlam]	P		
GE10159	99% <i>M. gordonae</i> ^a [Myco]	qP	100% <i>L. waltersii</i> [Legio]	P
	<i>Parachlamydiaceae</i> [Chlam]	P		
GE10160	100% <i>H. vermiformis</i> [Amoeba]	P		
GE10170	100% uncultured <i>Legionella</i> sp. [Legio]	qP	<i>Parachlamydiaceae</i> [Chlam]	qP
	<i>Parachlamydiaceae</i> [Chlam]	C	100% <i>H. vermiformis</i> [Amoeba]	P+C
GE10174	100% <i>L. pneumophila</i> [Legio]	C		
	97% <i>M. tusciae</i> ^a [Myco]	qP	<i>Parachlamydiaceae</i> [Chlam]	qP
GE10175	Failed sequencing [Chlam]	C	99% <i>L. waltersii</i> [Legio]	P
	100% <i>L. waltersii</i> [Legio]	P	99% uncultured eukaryote clone TKR07M.106 [Amoeba]	C
GE10179	98% <i>L. gratiana</i> [Legio]	C	100% <i>H. vermiformis</i> [Amoeba]	P
GE11050		P	99% <i>L. beliardensis</i> [Legio]	P
GE11064	99% <i>H. vermiformis</i> [Amoeba]	qP	97% <i>M. tusciae</i> ^a [Myco]	P+C
	<i>Chlamydiales</i> (failed sequencing) [Chlam]	P	100% <i>H. vermiformis</i> [Amoeba]	P
GE11093	94% <i>M. neoaurum</i> ^a [Myco]	P		
	100% <i>H. vermiformis</i> [Amoeba]	P+C		
GE11103	<i>Criblamydiaceae</i> [Chlam]	qP	<i>Parachlamydiaceae</i> putative species 1 [Chlam]	qP
	98% uncultured bacterium clone IC227246 [Legio]	P	<i>Parachlamydiaceae</i> putative species 2 [Chlam]	qP
GE11112	<i>Parachlamydiaceae</i> [Chlam]	qP		
	98% <i>L. fallonii</i> strain LLAP10 [Legio]	P	<i>Chlamydiales</i> (failed sequencing) [Chlam]	qP
HE20032	99% <i>M. abscessus</i> subsp. <i>bolletii</i> 50594 [Myco]	C	99% <i>Stenamoeba</i> CRIB 68 [Amoeba]	P+C
	<i>Parachlamydiaceae</i> putative species 1 [Chlam]	qP		
HE20036	100% <i>L. pneumophila</i> [Legio]	C		
	98% uncultured bacterium clone nbu179b03c1 [Legio]	P		
HE21001	99% <i>H. vermiformis</i> [Amoeba]	P+C		
HE21011	<i>Parachlamydiaceae</i> [Chlam]	qP	100% <i>L. waltersii</i> [Legio]	P
HE21012	<i>Criblamydiaceae</i> [Chlam]	qP	100% <i>M. chelonae</i> ^a [Myco]	P
HE21023	<i>Waddliaceae</i> [Chlam]	qP	<i>Parachlamydiaceae</i> species 2 [Chlam]	qP
HE21032	<i>Criblamydiaceae</i> [Chlam]	qP	100% <i>L. pneumophila</i> [Legio]	C
VS30003	<i>Parachlamydiaceae</i> [Chlam]	qP	100% <i>H. vermiformis</i> [Amoeba]	P+C
VS30013	100% <i>L. waltersii</i> [Legio]	C	<i>Waddliaceae</i> [Chlam]	qP
VS30044	<i>Criblamydiaceae</i> [Chlam]	qP	<i>Criblamydiaceae</i> [Chlam]	qP
	<i>Parachlamydiaceae</i> putative species 1 [Chlam]	qP	<i>Criblamydiaceae</i> [Chlam]	qP
VS30055	<i>Parachlamydiaceae</i> putative species 1 [Chlam]	qP	88% <i>Stenamoeba amazonica</i> strain P119 [Amoeba]	C
			<i>Simkaniaceae</i> [Chlam]	qP
			<i>Parachlamydiaceae</i> [Chlam]	qP
			<i>Chlamydiales</i> (failed sequencing) [Chlam]	qP
			<i>Parachlamydiaceae</i> putative species 2 [Chlam]	qP
			100% <i>Mycobacterium</i> sp. [Myco]	C

TABLE I. Continued

Household ID	Water		Biofilm	
	Species	Detection	Species	Detection
VS31006	<i>Parachlamydiaceae</i> [Chlam]	qP	100% <i>H. vermiformis</i> [Amoeba] <i>Chlamydiales</i> (failed sequencing) [Chlam]	C qP

Percentages of sequence identity with most similar GenBank sequence (for *Legionella*, mycobacteria or amoebae) or classification at the family or family-level lineage (for *Chlamydiales*) are indicated.
 C, culture; P, PCR; qP, quantitative real-time PCR; [Chlam], *Chlamydiales*; [Legio], *Legionella*; [Myco], *Mycobacterium*.
^aSpecies further identified by sequencing *rpoB* gene.
^bSpecies found within amoeba by nonnutrient agar screening.

lineages if fully characterized because the sequences exhibit a similarity with a previously reported species below 97% [17]. Figure 2 illustrates the number of bacteria detected in biofilm or water samples, based on the number of 16S rRNA gene copies quantified by the *Chlamydiales*-specific rtPCR. The majority of the sequences corresponded to members of the *Parachlamydiaceae* family ($n = 30$ sequences), which were detected in 20 different water systems. *Criblamydiaceae* DNAs were also amplified (18 sequences from 14 different domestic water systems) as well as two sequences of the *Waddliaceae* family and one sequence from the *Simkaniaceae* family. The highest number of bacteria was detected in biofilms and corresponded to members of the *Parachlamydiaceae* family (Fig. 2).




FIG. 2. *Chlamydiales* 16S ribosomal RNA gene copy number detected by *Chlamydiales*-specific real-time PCR in water and biofilm samples. Each symbol represents sample positive for *Chlamydiales* detected by specific quantitative PCR and its corresponding gene copy number expressed per litre of sample.

Legionella species

In total, the presence of *Legionella* was found in 21 (43.8%) drinking water systems. *Legionella* was detected by PCR and/or by amoebal co-culture (but never as an amoebal endosymbiont of amoebae grown using the amoebal enrichment method). The

results are shown in Table I, and the identification of *Legionella* species is detailed in Supplementary Table S2. By PCR and/or amoebal co-culture, *Legionella* was detected in 29 samples (ten biofilms and 19 waters); it corresponded to 15 different species (Table I and Supplementary Table S2). The most common species were *Legionella waltersii* (present in eight water systems) and *L. pneumophila* (present in three water systems).

Mycobacterium species

Using PCR and amoebal methods, 15 (31.3%) domestic water systems were positive for *Mycobacterium* species such as *Mycobacterium gordonae*, *chelonae* or *mucogenicum*. The results are summarized in Table I, and complete identification can be found in Supplementary Table S3. Of particular note, two different mycobacteria (*M. iranicum* strain CCUG52297 and *M. phocaicum*) were found within the amoeba *Hartmannella vermiformis*, recovered from water and biofilm samples of the same domestic water system (GEI0037).

Amoebae isolated by amoebal enrichment and/or detected by PCR

Using both PCR and amoebal enrichment, the presence of amoebae was documented in 18 (37.5%) domestic water systems (Fig. 1). Amoebae were present in water and/or biofilm samples (Fig. 1), with *Hartmannella vermiformis* being predominantly detected in 16 water systems (Table I). Two *Stenamoeba* species were also isolated from two different biofilms, one being a potential new amoebal species. Finally, in a biofilm already positive by PCR for *H. vermiformis*, an uncultured eukaryote strain related to the *Prostelium nocturnum* amoeba was isolated by culture (water system GEI0174). The complete identification of amoebae per type of sample can be found in Supplementary Table S4.

Discussion

In this study, the presence of ARB belonging to the *Chlamydiales* order as well as to the *Legionellaceae* and *Mycobacteriaceae*

families was investigated using amoebal culture methods and PCR on water and biofilm samples collected from domestic water systems of 48 different households. Overall, 39 (81.3%) of the investigated domestic water systems were positive for the presence of a *Chlamydiales*, a *Legionellaceae* and/or a *Mycobacteriaceae*. In 18 (46.2%) of these systems, the bacterium was detected by culture. In the other systems, the bacteria were only detected by PCR.

A *Chlamydiales*-specific rtPCR was used and allowed for the first time to observe such a high prevalence and diversity of *Chlamydiales* in domestic drinking water. The high sensitivity of the rtPCR allowed the detection of a *Chlamydiales* in 35 (72.9%) different domestic water systems, corresponding to members of at least four different family-level lineages of the *Chlamydiales* order. The dominant family-level lineage was the *Parachlamydiaceae* family. Members of the *Parachlamydiaceae* family have been frequently isolated from environmental samples [30,32,33]. The high prevalence of strains belonging to this family compared to other *Chlamydia*-related bacteria was also previously observed when using the same *Chlamydiales*-specific

rtPCR on nasopharyngeal swabs taken from children [21]. The second family detected in 14 water systems was the *Criblamydiaceae*. The presence of *Criblamydiaceae* species in water and biofilm samples was not surprising because these bacteria have been previously isolated from water and/or sediment samples [33–35]. This result is particularly interesting because serologic evidence indicates that *Criblamydiaceae* may be associated with cases of pneumonia (Lienard et al., personal communication).

We also detected two members of the *Waddliaceae* family. To our knowledge, this is the first documentation of *Waddliaceae* in drinking water systems. Although the bacterium *Waddlia chondrophila* was previously associated with human and bovine hosts [36–39], its potential presence in water was suggested by its ability to also grow and survive in amoebae [40,41]. Only one sequence corresponding to the *Simkaniaceae* family was detected, which did not correspond to the species *Simkania negevensis*. This result contrasts with a previous work where *S. negevensis* was detected by PCR in the majority of tap water samples [42]. However, this latter study was performed




FIG. 3. Protocol of collection and processing of water and biofilm samples. Samples were collected from distal water conduit after removal of shower head. Cold water was concentrated 100× by filtration, and biofilms swabs were resuspended in 3 mL of collected shower water. Samples were then directly inoculated in culture, or DNA was extracted for direct PCR approaches. After amoebal co-culture or enrichment methods, total DNA was extracted from culture well and PCR performed. For positive results, bacterial or amoebal strains were identified by sequencing.

in Israel, where the microbial ecology and drinking water treatment processes may be different from those in Switzerland.

Among the 55 sequences of *Chlamydiales* bacteria obtained in this work, only two corresponded to bacteria currently grown in our laboratory, which indicated that the sequences obtained here did not result from a PCR contamination. Overall, 28 different sequences showed less than 97% similarity with a previously reported species. Considering this 97% cutoff [17,29], these latter 28 sequences may correspond to putative new species, highlighting the broad and underestimated biodiversity of the *Chlamydiales* order [21]. This report suggests that man-made drinking water could represent an important ecological niche for *Chlamydiales* bacteria.

No *Chlamydiales* bacterium was recovered by amoebal co-culture in this study. Another study on drinking water failed to detect any *Chlamydiales*, either by amoebal co-culture with *A. castellanii* or by classical PCR [43]. Kahane et al. [42] detected *Simkania negevensis* in tap water but only by PCR and membrane immunoassay. In the present work, the *Chlamydiales*-specific quantitative PCR, which is more sensitive than regular PCR, revealed the common occurrence of *Chlamydiales* DNA in domestic drinking water systems. The growth of *Chlamydiales* bacteria from environmental samples could have been restricted here by the overgrowth of other environmental bacteria within the co-cultures in *A. castellanii*. Furthermore, in some cases, *Chlamydiales* bacteria were probably initially dead or not cultivable. The amoebal co-culture using *A. castellanii* was previously shown to be effective to recover *Chlamydiales*, including *Criblamydiaceae* and *Parachlamydiaceae* [33,34], but is clearly inadequate to grow all *Chlamydiales*. Indeed, considering the large biodiversity of the *Chlamydiales* order highlighted in the present study, only a few members have been isolated by amoebal co-culture [27,30,33,34,44]. In addition, a restricted amoebal host spectrum has already been shown for several *Chlamydiales* bacteria [20,40,45,46], which suggests that multiple amoebal strains should ideally be used to recover a higher biodiversity of these strictly intracellular bacteria in culture. In this work, an *Acanthamoeba* species was used, which is more suitable for the amoebal co-culture method, as it is less prone to encystment compared to *Hartmannella* spp. Furthermore, *Acanthamoeba* spp. are known to be permissive to a large number of bacteria [7,8,43,44,47,48]. Thus, other amoebae such as *Hartmannella* and *Naegleria* should also be included in future studies. Finally, several growth parameters such as temperature and media can also be optimized to increase the number of recovered ARB.

Legionella waltersii, which was previously associated with severe pneumonia [49], was the most prevalent species, followed

by *L. pneumophila*, among all *Legionella* found in this study. In addition, *Legionella* species considered as potential respiratory pathogens such as *L. anisa* [50–52], *L. longbeachae* [51,53,54] or *L. fallonii* [50] were also recovered. In all water systems positive for *L. pneumophila*, the amoeba *H. vermiformis* was systematically isolated by amoebal enrichment, supporting the importance of this amoeba as a reservoir for *L. pneumophila*.

In addition, various nontuberculous mycobacteria have been recovered using amoebal co-culture and amoebal enrichment, including several human pathogens, such as *M. mucogenicum* [55] and *M. chelonae*, which have mainly been shown to cause respiratory [56,57] and soft tissue [58] infection. *M. gordonae*, which is also sometimes considered pathogenic [59–63], has been previously isolated from drinking water [43,64] and was isolated in our study from water and biofilm samples. Other nontuberculous mycobacteria were also recovered in the present work, including *M. conceptionense* [65–68], *M. barrassiae* [69] and *M. neoaurum* [70–72]. Finally, one of the two mycobacteria recovered within the amoeba *H. vermiformis* was *M. iranicum*. This species was recently described as a new human pathogen; it was isolated from clinical samples such as cerebrospinal fluid and sputum samples from patients from different continents [73,74]. However, the source of infection has not been determined for these previously reported cases; drinking water should thus be considered.

Using amoebal enrichment and PCR, amoebae were documented in 18 systems (37.5%). Although the number of recovered amoebae is particularly variable between studies [75], the number of amoebae cultivated in this study ($n = 15$) is higher compared to a previous study using the same culture method [43]. However, the difference of water temperatures between the present and the previous study, with mean temperatures of 20.6°C and 56°C, respectively, may explain these results. Most of the amoebae isolated in this work corresponded to *H. vermiformis*, which is congruent with a previous investigation of drinking water by amoebal enrichment [43].

In conclusion, the current study highlighted the large colonization of drinking water points of use by ARB and amoebae. This work also demonstrated the common occurrence and large biodiversity of *Chlamydiales* bacteria in drinking water. Thus, drinking water represents a potential infection source for some *Chlamydia*-related bacteria. Because *Parachlamydia acanthamoebae* is associated with respiratory infections [17], the common occurrence of *Parachlamydiaceae* observed here is important in terms of public health. Larger prospective studies including different settings are needed to better investigate the role of domestic water systems or other systems generating aerosols, such as cooling towers, in the transmission of *Chlamydiales* to humans and other susceptible hosts.

Acknowledgements

We thank S. Aeby for technical help. We also thank A. L'Huillier, A. Hagerman, C. Salomon, S. Meier and S. Heiniger for their invaluable help in collecting water and biofilm samples. Supported in part by Suez-Environnement and the Association Nationale de la Recherche Technologique (ANRT) (CIFRE 680/2008).

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.nmni.2016.10.003>.

Conflict of Interest

None declared.

References

- [1] Coulon C, Collignon A, McDonnell G, Thomas V. Resistance of *Acanthamoeba* cysts to disinfection treatments used in health care settings. *J Clin Microbiol* 2010;48:2689–97.
- [2] Lloyd D, Turner NA, Khunkitti W, Hann AC, Furr JR, Russell AD. Encystation in *Acanthamoeba castellanii*: development of biocide resistance. *J Eukaryot Microbiol* 2001;48:11–6.
- [3] Mogoia E, Bodet C, Morel F, Rodier MH, Legube B, Hechard Y. Cellular response of the amoeba *Acanthamoeba castellanii* to chlorine, chlorine dioxide, and monochloramine treatments. *Appl Environ Microbiol* 2011;77:4974–80.
- [4] Loret JF, Greub G. Free-living amoebae: biological by-passes in water treatment. *Int J Hyg Environ Health* 2010;213:167–75.
- [5] Greub G, Raoult D. Microorganisms resistant to free-living amoebae. *Clin Microbiol Rev* 2004;17:413–33.
- [6] Kilvington S, Price J. Survival of *Legionella pneumophila* within cysts of *Acanthamoeba polyphaga* following chlorine exposure. *J Appl Bacteriol* 1990;68:519–25.
- [7] Kahane S, Dvoskin B, Mathias M, Friedman MG. Infection of *Acanthamoeba polyphaga* with *Simkania negevensis* and *S. negevensis* survival within amoebal cysts. *Appl Environ Microbiol* 2001;67:4789–95.
- [8] Steinert M, Birkness K, White E, Fields B, Quinn F. *Mycobacterium avium* bacilli grow saprozoically in coculture with *Acanthamoeba polyphaga* and survive within cyst walls. *Appl Environ Microbiol* 1998;64:2256–61.
- [9] Cirillo JD, Falkow S, Tompkins LS. Growth of *Legionella pneumophila* in *Acanthamoeba castellanii* enhances invasion. *Infect Immun* 1994;62:3254–61.
- [10] Cirillo JD, Falkow S, Tompkins LS, Bermudez LE. Interaction of *Mycobacterium avium* with environmental amoebae enhances virulence. *Infect Immun* 1997;65:3759–67.
- [11] Molmeret M, Horn M, Wagner M, Santic M, Abu Kwaik Y. Amoebae as training grounds for intracellular bacterial pathogens. *Appl Environ Microbiol* 2005;71:20–8.
- [12] Salah IB, Ghigo E, Drancourt M. Free-living amoebae, a training field for macrophage resistance of mycobacteria. *Clin Microbiol Infect* 2009;15:894–905.
- [13] Goy G, Croxatto A, Greub G. *Waddlia chondrophila* enters and multiplies within human macrophages. *Microbes Infect* 2008;10:556–62.
- [14] Horwitz MA, Silverstein SC. Legionnaires' disease bacterium (*Legionella pneumophila*) multiples intracellularly in human monocytes. *J Clin Invest* 1980;66:441–50.
- [15] Fraser DW, Tsai TR, Orenstein W, Parkin WE, Beecham HJ, Sharrar RG, et al. Legionnaires' disease: description of an epidemic of pneumonia. *N Engl J Med* 1977;297:1189–97.
- [16] Breiman RF, Fields BS, Sanden GN, Volmer L, Meier A, Spika JS. Association of shower use with Legionnaires' disease. Possible role of amoebae. *JAMA* 1990;263:2924–6.
- [17] Greub G. *Parachlamydia acanthamoebae*, an emerging agent of pneumonia. *Clin Microbiol Infect* 2009;15:18–28.
- [18] Friedman MG, Dvoskin B, Kahane S. Infections with the *Chlamydia*-like microorganism *Simkania negevensis*, a possible emerging pathogen. *Microbes Infect* 2003;5:1013–21.
- [19] Greub G, La Scola B, Raoult D. *Parachlamydia acanthamoeba* is endosymbiotic or lytic for *Acanthamoeba polyphaga* depending on the incubation temperature. *Ann N Y Acad Sci* 2003;990:628–34.
- [20] Hayashi Y, Nakamura S, Matsuo J, Fukumoto T, Yoshida M, Takahashi K, et al. Host range of obligate intracellular bacterium *Parachlamydia acanthamoebae*. *Microbiol Immunol* 2010;54:707–13.
- [21] Lienard J, Croxatto A, Aeby S, Jaton K, Posfay-Barbe K, Gervais A, et al. Development of a new *Chlamydiales*-specific real-time PCR and its application to respiratory clinical samples. *J Clin Microbiol* 2011;49:2637–42.
- [22] Greub G, Raoult D. Crescent bodies of *Parachlamydia acanthamoeba* and its life cycle within *Acanthamoeba polyphaga*: an electron micrograph study. *Appl Environ Microbiol* 2002;68:3076–84.
- [23] Rowbotham TJ. Isolation of *Legionella pneumophila* from clinical specimens via amoebae, and the interaction of those and other isolates with amoebae. *J Clin Pathol* 1983;36:978–86.
- [24] Lienard J, Greub G. Discovering new pathogens: amoebae as tools to isolate amoeba-resisting microorganisms from environmental samples. In: Sen K, Ashbolt NJ, editors. *Environmental microbiology: current technology and water applications*. Norfolk, UK: Caister Academic Press; 2011. p. 143–62.
- [25] Miyamoto H, Yamamoto H, Arima K, Fujii J, Maruta K, Izu K, et al. Development of a new seminested PCR method for detection of *Legionella* species and its application to surveillance of legionellae in hospital cooling tower water. *Appl Environ Microbiol* 1997;63:2489–94.
- [26] Kirschner P, Springer B, Vogel U, Meier A, Wrede A, Kiekenbeck M, et al. Genotypic identification of mycobacteria by nucleic acid sequence determination: report of a 2-year experience in a clinical laboratory. *J Clin Microbiol* 1993;31:2882–9.
- [27] Lienard J, Croxatto A, Prod'hom G, Greub G. *Estrella lausannensis*, a new star in the *Chlamydiales* order. *Microbes Infect* 2011;13:1232–41.
- [28] Ben Salah I, Adekambi T, Raoult D, Drancourt M. *rpoB* sequence-based identification of *Mycobacterium avium* complex species. *Microbiology* 2008;154(Pt 12):3715–23.
- [29] Everett KD, Bush RM, Andersen AA. Emended description of the order *Chlamydiales*, proposal of *Parachlamydiaceae* fam. nov. and *Simkaniaceae* fam. nov., each containing one monotypic genus, revised taxonomy of the family *Chlamydiaceae*, including a new genus and five new species, and standards for the identification of organisms. *Int J Syst Bacteriol* 1999;49:415–40.
- [30] Birtles RJ, Rowbotham TJ, Storey C, Marrie TJ, Raoult D. *Chlamydia*-like obligate parasite of free-living amoebae. *Lancet* 1997;349(9056):925–6.

- [32] Fritsche TR, Gautom RK, Seyedirashti S, Bergeron DL, Lindquist TD. Occurrence of bacterial endosymbionts in *Acanthamoeba* spp. isolated from corneal and environmental specimens and contact lenses. *J Clin Microbiol* 1993;31:1122–6.
- [33] Corsaro D, Feroldi V, Saucedo G, Ribas F, Loret JF, Greub G. Novel *Chlamydiales* strains isolated from a water treatment plant. *Environ Microbiol* 2009;11:188–200.
- [34] Thomas V, Casson N, Greub G. *Criblamydia sequanensis*, a new intracellular *Chlamydiales* isolated from Seine river water using amoebal coculture. *Environ Microbiol* 2006;8:2125–35.
- [35] Corsaro D, Venditti D. Detection of *Chlamydiae* from freshwater environments by PCR, amoeba coculture and mixed coculture. *Res Microbiol* 2009;160:547–52.
- [36] Dilbeck PM, Evermann JF, Crawford TB, Ward AC, Leathers CW, Holland CJ, et al. Isolation of a previously undescribed rickettsia from an aborted bovine fetus. *J Clin Microbiol* 1990;28:814–6.
- [37] Rurangirwa FR, Dilbeck PM, Crawford TB, McGuire TC, McElwain TF. Analysis of the 16S rRNA gene of micro-organism WSU 86-1044 from an aborted bovine foetus reveals that it is a member of the order *Chlamydiales*: proposal of *Waddliaceae* fam. nov., *Waddlia chondrophila* gen. nov., sp. nov. *Int J Syst Bacteriol* 1999;49:577–81.
- [38] Haider S, Collingro A, Walochnik J, Wagner M, Horn M. *Chlamydia*-like bacteria in respiratory samples of community-acquired pneumonia patients. *FEMS Microbiol Lett* 2008;281:198–202.
- [39] Goy G, Croxatto A, Posfay-Barbe KM, Gervais A, Greub G. Development of a real-time PCR for the specific detection of *Waddlia chondrophila* in clinical samples. *Eur J Clin Microbiol Infect Dis* 2009;28:1483–6.
- [40] Michel R, Steinert M, Zöller L, Hauröder B, Henning K. Free-living amoebae may serve as hosts for the *Chlamydia*-like bacterium *Waddlia chondrophila* isolated from an aborted bovine foetus. *Acta Protozoologica* 2004;43:37–42.
- [41] Goy G, Greub G. Antibiotic susceptibility of *Waddlia chondrophila* in *Acanthamoeba castellanii* amoebae. *Antimicrob Agents Chemother* 2009;53:2663–6.
- [42] Kahane S, Platzner N, Dvoskin B, Itzhaki A, Friedman MG. Evidence for the presence of *Simkania negevensis* in drinking water and in reclaimed wastewater in Israel. *Appl Environ Microbiol* 2004;70:3346–51.
- [43] Thomas V, Herrera-Rimann K, Blanc DS, Greub G. Biodiversity of amoebae and amoeba-resisting bacteria in a hospital water network. *Appl Environ Microbiol* 2006;72:2428–38.
- [44] Collingro A, Poppert S, Heinz E, Schmitz-Esser S, Essig A, Schweikert M, et al. Recovery of an environmental *Chlamydia* strain from activated sludge by co-cultivation with *Acanthamoeba* sp. *Microbiology* 2005;151(Pt 1):301–9.
- [45] Michel R, Müller KD, Zöller L, Walochnik J, Hartmann M, Schmid EN. Free-living amoebae serve as a host for the *Chlamydia*-like bacterium *Simkania negevensis*. *Acta Protozoologica* 2005;44:113–21.
- [46] Coulon C, Eterpi M, Greub G, Collignon A, McDonnell G, Thomas V. Amoebal host range, host-free survival and disinfection susceptibility of environmental *Chlamydiae* as compared to *Chlamydia trachomatis*. *FEMS Immunol Med Microbiol* 2012;64:364–73.
- [47] Thomas V, Loret JF, Jousset M, Greub G. Biodiversity of amoebae and amoeba-resisting bacteria in a drinking water treatment plant. *Environ Microbiol* 2008;10:2728–45.
- [48] Corsaro D, Pages GS, Catalan V, Loret JF, Greub G. Biodiversity of amoebae and amoeba-associated bacteria in water treatment plants. *Int J Hyg Environ Health* 2010;213:158–66.
- [49] König C, Hebestreit H, Valenza G, Abele-Horn M, Speer CP. *Legionella waltersii*—a novel cause of pneumonia? *Acta Paediatr* 2005;94:1505–7.
- [50] McNally C, Hackman B, Fields BS, Plouffe JF. Potential importance of *Legionella* species as etiologies in community acquired pneumonia (CAP). *Diagn Microbiol Infect Dis* 2000;38:79–82.
- [51] Yu VL, Plouffe JF, Pastoris MC, Stout JE, Schousboe M, Widmer A, et al. Distribution of *Legionella* species and serogroups isolated by culture in patients with sporadic community-acquired legionellosis: an international collaborative survey. *J Infect Dis* 2002;186:127–8.
- [52] Fenstersheib MD, Miller M, Diggins C, Liska S, Detwiler L, Werner SB, et al. Outbreak of Pontiac fever due to *Legionella anisa*. *Lancet* 1990;336(8706):35–7.
- [53] Grove DI, Lawson PJ, Burgess JS, Moran JL, O'Fathartaigh MS, Winslow WE. An outbreak of *Legionella longbeachae* infection in an intensive care unit? *J Hosp Infect* 2002;52:250–8.
- [54] Phares CR, Wangroongsarb P, Chantra S, Paveenkitiporn W, Tondella ML, Benson RF, et al. Epidemiology of severe pneumonia caused by *Legionella longbeachae*, *Mycoplasma pneumoniae*, and *Chlamydia pneumoniae*: 1-year, population-based surveillance for severe pneumonia in Thailand. *Clin Infect Dis* 2007;45:e147–55.
- [55] Adekambi T. *Mycobacterium mucogenicum* group infections: a review. *Clin Microbiol Infect* 2009;15:911–8.
- [56] Goto T, Hamaguchi R, Maeshima A, Oyama Y, Kato R. Pulmonary resection for *Mycobacterium chelonae* infection. *Ann Thorac Cardiovasc Surg* 2012;18:128–31.
- [57] Singh N, Yu VL. Successful treatment of pulmonary infection due to *Mycobacterium chelonae*: case report and review. *Clin Infect Dis* 1992;14:156–61.
- [58] Mullin D, Jothi S, Healy D. *Mycobacterium chelonae* infections involving the head and neck. *Ann Otol Rhinol Laryngol* 2009;118:714–20.
- [59] Weinberger M, Berg SL, Feuerstein IM, Pizzo PA, Witebsky FG. Disseminated infection with *Mycobacterium gordonae*: report of a case and critical review of the literature. *Clin Infect Dis* 1992;14:1229–39.
- [60] Jarikre LN. Case report: disseminated *Mycobacterium gordonae* infection in a nonimmunocompromised host. *Am J Med Sci* 1991;302:382–4.
- [61] Asija A, Prasad A, Eskridge E. Disseminated *Mycobacterium gordonae* infection in an immunocompetent host. *Am J Ther* 2011;18:e75–7.
- [62] Pinho L, Santos J, Oliveira G, Pestana M. *Mycobacterium gordonae* urinary infection in a renal transplant recipient. *Transpl Infect Dis* 2009;11:253–6.
- [63] den Broeder AA, Vervoort G, van Assen S, Verduyn Lunel F, de Lange WC, de Sevaux RG. Disseminated *Mycobacterium gordonae* infection in a renal transplant recipient. *Transpl Infect Dis* 2003;5:151–5.
- [64] Fischeder R, Schulze-Robbecke R, Weber A. Occurrence of mycobacteria in drinking water samples. *Zentralbl Hyg Umweltmed* 1991;192:154–8.
- [65] Adekambi T, Stein A, Carvajal J, Raoult D, Drancourt M. Description of *Mycobacterium conceptionense* sp. nov., a *Mycobacterium fortuitum* group organism isolated from a posttraumatic osteitis inflammation. *J Clin Microbiol* 2006;44:1268–73.
- [66] Shojaei H, Hashemi A, Heidarieh P, Ataei B, Naser AD. Pulmonary and extrapulmonary infection caused by *Mycobacterium conceptionense*: the first report from Iran. *JRSM Short Rep* 2011;2:31.
- [67] Thibeaut S, Levy PY, Pelletier ML, Drancourt M. *Mycobacterium conceptionense* infection after breast implant surgery, France. *Emerg Infect Dis* 2010;16:1180–1.
- [68] Liao CH, Lai CC, Huang YT, Chou CH, Hsu HL, Hsueh PR. Subcutaneous abscess caused by *Mycobacterium conceptionense* in an immunocompetent patient. *J Infect* 2009;58:308–9.
- [69] Adekambi T, Raoult D, Drancourt M. *Mycobacterium barrassiae* sp. nov., a *Mycobacterium moriokaense* group species associated with chronic pneumonia. *J Clin Microbiol* 2006;44:3493–8.
- [70] Morimoto Y, Chan ED, Heifets L, Routes JM. Pulmonary infection with *Mycobacterium neoaurum* identified by 16S ribosomal DNA sequence. *J Infect* 2007;54:e227–31.
- [71] Zanetti S, Faeda R, Fadda G, Dupre I, Mollicotti P, Ortu S, et al. Isolation and identification of *Mycobacterium neoaurum* from a patient with urinary infection. *New Microbiol* 2001;24:189–92.

- [72] Washer LL, Riddell Jt, Rider J, Chenoweth CE. *Mycobacterium neoaurum* bloodstream infection: report of 4 cases and review of the literature. *Clin Infect Dis* 2007;45:e10–3.
- [73] Shojaei H, Daley C, Gitti Z, Hashemi A, Heidarieh P, Moore ER, et al. *Mycobacterium iranicum* sp. nov., a rapidly growing scotochromogenic species isolated from clinical specimens on three different continents. *Int J Syst Evol Microbiol* 2013;63(Pt 4):1383–9.
- [74] Balakrishnan N, Tortoli E, Engel SL, Breitschwerdt EB. Isolation of a novel strain of *Mycobacterium iranicum* from a woman in the United States. *J Clin Microbiol* 2013;51:705–7.
- [75] Thomas JM, Ashbolt NJ. Do free-living amoebae in treated drinking water systems present an emerging health risk? *Environ Sci Technol* 2011;45:860–9.