

Assessment of Diesel exhaust particulate exposure and surface characteristics in association with levels of oxidative stress biomarkers

¹Setyan A., ¹Sauvain J.-J., ¹Riediker M., ²Rossi M., ¹Guillemin M.

Institut universitaire romand de Santé au Travail¹, Ecole Polytechnique Fédérale de Lausanne²

Exposure to PM₁₀ and PM_{2.5} (particulate matter with aerodynamic diameter smaller than 10 µm and 2.5 µm, respectively) is associated with a range of adverse health effects, including cancer, pulmonary and cardiovascular diseases. Surface characteristics (chemical reactivity, surface area) are considered of prime importance to understand the mechanisms which lead to harmful effects. A hypothetical mechanism to explain these adverse effects is the ability of components (organics, metal ions) adsorbed on these particles to generate Reactive Oxygen Species (ROS), and thereby to cause oxidative stress in biological systems (Donaldson *et al.*, 2003). ROS can attack almost any cellular structure, like DNA or cellular membrane, leading to the formation of a wide variety of degradation products which can be used as a biomarker of oxidative stress.

The aim of the present research project is to test whether there is a correlation between the exposure to Diesel Exhaust Particulate (DEP) and the oxidative stress status. For that purpose, a survey has been conducted in real occupational situations where workers were exposed to DEP (bus depots).

Different exposure variables have been considered:

- particulate number, size distribution and surface area (SMPS);
- particulate mass - PM_{2.5} and PM₄ (gravimetry);
- elemental and organic carbon (coulometry);
- total adsorbed heavy metals - iron, copper, manganese (atomic adsorption);
- surface functional groups present on aerosols (Knudsen flow reactor). (Demirdjian *et al.*, 2005).

Several biomarkers of oxidative stress (8-hydroxy-2'-deoxyguanosine and several aldehydes) have been determined either in urine or serum of volunteers.

Results obtained during the sampling campaign in several bus depots indicated that the occupational exposure to particulates in these places was rather low (40-50 µg/m³ for PM₄). Size distributions indicated that particles are within the nanometric range. Surface characteristics of sampled particles varied strongly, depending on the bus depot. They were usually characterized by high carbonyl and low acidic sites content.

Among the different biomarkers which have been analyzed within the framework of this study, mean levels of 8-hydroxy-2'-deoxyguanosine and several aldehydes (hexanal, heptanal, octanal, nonanal) increased during two consecutive days of exposure for non-smokers.

In order to bring some insight into the relation between the particulate characteristics and the formation of ROS by-products, biomarkers levels will be discussed in relation with exposure variables.

This project is financed by the Swiss State Secretariat for Education and Research. It is conducted within the framework of the COST Action 633 "Particulate Matter – Properties Related to Health Effects".

Demirdjian B., Rossi M. J. (2005). *Atmos. Chem. Phys. Discuss.*, 5, 607 – 654.

Donaldson K., Stone V., Borm P. J., Jimenez L. A., Gilmour P. S., Schins R. P., Knaapen A. M., Rahman I., Faux S. P., Brown D. M., MacNee W. (2003). *Free Radical Biol. Med.*, 34, 1369-1382.



Comité d'organisation 2008

Yvan Arsenijevic, Thérapie génique et biologie des cellules souches, Hôp. Ophtal.

Yann Barrandon, Dynamique des cellules souches, EPFL

Vassily Hatzimanikatis, Biotechnologie computationnelle des systèmes, EPFL

Jeffrey Hubbell, Médecine régénérative et de pharmacobiologie, EPFL

Brigitte Jolles-Haeberli, Hôpital Orthopédique

Thierry Pedrazzini, Médecine interne, CHUV

Ivan Stamenkovic, Institut de pathologie, CHUV

Melody Swartz, Mécanobiologie et de morphogenèse, EPFL

Administration de la Recherche :

Jovan Mirkovitch

Anne Tricot

Coraline Fraga