

Article

Intuitive Eating Behavior, Diet Quality and Metabolic Health in the Postpartum in Women with Gestational Diabetes

Dan Yedu Quansah ^{1,*}, Sybille Schenk ^{1,2}, Leah Gilbert ¹, Amar Arhab ¹, Justine Gross ², Pedro-Manuel Marques-Vidal ³, Elena Gonzalez Rodriguez ⁴, Didier Hans ⁴, Antje Horsch ^{5,6} and Jardena J. Puder ¹

- ¹ Obstetric Service, Department Woman-Mother-Child, Lausanne University Hospital and University of Lausanne, 1011 Lausanne, Switzerland
 - ² Service of Endocrinology, Diabetology and Metabolism, Department of Medicine, Lausanne University Hospital and University of Lausanne, 1005 Lausanne, Switzerland
 - ³ Department of Medicine, Internal Medicine, Lausanne University Hospital and University of Lausanne, 1011 Lausanne, Switzerland
 - ⁴ Interdisciplinary Center of Bone Diseases, Bone & Joint Department, Lausanne University Hospital and University of Lausanne, 1011 Lausanne, Switzerland
 - ⁵ Institute of Higher Education and Research in Healthcare (IUFERS), University of Lausanne, 1010 Lausanne, Switzerland
 - ⁶ Neonatology service, Department Woman-Mother-Child, Lausanne University Hospital and University of Lausanne, 1011 Lausanne, Switzerland
- * Correspondence: dan.quansah@chuv.ch; Tel.: +41-21-314-0638; Fax: +41-21-314-8031



Citation: Quansah, D.Y.; Schenk, S.; Gilbert, L.; Arhab, A.; Gross, J.; Marques-Vidal, P.-M.; Gonzalez Rodriguez, E.; Hans, D.; Horsch, A.; Puder, J.J. Intuitive Eating Behavior, Diet Quality and Metabolic Health in the Postpartum in Women with Gestational Diabetes. *Nutrients* **2022**, *14*, 4272. <https://doi.org/10.3390/nu14204272>

Academic Editor: Antonio Brunetti

Received: 20 September 2022

Accepted: 11 October 2022

Published: 13 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Little is known regarding intuitive eating (IE), diet quality and adherence. We investigated the associations between IE, diet quality and metabolic health after gestational diabetes (GDM), who have an increased diabetes risk. Data from 179 women with GDM from MySweetheart trial (NCT02872974) were analyzed. IE was assessed using the eating for physical rather than emotional reasons (EPR) and reliance on hunger and satiety cues (RHSC) subscales of the French Intuitive Eating Scale-2. Metabolic outcomes included weight, central body fat and insulin resistance. Diet quality was calculated using the Alternative Health Eating Index (AHEI) and compliance with national recommendations was evaluated. Both IE subscales were associated with lower BMI and fat mass (BIA) at 1-year postpartum (all $p \leq 0.034$). The EPR subscale inversely correlated with fat mass (DXA) and visceral adipose tissue (both $p \leq 0.028$), whereas RHSC with higher insulin sensitivity (Matsuda, $p = 0.034$). RHSC during pregnancy predicted increased AHEI ($p = 0.043$) at 1-year postpartum, whilst EPR predicted lower fat mass and insulin resistance (HOMA-IR) (all $p \leq 0.04$). In longitudinal analyses, both subscales were associated with increased adherence to dairy and fiber intake recommendations (both $p \leq 0.023$). These data suggest IE may be an interesting approach to improve diet quality and metabolic outcomes in women with GDM.

Keywords: intuitive eating; diet quality; dietary adherence; metabolic health; postpartum; insulin resistance

1. Introduction

Eating behavior is linked to food choices, eating practices, dieting, and eating-related problems. Intuitive Eating (IE) is an adaptive and regulated eating behavior that controls emotional eating and dietary restriction [1]. It is a non-dieting approach, which deals with the ability to interpret and adhere to instinctive feedback regarding the amount and timing of food intake. IE has three distinct characteristics: (1) reliance on internal hunger and satiety cues to determine when to start and stop eating; (2) nonrestrictive eating, and (3) eating for physical rather than emotional reasons [2,3].

Diet quality refers to a dietary pattern or an indicator of food variety across key food groups recommended in dietary guidelines [4]. The Alternate Healthy Eating Index (AHEI)

is a quantitative score measuring diet quality [5]. An updated review of 34 observational cohorts revealed that higher dietary quality is associated with adherence to dietary recommendations and lower all-cause mortality, metabolic and cardiovascular risks [6]. Diet quality can influence morbidity and mortality beyond weight (or glucose control) and both quality and adherence have long-lasting benefits for metabolic health [6,7].

IE can potentially improve dietary adherence, and encourage self-responsibility with food [1]. Although cross-sectional studies outside of pregnancy found no association between IE and diet quality [8,9], an IE intervention study in the general population led to an improved diet quality [10]. In pregnancy, one cross-sectional study suggested an association between IE and diet quality in the second and third trimesters of pregnancy [11], but longitudinal studies and studies in the postpartum are lacking. We are not aware of studies relating IE to adherence with national or international dietary recommendations.

Relationships between IE and body mass index (BMI), weight, gestational weight gain [12,13] in the general pregnant population and with weight [14] in the postpartum have been reported. We previously also showed both cross-sectional and longitudinal associations between IE with weight, BMI, weight retention and glucose control in the perinatal period up to 1-year postpartum in metabolically high-risk women with gestational diabetes (GDM; women with a glucose intolerance diagnosed in pregnancy) [15,16]. Data on IE with body composition including total and central body fat, and insulin resistance, both of which are correlates of adverse metabolic health outcomes in the postpartum period in women with GDM, are lacking. The perinatal period is an essential moment regarding maternal metabolic health. Existing recommendations aimed to improve metabolic health include women returning to their pre-pregnancy weight or a 5% decrease in weight if they are overweight or obese. However, at least two-thirds of women do not even return to pre-pregnancy weight at 1-year postpartum [17,18]. In women with GDM, postpartum weight is the most important predictor for future diabetes [19]. In view of the 7-fold increased risk for diabetes and cardiovascular outcomes [20,21], the postpartum period is especially a critical moment for women with GDM. Data on the relationship between IE, diet quality and metabolic health are lacking, and it may be promising to explore the potential benefits of IE as an alternative approach to promote healthy eating and to prevent short and long-term adverse metabolic health outcomes in the postpartum period in women after GDM.

We investigated the cross-sectional and longitudinal associations between IE during and after pregnancy with diet quality and metabolic health including weight, total and central body fat, and insulin resistance at 1-year postpartum in women with GDM. We also explored if a higher IE score during pregnancy was related with higher adherence with the Swiss Society of Nutrition (SSN) dietary recommendations at 1-year postpartum.

2. Methods

2.1. Study Design and Patient Population

This current study is based on data from the MySweetheart trial (trial registration: NCT02872974) data. The detailed study protocol has been previously described [22]. Briefly, the MySweetheart trial tested the effect of an interdisciplinary lifestyle and psychosocial intervention on improving metabolic and mental health outcomes in women with GDM up to 1-year postpartum [22]. General eligibility criteria included women ≥ 18 years, diagnosed with GDM between 24–32 weeks gestational age (GA) according to the International Association of Diabetes and Pregnancy Study Groups (IADPSG) and the American Diabetes Association (ADA) guidelines [23,24]. Of the 211 participants included at baseline (105 randomized to intervention and 106 to usual care, Table 1), 179 completed the 1-year postpartum follow-up (Figure 1) and were included in the analyses. Predictors and outcomes of interest in this analysis were similar in the intervention and usual care groups so we pooled all participants together and adjusted for group allocation (see statistical analysis).

Table 1. Baseline maternal socio-demographic and health characteristics ($n = 179$).

Variable	Mean \pm SD
Age (year)	33.6 \pm 5.0
GA at the first GDM visit (weeks)	28.9 \pm 2.2
Pre-pregnancy weight (kg)	69.0 \pm 14.9
Pre-pregnancy BMI (kg/m ²)	25.59 \pm 5.1
Weight at the first GDM visit (kg)	79.32 \pm 14.6
BMI at the first GDM visit (kg/m ²)	29.41 \pm 4.9
HbA1c at the first GDM visit	5.09 \pm 0.3
Fasting glucose at GDM visit	4.95 \pm 0.5
Ethnicity/Nationality, n (%)	
Switzerland	52 (28.9)
Rest of Europe and North America	68 (38.3)
Asia and Oceania	13 (7.2)
Africa	21 (11.7)
Latin America	7 (3.9)
Others	18 (10.0)
Education level ^a , n (%)	
Compulsory school incomplete ^b	2 (1.3)
Compulsory school achieved	21 (13.9)
High school	16 (10.6)
General and vocational education	32 (21.2)
University	80 (53.0)
Employment status	
Student	5 (2.8)
Professional worker	118 (65.9)
Housewife/unemployed	56 (31.3)
Glucose-lowering treatment in pregnancy, n (%)	
None	99 (55.6)
Insulin	70 (38.9)
Metformin	10 (5.6)
Parity, n (%)	
0	102 (57.2)
1	49 (27.2)
2	14 (7.8)
≥ 3	14 (7.8)
Gravida, n (%)	
1	78 (43.3)
2	40 (22.8)
≥ 3	61 (33.9)
GDM in previous pregnancy ^c , yes, n (%)	19 (10.6)
Family history of diabetes, yes, n (%)	109 (61.1)
Social support during pregnancy, yes, n (%)	12 (6.7)

GDM denotes gestational diabetes mellitus; GA denotes gestational age; BMI denotes body mass index; ^a 29 participants had missing data on education; ^b in Switzerland, compulsory schooling lasts eleven years including kindergarten; ^c only for women who had at least one previous pregnancy; All data are expressed as n , % for categorical variables or mean (standard deviation) for continuous variables.




Figure 1. Flow of study participants. Of the 211 women included at baseline, 179 completed the 1-year follow-up and were included in this analysis. Out of these 179 women with data at 1-year postpartum visit, 100 had valid dietary data and 162 had valid data for HOMA-IR.

2.2. GDM Management and Patient Follow-Up

Women in the usual care group were followed-up according to the current ADA and Endocrine Society guidelines [23,25]. Following the diagnosis of GDM, women were seen at 24–32 weeks GA by either a physician, or a diabetes-specialist nurse, and were followed until childbirth. During this visit, they received information on GDM, specific recommendations regarding lifestyle changes and gestational weight gain (GWG) based on the 2009 recommendations of the Institute of Medicine (IOM) [26]. We placed a strong focus on lifestyle behavioral changes. Women were taught how to perform self-control of blood glucose both fasting and 2 h postprandial. Women also had one appointment with a registered dietician in order to receive individualized dietary advice, which focused on distribution of carbohydrate intake over several meals and snacks, limiting the intake of free sugars to less than 10%, and increase fiber intake to up to 30 g per day [27]. Women were advised to reduce sedentary behavior and engage in physical activity ideally postprandial. Treatment with insulin, or rarely with metformin was introduced when glucose values remained above targets between two or more times during a 1 to 2-week period (fasting plasma glucose) (FPG) > 5.3 mmol/l, 1-h postprandial glucose > 8 mmol/l and/or 2-h postprandial glucose > 7 mmol/l) despite lifestyle changes according to Swiss guidelines [28]. At 1-year postpartum, patients underwent a 75 g oral glucose tolerance test (oGTT) and received general advice on lifestyle changes.

On top of the usual care, the intervention consisted of four clinical lifestyle sessions during pregnancy and four lifestyle and psychosocial visits in the postpartum, two peer support group workshop (one in pregnancy and one in the postpartum), and a bimonthly

lifestyle coach support, mostly through telemedicine. It focused on tailored behavioral and psychosocial strategies to improve diet, physical activity, mental health and social support, and to improve adherence to GWG and weight retention recommendations. Of the four additional sessions during pregnancy, women had two counseling sessions with a dietician that encouraged mindful eating, and prioritized higher quality fats, reducing the intake of red or processed meat or high-fat cheese. They also had two additional sessions with the physiotherapists to increase physical activity. In the postpartum, continuous breastfeeding for at least 6 months was encouraged and aerobic and resistance physical activity for 150 min a week and resistance physical activity twice a week, all at a moderate intensity, were recommended during one of the postpartum visits.

2.3. Measures

All outcome measures including IE, metabolic health (except those assessed at only 1-year; see below) and dietary variables including diet quality were assessed both at the first GDM visit during pregnancy (baseline) and at 1-year postpartum.

2.3.1. Baseline Demographic and Health Characteristics

Data on maternal socio-demographic characteristics including age, nationality/ethnic origin and educational level were collected during the first GDM visit. Information on medical characteristics including previous history of GDM, family history of diabetes, gravida, parity and family social support during pregnancy (living with partner or with support, yes/no) were extracted from participants' medical charts.

2.3.2. Intuitive Eating Assessment

We assessed IE with a 14-item self-report questionnaire consisting of "eating for physical rather than emotional reasons" (EPR, 8 items) and the "reliance on hunger and satiety cues" (RHSC, 6 items) subscales of the French Intuitive Eating Scale-2 (IES-2) [29]. Originally, the French IES-2 contains three (3) subscales including the EPR, the RHSC and the unconditional permission to eat (UPE) subscale. Although we measured IE before the diet visit, we did not include the UPE subscale in this study, because the diagnosis of GDM itself and subsequent dietary counselling could significantly influence responses to the UPE subscale questions, such as "I try to avoid certain foods high in fat, carbohydrates, or calories". However, we measured metabolic health and dietary intakes before the dietician visit to ensure that, diet counselling did not influence study outcomes.

Women completed the EPR and RHSC subscales by responding to a 5-point Likert scale response ranging from one ("strongly disagree") to five ("strongly agree") to each item in both subscales. Possible scores for each subscale ranged from 1 to 5. We then calculated the EPR and RHSC subscale scores as recommended [29]. A higher score of the EPR subscale reflects eating as an answer to hunger and a lower score meant eating to cope with emotional distress, whereas a higher score of the RHSC subscale signifies trust in internal cues, and a lower score reflects less ability to regulate food intake.

2.3.3. Dietary Intake, Diet Quality (AHEI) and Dietary Adherence

Dietary intake was assessed with a validated food-frequency questionnaire (FFQ) for French-speaking Swiss adults. Details of this FFQ are described elsewhere [30]. This FFQ assesses the intakes of 97 different food items in the previous 4 weeks. For each food item, consumption frequencies ranged from "less than once during the last 4 weeks" to "2 or more times per day". We compared participants' average portion sizes (smaller, equal or bigger) to a reference size. Reported consumption frequencies were converted into daily or weekly consumptions including: "never these last 4 weeks" = 0; "once/month" = 1/28; "2-3/month" = 2.5/28; "1-2/week" = 1.5/7; "3-4times/week" = 3.5/7; "once/day" = 1 and "2+/day" = 2.5. We summed all frequencies of foods in a food category to obtain the consumption frequency of that food category. Conversion into food nutrients were based

on the French CIQUAL food composition table. We expressed total protein, carbohydrate and fat as percentage of total energy intake (TEI) (alcohol excluded).

We calculated Alternative Health Eating Index (AHEI) as a measure of diet quality. The AHEI was adapted from McCullough et al. [5]. In our study, we could not assess the amount of trans fat. The modified AHEI score ranged between 2.5 and 77.5 instead of 2.5 and 87.5 for the original AHEI score [5]. Higher AHEI scores represented a higher diet quality and a healthier diet. At 1-year postpartum, compliance with the national Swiss Society of Nutrition (SSN) dietary recommendations were computed as recommended [31]. The recommendations include ≥ 2 fruit portions/day; ≥ 3 vegetable portions/day; ≤ 5 portions meat/week; ≥ 1 portion fish/week and ≥ 3 portions dairy products/day and fiber ≥ 30 g/day [32]. For each recommendation, a binary variable (1 = yes, 0 = no) was computed.

2.3.4. Metabolic Health Outcomes

Pre-pregnancy weight was extracted from participants' medical charts or, if missing, was self-reported. We measured height at the first GDM visit during pregnancy and weight at the same moment and at 1-year postpartum to the nearest 0.1 cm and 0.1 kg with electronic scales (Seca® model 220, Hamburg, Germany). BMI was expressed as a ratio of weight in kilograms to the square of height in meters (kg/m^2). Total GWG during pregnancy was defined as the difference in pre-pregnancy weight and weight at the end of pregnancy. Fat mass was estimated using Kyle equation based on reactance and resistance values from Bioelectrical Impedance Analysis (BIA) (Akern BIA 101, Akern Srl, Pontassieve, Italy) measures performed during the first visit and at 1-year postpartum [33]. We also assessed fat mass and visceral adipose tissue at 1-year postpartum using Dual-Energy X-ray Absorptiometry (DXA; iDXA device, GEHC-Lunar, Madison, WI, USA) in 109 women who signed an additional consent form for this procedure.

We extracted data on the need for glucose-lowering medical treatment during pregnancy (use of insulin and/or metformin) from maternal medical records. We measured fasting glucose, HbA1c and calculated Homeostatic Model Assessment for Insulin Resistance (HOMA-IR) at the first GDM visit. At 1-year postpartum, women underwent a 75 g oGTT with glucose and insulin sampling at 30 min intervals for 2 h. We calculated insulin resistance/sensitivity, both by HOMA-IR (mostly hepatic insulin resistance) and MATSUDA (mostly total body insulin sensitivity) [34,35].

2.4. Statistical Analyses

All statistical analyses were performed with Stata/SE 15.1 (StataCorp LLC, College Station, TX, USA) [36]. We presented demographic and other descriptive variables as means (\pm standard deviation) or percentages (%) where appropriate. Predictors (EPR and RHSC subscales) and outcomes including diet quality (AHEI), all metabolic health variables (weight, BMI, GWG, total fat mass (BIA and DXA), visceral adipose tissue (DXA) and measures of insulin resistance (HOMA-IR, MATSUDA)) were normally distributed. We used a paired *t*-test to determine the changes in EPR and RHSC, AHEI and metabolic health variables between the first GDM visit and the 1-year postpartum. We performed linear regression analyses to determine the associations between EPR and RHSC with AHEI and metabolic health, namely the cross-sectional association between EPR and RHSC with AHEI and metabolic health at 1-year postpartum, and longitudinal associations between EPR and RHSC at the first GDM visit during pregnancy with AHEI and metabolic health at 1-year postpartum.

We used a modified Poisson regression for dichotomous variables to determine the longitudinal relationship between EPR and RHSC at the first GDM visit during pregnancy and adherence to SSN dietary recommendations for fruits, vegetables, dairy, protein and fiber intakes at 1-year postpartum. In all analyses, predictors and outcomes were similar in both groups (intervention vs. usual care), and the results, particularly the effect sizes, were similar when the regression analyses were restricted only to the usual care group or both

groups together. Therefore, and to increase the sample size, we pooled both groups and adjusted for group allocation in all analyses. All reported beta-coefficients for all regression estimates were standardized. All statistical significances were two-sided and accepted at $p < 0.05$.

3. Results

3.1. Baseline Characteristics of Participants

Among the 211 women included at the first GDM visit, 179 (84.8%) completed the 1-year follow-up visit (Figure 1). Mean gestational age, pre-pregnancy weight and BMI at baseline were 29.0 ± 2.2 weeks, 69.0 ± 14.8 kg and 25.6 ± 5.1 kg/m² respectively (Table 1).

Table 2 shows the change in IE, diet quality and metabolic health variables between the first GDM visit and 1-year postpartum. Whereas the EPR subscale remained stable (0.01 ± 0.8), the RHSC increased by 0.13 ± 0.9 at 1-year postpartum ($p = 0.041$). Nationality/ethnic origin of women did not influence EPR and RHSC scores (data not shown). Dietary intake variables including AHEI scores (Table 2), TEI, total carbohydrate, protein, fat and fiber intakes (results not shown) were similar at both time points (all $p \geq 0.07$), whereas all measures of metabolic health including BMI, weight, fat mass (BIA), and insulin resistance (HOMA-IR), significantly decreased at 1-year postpartum (all $p \leq 0.007$).

Table 2. Frequency, mean and standard deviation of study variables at baseline and end of the study.

Variable	<i>n</i>	First GDM Visit Mean \pm SD	At 1-Year pp Mean \pm SD	Mean Difference Mean \pm SD	<i>p</i> -Value
Intuitive eating behaviors					
EPR subscale	179	3.8 ± 0.8	3.8 ± 0.9	0.01 ± 0.8	0.846
RHSC subscale	179	3.5 ± 0.8	3.6 ± 0.8	0.13 ± 0.9	0.041
Diet quality					
AHEI	100	31.7 ± 9.7	30.6 ± 9.5	-1.07 ± 11.0	0.395
Metabolic health variables					
BMI (kg/m ²)	179	29.4 ± 4.9	26.8 ± 5.6	-2.52 ± 2.1	<0.001
Weight (kg)	179	79.3 ± 14.5	72.4 ± 16.1	-6.85 ± 5.7	<0.001
Total fat mass (BIA) (kg)	179	31.5 ± 9.2	26.6 ± 10.6	-4.91 ± 4.2	<0.001
HOMA-IR	162	3.6 ± 2.0	3.2 ± 2.3	-0.34 ± 1.5	0.007

GDM denotes gestational diabetes mellitus; SD denotes standard deviation; BMI denotes body mass index, EPR denotes Eating for Emotional Rather than Physical reasons of the French Intuitive eating Scale-2 (IES-2) questionnaire; RHSC denotes Reliance on Hunger and Satiety subscale of the IES-2; AHEI denotes Alternative Healthy Eating Index, BIA denotes Bioelectrical Impedance Analysis. HOMA-IR denotes Homeostatic Model Assessment for Insulin Resistance; *p*-value derived from paired *t*-test; bold *p*-values are significant ($p < 0.05$).

3.2. Cross-Sectional Associations between IE, Diet Quality and Metabolic Health

In the cross-sectional analyses at 1-year postpartum, none of the subscales of IE was associated with AHEI (Table 3). Similarly, they were not associated with HOMA-IR. However, they were inversely associated with lower BMI (EPR: $\beta = -1.17$, 95% CI: -2.13 , -0.21 ; RHSC: $\beta = -1.20$, 95% CI: -2.14 , -0.22) and fat mass (EPR: $\beta = -2.01$, 95% CI: -3.85 , -0.17 ; RHSC: $\beta = -2.02$, 95% CI: -3.97 , -0.08). Additionally, the EPR subscale correlated with lower weight ($\beta = -2.82$, 95% CI: -5.59 , -0.04), fat mass (DXA) ($\beta = -4.44$, 95% CI: -7.08 , -1.80) and visceral adipose tissue ($\beta = -0.12$, 95% CI: -0.23 , -0.02) whereas the RHSC was associated with higher total body insulin sensitivity (MATSUDA, $\beta = 0.18$, 95% CI: 0.006 , 0.128).

Table 3. Cross-sectional associations between the two subscales of intuitive eating with diet quality and metabolic health outcomes at 1-year postpartum.

Variable	Effect Estimate		
	Standardized Beta Coefficient	β (95% CI)	<i>p</i> Value
EPR Subscale at 1-year pp			
Diet quality at 1-year pp			
AHEI	0.04	0.56 (−1.38, 2.50)	0.548
Metabolic health at 1-year pp			
Weight (kg)	−0.15	−2.82 (−5.59, −0.04)	0.040
BMI (kg/m ²)	−0.18	−1.17 (−2.13, −0.21)	0.013
Fat mass (BIA) (kg)	−0.16	−2.01 (−3.85, −0.177)	0.027
Fat mass (DXA) (kg)	−0.32	−4.44 (−7.08, −1.80)	0.001
Visceral adipose tissue (DXA) (kg)	−0.23	−0.12 (−0.23, −0.02)	0.028
HOMA-IR	−0.09	−0.25 (−0.68, 0.16)	0.184
MATSUDA	0.03	0.11 (−0.48, 0.71)	0.595
RHSC Subscale at 1-year pp			
Diet quality at 1-year pp			
AHEI	0.009	0.11 (−1.94, 2.17)	0.337
Metabolic health at 1-year pp			
Weight (kg)	−0.11	−2.30 (−5.26, 0.66)	0.109
BMI (kg/m ²)	−0.09	−1.20 (−2.14, −0.22)	0.023
Fat mass (BIA) (kg)	−0.15	−2.02 (−3.97, −0.08)	0.034
Fat mass (DXA) (kg)	−0.16	−2.34 (−5.16, 0.47)	0.101
Visceral adipose tissue (DXA) (kg)	−0.12	−0.06 (−0.17, 0.04)	0.319
HOMA-IR	−0.10	−0.30 (−0.75, 0.14)	0.142
MATSUDA	0.18	0.64 (−0.006, 1.28)	0.034

pp denotes postpartum; EPR denotes Eating for Emotional Rather than Physical reasons of the French Intuitive eating Scale-2 (IES-2) questionnaire; RHSC denotes Reliance on Hunger and Satiety subscale of the IES-2; AHEI denotes Alternative Healthy Eating Index, BIA denotes Bioelectrical Impedance Analysis; BMI denotes body mass index, DXA denotes Dual-energy X-ray Absorptiometry. *p*-values are based on linear regression estimates adjusted for group allocation; bold *p*-values are significant ($p < 0.05$).

3.3. Prospective Associations between IE, Diet Quality, Metabolic Health and Dietary Adherence

In the longitudinal analyses (Table 4), the EPR subscale at the first GDM visit was associated with a decreased BMI ($\beta = -1.42$, 95% CI: −4.21, −0.03), fat mass (DXA) ($\beta = -2.95$, 95% CI: −5.95, −0.02) and lower insulin resistance (HOMA-IR; $\beta = -1.61$, 95% CI: −3.21, −0.09) at 1-year postpartum but not with AHEI. On the other hand, the RHSC subscale during pregnancy predicted an improved AHEI score ($\beta = 2.03$, 95% CI: 0.09, 3.97) at 1-year postpartum. RHSC was also associated with both lower GWG during pregnancy and weight at 1-year postpartum (both $p \leq 0.041$).

We also investigated the relationship between IE at the first GDM visit and adherence to the SSN dietary recommendations for adults at 1-year postpartum (Table 5). Both subscales predicted an increased adherence to dairy and fiber intakes recommendations ($p \leq 0.023$). Additionally, the RHSC subscale was associated with an increased adherence to the fruit intake recommendation ($p = 0.024$).

Table 4. Longitudinal associations between the two subscales of intuitive eating during pregnancy with diet quality and metabolic health outcomes up to 1-year postpartum.

Variable	Effect Estimate		
	Standardized Beta Coefficient	β (95% CI)	<i>p</i> Value
EPR Subscale the First GDM Visit			
Diet quality at 1-year pp			
AHEI	0.14	1.96 (−1.70, 5.63)	0.279
Metabolic health during pregnancy			
Total GWG during pregnancy (kg)	0.10	0.79 (−0.26, 1.86)	0.157
Metabolic health at 1-year pp			
Weight (kg)	−0.10	−2.17 (−5.12, 0.76)	0.146
BMI (kg/m ²)	−0.11	−1.42 (−4.21, −0.03)	0.016
Fat mass (BIA) (kg)	−0.09	−1.30 (−3.27, 0.66)	0.193
Fat mass (DXA) (kg)	−0.18	−2.95 (−5.95, −0.02)	0.043
Visceral adipose tissue (DXA) (kg)	−0.13	−0.08 (−0.20, 0.03)	0.172
HOMA-IR	−0.20	−1.61 (−3.21, −0.09)	0.032
MATSUDA	0.04	0.16 (−0.49, 0.81)	0.640
RHSC Subscale the First GDM Visit			
Diet quality at 1-year pp			
AHEI	0.16	2.03 (0.09, 3.97)	0.043
Metabolic health during pregnancy			
Total GWG during pregnancy (kg)	−0.12	−1.42 (−4.20, −0.03)	0.041
Metabolic health at 1-year pp			
Weight (kg)	−0.14	−2.90 (−5.81, −0.04)	0.025
BMI (kg/m ²)	−0.11	−0.81 (−1.84, 0.20)	0.124
Fat mass (BIA) (kg)	−0.14	−1.82 (−3.75, 0.10)	0.067
Fat mass (DXA) (kg)	−0.12	−1.63 (−4.20, 0.92)	0.206
Visceral adipose tissue (DXA) (kg)	−0.05	−0.02 (−0.12, 0.07)	0.538
HOMA-IR	−0.10	−0.30 (−0.74, 0.14)	0.203
MATSUDA	0.06	0.22 (−0.43, 0.87)	0.537

EPR denotes Eating for Emotional Rather than Physical reasons of the French Intuitive eating Scale-2 (IES-2) questionnaire; RHSC denotes Reliance on Hunger and Satiety subscale of the IES-2; GDM denotes gestational diabetes mellitus; AHEI denotes Alternative Healthy Eating Index, GWG denotes gestational weight gain; BIA denotes Bioelectrical Impedance Analysis; BMI denotes body mass index, DXA denotes Dual-energy X-ray Absorptiometry. *p*-values are based on linear regression estimates adjusted for group allocation; bold *p*-values are significant (*p* < 0.05).

Table 5. Longitudinal relationship between intuitive eating subscales during pregnancy and adherence to Swiss society for nutrition recommendation at 1-year postpartum.

Variable	Mean ± SD	β (95% CI)	<i>p</i> Value
EPR Subscale the First GDM Visit			
Fruits intake			
Below	3.75 ± 0.98	Ref	
Adhere	3.84 ± 0.78	0.09 (−0.39, 0.59)	0.699
Vegetable intake			
Below	3.75 ± 0.96	Ref	
Adhere	3.83 ± 0.76	−0.08 (−0.47, 0.32)	0.700
Dairy intake			
Above	3.79 ± 0.77	Ref	
Adhere	4.83 ± 0.28	1.03 (0.14, 1.93)	0.023

Table 5. Cont.

Variable	Mean \pm SD	β (95% CI)	<i>p</i> Value
Protein (non-fried fish) intake			
Below	3.80 \pm 0.75	Ref	
Adhere	3.96 \pm 0.85	−0.14 (−0.39, 0.12)	0.288
Fiber intake			
Below	3.83 \pm 0.79	Ref	
Adhere	4.25 \pm 0.85	1.42 (1.07, 3.23)	0.024
RHSC Subscale the First GDM Visit			
Fruits intake			
Below	3.20 \pm 0.68	Ref	
Adhere	3.58 \pm 0.77	1.37 (0.42, 2.10)	0.031
Vegetable intake			
Below	3.53 \pm 0.80	Ref	
Adhere	3.67 \pm 0.91	0.14 (−0.27, 0.55)	0.505
Dairy intake			
Above	3.51 \pm 0.81	Ref	
Adhere	4.24 \pm 0.81	1.87 (0.56, 2.73)	0.003
Protein (non-fried fish) intake			
Below	3.56 \pm 0.82	Ref	
Adhere	3.57 \pm 0.78	0.17 (−0.25, 0.58)	0.901
Fiber intake			
Below	3.25 \pm 0.60	Ref	
Adhere	3.56 \pm 0.81	2.31 (1.98, 3.35)	0.021

EPR denotes Eating for Emotional Rather than Physical reasons of the French Intuitive eating Scale-2 (IES-2) questionnaire; RHSC denotes Reliance on Hunger and Satiety subscale of the IES-2; GDM denotes gestational diabetes mellitus; Ref denotes reference. Bold *p*-values are significant ($p < 0.05$). Analyses were adjusted for group allocation (intervention/control).

4. Discussion

This study found cross-sectional and longitudinal associations between intuitive eating (IE) with increased diet quality, healthier metabolic outcomes and increased adherence to dietary recommendations in women after GDM. Specifically, the cross-sectional analyses at 1-year postpartum showed an inverse association between the two subscales of IE (eating for physical rather than emotional (EPR) and reliance on hunger and satiety cues (RHSC) subscales) with BMI and fat mass (BIA). Additionally, the EPR subscale correlated with lower fat mass (DXA) and visceral adipose tissue, whereas the RHSC was associated with higher total body insulin sensitivity (MATSUDA). In the longitudinal analyses, RHSC in pregnancy predicted increased dietary quality (AHEI scores) and lower weight at 1-year postpartum, whilst EPR was inversely associated with BMI, fat mass (DXA) and insulin resistance (HOMA-IR). Higher scores of both subscales predicted an increased adherence to dairy and fiber intake and RHSC further predicted fruit intake recommendations at 1-year postpartum. These results shed new light on the importance of IE for nutrition and metabolic health in the perinatal period in these women.

Although the postpartum period is a critical and important public health period for metabolic health, particularly in women with GDM, there has been a lack of studies of IE and diet quality. Furthermore, there is a general need for longitudinal studies regarding IE. The longitudinal relationship between RHSC and increased AHEI scores in our study suggests that IE can promote diet quality [2,3]. IE emphasizes/focuses on internal cues of hunger and satiety, which are linked to diet quality by developing a positive relationship with food and instinctively recognizing food variety to promote nutritional balance [8]. Our results are concordant with a cross-sectional study that showed positive association between IE and diet quality during pregnancy in a general pregnant population [11]. The findings of this study are in agreement with several studies performed outside of the perinatal setting. This includes an IE intervention that increased diet quality [10], an IE program that reduced restrictive dieting and emotional eating towards a healthy eating

approach, ultimately promoting better diet quality [37] and a Swiss study that reported positive associations between EPR and RHSC with diet quality in women [38].

IE principles including honoring hunger and health [39] overlap with the key concepts of dietary adherence including making healthier food choices, healthy eating patterns and food variety [40]. It is therefore not surprising that higher scores of EPR and RHSC subscales during pregnancy predicted increased adherence to dairy and fiber intake recommendations at 1-year postpartum in our study. The latter also predicted an increased adherence to fruit recommendations. The relationship between IE practices and self-responsibility with food [1] leads to appropriate consumption of diverse food groups that influence dietary adherence [41]. Studies in the general population that showed associations between EPR subscale with lower intakes of dairy, meat and fish [12], lower intakes of sweets and fast foods [38] and with higher intakes of fruits, vegetables and whole grains [8] are in general consistent with our results. However, in contrast to our study, these studies rather demonstrate a relationship with healthy food choices, but do not investigate the relationship between IE and adherence to dietary recommendation of (inter) national scientific organizations. Adherence to dietary recommendations is an important predictor of long-term weight loss and improved metabolic outcomes [42].

In this study, the RHSC subscale predicted lower weight, whereas the EPR was inversely associated with BMI, fat mass (DXA) and insulin resistance (HOMA-IR). Increased IE practices reduce emotional eating triggers (anxiety and depression) that are often associated with increased weight gain [43] and its associated metabolic problems that are accentuated in women with a history of GDM. We also speculate that a significant positive relationship between IE with diet quality and adherence might play a role, as both diet quality and adherence have short- and long-term metabolic benefits [44].

Our finding of cross-sectional associations between the two subscales of IE and lower weight and BMI at 1-year postpartum in women after GDM has been previously reported by our research group [15,16]. However, this is the first to investigate and show the inverse association between IE with fat mass, visceral adipose tissue and with higher total body insulin sensitivity. Both visceral adipose tissue and insulin sensitivity have important clinical implications and are independently associated with cardio-metabolic outcomes in women after GDM [45]. The cross-sectional and longitudinal data of our study show the relationship between IE with central body fat and insulin resistance in women with GDM.

The strengths of this study include its design and longitudinal follow-up. It is the first study to investigate the relationship between IE with diet quality and adherence to dietary recommendations, particularly in women with GDM. It is also the first to show an association between IE with lower weight and metabolic parameters in the postpartum, notably fat mass and its distribution. Limitations include the use of self-reported data regarding IE and dietary intakes. Despite being self-reported, we used validated tools to measure IE, diet quality and compliance with SSN dietary recommendations. Another potential limitation is the lack of total IES-2 scores, as we did not include the UPE subscale because of the potential response bias described above. At 1-year postpartum, 15% ($n = 32/211$) of women were lost to follow-up, but this proportion is lower compared to other studies in women with GDM.

In the future, a more objective outcome measure for IE would be helpful. In addition, future research that utilizes IE intervention in women with GDM, starting early in pregnancy up to 1-year postpartum to increase diet quality, improve dietary adherence and metabolic outcomes both in pregnancy and in the postpartum are needed to determine the cause and effect of our associations.

5. Conclusions

We found cross-sectional and longitudinal associations between IE with increased diet quality (AHEI) and healthier metabolic outcomes at 1-year postpartum in women after GDM. Thus, the subscales of IES-2, RHSC and/or EPR, inversely correlated with weight, body fat and insulin resistance. Higher scores of both subscales predicted an increased

adherence to national dietary recommendations. Particularly the longitudinal associations imply that IE during and after pregnancy can be an alternative healthy eating approach that can increase diet quality, improve adherence, and reduce the risk of adverse metabolic outcomes in the postpartum in women after GDM who have an increased metabolic risk in the postpartum period.

Author Contributions: D.Y.Q. and J.J.P. conceived the study. D.Y.Q., A.H., L.G. and J.J.P. designed the study. D.Y.Q. and J.J.P. performed the data analysis. All authors (D.Y.Q., S.S., L.G., J.G., A.A., P.-M.M.-V., E.G.R., D.H., A.H. and J.J.P.) were involved in the interpretation of data. D.Y.Q. and J.J.P. wrote the draft manuscript. Authors S.S., L.G., J.G., A.A., P.-M.M.-V., E.G.R., D.H., A.H. revised the manuscript for important intellectual content and gave final approval for the version to be published. J.J.P. supervised all the work. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by a project grant from the Swiss National Science Foundation (SNF 32003B_176119) and by an unrestricted educational grant from Novo Nordisk. The SNF and Novo Nordisk had no role regarding the content of the original data or analyses or in the drafting of this manuscript.

Institutional Review Board Statement: The Human Research Ethics Committee of the Canton de Vaud (No. 2016-00745) approved the study protocol (Date: 21 July 2016). All procedures were performed in compliance with the Helsinki declaration. Participation in the study did not interfere with patients' routine clinical care during pregnancy and after delivery.

Informed Consent Statement: Written informed consent and consent for publication were obtained from all participants.

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The data are not publicly available because it is a clinical data maintained and kept in a secure server at the Lausanne University Hospital.

Acknowledgments: Our sincere appreciation goes to our study participants for their time and participation. We thank Deborah Degen, Dominique Stulz and Isabelle Cohen-Salmon who helped with data collection and Alain Lacroix and Bobby Stuijtzand for their contribution to data management.

Conflicts of Interest: The authors declare no conflict to interest.

References

1. Tylka, T.L. Development and psychometric evaluation of a measure of intuitive eating. *J. Couns. Psychol.* **2006**, *53*, 226–240. [[CrossRef](#)]
2. Tylka, T.L.; Calogero, R.M.; Danielsdóttir, S. Intuitive eating is connected to self-reported weight stability in community women and men. *Eat. Disord.* **2020**, *28*, 256–264. [[CrossRef](#)] [[PubMed](#)]
3. Tylka, T.L.; Kroon Van Diest, A.M. The Intuitive Eating Scale-2: Item refinement and psychometric evaluation with college women and men. *J. Couns. Psychol.* **2013**, *60*, 137–153. [[CrossRef](#)] [[PubMed](#)]
4. Wirt, A.; Collins, C.E. Diet quality—What is it and does it matter? *Public Health Nutr.* **2009**, *12*, 2473–2492. [[CrossRef](#)]
5. McCullough, M.L.; Feskanich, D.; Stampfer, M.J.; Giovannucci, E.L.; Rimm, E.B.; Hu, F.B.; Spiegelman, D.; Hunter, D.J.; Colditz, G.A.; Willett, W.C. Diet quality and major chronic disease risk in men and women: Moving toward improved dietary guidance. *Am. J. Clin. Nutr.* **2002**, *76*, 1261–1271. [[CrossRef](#)]
6. Schwingshackl, L.; Bogensberger, B.; Hoffmann, G. Diet Quality as Assessed by the Healthy Eating Index, Alternate Healthy Eating Index, Dietary Approaches to Stop Hypertension Score, and Health Outcomes: An Updated Systematic Review and Meta-Analysis of Cohort Studies. *J. Acad. Nutr. Diet.* **2018**, *118*, 74–100.e11. [[CrossRef](#)]
7. Patel, Y.R.; Robbins, J.M.; Gaziano, J.M.; Djoussé, L. Mediterranean, dash, and alternate healthy eating index dietary patterns and risk of death in the physicians' health study. *Nutrients* **2021**, *13*, 1893. [[CrossRef](#)]
8. Jackson, A.; Sano, Y.; Parker, L.; Cox, A.E.; Lanigan, J. Intuitive eating and dietary intake. *Eat Behav.* **2022**, *45*, 101606. [[CrossRef](#)]
9. Lopez, T.D.; Hernandez, D.; Bode, S.; Ledoux, T. A complex relationship between intuitive eating and diet quality among university students. *J. Am. Coll. Health* **2021**, 1–7. [[CrossRef](#)]
10. Carbonneau, E.; Bégin, C.; Lemieux, S.; Mongeau, L.; Paquette, M.C.; Turcotte, M.; Labonté, M.-È.; Provencher, V. A Health at Every Size intervention improves intuitive eating and diet quality in Canadian women. *Clin. Nutr.* **2017**, *36*, 747–754. [[CrossRef](#)]
11. Plante, A.S.; Savard, C.; Lemieux, S.; Carbonneau, É.; Robitaille, J.; Provencher, V.; Morisset, A.S. Trimester-Specific Intuitive Eating in Association With Gestational Weight Gain and Diet Quality. *J. Nutr. Educ. Behav.* **2019**, *51*, 677–683. [[CrossRef](#)]

12. Camilleri, G.M.; Méjean, C.; Bellisle, F.; Andreeva, V.A.; Kesse-Guyot, E.; Hercberg, S.; Péneau, S. Intuitive eating dimensions were differently associated with food intake in the general population-based nutrinet-santé study. *J. Nutr.* **2017**, *147*, 61–69. [[CrossRef](#)]
13. Paterson, H.; Treharne, G.J.; Horwath, C.; Haszard, J.J.; Herbison, P.; Hay-Smith, E.J.C. Intuitive eating and gestational weight gain. *Eat. Behav.* **2019**, *34*, 101311. [[CrossRef](#)]
14. Leahy, K.; Berlin, K.S.; Banks, G.G.; Bachman, J. The Relationship Between Intuitive Eating and Postpartum Weight Loss. *Matern. Child Health J.* **2017**, *21*, 1591–1597. [[CrossRef](#)]
15. Quansah, D.Y.; Gilbert, L.; Gross, J.; Horsch, A.; Puder, J.J. Intuitive eating is associated with improved health indicators at 1-year postpartum in women with gestational diabetes mellitus (GDM). *J. Health Psychol.* **2019**, *26*, 1168–1184. [[CrossRef](#)]
16. Quansah, D.Y.; Gross, J.; Gilbert, L.; Helbling, C.; Horsch, A.; Puder, J.J. Intuitive eating is associated with weight and glucose control during pregnancy and in the early postpartum period in women with gestational diabetes mellitus (GDM): A clinical cohort study. *Eat. Behav.* **2019**, *34*, 101304. Available online: <https://linkinghub.elsevier.com/retrieve/pii/S1471015319300558> (accessed on 25 May 2019). [[CrossRef](#)]
17. Walker, L.O.; Sterling, B.S.; Timmerman, G.M. Retention of pregnancy-related weight in the early postpartum period: Implications for women's health services. *J. Obstet. Gynecol. Neonatal Nurs.* **2005**, *34*, 418–427. [[CrossRef](#)]
18. Nicklas, J.M.; Zera, C.A.; Seely, E.W. Predictors of very early postpartum weight loss in women with recent gestational diabetes mellitus. *J. Matern. Neonatal Med.* **2018**, *33*, 120–126. [[CrossRef](#)]
19. Bao, W.; Yeung, E.; Tobias, D.K.; Hu, F.B.; Vaag, A.A.; Chavarro, J.E.; Mills, J.L.; Grunnet, L.G.; Bowers, K.; Ley, S.H.; et al. Long-term risk of type 2 diabetes mellitus in relation to BMI and weight change among women with a history of gestational diabetes mellitus: A prospective cohort study. *Diabetologia* **2015**, *58*, 1212–1219. [[CrossRef](#)]
20. Kramer, C.K.; Campbell, S.; Retnakaran, R. Gestational diabetes and the risk of cardiovascular disease in women: A systematic review and meta-analysis. *Diabetologia* **2019**, *62*, 905–914. [[CrossRef](#)]
21. Gunderson, E.P.; Sun, B.; Catov, J.M.; Carnethon, M.; Lewis, C.E.; Allen, N.B.; Sidney, S.; Wellons, M.; Rana, J.S.; Hou, L.; et al. Gestational Diabetes History and Glucose Tolerance after Pregnancy Associated with Coronary Artery Calcium in Women during Midlife: The CARDIA Study. *Circulation* **2021**, *143*, 974–987. [[CrossRef](#)]
22. Horsch, A.; Gilbert, L.; Lanzi, S.; Gross, J.; Kayser, B.; Vial, Y.; Simeoni, U.; Hans, D.; Berney, A.; Scholz, U.; et al. Improving cardiometabolic and mental health in women with gestational diabetes mellitus and their offspring: Study protocol for MySweetHeart Trial, a randomised controlled trial. *BMJ Open* **2018**, *8*, e020462. [[CrossRef](#)]
23. American Diabetes Association (ADA). Classification and diagnosis of diabetes: Standards of medical care in diabetes—2021. *Diabetes Care* **2021**, *44*, S15–S33. [[CrossRef](#)]
24. Metzger, B.E. International Association of Diabetes and Pregnancy Study Groups recommendations on the diagnosis and classification of hyperglycemia in pregnancy: Response to Weinert. *Diabetes Care* **2010**, *33*, e98. [[CrossRef](#)]
25. Blumer, I.; Hadar, E.; Hadden, D.R.; Jovanovic, L.; Mestman, J.H.; Murad, M.H.; Yogeve, Y. Diabetes and pregnancy: An Endocrine Society clinical practice guideline. *J. Clin. Endocrinol. Metab. Erratum in: J. Clin. Endocrinol. Metab.* **2022**, *107*, e3972. **2013**, *98*, 4227–4249. [[CrossRef](#)]
26. Institute of Medicine (IOM). *Weight Gain During Pregnancy: Reexamining the Guidelines*; Committee to Reexamine IOM Pregnancy Weight Guidelines: Washington, DC, USA, 2009.
27. Office fédéral de la Santé Publique OFSP. Alimentation Durant la Grossesse et la Période d'Allaitement. Berne. Available online: www.bag.admin.ch (accessed on 20 July 2022).
28. Arditi, C.; Puder, J.; Vial, Y.; Hagon-Traub, I.; Burnand, B. Pied diabétique Prise en charge multidisciplinaire du diabète: Recommandations pour la pratique clinique. *Rev. Med. Suisse* **2019**, *15*, 929. [[CrossRef](#)]
29. Camilleri, G.M.; Méjean, C.; Bellisle, F.; Andreeva, V.A.; Sautron, V.; Hercberg, S.; Péneau, S. Cross-cultural validity of the Intuitive Eating Scale—Psychometric evaluation in a sample of the general French population. *Appetite* **2015**, *84*, 34–42. [[CrossRef](#)]
30. Marques-Vidal, P.; Ross, A.; Wynn, E.; Rezzi, S.; Paccaud, F.; Decarli, B. Reproducibility and relative validity of a food-frequency questionnaire for French-speaking Swiss adults. *Food Nutr. Res.* **2011**, *55*, 5905. [[CrossRef](#)]
31. De Abreu, D.; Guessous, I.; Vaucher, J.; Preisig, M.; Waeber, G.; Vollenweider, P.; Marques-Vidal, P. Low compliance with dietary recommendations for food intake among adults. *Clin. Nutr.* **2013**, *32*, 783–788. [[CrossRef](#)]
32. Swiss Society of Nutrition. Substances Nutritives. 2011. Available online: www.sge-ssn.ch/media/sge_pyramid_E_basic_20161.pdf (accessed on 20 July 2022).
33. Kyle, U.G.; Genton, L.; Slosman, D.O.; Pichard, C. Fat-free and fat mass percentiles in 5225 healthy subjects aged 15 to 98 years. *Nutrition* **2001**, *17*, 534–541. [[CrossRef](#)]
34. Matsuda, M.; DeFronzo, R.A. Insulin sensitivity indices obtained from oral glucose tolerance testing: Comparison with the euglycemic insulin clamp. *Diabetes Care* **1999**, *22*, 1462–1470. [[CrossRef](#)] [[PubMed](#)]
35. Matthews, D.R.; Hosker, J.P.; Rudenski, A.S.; Naylor, B.A.; Treacher, D.F.; Turner, R.C. Homeostasis model assessment: Insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* **1985**, *28*, 412–419. [[CrossRef](#)] [[PubMed](#)]
36. StataCorp LP. *Stata Data Analysis and Statistical Software*; StataCorp LP: College Station, TX, USA, 2017.
37. Cole, R.E.; Horacek, T. Effectiveness of the “my body knows when” intuitive-eating pilot program. *Am. J. Health Behav.* **2010**, *34*, 286–297. [[CrossRef](#)] [[PubMed](#)]

38. Horwath, C.; Hagmann, D.; Hartmann, C. Intuitive eating and food intake in men and women: Results from the Swiss food panel study. *Appetite* **2019**, *135*, 61–71. [[CrossRef](#)] [[PubMed](#)]
39. Resch, E.; Tylka, T.L. *Handbook of Positive Body Image and Embodiment: Constructs, Protective Factors, and Interventions*; Oxford University Press: Oxford, UK, 2019.
40. National Academies of Sciences, Engineering, and Medicine; Health and Medicine Division; Food and Nutrition Board; Committee to Review the Process to Update the Dietary Guidelines for Americans. *Redesigning the Process for Establishing the Dietary Guidelines for Americans*; The National Academies Press: Washington, DC, USA, 2017.
41. Borelli, J.; Pignotti, G.P.; Widaman, A. Examining the Relationship Between Intuitive Eating and Diet Quality Among College Students. *Curr. Dev. Nutr.* **2020**, *4*, 1290. [[CrossRef](#)]
42. Gibson, A.A.; Sainsbury, A. Strategies to improve adherence to dietaryweight loss interventions in research and real-world settings. *Behav. Sci.* **2017**, *7*, 44. [[CrossRef](#)] [[PubMed](#)]
43. De Lauzon-Guillain, B.; Basdevant, A.; Romon, M.; Karlsson, J.; Borys, J.M.; Charles, M.A. Is restrained eating a risk factor for weight gain in a general population? *Am. J. Clin. Nutr.* **2006**, *83*, 132–138. [[CrossRef](#)]
44. Mattei, J.; Sotres-Alvarez, D.; Daviglus, M.L.; Gallo, L.C.; Gellman, M.; Hu, F.B.; Tucker, K.L.; Willett, W.C.; Siega-Riz, A.M.; Van Horn, L.; et al. Diet quality and its association with cardiometabolic risk factors vary by hispanic and latino ethnic background in the Hispanic community health study /study of Latinos. *J. Nutr.* **2016**, *146*, 2035–2044. [[CrossRef](#)]
45. Hogling, D.E.; Rydén, M.; Bäckdahl, J.; Thorell, A.; Arner, P.; Andersson, D.P. Body fat mass and distribution as predictors of metabolic outcome and weight loss after Roux-en-Y gastric bypass. *Surg. Obes. Relat. Dis.* **2018**, *14*, 936–942. [[CrossRef](#)]