

Understanding sexual health indicators during adolescence: a study to consider time since sexual debut when exploring multiple sexual partners

Yara Barrense-Dias^a, Richard E. Bélanger^{ab}, François Desbiens^{ab}, Scott T. Leatherdale^c, Slim Haddad^{ab}

Affiliations: ^aCentre de recherche sur les soins et les services de première ligne de l'Université Laval, 2525 Chemin de la Canardière, Québec, QC G1J 0A4, Canada; ^bCentre intégré universitaire de santé et de services sociaux de la Capitale-Nationale, Québec, Canada; ^cSchool of Public Health and Health Systems, University of Waterloo, Ontario, Canada.

Corresponding author: Yara Barrense-Dias, Yara.Barrense-Dias@unisante.ch, tel. +4121 314 69 46, Unisanté, Route de la Corniche 10, 1010 Lausanne (Switzerland)

Sources of funding: The COMPASS study was supported by a bridge grant from the Canadian Institutes of Health Research (CIHR) Institute of Nutrition, Metabolism and Diabetes (INMD) through the “Obesity – Interventions to Prevent or Treat” priority funding awards (OOP-110788; grant awarded to S. Leatherdale) and an operating grant from the Canadian Institutes of Health Research (CIHR) Institute of Population and Public Health (IPPH) (MOP-114875; grant awarded to S. Leatherdale). The COMPASS-Quebec project additionally benefits from funding from the Ministère de la Santé et des Services sociaux of the province of Québec, and the Direction régionale de santé publique du CIUSSS de la Capitale-Nationale. The participation of Yara Barrense-Dias was made possible thanks to a research grant (# IZSEZ0_191187 / 1) granted by the Swiss National Fund for Scientific Research.

Acknowledgements: The authors wish to thank the Direction régionale de santé publique du CIUSSS de la Capitale-Nationale, the COMPASS team, especially Mireille Desrochers-Couture

and Claude Bacque-Dion, the relevant school boards, the administrations of participating schools, and all participants, for their continued confidence in our team.

Conflict of interest: All authors have indicated they have no potential conflicts of interest to disclose.

ACCEPTED VERSION

ABSTRACT

Understanding risky sexual behaviors among adolescents is key in efforts devoted to reducing the health burden related to sexually transmitted infections and unintended or unplanned pregnancies. The aims of this study were to understand the association between number of lifetime sexual partners and time since sexual debut (TSSD) among adolescents and to determine whether sex modifies this association. Data were drawn from the 2018–2019 COMPASS-Quebec study, a cohort study conducted in secondary schools in the province of Quebec, Canada. Of 18,467 respondents aged 14 years and older, 6991 (37.9%; mean age 15.3) reported consensual sexual intercourse and answered questions on their age at sexual initiation and number of lifetime sexual partners. Multilevel Poisson regressions with robust standard errors were estimated to adjust for covariates and produce adjusted group mean differences. The adjusted mean number of lifetime sexual partners ranged from 1.5 for those who had recently begun sexual activity (<12 months) to 4.0 for those who had been active for >35 months, an average rise of about 0.6 per year. Females-to-males adjusted mean differences showed that males reported more sexual partners than females at all time points, but the differences were only significant at the shorter (<12 months) and longer (>35 months) time spans. This study highlights the importance of taking into account TSSD when using and interpreting the number of lifetime sexual partners as risky sexual behavior among adolescents. Sex did not have a significant modifying effect on the relationship between number of lifetime sexual partners and TSSD.

Keywords: Adolescence; Sexual debut; Sexual partners; Sex differences.

INTRODUCTION

Sexuality and sexual health are recognized as important dimensions in the development and well-being of young people (World Health Organization, 2010). Initiating sexual activity and experiencing intimate relationships are thus expected during adolescence. However, sexual initiation and experimentation can also be associated with risk-taking, potentially leading to negative health outcomes, such as sexually transmitted infections (STI) and unwanted or unplanned pregnancies. Research to identify risky sexual behaviors is therefore useful to inform prevention and education programs aimed at reducing negative health outcomes (Turchik & Garske, 2009).

Risky sexual behaviors among youth have been defined in various ways. The most commonly used variables are unprotected sex, early sexual initiation, and multiple sexual partners (Camirand, Traoré, & Baulne, 2016; Public Health Agency of Canada, 2019; Ritchwood, Ford, DeCoster, Sutton, & Lochman, 2015; Sales, Brown, Vissman, & DiClemente, 2012). The multiple sexual partners variable has been widely explored in relation to STIs, as it increases the probability of being exposed to pathogens through multiple contacts (Ashenhurst, Wilhite, Harden, & Fromme, 2017; Dickson, Paul, Herbison, McNoe, & Silva, 1996; Dimbuene, Emina, & Sankoh, 2014; Valois, Oeltmann, Waller, & Hussey, 1999).

The literature is inconsistent regarding both the number of sexual partners to be defined as a risk during adolescence (Baumann, Bélanger, Akre, & Suris, 2011; Eaton et al., 2008; Kann et al., 2018; Lehavot et al., 2014; Lehrer, Shrier, Gortmaker, & Buka, 2006; Olesen et al., 2012; Santelli, Brener, Lowry, Bhatt, & Zabin, 1998; Traoré et al., 2018; United Nations Programme on HIV/AIDS (UNAIDS), 2013) and the time period used to assess this number. Some studies used a lifetime period (Baumann et al., 2011; Bruederle, Delany-Moretlwe, Mmari, &

Brahmbhatt, 2019; Eaton et al., 2008; Grossbard, Lee, Neighbors, Hendershot, & Larimer, 2007; Kann et al., 2018; Santelli et al., 1998), while others focused on a limited time period (Ashenurst et al., 2017; Olesen et al., 2012; Scott et al., 2011; United Nations Programme on HIV/AIDS (UNAIDS), 2013; Ybarra, Rosario, Saewyc, & Goodenow, 2016). For example, some studies found that a higher number of lifetime sexual partners was more prevalent in upper grades (Eaton et al., 2008; Kann et al., 2018; Valois et al., 1999), and others, that early sexual debut was found to be associated with multiple sexual partners (Albert, Brown, & Flanigan, 2003; Kastbom, Sydsjö, Bladh, Priebe, & Svedin, 2015; Ma et al., 2009; Sandfort, Orr, Hirsch, & Santelli, 2008; Tuoyire, Anku, Alidu, & Amo-Adjei, 2018). A logical explanation for such associations would seem to be the time since sexual debut (TSSD): the earlier sexual activity begins, the greater the likelihood of having multiple sexual partners due to longer exposure. This issue has been highlighted for substance use, with early initiators having higher rates of substance use because of their longer exposure to risk (Patton et al., 2004). However, even though some studies have pointed out this limitation (Baumann et al., 2011; Kastbom et al., 2015; Lehrer et al., 2006), the use of TSSD in sexual health studies among youth remains sparse. Moreover, the natural history of multiple sexual partnering among adolescents remains to be described.

The aims of this study were, first, to understand the association between number of lifetime sexual partners and TSSD among a sample of sexually active adolescents and, second, to determine whether sex modifies this association. Some studies have identified a sex difference, with males reporting more sexual partners than females (Bruederle et al., 2019; Lehrer et al., 2006; Romero-Estudillo, González-Jiménez, Mesa-Franco, & García-García, 2014; Traoré et al.,

2018), while others have found no significant differences in adolescence (Vasilenko & Lanza, 2014).

METHOD

Survey design and study population

The COMPASS Study (Leatherdale et al., 2014) is a prospective cohort study (2012–2021) designed to collect hierarchical and longitudinal data annually from a convenience sample of secondary school students and the schools they attend. In accordance with an active-information passive-consent parental permission protocol, students are eligible if their parent(s) or guardian(s) do not withdraw their child from the survey. Eligible students who agree to participate answer a pen-and-pencil self-administered anonymous questionnaire that covers a broad spectrum of topics, such as substance use, bullying, violence, mental health, and other health-related behaviors. A full description of the COMPASS host study is available online (www.compass.uwaterloo.ca) and in print (Leatherdale et al., 2014).

In 2016, a sample of schools in Quebec City (COMPASS-Quebec) was voluntarily recruited to join the COMPASS host study. The recruitment and data collection protocols remained the same as the COMPASS host study; however, the recruitment and data collection tools were translated into French and additional questions related to student sexual health were added to the COMPASS-Quebec student questionnaire.

For this paper, data were drawn from the COMPASS-Quebec project. The study population includes all students in the five years of the secondary level.¹ Students aged 14 years

¹ In the province of Quebec, secondary schools cover five years, grades 1 to 5, equivalent to grades 7 to 11 in the U.S. and the rest of Canada.

and older are invited to respond to a subset of questions related to their behaviors and perceptions in relation to sexuality.

The data used in the present study were drawn from the Year 3 data collection, conducted during the 2018–2019 school year. Overall, parents' refusal rate was 0.1% (32 cases), and about 8% of students were not reached, either because of personal refusal or absence on the day of the survey. Of 18,467 respondents aged 14 years and older, 6991 (37.9%; mean age 15.3; 51.3% females) answered yes to the filter question, *Have you ever had sexual intercourse (oral, vaginal, or anal) with your consent?* of the sub-questionnaire on sexuality, and were invited to answer additional questions.

Variables

The number of lifetime sexual partners was obtained from the response to the question, *With how many different people have you had sexual intercourse (oral, vaginal, or anal) with your consent?* Response options ranged from one person to nine or more. This variable was used as a continuous one ranging from 1 to 9.

TSSD was obtained by subtracting age at first consensual sexual intercourse from age at time of survey and then recoded into four subcategories: less than 12 months, 12–23 months, 24–35 months, and 36 months or more. Age at first consensual sexual intercourse was measured by asking, *How old were you when you first had sex (with your consent)?* Seven response options were offered, ranging from 12 years or younger to 18 years or older.

Sex was used as a potential modifier of the relationship between number of lifetime sexual partners and TSSD. Covariates used for adjustments in statistical analyses were: age at first consensual sexual intercourse, ethnicity (non-white, white), family deprivation, and depression symptoms. These were chosen on the basis of their known association with risky

sexual behaviors, and especially with multiple sexual partners (Cavazos-Rehg et al., 2011; Valois et al., 1999; Vasilenko & Lanza, 2014). Early sexual initiation was also defined according to the answer to the question on the age at first consensual sexual intercourse. The seven response options were dichotomized by defining early sexual initiation as occurring before the age of 14 (Kastbom et al., 2015; Young, Burke, & Nic Gabhainn, 2018). Age at first consensual sex allows us to create TSSD and early sexual initiation. In a developmental and behavioral approach, we hypothesized that a TSSD of 2 years before the age of 14 and a TSSD of 2 years at the age of 16, for example, will not have the same influence on the number of sexual partners.

Family deprivation was based on a proxy variable whose value was 1 (0 otherwise) in the presence of at least one of three situations: 1) having less than \$6 a week for personal spending or savings; 2) sometimes skipping breakfast because there is nothing to eat at home; or 3) sometimes going to bed hungry at night because there is not enough money to buy food.

Depression symptoms were drawn from the *Center for Epidemiologic Studies Depression Scale Revised* (Eaton, Smith, Ybarra, Muntaner, & Tien, 2004). This scale includes 10 items referring to the past seven days (e.g. *I had trouble keeping my mind on what I was doing*). Total score could range from 0 to 30, with a score of 10 or higher being considered a threshold signaling the presence of significant depressive symptoms.

Data Analysis

Multilevel Poisson regressions with robust standard errors were estimated to adjust for covariates and produce adjusted group mean differences. Possible modifying effects were explored through double group mean differences (margins command in Stata®). There was no attempt to impute missing values of the dependent variables (rate of missing values = 2.7%) or

the covariates (overall rate of missing values = 6.5%). All analyses were performed with Stata[®] 16.0 (StataCorp, College Station, Texas).

RESULTS

More than one-third (38%) of respondents reported being sexually active. Table 1 presents the characteristics of those students. The prevalence of early sexual initiation (before age 14) was about 20%. Two out of three respondents reported TSSD of less than 24 months. Mean number of lifetime sexual partners was 2.4 (males 2.5, females 2.2).

[Insert Table 1 here]

The adjusted mean number of lifetime sexual partners ranged from 1.5 for those who had recently begun sexual activity (<12 months) to 4.0 for those who had been active for 36 months or more, an average rise of about 0.6 sexual partners a year.

[Insert Table 2 here]

The females-to-males adjusted mean differences showed significant differences in numbers of partners in the first stratum (less than 12 months since sexual debut), with a difference of 0.2 partners between females and males, and a greater differential number of partners appearing in the 4th stratum (36 months or more since sexual debut), with a difference of 0.5 partners. Between these two durations, the difference between females and males disappeared. Significant differences remained slight (-0.21 for <12 months, -0.54 for >35 months since sexual debut).

[Insert Figure 1 here]

The number of partners predicted after adjustment increased gradually with duration of sexual activity, reaching a mean of 3.74 [CI 3.53, 3.95] for females and 4.27 [CI 3.95, 4.59] for

males among respondents who had been active for at least 36 months. Difference-in-difference analysis, however, did not rule out the hypothesis of equal progression in both sexes (Table 3).

[Insert Table 3 here]

DISCUSSION

This study highlights the importance of considering TSSD when using and interpreting the number of lifetime sexual partners among adolescents. Previous studies (Albert et al., 2003; Eaton et al., 2008; Kann et al., 2018; Kastbom et al., 2015; Ma et al., 2009; Sandfort et al., 2008; Tuoyire et al., 2018; Valois et al., 1999) have found associations between multiple sexual partners and upper grades, older age, and/or early sexual initiation. However, TSSD has rarely been considered, whereas the number of lifetime sexual partners would appear to depend more on this variable than on those three (Baumann et al., 2011).

The present study sheds light on the association between number of lifetime sexual partners and TSSD in the first years of sexual activity among adolescents. Regardless of age at sexual initiation, over the subsequent four-year period the mean number of lifetime sexual partners was found to be 2.5, representing an average increase of 0.6 per year. These results show an average progression in the number of sexual partners that is relatively linear and regular in the first years of sexual activity. When number of sexual partners is used as a risk factor without considering TSSD, some might conclude that adolescents have a lot of casual sexual partners or one-night stands. Therefore, such linear results when TSSD is taken into account could challenge public opinion and societal perception about adolescent sexuality. Indeed, As such, one way to look at risky behaviors would be to define them in terms of where they fit in relation to norms, rather than applying a predefined number or threshold. In this vein, previous studies (Huang, Murphy, & Hser, 2012; Lansford et al., 2010; Moilanen, Crockett, Raffaelli, &

Jones, 2010; Rossi, Poulin, & Boislard, 2017) have aimed to define normative and deviant pathways in relation to sexual behaviors. For example, Rossi et al. (Rossi et al., 2017) identified four trajectory groups among young people aged 16 to 22 years regarding number of sexual partners: abstainers, low-increasing, medium-increasing, and multiple-partners. However, none of these studies considered TSSD in their analyses, with the period assessed for sexual partners being limited to the past year. The challenge that therefore persists is to define thresholds and healthy trajectories. In the current paradigm, an adolescent who does not conform to the trend and reports a significantly higher number of sexual partners could be regarded as representing a potential risk in terms of multiple partners. However, having multiple sexual partners is not necessarily an indicator of unhealthy sexual behavior unless it is done without protection. So, in addition to TSSD, other variables must be considered when trying to understand risky sexual behavior, particularly protection and contraception use, among adolescents.

As with other studies (Albert et al., 2003; Kastbom et al., 2015; Ma et al., 2009; Sandfort et al., 2008; Tuoyire et al., 2018), the present study found that earlier debut of sexual activity was associated with a higher number of lifetime sexual partners. Sexual initiation at an early age is also considered a risk because of other factors, such as adolescents' ongoing cognitive development, sensitivity to peer pressure, and greater impulsivity (Rossi et al., 2017; Steinberg et al., 2008; Sumter, Bokhorst, Steinberg, & Westenberg, 2009). However, it seems more appropriate to consider and interpret the association between early sexual initiation and number of sexual partners in terms of duration of risk exposure rather than a particular age or pre-set threshold.

In the initial years of sexual activity, females and males tend to follow a similar trajectory regarding number of sexual partners when TSSD is taken into account. In the present study, sex

did not have a significant modifying effect on the relationship between the numbers of lifetime sexual partners and TSSD. Furthermore, females and males followed similar trajectories even when they initiated sexual activity at different ages, such that the number of sexual partners in the first four years was not sex-related. This result is inconsistent with several studies (Bruederle et al., 2019; Lehrer et al., 2006; Romero-Estudillo et al., 2014; Traoré et al., 2018) that found males reporting higher numbers of sexual partners. Other studies have, however, challenged this difference as being potentially attributable to a bias in males' responses (Mitchell et al., 2019).

This study presents certain limitations that should be kept in mind. First, as with all population-based surveys of youth, sexual practices are documented based on respondents' self-reports, which may lead to underestimation of certain risky behaviors. This risk is mitigated, however, by the anonymity of the questionnaires (Brener, Billy, & Grady, 2003). Moreover, the high response rate indicated that adolescents were not embarrassed by the topic of sexual activity, which was addressed only at the end of the COMPASS-Quebec questionnaire, by which time some respondent fatigue might have been expected. Second, the results are not fully generalizable to other contexts, including the province of Quebec or its regions, as it is a convenience sample and school participation in the COMPASS-Quebec study is voluntary. The solicitation was carried out on convenience basis by the school boards; there is no reason to believe there is a serious selection bias at this level, since 90% of the schools solicited gave their consent. Moreover, student participation is probably free of the risk of selection bias, as: 1) less than half a percent of the parents asked to withdraw their children from the study; 2) the response rate in the participating schools varied between 80% and 90%, proportions that correspond to the rates of absenteeism observed. The results should nevertheless be interpreted with caution and not generalized. Third, no information was collected on relationships, simultaneous sexual

activity, or protection/contraception (only on condom use at last intercourse) (Dimbuene et al., 2014; Grossbard et al., 2007). Lastly, we used covariates that were available in the COMPASS study, but other confounders may be included.

Further studies are needed to determine whether questions on recent sexual partners with a time-limited assessment are more appropriate for studying risky sexual behaviors among adolescents. However, given a recent study (Haderxhanaj, Leichliter, Aral, & Chesson, 2014) found that number of lifetime sexual partners was associated with the number of recent sexual partners, showing that lifetime assessment is also important to understand recent risky sexual behaviors. How time since sexual debut may impact other risky sexual behaviors, such as non-use of condoms, remains a question of interest. Finally, as mentioned, having multiple sexual partners is not necessarily unhealthy unless it is done without protection. Therefore, future research should consider protection and contraception use in their interpretation of risky sexual behaviors among adolescents.

COMPLIANCE WITH ETHICAL STANDARDS

Conflict of Interest

The authors declare that they have no conflict of interest.

Ethical Approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or provincial research committee and with the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards. All procedures in the COMPASS study received ethics approval from the University of Waterloo Research Ethics Board (ORE 30118), as well from the Research Ethics Review Board of the

Centre intégré universitaire de santé et de services sociaux de la Capitale-Nationale [#MP-13-2017-1264] and participating school board review panels.

Informed Consent

The protocol for COMPASS involves active-information passive-consent parental permission procedures. Students could decline to participate at any time.

ACCEPTED VERSION

REFERENCES

- Albert, B., Brown, S., & Flanigan, C. M. (Eds.) (2003). *14 and younger: The sexual behavior of young adolescents*. Washington DC: National Campaign to Prevent Teen Pregnancy. Retrieved April 20, 2020, from: <https://eric.ed.gov/?id=ED477795>
- Ashenhurst, J. R., Wilhite, E. R., Harden, K. P., & Fromme, K. (2017). Number of sexual partners and relationship status are associated with unprotected sex across emerging adulthood. *Archives of Sexual Behavior*, *46*, 419–432. <https://doi.org/10.1007/s10508-016-0692-8>
- Baumann, P., Bélanger, R. E., Akre, C., & Suris, J.-C. (2011). Increased risks of early sexual initiators: Time makes a difference. *Sexual Health*, *8*, 431–435. <https://doi.org/10.1071/SH10103>
- Brener, N. D., Billy, J. O. G., & Grady, W. R. (2003). Assessment of factors affecting the validity of self-reported health-risk behavior among adolescents: Evidence from the scientific literature. *Journal of Adolescent Health*, *33*, 436-457. [https://doi.org/10.1016/S1054-139X\(03\)00052-1](https://doi.org/10.1016/S1054-139X(03)00052-1)
- Bruederle, A., Delany-Moretlwe, S., Mmari, K., & Brahmhbhatt, H. (2019). Social support and its effects on adolescent sexual risk taking: A look at vulnerable populations in Baltimore and Johannesburg. *Journal of Adolescent Health*, *64*, 56-62. <https://doi.org/10.1016/j.jadohealth.2018.07.004>
- Camirand, H., Traoré, I., & Baulne, J. (2016). *L'Enquête québécoise sur la santé de la population, 2014–2015 : Pour en savoir plus sur la santé des Québécois. Résultats de la deuxième édition*. Montreal: Institut de la statistique du Québec. Retrieved April 20,

2020, from: <https://www.stat.gouv.qc.ca/statistiques/sante/etat-sante/sante-globale/sante-quebecois-2014-2015.pdf>

- Cavazos-Rehg, P. A., Krauss, M. J., Spitznagel, E. L., Schootman, M., Cottler, L. B., & Bierut, L. J. (2011). Number of sexual partners and associations with initiation and intensity of substance use. *AIDS and Behavior*, *15*, 869–874. <https://doi.org/10.1007/s10461-010-9669-0>
- Dickson, N., Paul, C., Herbison, P., McNoe, B., & Silva, P. (1996). The lifetime occurrence of sexually transmitted diseases among a cohort aged 21. *The New Zealand Medical Journal*, *109*(1028), 308–312.
- Dimbuene, Z. T., Emina, J. B. O., & Sankoh, O. (2014). UNAIDS ‘multiple sexual partners’ core indicator: Promoting sexual networks to reduce potential biases. *Global Health Action*, *7*(23103), 1–6. <https://dx.doi.org/10.3402%2Fgha.v7.23103>
- Eaton, D. K., Kann, L., Kinchen, S., Shanklin, S., Ross, J., Hawkins, J., . . . Wechsler, H. (2008). Youth risk behavior surveillance – United States, 2007. *Morbidity and Mortality Weekly Report. Surveillance Summaries*, *57*(SS04), 1–131. Retrieved April 20, 2020 from: <https://www.cdc.gov>
- Eaton, W. W., Smith, C., Ybarra, M., Muntaner, C., & Tien, A. (2004). Center for Epidemiologic Studies Depression Scale: Review and Revision (CESD and CESD-R). In M. E. Maruish (Ed.), *The use of psychological testing for treatment planning and outcomes assessment: Instruments for adults (3rd ed.)* (pp. 363–377): Lawrence Erlbaum Associates Publishers.
- Grossbard, J. R., Lee, C. M., Neighbors, C., Hendershot, C. S., & Larimer, M. E. (2007). Alcohol and risky sex in athletes and nonathletes: What roles do sex motives play?

Journal of Studies on Alcohol and Drugs, 68, 566–574.

<https://doi.org/10.15288/jsad.2007.68.566>

Haderxhanaj, L. T., Leichter, J. S., Aral, S. O., & Chesson, H. W. (2014). Sex in a lifetime: Sexual behaviors in the United States by lifetime number of sex partners, 2006-2010. *Sexually Transmitted Diseases*, 41, 345-352.

<https://doi.org/10.1097/OLQ.0000000000000132>

Huang, D. Y. C., Murphy, D. A., & Hser, Y.-I. (2012). Developmental trajectory of sexual risk behaviors from adolescence to young adulthood. *Youth & Society*, 44, 479–499.

<https://doi.org/10.1177/0044118X11406747>

Kann, L., McManus, T., Harris, W. A., Shanklin, S. L., Flint, K. H., Queen, B., . . . Ethier, K. A. (2018). Youth risk behavior surveillance – United States, 2017. *Morbidity and Mortality Weekly Report. Surveillance Summaries*, 67(8), 1-114.

<http://dx.doi.org/10.15585/mmwr.ss6708a1>

Kastbom, Å. A., Sydsjö, G., Bladh, M., Priebe, G., & Svedin, C.-G. (2015). Sexual debut before the age of 14 leads to poorer psychosocial health and risky behaviour in later life. *Acta Paediatrica*, 104, 91–100. <https://doi.org/10.1111/apa.12803>

Lansford, J. E., Yu, T., Erath, S. A., Pettit, G. S., Bates, J. E., & Dodge, K. A. (2010).

Developmental precursors of number of sexual partners from ages 16 to 22. *Journal of Research on Adolescence*, 20, 651–677. <https://doi.org/10.1111/j.1532-7795.2010.00654.x>

Leatherdale, S. T., Brown, K. S., Carson, V., Childs, R. A., Dubin, J. A., Elliott, S. J., . . .

Thompson-Haile, A. (2014). The COMPASS study: a longitudinal hierarchical research platform for evaluating natural experiments related to changes in school-level programs,

policies and built environment resources. *BMC Public Health*, 14, article 331.

<https://doi.org/10.1186/1471-2458-14-331>

- Lehavot, K., Katon, J. G., Williams, E. C., Nelson, K. M., Gardella, C. M., Reiber, G. E., & Simpson, T. L. (2014). Sexual behaviors and sexually transmitted infections in a nationally representative sample of women veterans and nonveterans. *Journal of Women's Health*, 23, 246–252. <https://doi.org/10.1089/jwh.2013.4327>
- Lehrer, J. A., Shrier, L. A., Gortmaker, S., & Buka, S. (2006). Depressive symptoms as a longitudinal predictor of sexual risk behaviors among US middle and high school students. *Pediatrics*, 118, 189–200. <https://doi.org/10.1542/peds.2005-1320>
- Ma, Q., Ono-Kihara, M., Cong, L., Xu, G., Pan, X., Zamani, S., . . . Kihara, M. (2009). Early initiation of sexual activity: A risk factor for sexually transmitted diseases, HIV infection, and unwanted pregnancy among university students in China. *BMC Public Health*, 9, article 111. <https://doi.org/10.1186/1471-2458-9-111>
- Mitchell, K. R., Mercer, C. H., Prah, P., Clifton, S., Tanton, C., Wellings, K., & Copas, A. (2019). Why do men report more opposite-sex sexual partners than women? Analysis of the gender discrepancy in a british national probability survey. *The Journal of Sex Research*, 56, 1–8. <https://doi.org/10.1080/00224499.2018.1481193>
- Moilanen, K. L., Crockett, L. J., Raffaelli, M., & Jones, B. L. (2010). Trajectories of sexual risk from middle adolescence to early adulthood. *Journal of Research on Adolescence*, 20, 114–139. <https://doi.org/10.1111/j.1532-7795.2009.00628.x>
- Olesen, T. B., Jensen, K. E., Nygård, M., Tryggvadottir, L., Sparén, P., Hansen, B. T., . . . Kjær, S. K. (2012). Young age at first intercourse and risk-taking behaviours – a study of nearly

- 65 000 women in four Nordic countries. *European Journal of Public Health*, 22, 220–224. <https://doi.org/10.1093/eurpub/ckr055>
- Patton, G. C., McMorris, B. J., Toumbourou, J. W., Hemphill, S. A., Donath, S., & Catalano, R. F. (2004). Puberty and the onset of substance use and abuse. *Pediatrics*, 114, e300-306. <https://doi.org/10.1542/peds.2003-0626-F>
- Public Health Agency of Canada. (2019). *Report on sexually transmitted infections in Canada, 2017*. Retrieved April 20, 2020, from <https://www.canada.ca/en/public-health/services/publications/diseases-conditions/report-sexually-transmitted-infections-canada-2017.html>
- Ritchwood, T. D., Ford, H., DeCoster, J., Sutton, M., & Lochman, J. E. (2015). Risky sexual behavior and substance use among adolescents: A meta-analysis. *Children and Youth Services Review*, 52, 74–88. <https://doi.org/10.1016/j.childyouth.2015.03.005>
- Romero-Estudillo, E., González-Jiménez, E., Mesa-Franco, M. C., & García-García, I. (2014). Gender-based differences in the high-risk sexual behaviours of young people aged 15-29 in Melilla (Spain): A cross-sectional study. *BMC Public Health*, 14, article 745. <https://dx.doi.org/10.1186%2F1471-2458-14-745>
- Rossi, E., Poulin, F., & Boislard, M.-A. (2017). Trajectories of annual number of sexual partners from adolescence to emerging adulthood: Individual and family predictors. *Journal of Youth and Adolescence*, 46, 995–1008. <https://doi.org/10.1007/s10964-016-0571-6>
- Sales, J. M., Brown, J. L., Vissman, A. T., & DiClemente, R. J. (2012). The association between alcohol use and sexual risk behaviors among African American women across three developmental periods: a review. *Current Drug Abuse Reviews*, 5, 117–128. <https://doi.org/10.2174/1874473711205020117>

- Sandfort, T. G. M., Orr, M., Hirsch, J. S., & Santelli, J. (2008). Long-term health correlates of timing of sexual debut: Results from a national US study. *American Journal of Public Health, 98*, 155–161. <https://doi.org/10.2105/AJPH.2006.097444>
- Santelli, J. S., Brener, N. D., Lowry, R., Bhatt, A., & Zabin, L. S. (1998). Multiple sexual partners among U.S. adolescents and young adults. *Family Planning Perspectives, 30*, 271–275. <https://doi.org/10.2307/2991502>
- Scott, M. E., Wildsmith, E., Welti, K., Ryan, S., Schelar, E., & Steward-Streng, N. R. (2011). Risky adolescent sexual behaviors and reproductive health in young adulthood. *Perspectives on Sexual and Reproductive Health, 43*, 110–118. <https://doi.org/10.1363/4311011>
- Steinberg, L., Albert, D., Cauffman, E., Banich, M., Graham, S., & Woolard, J. (2008). Age differences in sensation seeking and impulsivity as indexed by behavior and self-report: Evidence for a dual systems model. *Developmental Psychology, 44*, 1764–1778. <https://doi.org/10.1037/a0012955>
- Sumter, S. R., Bokhorst, C. L., Steinberg, L., & Westenberg, P. M. (2009). The developmental pattern of resistance to peer influence in adolescence: Will the teenager ever be able to resist? *Journal of Adolescence, 32*, 1009–1021. <https://doi.org/10.1016/j.adolescence.2008.08.010>
- Traoré, I., Street, M.-C., Camirand, H., Julien, D., Joubert, K., & Berthelot, M. (2018). *Enquête québécoise sur la santé des jeunes du secondaire de 2016-2017. Résultats de la deuxième édition. La santé physique et les habitudes de vie des jeunes, Tome 3*. Montreal: Institut de la statistique du Québec. Retrieved April 20, 2020, from:

<https://www.stat.gouv.qc.ca/statistiques/sante/enfants-ados/alimentation/sante-jeunes-secondaire-2016-2017-t3.pdf>

Tuoyire, D. A., Anku, P. J., Alidu, L., & Amo-Adjei, J. (2018). Timing of first sexual intercourse and number of lifetime sexual partners in sub-Saharan Africa. *Sexuality & Culture*, 22, 651–668. <https://doi.org/10.1007/s12119-017-9488-9>

Turchik, J. A., & Garske, J. P. (2009). Measurement of sexual risk taking among college students. *Archives of Sexual Behavior*, 38, 936–948. <https://doi.org/10.1007/s10508-008-9388-z>

United Nations Programme on HIV/AIDS (UNAIDS). (2013). *Global AIDS response progress reporting 2013: Construction of Core Indicators for monitoring the 2011 UN Political Declaration on HIV/AIDS*. Retrieved April 20, 2020, from https://www.unaids.org/sites/default/files/media_asset/GARPR_2013_guidelines_en_0.pdf

Valois, R. F., Oeltmann, J. E., Waller, J., & Hussey, J. R. (1999). Relationship between number of sexual intercourse partners and selected health risk behaviors among public high school adolescents. *Journal of Adolescent Health*, 25, 328–335. [https://doi.org/10.1016/S1054-139X\(99\)00051-8](https://doi.org/10.1016/S1054-139X(99)00051-8)

Vasilenko, S. A., & Lanza, S. T. (2014). Predictors of multiple sexual partners from adolescence through young adulthood. *Journal of Adolescent Health*, 55, 491–497. <https://doi.org/10.1016/j.jadohealth.2013.12.025>

World Health Organization. (2010). *Measuring sexual health: Conceptual and practical considerations and related indicators*. Retrieved April 20, 2020, from <https://apps.who.int/iris/rest/bitstreams/66657/retrieve>

Ybarra, M. L., Rosario, M., Saewyc, E., & Goodenow, C. (2016). Sexual behaviors and partner characteristics by sexual identity among adolescent girls. *Journal of Adolescent Health, 58*, 310–316. <https://doi.org/10.1016/j.jadohealth.2015.11.001>

Young, H., Burke, L., & Nic Gabhainn, S. (2018). Sexual intercourse, age of initiation and contraception among adolescents in Ireland: Findings from the Health Behaviour in School-aged Children (HBSC) Ireland study. *BMC Public Health, 18*, article 362. <https://doi.org/10.1186/s12889-018-5217-z>

ACCEPTED VERSION

Table 1. Characteristics of sexually active respondents and number of lifetime sexual partners, crude prevalence and mean

| | Sexually active | | Number of lifetime sexual partners | |
|--------------------------------|-----------------|-------|------------------------------------|---------|
| | N | %* | N | Mean±SD |
| All respondents | 6991 | 100.0 | 6800 | 2.4±.03 |
| Sex | | | | |
| <i>Male</i> | 3388 | 48.7 | 3286 | 2.5±.04 |
| <i>Female</i> | 3573 | 51.3 | 3485 | 2.2±.03 |
| Age at time of survey (years) | | | | |
| 14 | 1084 | 15.5 | 1047 | 2.2±.07 |
| 15 | 1819 | 26.0 | 1178 | 2.1±.04 |
| 16 | 2455 | 35.1 | 2380 | 2.3±.04 |
| 17 or older | 1633 | 23.4 | 1595 | 2.9±.06 |
| Grades (secondary level) | | | | |
| 1–2 | 719 | 10.7 | 691 | 2.5±.09 |
| 3 | 1493 | 22.2 | 1456 | 2.2±.05 |
| 4 | 1964 | 29.2 | 1910 | 2.2±.04 |
| 5 | 2550 | 37.9 | 2487 | 2.5±.04 |
| Ethnicity | | | | |
| White | 5948 | 85.2 | 5798 | 2.2±.02 |
| Non-White | 1035 | 14.8 | 994 | 3.3±.09 |
| Family deprivation | | | | |
| Yes | 960 | 13.7 | 932 | 2.9±.09 |
| No | 6028 | 86.3 | 5867 | 2.3±.03 |
| Depression | | | | |
| Yes | 2833 | 41.5 | 2757 | 2.5±.04 |
| No | 3992 | 58.5 | 3889 | 2.3±.03 |
| Early sexual initiation | | | | |
| <14 years old | 1392 | 20.2 | 1357 | 3.8±.08 |
| 14 years old or older | 5486 | 79.8 | 5365 | 2.0±.02 |
| Months since sexual initiation | | | | |
| 0–11 | 1937 | 28.2 | 1884 | 1.4±.03 |
| 12–23 | 2564 | 37.3 | 2508 | 2.0±.03 |
| 24–35 | 1437 | 20.9 | 1407 | 3.0±.06 |
| ≥36 | 940 | 13.7 | 923 | 4.6±.09 |

* Percentages are proportions among all respondents aged 14 years or older who reported being sexually active.

Table 2. Adjusted mean number of lifetime sexual partners among sexually active participants

| | | Adjusted mean±SD | Mean difference [95 CI] | Risk ratio [95 CI] |
|----------------------------------|-------------------------|-------------------------|------------------------------------|-------------------------------|
| Sex | | | | |
| | <i>Male</i> | 2.46±.05 | (ref) | (ref) |
| | <i>Female</i> | 2.25±.03 | -0.22 [-0.31, -0.12] | 0.87 [0.81, 0.93] |
| Sexual initiation | | | | |
| | <i><14 years old</i> | 2.68±.07 | 0.45 [0.31, 0.60] | 1.20 [1.14, 1.27] |
| | <i>≥14 years old</i> | 2.23±.03 | (ref) | (ref) |
| Time since sexual debut (months) | | | | |
| | <i>0–11</i> | 1.48±.03 | (ref) | (ref) |
| | <i>12–23</i> | 2.00±.04 | 0.53 [0.43, 0.63] | 1.30 [1.19, 1.43] |
| | <i>24–35</i> | 2.91±.07 | 1.43 [1.29, 1.57] | 1.89 [1.74, 2.05] |
| | <i>≥36</i> | 4.00±.11 | 2.52 [2.29, 2.75] | 2.69 [2.44, 2.96] |
| Ethnicity | | | | |
| | <i>White</i> | 2.27±.04 | (ref) | (ref) |
| | <i>Non-White</i> | 2.79±.08 | 0.52 [0.33, 0.71] | 1.22 [1.14, 1.32] |
| Depression | | | | |
| | <i>Yes</i> | 2.45±.04 | 0.17 [0.09, 0.25] | 1.07 [1.04, 1.11] |
| | <i>No</i> | 2.28±.04 | (ref) | (ref) |
| Family deprivation | | | | |
| | <i>Yes</i> | 2.57±.07 | 0.25 [0.12, 0.38] | 1.11 [1.05, 1.17] |
| | <i>No</i> | 2.32±.03 | (ref) | (ref) |

Table 3. Adjusted mean number of lifetime sexual partners and effect of sex on number of lifetime sexual partners by time since sexual debut

| Modality | Females | | | Males | | | Difference in mean differences |
|------------------------------|----------------------|-----------------|----------------------|----------------------|-----------------|----------------------|--------------------------------|
| | Mean CI | Mean Difference | | Mean CI | Mean Difference | | |
| | | Contrast | Mean Diff CI | | Contrast | Mean Diff CI | |
| (1) T1: 0–11 months | 1.37 [1.31, 1.43] | Ref. | | 1.59 [1.49, 1.69] | Ref. | | |
| (2) T2: 12–23 months | 1.95 [1.87, 2.02] | (2)-(1) | 0.57 [0.49, 0.66] | 2.07 [1.94, 2.19] | (2)-(1) | 0.48 [0.32, 0.65] | 0.09 [-0.07, 0.25] |
| (3) T3: 24–35 months | 2.82 [2.67, 2.98] | (3)-(1) | 1.45 [1.29, 1.60] | 3.00 [2.80, 3.20] | (3)-(1) | 1.41 [1.21, 1.62] | 0.03 [-0.20, 0.27] |
| (4) T4: 36 months or more | 3.74 [3.53, 3.95] | (4)-(1) | 2.37 [2.15, 2.59] | 4.27 [3.95, 4.59] | (4)-(1) | 2.68 [2.35, 3.02] | -0.31 [-0.63, 0.00] |

Figure 1. Adjusted mean number of lifetime sexual partners according to time since sexual debut, overall and by sex

