



Neurite Exchange Imaging (NEXI): A minimal model of diffusion in gray matter with inter-compartment water exchange

Ileana O. Jelescu^{a,b,c,*}, Alexandre de Skowronski^a, Françoise Geffroy^d, Marco Palombo^{e,f,g,1}, Dmitry S. Novikov^{h,1}

^a CIBM Center for Biomedical Imaging, Animal Imaging and Technology, Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

^b Department of Radiology, Lausanne University Hospital (CHUV), Lausanne, Switzerland

^c School of Biology and Medicine, University of Lausanne (UNIL), Lausanne, Switzerland

^d NeuroSpin, CEA, Saclay, France

^e School of Psychology, Cardiff University Brain Research Imaging Centre (CUBRIC), Cardiff University, Cardiff, UK

^f School of Computer Science and Informatics, Cardiff University, Cardiff, UK

^g Department of Computer Science, Centre for Medical Image Computing, University College London, London, UK

^h Department of Radiology, Center for Biomedical Imaging, New York University School of Medicine, New York, NY, USA

ARTICLE INFO

Keywords:

Diffusion MRI

Microstructure

Gray matter

Cortex

Exchange

Cell membrane permeability

ABSTRACT

Biophysical models of diffusion in white matter have been center-stage over the past two decades and are essentially based on what is now commonly referred to as the “Standard Model” (SM) of non-exchanging anisotropic compartments with Gaussian diffusion. In this work, we focus on diffusion MRI in gray matter, which requires rethinking basic microstructure modeling blocks. In particular, at least three contributions beyond the SM need to be considered for gray matter: water exchange across the cell membrane – between neurites and the extracellular space; non-Gaussian diffusion along neuronal and glial processes – resulting from structural disorder; and signal contribution from soma. For the first contribution, we propose Neurite Exchange Imaging (NEXI) as an extension of the SM of diffusion, which builds on the anisotropic Kärger model of two exchanging compartments. Using datasets acquired at multiple diffusion weightings (b) and diffusion times (t) in the rat brain *in vivo*, we investigate the suitability of NEXI to describe the diffusion signal in the gray matter, compared to the other two possible contributions. Our results for the diffusion time window 20–45 ms show minimal diffusivity time-dependence and more pronounced kurtosis decay with time, which is well fit by the exchange model. Moreover, we observe lower signal for longer diffusion times at high b . In light of these observations, we identify exchange as the mechanism that best explains these signal signatures in both low- b and high- b regime, and thereby propose NEXI as the minimal model for gray matter microstructure mapping. We finally highlight multi- b multi- t acquisition protocols as being best suited to estimate NEXI model parameters reliably. Using this approach, we estimate the inter-compartment water exchange time to be 15 – 60 ms in the rat cortex and hippocampus *in vivo*, which is of the same order or shorter than the diffusion time in typical diffusion MRI acquisitions. This suggests water exchange as an essential component for interpreting diffusion MRI measurements in gray matter.

1. Introduction

The bedrock of biophysical models of diffusion MRI is water compartmentalization. Morphologically, there are at least three compartments in brain tissue that are essential for interpreting an MRI measurement. The first one is a collection of micron-thin long and often branch-like cellular structures, referred to “cellular processes or projections” – either axons, dendrites or glial cell processes. There, water diffusion is locally unidirectional, and is typically modeled in terms of

the so-called “sticks”, i.e., zero-radius cylinders (Jespersen et al., 2007; Kroenke et al., 2004). The second compartment is cell bodies (soma), which are roughly spherical and of $\sim 15\mu\text{m}$ in diameter (Palombo et al., 2021). Their size is generally comparable with the typical mean squared displacement (the diffusion length) of a water molecule during measurements. The third compartment is the extra-cellular space in which the first two are embedded.

In the white matter (WM), soma are typically neglected due to their relatively small density (5–10% *ex vivo*) (Andersson et al., 2020;

* Corresponding author at: Department of Radiology, CHUV, Lausanne, Switzerland.

E-mail address: ileana.jelescu@chuv.ch (I.O. Jelescu).

¹ These authors contributed equally to this work.

Veraart et al., 2020). Furthermore, the myelin sheath around axons contributes to impermeability (i.e., negligible exchange with the extra-axonal water) over the diffusion MRI-relevant timescales, and thus the sought compartmentalization. Biophysical models of diffusion in WM have therefore gained a lot of traction and are essentially based on what is now commonly referred to as the “Standard Model” (SM) (Novikov et al., 2018a, 2019) of non-exchanging compartments with Gaussian diffusion: a collection of sticks (axons) with some orientation distribution function (ODF); an anisotropic extra-axonal space surrounding each local fascicle (a bundle of sticks) aligned in a given direction; and, if relevant, the “free water” compartment describing the partial volume contribution of the cerebrospinal fluid (CSF), free from the hindrances of the extra-axonal space. Within the SM family, a constellation of implementations has been proposed, each with its own acronym and its own further simplifying assumptions, e.g., on the shape of the ODF for fiber fascicles, or on the relations between the compartmental diffusivities and volume fractions (Fieremans et al., 2011; Jespersen et al., 2010; Novikov et al., 2018b; Reisert et al., 2017; Wang et al., 2011; Zhang et al., 2012). These models are widely used to characterize WM microstructure, and are occasionally applied in gray matter (GM). Physics beyond SM has been revealed in WM, such as the residual non-Gaussian diffusion along sticks (axons) (Arbabi et al., 2020; Fieremans et al., 2016; Lee et al., 2020a).

In this work, we focus on diffusion MRI in GM, which is sufficiently distinct from WM morphologically. This implies rethinking basic microstructure modeling blocks, leading to a different simplified picture of diffusion MRI-relevant microgeometry (Jelescu et al., 2020; Novikov, 2021; Palombo et al., 2020). In particular, at least three contributions beyond the SM need to be considered:

- (i) Exchange across the membrane of cellular processes.
- (ii) Non-Gaussian diffusion along cellular processes – resulting from structural disorder.
- (iii) Signal contribution from cell bodies (soma).

For (i), as myelin content is limited in GM, there is growing evidence that water exchange across the neurite membrane cannot be neglected for typical clinical diffusion times ($20 < t < 80$ ms). Evidence for the deviation from the impermeable stick model for neurites in GM and its relationship to exchange has been highlighted in human cortex, with an estimated characteristic *in vivo* exchange time of 10–30 ms (Veraart et al., 2018a, 2020). Similar exchange time ranges have been reported for perfused neonatal mouse spinal cords (Williamson et al., 2019), while other groups have reported longer exchange times of 100–150 ms in astrocyte and neuron cultures (Yang et al., 2018), rat brain (Quirk et al., 2003) and rat brain cortical cultures (Bai et al., 2018). Much shorter exchange times (3–5 ms) have been recently reported in the rat brain *ex vivo* (Olesen et al., 2022).

For (ii), the intra-compartment structural disorder along the effectively one-dimensional neurite has been suggested in the rat brain *in vivo* based on the power law exponent $\vartheta = 1/2$ (Novikov et al., 2014) in oscillating-gradient diffusion MRI data of (Does et al., 2003), at time scales (0.4–10 ms) long relative to the correlation length, but shorter than typical PGSE (Pulsed Gradient Spin Echo) diffusion times. Recently, a related kurtosis time-dependence with the same exponent, $K(t) \propto t^{-1/2}$, was found in the human cortex at long t (Lee et al., 2020b). However, in that work, the inter-compartment exchange could not be ruled out (generating the sub-leading, faster kurtosis decay $K(t) \propto t^{-1}$ at long t), especially in the face of weak-to-absent time-dependent diffusivity within the diffusion time range accessible with PGSE on a clinical scanner (20–100 ms). Notably, structural disorder in the extra-cellular space is naturally provided by the embedded neurites, which, given the three-dimensional nature of diffusion in this compartment, would lead to diffusion and kurtosis time-dependence as $(\ln t)/t$ at long t (Novikov et al., 2014).

Finally, for (iii), cell bodies (soma) occupy ~10–20% of gray matter by volume (Bondareff and Pysh, 1968; Motta et al., 2019;

Shapson-Coe et al., 2021; Spocter et al., 2012) and may need to be modeled, as proposed in a recent three-compartment model (SANDI) that accounted for this tissue component by representing soma as impermeable spheres (Palombo et al., 2020). The potential issue of inter-compartment exchange was partially circumvented by the use of short diffusion times ($t < 20$ ms). Diffusion in soma was modeled in the Gaussian phase approximation, whereby diffusivity time-dependence in this compartment was retained, but higher-order terms (kurtosis and above) were neglected.

Fig. 1 summarizes the picture underlying each model, the expected functional forms of time-dependent diffusion and kurtosis in the long-time limit as well as the functional form of the powder-averaged signal at high b . Model details and full expressions of the functional forms are provided in the Theory Section.

This work is organized in two main parts.

First, we examine the diffusion and kurtosis time-dependence as well as powder-average signal signature at high b -value in the rat brain *in vivo* in order to assess the importance of the effects (i)–(iii) above. Within the ranges of diffusion weightings (b) and diffusion times (t) explored, we highlight negligible diffusivity time-dependence and a time-dependent kurtosis which can be consistent with the t^{-1} power-law. The two trends combined are compatible with exchange, whereby structural disorder may be less relevant in modeling rat cortex in this diffusion time range. Signal decay curves at high b -values and for different diffusion times are also better accounted for by an exchange term as compared with adding the soma compartment, as also recently shown in *ex vivo* rat cortex (Olesen et al., 2022).

Second, we therefore propose NEXI (Neurite Exchange Imaging), an implementation of the anisotropic Kärger model of exchange accounting only for the effect (i) above, as the minimal model suitable for gray matter. We show in simulations that the availability of multi- b multi- t data is critical for the reliable estimation of model parameters, in particular of the exchange time. We further demonstrate NEXI performance in experimental data. Fit stability permitting, the model could be extended to account for a soma fraction as well (Olesen et al., 2022; Palombo et al., 2020).

For our experiments, we exploit the potential of strong preclinical gradients (1 T/m) to probe a range of short to intermediate diffusion times (10–45 ms) with strong diffusion-weighting (up to $b = 10$ ms/ μm^2). The GM regions of interest (ROIs) are cortex and hippocampus, with corpus callosum, internal capsule and cingulum serving as reference WM ROIs. We take advantage of the rat brain anatomy where cortex is much less affected by partial volume effects with neighboring white matter or CSF than its human counterpart.

2. Methods

2.1. Theory

Let us first provide a theoretical description of the models sketched in Fig. 1, that will be compared throughout this work.

2.1.1. Standard model

To build a GM model, we begin with the Standard Model (Novikov et al., 2019) of brain tissue composed of two compartments (we will neglect the CSF contribution in what follows, assuming voxels are devoid of the CSF contamination). In SM, the intra-neurite compartment – occupying a relative signal fraction f – is modeled as a collection of “sticks” (zero-radius cylinders) where diffusion is unidirectional with diffusivity $D_{i,\parallel}$. The extra-neurite compartment – the immediate environment of sticks – is modeled as a Gaussian anisotropic medium with characteristic diffusivities $D_{e,\parallel}$ and $D_{e,\perp}$ parallel and perpendicular to the local orientation of neurites, respectively. For a neurite ensemble coherently oriented along the unit direction \mathbf{n} , its signal (response) in the unit direction \mathbf{g} is:




Fig. 1. Sketch of relevant features and parameters in the Standard Model, NEXI, SANDI and structural disorder models, along with the associated functional form for time-dependence in diffusivity D and kurtosis K , as well as the functional form of the signal decay in the high- b regime. Note that parameters A and B stand for constants that are different in each instance. The Standard Model considers a collection of impermeable sticks – occupying a relative signal fraction f – where diffusion is Gaussian and unidirectional with diffusivity $D_{i,\parallel}$ and an extra-neurite Gaussian anisotropic compartment with characteristic diffusivities $D_{e,\parallel}$ and $D_{e,\perp}$ parallel and perpendicular to the local neurite orientation, respectively. ODF anisotropy can be characterized by the $l = 2$ order rotational invariant p_2 or its derived dispersion angle $c_2 \equiv \langle \cos^2 \psi \rangle = \frac{2p_2 + 1}{3}$. NEXI considers a collection of randomly-oriented sticks – occupying a relative signal fraction f – where diffusion is unidirectional with diffusivity $D_{i,\parallel}$ and an extra-neurite Gaussian isotropic compartment with characteristic diffusivity D_e . The two compartments exchange with a characteristic time t_{ex} . SANDI considers a similar picture as NEXI, but accounts for a third compartment of spheres of radius R_s (occupying a relative fraction f_s) and neglects inter-compartment exchange. The structural disorder model assumes a certain type of disorder (here short-range disorder) and its signature in diffusion and kurtosis time-dependence, also as a function of the spatial dimensionality in which the disorder is manifest (1D for intra-neurite water, 2D or 3D for extracellular water).

$$\begin{aligned} \mathcal{K}_{SM}(q, t, \mathbf{g} \cdot \mathbf{n}; f, D_{i,\parallel}, D_{e,\parallel}, D_{e,\perp}) \\ = f e^{-q^2 t} D_{i,\parallel}(\mathbf{g} \cdot \mathbf{n})^2 + (1-f) e^{-q^2 t} (D_{e,\parallel}(\mathbf{g} \cdot \mathbf{n})^2 + D_{e,\perp}(1 - (\mathbf{g} \cdot \mathbf{n})^2)) \end{aligned} \quad (1)$$

The signal attenuation (1) for the elementary neurite ensemble and its immediate extracellular space is then convolved on the unit sphere $|\mathbf{n}| = 1$ with the orientation distribution function (ODF) $P(\mathbf{n})$ for the neurites to give the overall SM signal

$$S_{SM}(q, \mathbf{g}, t) = \int \mathcal{K}_{SM}(q, t, \mathbf{g} \cdot \mathbf{n}; f, D_{i,\parallel}, D_{e,\parallel}, D_{e,\perp}) P(\mathbf{n}) d\mathbf{n}. \quad (2)$$

The SM is suitable for white matter, where neurites – mainly myelinated axons – are the dominant structure (vs negligible soma), and the accepted assumption is to neglect the exchange between intra- and extra-neurite spaces due to the myelin sheath.

2.1.2. NEXI: adding exchange to SM

In the GM, most neurites are unmyelinated and inter-compartment water exchange across the cell membrane may be non-negligible for clinical diffusion times ($t > 20$ ms). A model of two exchanging compartments, one being the collection of isotropically oriented neurites (sticks) and the other being the extra-neurite space, is built based on the anisotropic Kärger model for a coherent fiber tract (Fieremans et al., 2010; Kärger, 1985). Namely, the signal (response) from an elementary coherent ensemble of “neurite + its proximal extracellular space” is the result of mixing two anisotropic Gaussian compartments in Eq. (1) by the barrier-limited exchange, with the rates r_{ie} and r_{ei} related by the detailed balance condition $f r_{ie} = (1-f) r_{ei}$:

$$\mathcal{K}(q, t, \mathbf{g} \cdot \mathbf{n}; f, D_{i,\parallel}, D_{e,\parallel}, D_{e,\perp}, t_{ex}) = f' e^{-q^2 t D'_i} + (1-f') e^{-q^2 t D'_e} \quad (3)$$

$$D'_{i/e} = \frac{1}{2} \left\{ D_i + D_e + \frac{1}{q^2 t_{ex}} \mp \left[\left(D_e - D_i + \frac{2f-1}{q^2 t_{ex}} \right)^2 + \frac{4f(1-f)}{q^4 t_{ex}^2} \right]^{1/2} \right\} \quad (4)$$

$$f' = \frac{1}{D'_i - D'_e} [f D_i + (1-f) D_e - D'_e] \quad (5)$$

where $D_i \equiv D_{i,\parallel}(\mathbf{g} \cdot \mathbf{n})^2$, $D_e \equiv D_{e,\parallel}(\mathbf{g} \cdot \mathbf{n})^2 + D_{e,\perp}(1 - (\mathbf{g} \cdot \mathbf{n})^2)$ and $t_{ex} = 1/r$, with $r = r_{ie} + r_{ei}$. In other words, the bi-exponential expression in Eq. (3) is reminiscent of the SM expression in Eq. (1), but with *apparent* fractions and diffusivities that depend on all model parameters as well as on diffusion wavenumber q and on the characteristic exchange time t_{ex} . Eqs. (3)–(5) are valid in the narrow pulse approximation. In this work this approximation is *a posteriori* justified with estimated values of t_{ex} notably exceeding pulse width δ .

In principle, the response function (3) should be convolved with the GM neurite ODF to get the overall signal in direction \mathbf{g} , as in Eq. (2). Since diffusion anisotropy in GM can be variable across cortical layers, though overall negligible (FA~0.15), we choose to consider the orientational average (the so-called powder average) of the signal instead, which is independent of the ODF. By the same token, we further assume the extra-neurite space to be isotropic $D_{e,\parallel} = D_{e,\perp} \equiv D_e$. This approximation helps reduce the number of parameters to be estimated and raises the precision on the remaining ones. The model parameters are therefore $\mathbf{p} = [f, D_{i,\parallel}, D_e, t_{ex}]$ (Fig. 1). Hence, the model signal equation, to be fit to the powder-averaged measured signal \bar{S} , is:

$$\bar{S}(q, t) = S|_{q=0} \cdot \int_0^1 \mathcal{K}(q, t, \mathbf{g} \cdot \mathbf{n}; \mathbf{p}) d(\mathbf{g} \cdot \mathbf{n}) \quad (6)$$

Technically, this means that we are only using the $l = 0$ rotational invariant of the overall directional signal, and discarding potential information residing in $l = 2, 4, \dots$ invariants. Nonetheless, the latter are small in GM, and are not expected to contribute much information to the original directional signal.

We call this implementation, Eqs. (3)–(6), NEXI.

NEXI kurtosis. One of the assumptions behind the Kärger model, and thus behind NEXI, is time-independent diffusivity. If this condition is met, the NEXI kurtosis of Eq. (6) (see Appendix for the derivation) is:

$$K(t) = K_0 \cdot \frac{2t_{ex}}{t} \left[1 - \frac{t_{ex}}{t} (1 - e^{-t/t_{ex}}) \right] + K_\infty. \quad (7)$$

It has two contributions: the inter-compartment heterogeneity $\sim K_0$ which decays to zero as $1/t$ at long times $t \gg t_{ex}$ as a result of exchange, identical to that found in Fieremans et al. (2010), Jensen et al. (2005);

and the constant offset K_∞ that captures potential residual sources of kurtosis in the limit $t \gg t_{ex}$. Residual kurtosis could stem from partial volume (macroscopic heterogeneity) within a voxel (in our experiment, this is deemed small due to homogeneity of rat GM at our imaging resolution) and, in the case of powder-averaged signal, from non-exchanging microscopic anisotropic structures (Szczepankiewicz et al., 2016). Indeed, the powder-averaged signal, Eq. (6), has residual kurtosis, as derived in the Appendix; it originates from the assumption that exchange happens only within a local neighborhood, between a single stick (neurite) and its accompanying extracellular space. The actual tissue geometry may differ from this latter assumption. If multiple sub-units of “neurite + extracellular space” with different orientations fall within the same volume $\sim L^3(t)$, where $L \sim \sqrt{6D_e t}$ is the diffusion length, then exchange can occur across multiple neurites. Since the extra-cellular space is a connected medium, at long times $t \gg t_{ex}$, exchange would coarse-grain the medium fully, yielding $K_\infty = 0$.

We note that the absence of exchange across multiple neurites is a mathematical construct to make the problem solvable from an analytical point of view. A plausible picture is naturally one with exchange across multiple neurites, given the random orientations of neurites and the tight network they constitute. This plausible picture led us to assume isotropic ECS diffusion and $K_\infty = 0$. At reasonably short diffusion times as the ones used here, the two pictures are compatible, but for longer diffusion times, exchange across multiple neurites is of course expected.

Finally, we also note that even for anisotropically oriented neurites, the kurtosis of the non-powder-averaged signal will have the functional form of Eq. (7), with the parameters K_0 and K_∞ depending on the ODF but the exchange time being ODF-independent. Hence, we will use Eq. (7) to analyze mean kurtosis. Fitting Eq. (7) to the measured mean kurtosis can provide a complementary estimate of t_{ex} stemming from the low- b regime (within the convergence radius of the cumulant expansion), to be compared to the one from the full NEXI model Eq. (6). We label this estimate $t_{ex}^{K(t)}$.

$$\bar{A}_s(b, D_s, R_s) \approx \exp \left\{ -\frac{2g^2 R_s^4}{D_s} \sum_{m=1}^{\infty} \frac{\alpha_m^{-4}}{\alpha_m^2 - 2} \cdot \left[2\delta - \frac{R_s^2}{\alpha_m^2 D_s} \left(2 + e^{-\alpha_m^2 D_s (\Delta - \delta) / R_s^2} - 2e^{-\alpha_m^2 D_s \delta / R_s^2} - 2e^{-\alpha_m^2 D_s \Delta / R_s^2} + e^{-\alpha_m^2 D_s (\Delta + \delta) / R_s^2} \right) \right] \right\}$$

High- b scaling. An independent hallmark of the model (3)–(6) is the functional form

$$\bar{S}|_{b \rightarrow \infty} = \sqrt{\frac{\pi}{4}} \frac{f}{\sqrt{bD_{i,\parallel}}} e^{-\frac{t(1-f)}{t_{ex}}} \times \left[1 + \frac{2(1-f)t/t_{ex} + f(1-f)(t/t_{ex})^2}{bD_{e,\perp}} + O\left(\frac{1}{b^2}\right) \right] \quad (8)$$

of its expansion in the inverse powers of the diffusion weighting parameter $b = q^2 t$. The first term in the square brackets, corresponding to $b^{-1/2}$ decrease, is the signature of impermeable sticks (Callaghan et al., 1979; McKinnon et al., 2017; Veraart et al., 2019). The subsequent $b^{-3/2}$ term arises due to slow exchange, such that $t/t_{ex} \ll bD$, where D is the smallest of the compartment diffusivities; the lower bound is practically set by $D_{e,\perp} = D_e$. The $\sim t/t_{ex}$ term was obtained by Veraart et al. (2020), and the $\sim (t/t_{ex})^2$ term by Olesen et al. (2022).

We note that the Kärger Model is treated in the narrow pulse approximation regime. This condition translates into $\delta \ll t_{ex}$, which will be justified a posteriori by comparing experimental values of $\delta = 4 - 4.5$ ms and $t_{ex} = 20 - 40$ ms. However, a numerical solution to the Kärger model in the finite pulse regime (Olesen et al., 2022) could be implemented in the case of longer diffusion pulses (or shorter exchange times).

2.1.3. Structural disorder

The assumption of Gaussian compartments may break in the presence of irregularities on length scales similar to the diffusion length, such as dendritic spines and neurite beading. While there is no analytical formula to describe the signal in this case exactly, the relative importance of non-Gaussian effects can be determined by examining the

diffusivity and kurtosis time-dependence at diffusion times $t \gg t_c$, the time t_c to diffuse past the disorder correlation length (Novikov et al., 2014). In particular, kurtosis should follow a $t^{-1/2}$ functional form in the case of 1-dimensional disorder (Dhital et al., 2018; Lee et al., 2020b):

$$K(t)|_{t \gg t_c} \simeq A \cdot t^{-\vartheta} + K_\infty, \quad \vartheta = 1/2 \quad (9)$$

or $(\ln t)/t$ in the case of 2-dimensional disorder (Burcaw et al., 2015; Lee et al., 2020b). The offset K_∞ arises in the case of residual voxel heterogeneity in the long-time limit, similar to that in Eq. (7). If the relevant correlation time t_c for diffusion across these structural irregularities is of the order of t_{ex} , the competing effects of coarse-graining over the structural disorder and of exchange are both contributing significantly to the time-dependence of the measured $K(t)$, which complicates the interpretation (Lee et al., 2020b).

2.1.4. SANDI: adding soma

A three-compartment model (SANDI) was proposed as an extension of the SM that models the total direction-averaged signal as the sum of three non-exchanging compartments (Palombo et al., 2020): (i) randomly oriented sticks with intra-stick axial diffusivity $D_{i,\parallel}$ and relative signal fraction f ; (ii) restriction in sphere of apparent radius R_s , fixed intra-sphere diffusivity $D_s = 3 \frac{\text{Lum}^2}{\text{ms}}$ and relative signal fraction f_s (modeled in the Gaussian phase approximation); (iii) Gaussian isotropic diffusion in the extracellular space with diffusivity D_e and relative signal fraction $f_e = 1 - f - f_s$. SANDI provides estimates for the five model parameters: $[f, f_s, D_{i,\parallel}, D_e, R_s]$ which by design should be independent of diffusion time. The direction-averaged SANDI signal is:

$$\frac{\bar{S}(b)}{S(0)} = f \cdot \sqrt{\frac{\pi}{4}} \frac{1}{\sqrt{bD_{i,\parallel}}} \text{erf}\left(\sqrt{bD_{i,\parallel}}\right) + f_s \cdot \bar{A}_s(b, D_s, R_s) + f_e \cdot e^{-bD_e} \quad (10)$$

where

with δ and Δ the diffusion gradient pulse duration and separation, respectively, g the product of diffusion gradient amplitude and gyromagnetic ratio, α_m the m -th root of $\frac{1}{2} J_{3/2}(\alpha) = \alpha J'_{3/2}(\alpha)$, and $J_n(x)$ the Bessel function of the first kind. In practice, summation up to $m = 20$ roots is sufficient for a good approximation.

2.2. Experimental

Animal experiments were approved by the Service for Veterinary Affairs of the canton of Vaud. Six Wistar rats (Charles River) weighing 250–300 g were scanned on a 14 T Bruker system equipped with 1 T/m gradients (Resonance Research Inc.) using a home-built surface quadrature transceiver. Rats were set up and maintained under isoflurane anesthesia, and body temperature was monitored and maintained around 38 °C for the duration of the experiment. Diffusion MRI data were acquired using a PGSE EPI sequence, with parameters provided in Table 1. All six datasets were used to assess the behavior of time-dependent diffusion and kurtosis, while four datasets (labeled 1–4) were used for high- b signal analysis, SANDI and NEXI estimations. In datasets 5 and 6 we prioritized a larger number of diffusion times over b -values to capture trends in $D(t)$ and $K(t)$.

Images were denoised using MP-PCA and corrected for Rician bias (Veraart et al., 2016b, 2016a), for Gibbs ringing (Kellner et al., 2016) and for motion (Jenkinson et al., 2002). No strong distortions due to eddy currents were observed.

ROIs in both white matter – internal capsule (IC), corpus callosum (CC) and cingulum (Cg) – and gray matter – cortex (CTX) and hippocampus (HPC) – were manually drawn.

Table 1
Acquisition parameters for the six datasets included in this study.

# Datasets	3	1	1	1
Dataset label	1–3	4	5	6
TE (ms)	50	58	52	58
δ (ms)	4.5	4.5	4	4
Δ (ms)	12, 20, 30, 40	11, 25, 45	10, 15, 20, 25, 30, 40	10, 15, 20, 25, 30, 35, 40, 45
b -values (ms/ μm^2)	1, 2.5, 4, 5.5, 7, 8.5, 10	1, 2.5, 5, 6, 7, 8, 9, 10	1, 1.8, 2.5	1, 1.4, 2.5
Dirs. per shell	24	24	24	24
TR (ms)	2500	3000	2500	3000
In-plane res (mm ²)	0.2 \times 0.2	0.25 \times 0.25	0.2 \times 0.2	0.25 \times 0.25
Slice thickness (mm)	0.5	0.8	0.5	0.8

2.3. Impact of each contribution (i)–(iii): exchange, structural disorder & soma

Standard Model: To examine potential time-dependence and b -range dependence of SM estimates in various brain regions, for the four datasets with $b_{\max} = 10$ ms/ μm^2 , the SM parameters were estimated for each diffusion time using likelihood maximization in the rotational invariant framework RotInv using up to $l = 4$ (Novikov et al., 2018b) on various data subsets ($b_{\max} = 2.5, 6$ or 10). RotInv was implemented in Matlab using non-linear least-squares minimization with a trust-region-reflective algorithm ('lsqnonlin' function). The $D_{i,\parallel} > D_{e,\parallel}$ solution was favored by choosing a random algorithm initialization that met this inequality. The time- and b_{\max} -dependence of model parameters were evaluated in the various brain ROIs. Notable time-dependence of SM parameter estimates was tested via the slope – and its uncertainty – of a simple linear regression.

Time-dependent diffusion and kurtosis: To determine the extent to which inter-compartment exchange and/or structural disorder are relevant in various brain regions, diffusion and kurtosis tensors were estimated for each diffusion time using shells up to $b = 2.5$ ms/ μm^2 and a weighted linear least-squares algorithm custom-written in Matlab (Veraart et al., 2013), from which mean diffusivity and kurtosis were derived. The time-dependence of these metrics was evaluated in the various brain ROIs. Notable time-dependence of diffusivities was first tested via the slope – and its uncertainty – of a simple linear regression. To establish the dominant power-law of $K(t)$ decay, Eq. (9) with variable ϑ was fit to the experimental $K(t)$ using the 'lsqnonlin' function a trust-region reflective algorithm in Matlab. The Kärger time-dependent kurtosis, Eq. (7), was also fit to the experimental $K(t)$, either allowing for nonzero K_{∞} or setting it to zero. The 1D structural disorder functional form (Eq. (9) with $\vartheta = 1/2$) was also fit to the measured $K(t)$ for comparison.

For SM estimates, as well as for $D(t)$ and $K(t)$, the uncertainty on fit parameters was estimated using a bootstrapping method where random noise with variance equal to the residual variance was added to the datapoints for $N = 1000$ realizations, from which mean and standard deviation of the estimated parameters were extracted.

Soma vs exchange: The signal was averaged over each shell $\bar{S}(b, t)$ to fit either the SANDI model, Eq. (10) (Palombo et al., 2020), for each diffusion time t separately, or the NEXI model, for all diffusion times t jointly (see Section 2.4 on NEXI parameter estimation below).

The SANDI fit was performed using its implementation in the accelerated microstructure imaging via convex optimization (AMICO) framework in Python 3.5 (Daducci et al., 2015), publicly available at : <https://github.com/daducci/AMICO/wiki/Fitting-the-SANDI-model>. Briefly, AMICO (Daducci et al., 2015) rewrites Eq. (10) as a linear system $Ax = y$, where $A = [A_{\text{stick}}, A_{\text{sphere}}, A_{\text{extra}}]$ is a matrix whose columns contain simulated signals of each compartment (stick, sphere or isotropic Gaussian), y is the vector of measured signals, and x the unknown contributions. To build A , we used a dictionary of signals simulated using: 5 values of $D_{i,\parallel}$ linearly spaced within the interval $[0.25, 3]$ $\mu\text{m}^2/\text{ms}$ (namely p_{stick}) for A_{stick} ; 5 values of R_s linearly spaced within the interval $[1, 12]$ μm (namely p_{sphere}) for A_{sphere} ; and 5 values of D_e linearly spaced

within the interval $[0.25, 3]$ $\mu\text{m}^2/\text{ms}$ (namely p_{extra}) for A_{extra} . The elements of x are then estimated using non-negative least squares with Tikhonov regularization (Efron et al., 2004) (regularization parameter $\lambda_2 = 0.005$) using the Lasso function implemented in the SPAMS optimization toolbox (<http://spams-devel.gforge.inria.fr>). From x , we then computed the SANDI model parameters as: $f = \frac{\sum_{i=1}^5 x_i}{\sum_{i=1}^{15} x_i}$; $f_s = \frac{\sum_{i=6}^{10} x_i}{\sum_{i=1}^{15} x_i}$; $D_{i,\parallel} = \frac{\sum_{i=1}^5 x_i p_{\text{stick},i}}{\sum_{i=1}^5 x_i}$; $R_s = \frac{\sum_{i=6}^{10} x_i p_{\text{sphere},i}}{\sum_{i=6}^{10} x_i}$; $D_e = \frac{\sum_{i=11}^{15} x_i p_{\text{extra},i}}{\sum_{i=11}^{15} x_i}$. The fit provided estimates for the five model parameters: $[f, f_s, D_{i,\parallel}, R_s, D_e]$ in the cortex and hippocampus at each investigated diffusion time. The dependence of SANDI model parameters on diffusion time was quantified by computing the mean percentage difference of parameter estimates at each time with respect to the shortest diffusion time ($t = 12$ ms), and, in parallel, by performing one-way ANOVA as a function of time and reporting the significant differences pairwise for available diffusion times. Finally, the significance of a linear trend of model parameters over time was also calculated.

The SANDI parameter estimates at the shortest diffusion time were also used to predict the signal decay in the cortex at longer diffusion times, and compared to experimental outcomes, as suggested in Olesen et al. (2022).

The NEXI fit was performed using a non-linear least-squares (NLLS) optimization based on a quasi-Newton algorithm without constraints, implemented as 'fminunc' function in Matlab.

The performance of SANDI and NEXI to capture the deviation from the stick model at high b -values and the qualitative signal decay curves across multiple diffusion times was evaluated and compared.

2.4. NEXI parameter estimation

Simulations: Synthetic signals ($N = 10^4$) were generated based on Eqs. (3)–(6) assuming a protocol of $b = 0$ and seven shells at $b = 1, 2.5, 4, 5.5, 7, 8.5$ and 10 ms/ μm^2 , four diffusion times ($t = 12, 20, 30, 40$ ms) and a realistic SNR level of 100 – as estimated from experimental data in cortex following MP-PCA denoising and powder-averaging – see Experimental paragraph below. The ground truth was either fixed to $[t_{\text{ex}}, D_{i,\parallel}, D_e, f] = [20, 2.5, 0.75, 0.34]$ with only the noise realization changing for each iteration, or randomly chosen within physical ranges, that is $t_{\text{ex}} \in [5, 120]$, $D_{i,\parallel} \in [1.5, 3]$, $D_e \in [0, 1.5]$ and $f \in [0.1, 0.9]$, thus enforcing the $D_{i,\parallel} > D_e$ solution of the NEXI model. The exploration of disjoint intervals with $D_{i,\parallel} > D_e$ was supported by experimental data where, when repeating the NEXI estimation by varying the algorithm initialization (within full ranges $D_{i,\parallel}, D_e \in [0, 3]$) the mode of the outcome distribution yielded $D_{i,\parallel} > D_e$ in both cortex and hippocampus (see Results and Fig. S9).

Parameter estimation was done either based on the signals for each diffusion time separately (as in standard multi-shell datasets) or jointly. NLLS used a trust-region-reflective algorithm with box constraints for multi-shell data and quasi-Newton algorithm without constraints for multi-shell multi- t data. We also tested the performance of a deep learning (DL) algorithm for parameter estimation, in terms of precision and accuracy with respect to the more widespread NLLS approach, in the




Fig. 2. Time-dependence of SM parameters, also as a function of maximum b -value available, in two GM ROIs: cortex and hippocampus. Symbols: mean \pm std across rats. Solid line: linear fits.

perspective of providing a fast implementation of NEXI with on-the-fly estimation of model parameter maps (Supplementary Methods).

Impact of b range. In the case of the joint diffusion times fit, we further evaluated the impact of the maximum b -value on NEXI estimates by retaining only subsets of the data for the estimation: $b_{\max} = 2.5$ (2 shells), 5.5 (4 shells), or 10 $\text{ms}/\mu\text{m}^2$ (7 shells). For completeness, we also compared the performance of the estimation for datasets with variable b_{\max} but same number of equally spaced shells ($N_{\text{shells}} = 7$).

Experimental: The signal was averaged over each shell $\bar{S}(b, t)$ to be used for NEXI parameter estimation. All four model parameters were estimated with NLLS (and DL) by using all shells and diffusion times jointly. This joint fit was performed on a voxel-wise basis to generate parametric maps. The t_{ex} estimate from NEXI was also compared to $t_{\text{ex}}^{K(t)}$.

The NLLS fit was unconstrained. To test the impact of algorithm initialization on the outcome, NEXI was fit to the average signal in cortex and hippocampus (separately) using $N = 100$ random initializations covering the entire range of physical values for each parameter ($D_{i,\parallel}, D_e \in [0, 3]$, $f \in [0, 1]$, $t_{\text{ex}} \in [0, 100]$). Based on this outcome, the range of NLLS fit initializations was further reduced to ($D_{i,\parallel} \in [1.5, 3]$, $D_e \in [0.5, 1.5]$, $f \in [0.1, 0.9]$, $t_{\text{ex}} \in [5, 60]$) to limit the impact of spurious noise-driven minima in voxel-wise fits.

The agreement between membrane permeability estimates derived from the experimental t_{ex} values and existing literature for physiologically relevant membrane permeability values in healthy cells were compared.

2.5. Histology

A fixed brain sample from a 6 month-old rat was cut into 30 μm -thick slices using a cryomicrotome, and positioned on glass slides. Then, immunohistochemical stainings were performed to label various microstructure features. After a step of blocking non-specific antigens with donkey serum 5% buffer, with detergent (Triton, Sigma Aldrich, X-100, 1% 2 h incubation), a quadruple staining was prepared. It included labeling for microglial cells (anti-Iba 1, AbCam ab5076, 1/500 dilution), astrocytes (anti-GFAP, AbCam, ab7260, 1/500 dilution), neuron microfilaments (anti-NF, AbCam ab4680, 1/2000 dilution) and neuron nuclei (anti-NeuN, Millipore, MAB377X Alexa488, 1/100 dilution). Each antibody was incubated for one hour, followed by two steps of washing with PBS. Secondary antibodies were incubated at the same time, with anti-NeuN already coupled with Alexa488. We used donkey anti-chicken Cy5 (Millipore, AP194C), donkey anti-rabbit Alexa350 (ThermoFisher, 1710039) and donkey anti-goat Alexa647 (AbCam, ab150135).

Slices were mounted with Permafluor (ThermoFisher, TA-030-Fr), then fluorescence microscopy images acquired with an Axio Vision Observer microscope at x20 magnification (Carl Zeiss).

The patterns of staining intensity across the brain (mainly cortex and hippocampus) were compared to patterns of NEXI model parameters, in particular neurite density f .

3. Results

3.1. Impact of exchange, structural disorder and soma

3.1.1. Time-dependent standard model parameters in GM and WM

To underline the limits of applicability of the Standard Model, Eqs. (1) and (2), we evaluated SM parameter estimates in GM vs WM ROIs at different diffusion times and b -value ranges.

Geometric parameters f and c_2 . The apparent intra-neurite water fraction f decreased with increasing diffusion time irrespective of b_{\max} in both GM (Fig. 2) and WM (Fig. S1). The decrease had a slow rate of $(2 - 3) \cdot 10^{-3} \text{ ms}^{-1}$ for all ROIs (Table S1), which would translate into an underestimation of the fraction by 0.2 points for a diffusion time of 100 ms. Neurite alignment c_2 increased with diffusion time, most markedly in GM.

Compartment diffusivities. The trends for compartment diffusivities as a function of diffusion time and b_{\max} were more complex. Significant decreases in parallel diffusivities (both intra- $D_{i,\parallel}$ and extra-neurite $D_{e,\parallel}$) with longer times were found in GM, but much more markedly beyond the second-order cumulant expansion regime ($b_{\max} \geq 6$ (Fig. 2, Table S1). The extra-axonal radial diffusivity $D_{e,\perp}$ increased with longer times only for $b_{\max} \geq 6$. Time-dependence of compartment diffusivities was less pronounced in WM ROIs than GM ROIs, with only cingulum showing a reliable trend (Fig. S1, Table S1).

3.1.2. Exchange vs structural disorder: time-dependent diffusion and kurtosis

As all SM parameter estimates significantly depended on diffusion time in GM ROIs, we explored whether inter-compartment exchange and/or intra-compartment non-Gaussian diffusion were relevant mechanisms in GM. To this end, we examined the time-dependence of mean diffusivity and kurtosis.

All tensor estimates were consistent across animals and displayed a reproducible ordering of rat brain structures from most coherent to least coherent: internal capsule had highest anisotropy and kurtosis, followed by corpus callosum, cingulum, cortex and finally hippocampus.

No significant time-dependence of diffusivities could be measured over the 10–45 ms range, based on slopes of linear fits (Table S2). Fitting the generic power-law formula (Eq. (9) with the exponent ϑ as a free parameter) to $\text{MD}(t)$ yielded unreliable estimates with the exception of D_{∞} (Fig. 3A for GM and Fig. S2 for WM).




Fig. 3. Mean diffusivity and kurtosis as a function of diffusion time, in the cortex (Ctx) and hippocampus (Hpc), averaged across animals. Fit parameters (mean \pm std) for each functional form are collected in the tables. A: Fitting the power-law to MD yielded very large exponent ϑ (with high variability), mainly driven by the diffusion times 10–20 ms. B: The behavior of MK was markedly different, with a decay throughout the 10–45 ms span. As a result, the power-law fit to MK yielded exponent ϑ close to 1 (and with reduced variability). C-D: The direct fitting to either the KM kurtosis (imposing $K_\infty = 0$) or the 1D structural disorder form ($\vartheta = 1/2$) showed both approaches fit the data similarly, though KM kurtosis captures the curve at the longest times (leftmost of x-axis) better. Releasing $K_\infty = 0$ in the KM results in a similar curve to 1D disorder but with poorer parameter estimates (3 free parameters instead of 2). [The number of datasets N averaged for each diffusion time t is variable: t(N) = 10(2), 11(1), 12(3), 15(2), 20(5), 25(3), 30(5), 35(1), 40(5), 45(2), see also Table 1].

Mean kurtosis on the other hand showed marked time dependence over the 10–45 ms range (Fig. 3 for GM and Fig. S2 for WM). The important observation is that the exponents of decay of MD(t) and MK(t) at long times are very different from each other, with MD at a plateau and MK still decreasing markedly. While MD(t) may still exhibit some time-dependence due to structural disorder at short times ($t < 20$ ms), this effect is practically fully coarse-grained at $t > 20$ ms such that each compartment can be approximated as Gaussian. In contrast, MK(t) has sustained sources of time-dependence throughout the 20–45 ms interval suggesting an additional mechanism: inter-compartment exchange.

We note however that the fit of a three-parameter power-law to the average MK(t) from all datasets yielded an exponent between 0.5 and 1, with large uncertainty for all three parameters (Fig. 3B). Presumably, a direct comparison of the two possible power laws governing MK(t) in the GM, between exchange ($\sim 1/t$) and 1D structural disorder ($\sim 1/\sqrt{t}$), was not conclusive as the long-time limit was likely not reached at the longest diffusion time (45 ms); e.g., in the Kärger model the sub-leading negative $1/t^2$ term in Eq. (7) still weighed in significantly (with a numerical value of 0.2–0.3 vs 0.5–0.7 for the leading term, at $t = 45$ ms and assuming $t_{ex} \sim 20$ ms), which may explain the curvature for KM kurtosis in Fig. 3C. Remarkably, fitting MK(t) in each dataset individually yielded a dominant apparent exponent of 0.3 for the large majority of datasets, which was further accompanied by a $K_\infty = 0$ estimate. Simulating time-dependent MK(t) in the diffusion time range 20 – 45 ms based on the Kärger model of exchange (with $K_0 = 0.75$; $t_{ex} = 25$; $K_\infty = 0$ as ground truth) and fitting a generic power-law to it also yielded an apparent exponent of 0.3, therefore consistent with our data and the exploration of an intermediate time regime (Supplementary Fig. S3).

In that regard, the Kärger kurtosis enforcing $K_\infty = 0$ was the functional form that captured best the decay of $K(t)$ at the longest diffusion times available (leftmost on the plots, Fig. 3C and D). Allowing for nonzero K_∞ when fitting the NEXI kurtosis (Eq. (7)) to the measured $K(t)$ yielded extremely large uncertainty on both K_0 and t_{ex} estimates – up to 1800% – obscuring their interpretation completely (Fig. 3). A finite $K_\infty \sim 0.3$ is associated with very short exchange time estimates (1 – 3 ms) which (i) are too short to be reliably estimated from our diffusion time range and (ii) correspond to a timescale where different mechanisms may also come into play, such as structural disorder.

Setting $K_\infty = 0$ enabled a more robust fit. The $K_\infty = 0$ approximation is justified in the context of the reasonable picture of a fully mixed (Gaussian) medium in a rat GM voxel at infinitely long diffusion times, due to intra-/extracellular exchange and a fully connected extracellular space. The GM yielded exchange times in the same range as the diffusion times explored and thus with better precision, e.g., $t_{ex} = 21 \pm 4$ ms in cortex and $t_{ex} = 16 \pm 3$ ms in hippocampus. Estimated exchange times were longer ($t_{ex} > 80$ ms) in the internal capsule and the corpus callosum but were also associated with a fairly large uncertainty ($\sim 35\%$) likely related to the inappropriate diffusion time range (10 – 45 ms) to estimate long exchange times. The cingulum (WM) displayed an intermediate behavior between GM and WM, with $t_{ex} = 43 \pm 24$ ms. This suggests that the myelin sheath plays a significant role in slowing down inter-compartment water exchange – the IC and CC are most myelinated, the GM the least, while CG may be affected by partial volume effects with neighboring GM due to its thinner structure.

In light of the possible MD(t) time-dependence for $t < 20$ ms, we further estimated the exchange time by fitting the Kärger kurtosis to MK(t) using $t \geq 20$ ms data only, which yielded longer yet consistent exchange times of $t_{ex} = 41 \pm 18$ ms in cortex and $t_{ex} = 28 \pm 15$ ms in hippocampus (Supplementary Fig. S3). As expected, the diffusion time range affects the exchange time estimation and excluding short diffusion times yields a longer exchange time estimate.

3.1.3. Exchange vs soma

To explore whether soma and/or exchange are relevant features to explain the diffusion signal in GM, we assessed whether the estimated SANDI model parameters showed any significant time dependence, and then compared the quality of fit and predictions of the SANDI and NEXI models at high b-values.

The SANDI model was applied to GM ROIs at different diffusion times (Fig. 4). We overall observed statistically significant time dependence of all SANDI model parameters, based on slopes of linear fits (Table S3). However, for all parameters except R_s and f , the mean absolute percentage differences of the values at each time point with respect to the first time point at $t = 12$ ms were within 10% at $t = 20$ ms, suggesting that estimates of those SANDI parameters are stable for $t \leq 20$ ms (see Fig. S4). In particular, for $D_{i,\parallel}$ and D_e , this variability drops further down to within $\pm 5\%$ for $t \leq 20$ ms.




Fig. 4. Time-dependence of SANDI model parameters for the ROI in the cortex and hippocampus. Open symbols: mean value; error bars: standard deviation over all the voxels within the ROI for each investigated rat. Note that data from Rat #1–3 were acquired at higher resolution (blue, red, yellow – voxel size $0.2 \times 0.2 \times 0.5 \text{ mm}^3$) and different diffusion times than Rat #4 (purple – voxel size $0.25 \times 0.25 \times 0.8 \text{ mm}^3$). See Table 1 for further details on the acquisition.




Fig. 5. A. Various models were fit to the average signal in the cortex (Rat #2) at $t = 12 \text{ ms}$: SANDI, Eq. (10), and NEXI, Eq. (6), covering the full b -value range; the impermeable stick approximation (Callaghan’s model); NEXI approximation at high b , Eq. (8); and the NEXI-derived diffusivity + kurtosis approximation at low b (Appendix). B. Zoom-in of the black framed region in panel A. Both SANDI and NEXI explain the data at a single diffusion time well. Callaghan’s model does not describe diffusion signal decay in the cortex appropriately due to the signal’s notable curvature with respect to $b^{-1/2}$, cf Eq. (8). The NEXI low- b and high- b approximations are reasonable in their respective regimes. It should be noted the low- b approximation is derived from NEXI parameter estimates obtained over the entire b -value range available hence some mismatch with the experimental datapoints. The mismatch is reduced for longer diffusion times, where the Gaussian compartment approximation may be more suitable (Fig. S10). Estimated model parameters, underlying the plotted curves: SANDI: $f = 0.22$; $f_s = 0.41$; $D_{i,||} = 2.3$; $R_s = 9.3$; $D_e = 0.54$; NEXI: $f = 0.35$; $D_{i,||} = 3$; $D_e = 0.73$; $t_{ex} = 20$; Sticks: $f = 0.25$; $D_{i,||} = 2.4$; High b NEXI approx.: $f = 0.29$; $D_{i,||} = 1.9$; $D_e = 0.35$; $t_{ex} = 12$.

A one-way ANOVA analysis with Bonferroni correction for multiple comparison further showed that the estimates of all SANDI model parameters (except R_s) were not statistically different between $t = 12$ and 20 ms (see Fig. S5). In contrast, R_s estimates showed significant increase with increasing diffusion times. This is to some extent expected: R_s is an MR apparent estimate of the sphere radius, weighted by the tail of the distribution, and such weighting depends on the pulse timings (Alexander et al., 2010). Therefore, our findings suggest limited bias due to exchange at $t < \sim 20 \text{ ms}$ for all SANDI model parameters, except R_s , *in vivo* in rat.

Accounting for either a soma compartment or exchange between neurites and the extracellular space captures well the curvature of the signal decay as function of $b^{-1/2}$, which distinguishes the diffusion behavior in gray matter from that in white matter, where $\bar{S}_{|b \rightarrow \infty} \propto b^{-1/2}$ (an

asymptotically straight line). The quality of the fit for SANDI, NEXI and other signal approximations at a single diffusion time is shown in an example dataset (Fig. 5).

Remarkably though, a SANDI fit at short diffusion time predicted a qualitative trend of higher diffusion signals at longer diffusion times, while the experimental trend was the opposite. In this respect, the NEXI model of exchange explained signal decay curves for multiple diffusion times better than SANDI (Fig. 6).

3.2. NEXI parameter estimation

Since inter-compartment exchange appears to be a relevant mechanism to explain the diffusion signal both in the low-order approximation and at high b -values, we assess the performance of NEXI – a biophysi-




Fig. 6. A. The SANDI model was fit to the average signal in the cortex (Rat #2) at $t = 12$ ms. Estimated model parameters [$f = 0.22$; $f_s = 0.41$; $D_{i,\parallel} = 2.3$; $R_s = 9.3$; $D_e = 0.54$] were used to predict the signal for longer diffusion times (solid lines), as suggested by Olesen et al. (2021). Qualitatively, SANDI predicted higher signal at longer diffusion times, which was opposite to the experimental pattern of increasingly reduced signal with longer diffusion time (dots). B. The NEXI model of exchange was fit to data from all diffusion times jointly (solid lines). The estimated model parameters were [$f = 0.29$; $D_{i,\parallel} = 2.5$; $D_e = 0.74$; $t_{ex} = 44$]. This model explained decay curves at different diffusion times well, though the agreement was poorer at the highest b -values, potentially due to an imperfect correction for Rician noise floor or to soma. All units in μm , ms and $\mu\text{m}^2/\text{ms}$.




Fig. 7. Simulation results fitting multi-shell data for each diffusion time separately using NLLS, without noise (A) or with $\text{SNR} = 100$ (B). Displayed is the ground truth (GT) vs estimation for 10^4 set of random parameters. Markers correspond to the median & IQR in the corresponding intervals. Black lines are the ideal estimation $\pm 10\%$ error. In all cases, the precision is good on D_e and acceptable on f . However, in a finite SNR case, $D_{i,\parallel}$ and t_{ex} cannot be estimated, irrespective of the diffusion time.

cal model of two compartments with exchange – in terms of accuracy and precision in simulations, as well as feasibility and sensibility of estimated microstructure parameters in experimental rat data *in vivo*.

3.2.1. Simulations

We first present the performance of NEXI in simulations, using Eqs. (3)–(6) to generate ground truth signal, and estimating the four model parameters using conventional NLLS (see Supplementary Material for DL-based fitting, Figs. S6 and S7).

Scenario a: Fitting the four model parameters for each diffusion time separately. The precision was good on D_e and acceptable on f . However, in a finite SNR case, $D_{i,\parallel}$ and t_{ex} could not be estimated, irrespective of the diffusion time (Fig. 7).

Scenario b: Fitting the four model parameters using all diffusion times jointly. In all cases, this approach significantly improved the precision on f and t_{ex} compared to Scenario a. Some sensitivity to $D_{i,\parallel}$ was also restored (Fig. 8).

Varying b_{max} showed that b -values larger than $2.5 \text{ ms}/\mu\text{m}^2$ are needed for accuracy (Fig. S8). Both accuracy and precision were fur-

ther improved for $b_{\text{max}} = 10$ vs $6 \text{ ms}/\mu\text{m}^2$ but the benefits were less substantial. When inspecting the impact of b_{max} given a constant number of shells, the performance of $b_{\text{max}} = 10$ was still superior to that of $b_{\text{max}} = 2.5$ in terms of accuracy and precision, confirming it is the b -value range that is critical for sensitivity to model parameters.

3.2.2. Experimental: *in vivo* rat GM

The time-dependence analysis of diffusion, kurtosis and SM metrics highlighted the sharp difference in behavior between highly myelinated white matter fibers such as the internal capsule and the corpus callosum, and GM. Results from the previous sections suggest non-negligible inter-compartment water exchange in GM, which should be accounted for by biophysical models of this type of tissue when working at relatively long diffusion times ($t > 20$ ms).

Simulation results for NEXI performance in turn suggested the use of multi-shell multi- t data for the reliable estimation of model parameters, and of the exchange time in particular, for a broad range of ground truth values.




Fig. 8. Simulation results fitting multi-shell multi- t_d data jointly using NLLS, for random GT (A) or fixed to $[t_{ex}^{th}, D_{i||}^{th}, D_e^{th}, f^{th}] = [20, 2.5, 0.75, 0.34]$ (B). A: Displayed are the medians & IQR in each bin. Black lines: ideal estimation $\pm 10\%$ error. Without noise, NLLS fits all parameters with high accuracy and precision. At SNR = 100, uncertainty increases primarily for $D_{i||}$ and t_{ex} and sensitivity to high t_{ex} values is lost but the performance is much improved compared to single t_d fits (Fig. 7). B: At SNR = 100, good accuracy is achieved for all NEXI parameters. For $D_{i||}$ the precision is poor. Black solid line: ground truth.




Fig. 9. A–D: Four coronal slices of NEXI parametric maps calculated using NLLS from a multi-shell multi- t dataset. The maps enable a good differentiation between GM & WM as well as between different cortical layers (white arrows) or hippocampal subfields (black arrow). E: Median & IQR of model parameters in the cortex and hippocampus ROIs across the four datasets. The exchange time estimate is also compared with $t_{ex}^{K(t)}$, Eq. (7). Experimental trends agree with the simulations. Regarding $t_{ex}^{K(t)}$, the estimation agrees with t_{ex} very well for Dataset #4, which had the highest SNR (larger voxels), and is otherwise shorter.

Given a dataset comprised of b -values up to $10 \text{ ms}/\mu\text{m}^2$ and three to four diffusion times, all NEXI model parameters could be estimated both at the ROI and at the single voxel level in the rat GM.

Testing 100 random initializations on ROI-averaged signal showed that the overwhelming mode of the distribution of outcomes corresponded to a solution where $D_{i||} > 2$ and $D_e < 1$ (Fig. S9).

Parametric maps of NEXI estimates were consistent with expected neuroanatomy of the rat brain (Fig. 9A–D). Simulations predicted the variability was largest for t_{ex} and $D_{i||}$. ROI-based analysis in the cortex and hippocampus confirmed these trends and also revealed good

between-subject consistency (Fig. 9E). On average, the intra-neurite diffusivity was $2.5 \mu\text{m}^2/\text{ms}$, the extra-neurite diffusivity was $0.75 \mu\text{m}^2/\text{ms}$ and the neurite fraction was around 0.3.

For Datasets 1–3, the estimated exchange time t_{ex} was higher with NEXI than with $t_{ex}^{K(t)}$: 30–60 ms vs 10–40 ms, respectively, in cortex and 25–65 ms vs 10–25 ms, respectively, in hippocampus. Remarkably, for Dataset 4 (x2.5 voxel volume compared to Datasets 1–3), the two approaches displayed better agreement and estimates for t_{ex} ranged 10–30 ms. This comparison is based on fits using the entire diffusion time range available. As shown for MK(t), excluding $t < 20 \text{ ms}$ datapoints




Fig. 10. Features of NEXI neurite density map features as compared to cellular components obtained from histological stainings: neurofilaments (orange), astrocytes (blue), neuron nuclei (green) and microglia (red). The WM is outlined in fine dotted lines for legibility; cortex lies above, hippocampus below. Higher NEXI neurite density in central cortical layers agrees with higher density of neurofilament staining (dashed lines). Higher NEXI neurite density in the central part of the hippocampus (dorsal dentate gyrus) agrees especially with higher density of astrocytes but also neurofilaments (long-dashed contour). Neuron soma and microglia do not seem to contribute to NEXI neurite density contrast.

yielded longer exchange times. This can also be expected for NEXI but was not implemented here for concerns of fit stability and precision.

The exchange time can be related to the cell membrane permeability of a cylinder via $P = \frac{d}{4(t_{ex} - \frac{d^2}{32 D_{i,\parallel}})}$, where d is the diameter of the cylinder (neurite) and P is the *diffusional water membrane permeability* (affected by the properties of the lipids in the membrane and by water-channel proteins embedded in the membrane and different from the osmotic permeability, generally larger and measured in the presence of an osmotic pressure gradient over the membrane) (Meier et al., 2003; Nilsson et al., 2013). Given the typical diameter and diffusivity values, the relationship between exchange time and permeability is very well approximated as $P \cong \frac{d}{4t_{ex}}$. Assuming $d \sim [0.5\text{--}2] \mu\text{m}$, a characteristic exchange time $t_{ex} = [15\text{--}60] \text{ms}$, as estimated here using NEXI, yields $P \sim [2.1\text{--}33] \times 10^{-3} \mu\text{m}/\text{ms}$.

The intra-neurite fraction map displayed substantially larger values in white matter than gray matter though the model does not in principle support white matter (the extra-neurite space cannot be assumed to be isotropic). This can be explained by the fact that myelin is MR-invisible in our diffusion MRI measurements (due to the long TE) and the physical space occupied by myelin is therefore not considered. Assuming neurite physical occupancy fractions are similar in GM and WM, the *relative neurite fraction* we estimate is higher in WM because the myelin space reduces the extracellular space. Different T_2 relaxation times in compartments between GM and WM could also account for the difference. The other NEXI parameters were also higher in WM, which is in good agreement with more aligned structures (enabling faster diffusivity) and longer exchange times in highly myelinated axons.

All NEXI parameters also showed contrast within GM structures, such as across cortical layers and hippocampal subfields. Notably, the neurite fraction was higher in central cortical layers, consistent with neurofilament staining in *ex vivo* rat cortex slices (Fig. 10). A higher NEXI neurite fraction in the central section of the hippocampus (dorsal dentate

gyrus) agreed especially with higher astrocyte density in that region, which suggests astrocytic processes also contribute to this parameter via their similar geometry to neurites, i.e., long, thin structures.

4. Discussion

In this work, we propose NEXI as an extension of the SM of diffusion suitable for GM. NEXI accounts for inter-compartment exchange between neurites and the extracellular space, building on the anisotropic Kärger model of two exchanging compartments. Using multi-shell multi- t datasets acquired in the rat brain *in vivo*, we investigate the suitability of NEXI to describe diffusion in the GM, compared to other approaches such as SM, structural disorder, or the addition of a soma compartment. We identify exchange as the mechanism that best explains diffusion-time-dependence of signal in both low- b and high- b regime, and thereby propose NEXI as the minimal model for GM microstructure mapping. We finally propose multi- b multi- t acquisitions schemes as best suited to estimate NEXI model parameters $[f, D_{i,\parallel}, D_e, t_{ex}]$ reliably.

SM applicability. The presence of exchange yields a spurious time-dependence of SM parameters. Unsurprisingly, the time-dependence of geometric parameters for the SM was most marked in GM and cingulum. The apparent intra-neurite fraction decreased with increasing diffusion time. Qualitatively, this can be interpreted as water molecules that leave the intra-neurite space developing a diffusion signature closer to hindered diffusion – as the extracellular space or large soma – rather than restricted and unidirectional along the neurite. The neurite alignment also increased with longer times. These results suggest that in the SM the diffusion time acts as a filter that attributes to the intra-neurite space only cellular processes (axons, dendrites and glial processes) that can be considered impermeable over that time scale. Arguably, only the more myelinated and aligned neurites are retained at longer times.

The time-dependence of SM compartment apparent diffusivities varied with b -value regime, with more pronounced trends for

$b_{\max} \geq 6 \text{ ms}/\mu\text{m}^2$ and for GM than for WM ROIs. A trend of apparent diffusivity decreasing with time can be a signature of either non-Gaussian diffusion, exchange with a slower compartment or both. A trend of apparent diffusivity increasing with time is a sign of model failure or insufficiency. Essentially, in GM the SM parameters become ill-defined and their interpretation in terms of microstructure becomes challenging. Since exchange is not accounted for, compartment diffusivities D_i and D_e should be interpreted as “apparent” compartment diffusivities D'_i and D'_e . At $t \rightarrow 0$, the two compartments have not mixed at all, a standard model fit would yield apparent $D'_i = D_i$ and $D'_e = D_e$, while $\bar{D} = fD_i + (1-f)D_e$. In the long-time limit $t \rightarrow \infty$, the mean diffusivity \bar{D} is unchanged but the compartments are well-mixed. It is unlikely that the mixture effect can be absorbed entirely by a decrease in f , and apparent compartment diffusivities will also be affected: they will each tend to converge towards \bar{D} , hence, assuming without loss of generality that $D_i > D_e$, $D'_i(t \rightarrow 0) > D'_i(t \rightarrow \infty)$ while $D'_e(t \rightarrow 0) < D'_e(t \rightarrow \infty)$.

Our analysis confirms that the SM is applicable in thick WM bundles. The quantitative estimate of compartment diffusivities may however depend on the b -value range (the anisotropy of the extra-neurite compartment in particular may be exacerbated) and the intra-neurite fraction may decrease with longer times likely by dropping unmyelinated axons in WM. The latter are more numerous in the rodent than in the human brain (Wang et al., 2008) so this effect may not impact human brain estimates as much. However, the SM assumptions are likely not met in GM and this could also impact WM bundles that have substantial partial volume with cortex, such as the cingulum. In particular, while in the cumulant regime ($b \lesssim 2.5$) $b \leq 2.5$) only a progressive filtering of neurites with increasing time was observed, for higher b -values apparent compartment diffusivities were also affected, particularly in the intra-neurite space.

Exchange vs structural disorder in cortex. Our data show negligible time-dependence of mean diffusivity, over the range of diffusion times 20–45 ms, in the rat GM. This is consistent with previous reports for the *in vivo* rat brain by Pyatigorskaya et al. (2014) and *ex vivo* mouse brain by Aggarwal et al. (2020). We note however that pronounced OGSE frequency dependence of diffusivities has been reported at shorter time scales in rat and mouse cortex (Aggarwal et al., 2020; Does et al., 2003; Pyatigorskaya et al., 2014), and, in the case of the work by Does et al. (2003), has been attributed to the neurites based on the exponent $\vartheta = 1/2$ by Novikov et al. (2014).

Weak diffusivity time-dependence has also been highlighted in human cortical gray matter for diffusion times 21 – 100 ms (Lee et al., 2020b). The human brain presents the additional challenge of thin cortical ribbons and thereby relatively strong partial volume with white matter and CSF in “cortical voxels”. The rat brain is, from this perspective, a well-suited model system for exploring cortical properties. We note however the weak time-dependence of diffusion is not necessarily transposable to other organs, *ex vivo* conditions and different diffusion time ranges. For example, (Jespersen et al., 2018) have shown significant time-dependence of the diffusion coefficient in fixed pig spinal cord, for diffusion times 6–350 ms. The negligible time-dependence of D in our context suggests/validates that, within the cumulant expansion regime, the tissue can be considered as a collection of Gaussian compartments, one of the main assumptions behind multi-compartment models of diffusion, and that structural disorder is therefore negligible at our diffusion time scales. A larger dynamic range of diffusion times, covering at least a logarithmic decade, may be necessary to detect diffusivity time-dependence.

On the other hand, kurtosis displayed marked time-dependence, also in agreement with findings of all afore-mentioned studies (Aggarwal et al., 2020; Jespersen et al., 2018; Lee et al., 2020b; Pyatigorskaya et al., 2014). Together with the absence of marked time-dependence of D , the decrease in $K(t)$ with t can be attributed to inter-compartment exchange, Eq. (7) (Fieremans et al., 2010; Jensen et al., 2005; Kärger, 1985) rather than structural disorder, with kurtosis decaying to zero at very long times when compartments are fully mixed and

appear as a single Gaussian compartment. The confirmation of dominant exchange by the analysis of the power-law exponent of the $K(t)$ decay was challenged by the fact that the long-time limit has not been reached in order to yield either a decay as $1/t$ which would support exchange, or a $1/\sqrt{t}$ functional form, which would favor structural disorder as in Lee et al. (2020b). Nevertheless, simulations of exchange-driven $K(t)$ in our experimental time range yielded an apparent power-law exponent of 0.3, which agreed with the experimental exponents estimated from $K(t)$ power law fits on each animal and GM ROI. The exploration of a broader range of diffusion times in future work may enable a more definite assessment of the most relevant power law of $K(t)$ decay and of the relative contribution of the competing effects of incomplete coarse-graining over the structural disorder, and inter-compartmental exchange.

It should be underlined that the functional form for structural disorder as $1/\sqrt{t}$ corresponds to one-dimensional short-range disorder which is potentially suited for intra-neurite diffusion (Novikov et al., 2014). In principle, structural disorder could also arise from extracellular water, and would in this case be expected to follow the functional form for 2d or 3d disorder, as $(\ln t)/t$ or $1/t$. This functional form should however be followed by both $D(t)$ and $K(t)$. Overall, the trend in $D(t)$ was flat and certainly did not support a decay as $(\ln t)/t$ which is more pronounced than $1/\sqrt{t}$. The 2d or 3d disorder was also not supported by $D(t)$ in human cortex (Lee et al., 2020b).

The estimation of inter-compartment exchange based on NEXI $K(t)$ yielded relatively long exchange times ($t_{\text{ex}}^{K(t)} = 80\text{--}130$ ms, exceeding our diffusion time range) in highly myelinated white matter bundles such as the corpus callosum and internal capsule, intermediate exchange times ($t_{\text{ex}}^{K(t)} \sim 40$ ms) in thinner bundles such as the cingulum that may experience partial volume effects with neighboring gray matter, and relatively short exchange times in the cortex and hippocampus ($t_{\text{ex}} \sim 15\text{--}20$ ms).

Estimates of exchange time in WM bundles were unsurprisingly imprecise due to the mismatch between probed timescales (10 – 45 ms) and the expected exchange time $t_{\text{ex}}^{K(t)} > 80$ ms. Nevertheless, this result validates *a posteriori* the assumption of non-exchanging compartments for white matter models at diffusion times typical for PGSE acquisitions: $t < 80$ ms. Our findings are also consistent with previous studies in the human WM reporting exchange times above 500 ms (Lampinen et al., 2017; Nedjati-Gilani et al., 2017), and 350–400 ms in mouse corpus callosum (Hill et al., 2021).

In the case of gray matter, the $t_{\text{ex}}^{K(t)}$ estimates of 15–20 ms were consistent with previous studies in human gray matter (Veraart et al., 2018a) and perfused pup rat spinal cord (Williamson et al., 2019). Other studies using relaxation-based methods suggested however longer exchange times of 100–150 ms in astrocyte and neuron cultures (Yang et al., 2018), in rat subcortical structures – presumably the striatum (Quirk et al., 2003) and in rat perfused cortical cultures (Bai et al., 2018). Filter-exchange imaging (FEXI), another diffusion-based method to estimate the exchange time between a slow and a fast water pool reported an exchange time on the order of 1 s in WM and 2.5 s in GM using a filtering block of $b_f = 0.9 \text{ ms}/\mu\text{m}^2$ (Lampinen et al., 2017; Nilsson et al., 2013). While the exchange times between different WM tracts agreed with expected myelination levels, e.g., up to 3 s in corpus callosum and 500 ms in anterior corona radiata, it is somewhat counterintuitive that the exchange time would be longest in GM – the authors suggested the latter was likely overestimated. Recent work using FEXI with a similar filter also yielded an exchange time of around 1 s in WM and 1.4 s in GM (Bai et al., 2020), while arguing that there is no direct evidence that what FEXI measures is the exchange between intra- and extra-cellular compartments. Conversely, shorter exchange times of 2 – 5 ms have recently been reported in fixed mouse gray matter using SMEX (Standard Model with EXchange), a similar approach to NEXI, as will be discussed in the more detail further on Olesen et al. (2022).

Related to FEXI, while bi-exponential functions typically fit diffusion decay in brain tissue well, the association of the slow and fast water pools to specific tissue compartments has never been straightforward (Kiselev and Il'yasov, 2007; Novikov et al., 2018a), in particular since

a distribution of non-parallel sticks – a single compartment, technically – also yields a characteristic decay that is well approximated by a bi-exponential function (Assaf and Cohen, 1998; Callaghan et al., 1979; Novikov et al., 2018a; Sehy et al., 2002). For this same reason, alternative approaches to the Kärger model, extracting the exchange time from the decay of the intra-cellular fraction – as estimated from a bi-exponential model – with increasing diffusion time (Moutal et al., 2018) were explored in yeast suspensions but are not suitable for brain tissue. Indeed, predominant stick-like geometries in both white and gray matter invalidate the approximation of a sum of two Gaussian compartments in any direction or for the powder-average signal.

Our analysis of $K(t)$ so far cannot provide direct information on the mechanisms of exchange, such as intra/extracellular, neurite/soma or neurite/neurite exchange. However, numerical simulations suggest that neurite/soma and neurite/neurite exchange within the same neuron occur at longer time scales (on the order of 100 ms or more) than those estimated here (~20 ms) (Ianus et al., 2020) and support the intra/extracellular exchange as the dominant mechanism in these experiments.

Exchange vs soma. Going a step further, we investigated the performance of a two-compartment model with exchange (NEXI) and of a three-compartment model accounting for soma (SANDI) to capture diffusion signal decay at high b -values, and for multiple diffusion times. SANDI extends the SM by adding an extra compartment for modeling explicitly diffusion restricted in soma and relies on the assumption of negligible exchange between the three tissue compartments: intra-neurite, intra-soma and extra-cellular. Our results do not challenge this assumption in the rat GM *in vivo* for relatively short diffusion times (≤ 20 ms), while challenging it for longer diffusion times (> 20 ms), where the SANDI model parameters show some time-dependence. This diffusion time cutoff is in line with $t_{\text{ex}}^{K(t)}$ and suggests that unaccounted exchange mechanisms between the three major tissue compartments in GM (cellular processes, soma and extra-cellular space) may bias SANDI parameter estimation at diffusion times longer than 20 ms. On the other hand, our results also suggest that SANDI model parameter estimation provides f , $D_{i,\parallel}$ and D_e estimates in good agreement with the equivalent counterpart from the SM and NEXI. The importance of modeling exchange in addition to soma was mostly evident in the ability of NEXI vs SANDI to predict signal decay curves for longer diffusion times based on model parameters estimated at short diffusion times. This result is consistent with findings in the rat cortex *ex vivo* (Olesen et al., 2022). However, based on the NEXI parameter estimation performance alone – discussed below – a larger q - t coverage and higher SNR would likely be needed *in vivo* to account for both soma and exchange in a model.

NEXI parameter estimation. To provide recommendations of minimum data and fitting procedures for NEXI, we first established its performance in simulations. Given a comprehensive protocol with 7 shells up to $b_{\text{max}} = 10 \text{ ms}/\mu\text{m}^2$ and high final SNR of 100 – boosted by the MP-PCA denoising procedure and the powder-averaging over directions – data at a single diffusion time were insufficient to estimate t_{ex} and $D_{i,\parallel}$. Noise was clearly the culprit as the noiseless simulations otherwise demonstrated good performance for all four model parameters. Fitting the NEXI model to joint data over four diffusion times dramatically improved the accuracy and precision for all four model parameters though $D_{i,\parallel}$ remained the most challenging parameter to estimate, consistent with other model frameworks (Jelescu et al., 2016; Novikov et al., 2018b; Palombo et al., 2020). The benefit of a broader b -value range was critical between 2.5 and 6 $\text{ms}/\mu\text{m}^2$, but only marginal beyond, which suggests a range $0 < b < 6 \text{ ms}/\mu\text{m}^2$ could be sufficient to estimate NEXI parameters.

Simulations also suggested the exchange time estimate \hat{t}_{ex} plateaus beyond ground truths $t_{\text{ex}} \geq 80 \text{ ms}$ approximately. This is likely related to the diffusion time range simulated 12–40 ms, which is too short to probe slow processes with longer characteristic exchange times. For tissues where longer exchange times are expected, the diffusion time range should be adjusted accordingly.

Importantly, the performance of NEXI on experimental data using a multi-shell multi- t protocol was consistent with simulations. On average, the intra-neurite diffusivity was 2.2–2.5 $\mu\text{m}^2/\text{ms}$, in agreement with its estimate from the SM and from SANDI at the shortest diffusion times, as well as with previous reports of intra-neurite/axonal diffusivity (Dhital et al., 2019; Kunz et al., 2018; Olesen et al., 2021). The extra-neurite diffusivity was 0.75 $\mu\text{m}^2/\text{ms}$ and remarkably also agreed with the SANDI estimate at the shortest times. We underline that the intra-neurite diffusivity corresponds to the parallel diffusivity, with $D_{i,\perp} = 0$ in the perpendicular direction (the stick picture). A three-fold ratio between $D_{i,\parallel}$ and D_e is consistent with previous literature that reported similar Apparent Diffusion Coefficient (ADC) between intra- and extra-cellular water in rat GM (Duong et al., 1998). Considering the picture of isotropically-oriented neurites, the ADC of intra-cellular water in any given direction would be estimated at $D_{i,\parallel}/3$ and thereby similar to D_e .

Compartment fractions: The neurite fraction was about 0.3, which is lower than estimates from *ex vivo* histology (~0.65) but nonetheless higher than the SANDI estimate for neurite fraction at the shortest diffusion time ($f \sim 0.25$). This suggests that even at short diffusion times, neurite fractions from models that do not account for exchange may be underestimated, possibly by “missing” fast-exchanging components. In parallel, by comparing NEXI and SANDI compartment fractions, it appears the soma is associated to extra-neurite space in NEXI. While accounting for exchange had the advantage of providing a time-independent estimate of the neurite fraction, thereby correcting for its decrease with longer times in models of non-exchanging compartments, the absolute value of the neurite fraction estimate remains lower than the 60–70% expected from histology (Bondareff and Pysh, 1968; Motta et al., 2019; Shapson-Coe et al., 2021; Spocter et al., 2012). Combined relaxation-diffusion measurements may help improve the quantification of the neurite fraction by correcting for relaxation time-weighting (Barakovic et al., 2021; Hutter et al., 2018; Tax et al., 2021; Veraart et al., 2018b). Exchange processes on a shorter scale than those explored here also cannot be excluded, as very short exchange times have been recently reported in rat brain *ex vivo* (Olesen et al., 2022).

Exchange time: Determining the accurate value of t_{ex} correctly is of utmost importance to mapping GM microstructure, as this value sets the relevant tissue features to be modeled, given that typically accessible diffusion times on clinical MRI systems are $t \gtrsim 10t \geq 10 \text{ ms}$. If, for instance, $t_{\text{ex}} \sim 1 \text{ ms}$ (well below 10 ms), the exchange can be practically classified as fast, i.e., there are no distinct intra- and extra-stick exchanging compartments; neurites and extra-neurite space form a homogenized effective medium. This is the picture suggested by recent results of Olesen et al. If, conversely, $t_{\text{ex}} \gtrsim 100 \text{ ms}$, as Lampinen et al. (2017), Yang et al. (2018) suggest, then exchange is slow and can be neglected, and a Standard Model-like approach is applicable. Finally, if exchange is intermediate, between ~10 ms and ~50–100 ms, as our present results and the 2-dimensional NMR measurements by Williamson et al. (2019) suggest, then exchange is relevant, must be explicitly modeled, and can be mapped. This possibility, albeit requiring more complex modeling, would open a tantalizing prospect of mapping *tissue function in vivo*, since permeability has been shown to be modulated by metabolism (Bai et al., 2019). Below we consider the confounding effects for t_{ex} .

The discrepancy in exchange time estimates between Olesen et al. and the current work, which are otherwise similar in approach, can be due to multiple sources. First, *in vivo* vs fixed tissue is an important driver of exchange time differences (Li et al., 1998; Shepherd et al., 2009; Thelwall et al., 2006). Recent additional work using NEXI on *ex vivo* rat brain at 20 °C also yielded shorter exchange times of 4–6 ms, than the ones reported here *in vivo* (Jelescu and Uhl, 2022). Another possible source of discrepancy is the range of diffusion times explored in either study: 7.5–16 ms in Olesen et al. vs 11–45 ms here. The diffusion times act as a filter for the exchange times that can be reliably estimated (see Fig. 8, where, in simulations, the estimation of t_{ex} plateaus beyond

Table 2

NEXI and NEXI_dot parameter estimates from the average signal in the rat cortex (as in Fig. 6). Mean \pm std as provided by the fits' confidence intervals.

	f_n	t_{ex}	D_i	D_e	f_{dot} / σ
NEXI	0.29 \pm 0.01	43 \pm 8	2.55 \pm 0.24	0.74 \pm 0.02	–
NEXI_dot	0.22 \pm 0.03	21 \pm 9	3.6 \pm 0.6	0.70 \pm 0.02	0.022 \pm 0.008
NEXI_rm	0.29 \pm 0.01	22 \pm 6	3.2 \pm 0.3	0.69 \pm 0.02	0.025 \pm 0.002

75 ms for simulated diffusion times of 12–40 ms). Slow exchange cannot be accurately captured with short diffusion times.

Finally, since the NEXI fit deviates from the data at high b -values (Fig. 6), we also tested an adapted model of NEXI including a “dot” compartment (i.e., water apparently immobile, as reported mainly *ex vivo* and in the cerebellum) and a model fitting for a Rician floor. In the first case, the NEXI_dot model is thus characterized by five parameters to be estimated, similar to the SMEX model by Olesen et al.: f_n , t_{ex} , D_i , D_e and f_{dot} . In the second case, the NEXI_rm (for Rician mean) model is also characterized by five parameters to be estimated: f_n , t_{ex} , D_i , D_e and the Rician noise parameter σ :

$$NEXI_{rm}(f, t_{ex}, D_i, D_e, \sigma) = \sqrt{\frac{\pi}{2}} \cdot \sigma \cdot L_{1/2} \left(-\frac{1}{2} \left(\frac{NEXI(f, t_{ex}, D_i, D_e)}{\sigma} \right)^2 \right)$$

Where $L_{\frac{1}{2}}(x) = {}_1F_1(-\frac{1}{2}, 1, x) = e^{x/2} [(1-x)I_0(-\frac{x}{2}) - xI_1(-\frac{x}{2})]$ is the generalization of Laguerre polynomial $L_n(x)$, which for non-integer n is given in terms of the confluent hypergeometric function. The above equation elevates NEXI(f, t_{ex}, D_i, D_e) based on the Rician expectation value.

NEXI, NEXI_dot and NEXI_rm were fit to the data in Fig. 6 within the ranges: f_n : [0, 1], t_{ex} : [0, 200], D_i : [0, 4], D_e : [0, 4], f_{dot} : [0, 0.2] (Table 2). NEXI_dot and NEXI_rm indeed yielded a shorter t_{ex} than NEXI, but not as short as reported in Jelescu and Uhl (2022), Olesen et al. (2022). Furthermore, the uncertainty in NEXI_dot estimates was largest among the three models, while the intra-neurite diffusivity largely exceeded 3 $\mu\text{m}^2/\text{ms}$ systematically. The NEXI_dot and NEXI_rm estimates were similar and the resulting fitted curves approximated the data well (Supplementary Fig. S11). This suggests both an actual dot compartment and an impact of non-zero Rician mean could explain the NEXI departure from experimental points at highest b -values.

The exchange time estimates depended on the underlying SNR of the data, on the diffusion time range and on the estimation approach. For a diffusion time range 10–45 ms, NEXI estimated an exchange time of 15 – 60 ms while the matching estimate from $K(t)$ was 10–40 ms. The latter became 18–60 ms when the diffusion time range was reduced to 20–45 ms (albeit with increased uncertainty). These exchange times, combined with intra-neurite realistic diameter values yielded a range of cell membrane permeability values on the order of $P \cong [2.1 - 33]$ $\mu\text{m}/\text{s}$. This range of permeability values is in agreement with previous reports of physiologically relevant membrane permeability values in healthy cells: $P \cong [6 - 30]$ $\mu\text{m}/\text{s}$ (Baylis, 1988; Harkins et al., 2009; Latour et al., 1994; Stanisiz et al., 1997; Vestergaard-Poulsen et al., 2007). Cell-specific membrane water permeability values have also been reported as [18–76] $\mu\text{m}/\text{s}$ in murine neurons, [41–112] $\mu\text{m}/\text{s}$ in murine astrocytes and [23–81] $\mu\text{m}/\text{s}$ in human red blood cells (Boss et al., 2013). The latter have been extensively studied in terms of water permeability, with variable reported ranges (Benga et al., 2009, 2000). Shorter exchange times of 3–5 ms reported *ex vivo* by Jelescu and Uhl (2022), Olesen et al. (2022) would translate into a permeability $P \cong [25 - 167]$ $\mu\text{m}/\text{s}$, consistent with chemical fixation increasing permeability over three-fold (Shepherd et al., 2009).

Parametric maps agreed with known rat brain structure, with clear delimitation between gray and white matter. Contrast between adjacent cortical layers and between hippocampal sub-fields was also apparent. Comparison with histological staining revealed that higher NEXI neu-

rite fraction in middle cortical layers corresponded to higher neurofilament density in that area, but that in hippocampus, abundant astrocytic processes could contribute to the higher NEXI neurite fraction. Indeed, water is ubiquitous and microstructural features with similar geometry are typically pooled together – here neurites and astrocytic processes are both thin elongated cylinders. This is expected to be the case for all water diffusion models proposed, though the balance of contributions between neurons and glial cells has never been firmly established.

The importance of high SNR (> 20 – 30 in an individual $b = 0$ image) for reliable parameter estimation was manifest throughout our data which had a strong SNR spatial gradient from cortex to deep brain due to the use of a surface transceiver. An experimental setup with a volume coil for transmission and a surface coil for reception would yield uniform SNR and enable estimates of NEXI parameters over the whole brain. DL approaches are increasingly replacing NLLS in biophysical model estimation (Diao and Jelescu, 2022; Hill et al., 2021; Nedjati-Gilani et al., 2017; Palombo et al., 2020). We also showed here an improved management of noise by DL vs NLLS, although bias towards the mean of the prior was also more pronounced and DL outputs should always be examined carefully (Coelho et al., 2021; Gyori et al., 2021b; Martins et al., 2021). Alternatively, a log-likelihood objective function could be considered for the NLLS optimization to account for Rician noise in a voxel-wise fashion. As there was already good agreement between sum-of-squares NLLS and DL on our data, we did not implement the log-likelihood minimization, but it may be worth considering for lower SNR data.

Limitations. We note that a multi-shell multi- t acquisition protocol may be difficult to implement when scan time is of the essence. Future work will focus on optimizing the protocol to the best compromise between minimal scan time and maximal accuracy and precision of model parameter estimates. The currently large uncertainty on t_{ex} and $D_{i||}$ estimates will also benefit from such an optimization, which may include schemes that combine multiple diffusion tensor encodings (Chakwizira et al., 2021).

After inspecting the mode of NEXI outcomes as a function of random algorithm initialization, and in line with recent evidence that $D_{i||} \cong 2 - 2.5$ $\mu\text{m}^2/\text{ms}$ (Dhital et al., 2019; Howard et al., 2020; Kunz et al., 2018; Olesen et al., 2021), we chose an algorithm initialization where $D_{i||} > D_e$ in NLLS, and trained the DL network on disjoint intervals $D_{i||} \geq 1.5$ and $D_e \leq 1.5$. These constraints would likely need to be reconsidered and relaxed, particularly when characterizing pathological conditions or diseases, such as Alzheimer's or Huntington's diseases, where tangles and/or proteins accumulate in the intracellular space, hence increasing the cytoplasm tortuosity and reducing $D_{i||}$, potentially to the level of being slower than D_e .

One substantial limitation of the NEXI model is that it does not account for soma as a third compartment, which are then artificially absorbed into the extra-neurite space as the relative fractions suggest. Recently, the evidence of curved boundaries in the cortex, attributed to soma, was presented by observing the localization regime of diffusion on a human Connectom scanner (Lee et al., 2021). An important line of future work is the extension of NEXI to three compartments, or, equivalently, the incorporation of exchange processes in the SANDI model, as has been implemented in fixed mouse gray matter (Olesen et al., 2022). For now, SANDI accounts for three non-exchanging compartments – soma, neurites, extracellular space – but thus requires data acquisition at relatively short diffusion times (≤ 20 ms) concomitantly with high b -values ($b_{\text{max}} \geq 6$ $\text{ms}/\mu\text{m}^2$) (Palombo et al., 2020) which can only be achieved on preclinical and Connectom scanners, but not on typical clinical scanners. Going forward, accounting for exchange in SANDI will make it translatable to clinical settings, by enabling the use of longer diffusion times. The feasibility of estimating a large number of model parameters on data limited in terms of SNR and (q, t) coverage by *in vivo* and/or clinical hardware settings is yet to be determined.

Furthermore, NEXI considers Gaussian compartments, an assumption which seems to break for the neurite compartment, as revealed at higher b -values, likely due to finite length of dendritic processes,

branching, etc. This poses a significant conundrum, as low b -values where non-Gaussian contributions in each compartment can be neglected are detrimental for estimation accuracy and precision, while higher b -values reveal non-Gaussian effects which bias the outcomes as well. Ultimately, computational models based on realistic simulations of neural cells (Callaghan et al., 2020; Ginsburger et al., 2019; Lee et al., 2020a; Palombo et al., 2019) and cortical substrate may be the best approach for characterizing gray matter microstructure *in vivo*. Analytical approaches able to incorporate intra-compartmental non-Gaussian time-dependent effects, such as Lee et al. (2018), are in need to faithfully quantify the structural disorder contributions.

Finally, built on the anisotropic KM, NEXI assumes exchange happening within each ensemble of “neurite + its immediate extracellular space” separately. This may differ from GM microanatomy, where neurites at different angles can be piercing a volume of the size of the diffusion length. Sequential exchange processes can bring a molecule from, e.g., one stick to the extra-stick space to a differently-oriented stick, and so on; such a model geometry has not been considered. We note, however, that in the limit of $bD_e \gg 1$, the distinction between this more general geometry and NEXI should vanish, since, once a spin enters the extra-neurite space, its contribution to the overall signal gets exponentially suppressed, as does the memory about the above sequential exchange processes with different orientations. Hence, we expect that t_{ex} estimated from the full protocol including strong diffusion weightings will be more accurate than that from the time-dependent kurtosis approximation of Eq. (6).

Value. Taken together, our results suggest that inter-compartment exchange is not negligible in gray matter at typical PGSE or clinical diffusion times ($t > 20$ ms) and should therefore be accounted for in biophysical models of gray matter and potentially even in thinner white matter tracts such as the cingulum (in rodents), and by extension to demyelinating WM as a result of disease. Our findings also highlight an additional challenge for approaches that use b -tensor encoding techniques to disentangle various tissue geometries or solve model degeneracy (Afzali et al., 2021; Coelho et al., 2019; Gyori et al., 2021a; Reisert et al., 2019). Since free gradient waveforms introduce by design a whole spectrum of relatively long diffusion times (> 20 ms), the ill-definition of the diffusion time may become problematic in a regime where exchange cannot be neglected.

NEXI constitutes an important first step in accounting for inter-compartment exchange in GM and developing a more realistic model of diffusion in gray matter. The estimate of the exchange time alone can be used as a proxy for membrane permeability, which is known to increase with injury or neurodegeneration (Nilsson et al., 2013; Pacheco et al., 2015), and could yield an original and valuable new biomarker of tissue integrity, metabolism and function (Bai et al., 2018). In myelinated structures, the exchange time could also become a strong proxy for the myelin thickness, which is at the heart of several “*in vivo* histology” efforts (Brusini et al., 2019; Hill et al., 2021; Lazari and Lipp, 2021; Mancini et al., 2020).

5. Conclusions

One fundamental challenge in brain microstructure is to establish the biophysical origin of effects beyond the “Standard Model” (SM) picture of non-exchanging Gaussian compartments. Here we showed that in the rat GM *in vivo*, the exchange dominates over structural disorder, and offer the picture of diffusion time effectively filtering out the contribution of unmyelinated neurites with stronger dispersion. At long times, this picture suggests that only the myelinated (non-exchanging) neurites contribute to the intra-neurite SM compartment, and the rest is asymptotically attributed to extra-neurite space. Exchange also explains signal decay curves across different diffusion times better than the addition of a soma compartment. If a choice is warranted, a two-compartment model with exchange – NEXI – is better suited than a three-compartment model with soma for characterizing cortical microstructure at diffusion times

$t > 20$ ms, while also yielding a valuable estimate of exchange time, which can be used as a proxy for membrane permeability. Going forward, both soma and exchange should ideally be accounted for, if the data support the estimation of a larger number of parameters.

Data and code availability statement

The data used for this manuscript were newly acquired for the purpose of this project and can be shared upon request by emailing the Corresponding Author (ileana.jelescu@chuv.ch). Data sharing will require a formal data sharing agreement, including co-authorship of the Corresponding Author and possibly approval from the requesting researcher’s local ethics committee.

The processing tools are freely or widely available (referenced in the Methods section where applicable) and all details have been provided to enable replication of the analysis.

Declaration of Competing Interest

None.

Credit authorship contribution statement

Ileana O. Jelescu: Conceptualization, Methodology, Software, Investigation, Resources, Writing – review & editing, Visualization, Visualization, Formal analysis, Writing – original draft. **Alexandre de Skowronski:** Software, Formal analysis, Writing – original draft. **Françoise Geffroy:** Investigation, Resources, Writing – original draft. **Marco Palombo:** Conceptualization, Formal analysis, Writing – original draft, Writing – review & editing. **Dmitry S. Novikov:** Conceptualization, Methodology, Writing – original draft, Writing – review & editing.

Acknowledgments

We thank reviewers, including Sune Jespersen and Jonas Olesen, for constructive comments on the manuscript. We thank Analina da Silva, Mario Lepore and Stefan Mitrea for assistance with animal monitoring. This work was made possible thanks to the CIBM Center for Biomedical Imaging, founded and supported by Lausanne University Hospital (CHUV), University of Lausanne (UNIL), Ecole Polytechnique Federale de Lausanne (EPFL), University of Geneva (UNIGE) and Geneva University Hospitals (HUG). I.O.J. is supported by the SNSF Eccellenza Fellowship PCEFP2_194260. M.P. is supported by the UKRI Future Leaders Fellowship MR/T020296/2. D.S.N. is supported by NIH under NINDS award R01 NS088040 and by the Center of Advanced Imaging Innovation and Research (CAI²R, www.cai2r.net), a NIBIB Biomedical Technology Resource Center: P41 EB017183.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.neuroimage.2022.119277.

Appendix: DKI(t) for the orientationally-averaged anisotropic KM

We begin from a familiar DKI representation of a standard (scalar) KM:

$$\ln S_{KM}(b) = -bD + \frac{b^2}{2} f(1-f)(D_1 - D_2)^2 \cdot F(\tau),$$

$$F(\tau) = \frac{2(e^{-\tau} - 1 + \tau)}{\tau^2}, \quad \tau = \frac{t}{t_{ex}}$$

corresponding to the kurtosis (Fieremans et al., 2010) $K(\tau) = K_0 F(\tau)$ with

$$K_0 = 3f(1-f) \cdot \frac{(D_1 - D_2)^2}{D^2}.$$

We now use the above expressions for a single ensemble of neurites and their immediate extracellular space, oriented at an angle θ to the gradient direction, as function of $x = \cos \theta$. This means that $D_1 = D_{i,\parallel} x^2$ and $D_2 = D_{e,\perp} + \Delta_e x^2$, where $\Delta_e = D_{e,\parallel} - D_{e,\perp}$. Note that $\Delta_e \equiv 0$ and $D_{e,\perp} \equiv D_e$ for the isotropic extra-cellular space assumed in the main text. To average over directions, we expand $S_{KM}(b, x)$ in moments up to b^2 , integrate term-by-term over the orientations and re-expand in the exponential:

$$\bar{S}(b) = \int_0^1 dx S_{KM}(b, x) \simeq e^{-b\bar{D} + \frac{b^2 \bar{D}^2 \bar{K}}{6}}.$$

In this way, after some algebra we obtain

$$\bar{D} = \frac{1}{3} [f D_{i,\parallel} + (1-f)(3D_{e,\perp} + \Delta_e)]$$

and

$$\bar{K} = K_0 F(\tau) + K_\infty,$$

where

$$K_0 = \frac{3f(1-f) \left[D_{e,\perp}^2 + \frac{2}{3} D_{e,\perp} (\Delta_e - D_{i,\parallel}) + \frac{1}{5} (\Delta_e - D_{i,\parallel})^2 \right]}{\bar{D}^2},$$

$$K_\infty = \frac{4}{15} \frac{[f D_{i,\parallel} + (1-f)\Delta_e]^2}{\bar{D}^2}.$$

For $D_{e,\perp} = 0$ or for $f = 1$ (only sticks), $\bar{K} = K_\infty = \frac{12}{5}$, the familiar kurtosis value for the Callaghan model. Since $F(0) = 1$, the initial value

$$K(t)|_{t=0} = K_0 + K_\infty.$$

The long-time asymptotic behavior is

$$K(t)|_{t \gg t_{ex}} \simeq K_\infty + 2K_0 \cdot \frac{t_{ex}}{t}.$$

Note that the residual kurtosis $K_\infty \equiv K(t)|_{t \rightarrow \infty}$ corresponds to that of the isotropic mixture of diffusion tensors with axial and radial diffusivities $f D_{i,\parallel} + (1-f) D_{e,\parallel}$ and $(1-f) D_{e,\perp}$, respectively.

In the main text we set $\Delta_e = 0$ and $D_{e,\perp} = D_{e,\parallel} = D_e$ in the above expressions, and end up neglecting K_∞ when analyzing data, since the above model does not adequately describe mixing between sticks with different orientations at sufficiently long times.

References

- Afzali, M., Nilsson, M., Palombo, M., Jones, D.K., 2021. SPHERIOUSLY? The challenges of estimating sphere radius non-invasively in the human brain from diffusion MRI. *Neuroimage* 237, 118183. doi:10.1016/j.neuroimage.2021.118183.
- Aggarwal, M., Smith, M.D., Calabresi, P.A., 2020. Diffusion-time dependence of diffusional kurtosis in the mouse brain. *Magn. Reson. Med.* 84, 1564–1578. doi:10.1002/mrm.28189.
- Alexander, D.C., Hubbard, P.L., Hall, M.G., Moore, E.A., Pitto, M., Parker, G.J.M., Dyrby, T.B., 2010. Orientationally invariant indices of axon diameter and density from diffusion MRI. *Neuroimage* 52, 1374–1389. doi:10.1016/j.neuroimage.2010.05.043.
- Andersson, M., Kjer, H.M., Rafael-Patino, J., Pacureanu, A., Pakkenberg, B., Thiran, J.P., Pitto, M., Bech, M., Dahl, A.B., Dahl, V.A., Dyrby, T.B., 2020. Axon morphology is modulated by the local environment and impacts the noninvasive investigation of its structure–function relationship. *PNAS* 117, 33649–33659. doi:10.1073/pnas.2012533117.
- Arbabi, A., Kai, J., Khan, A.R., Baron, C.A., 2020. Diffusion dispersion imaging: mapping oscillating gradient spin-echo frequency dependence in the human brain. *Magn. Reson. Med.* 83, 2197–2208. doi:10.1002/mrm.28083.
- Assaf, Y., Cohen, Y., 1998. Non-mono-exponential attenuation of water and N-acetyl aspartate signals due to diffusion in brain tissue. *J. Magn. Reson.* 131, 69–85. doi:10.1006/jmre.1997.1313.
- Bai, R., Li, Z., Sun, C., Hsu, Y.C., Liang, H., Basser, P., 2020. Feasibility of filter-exchange imaging (FEXI) in measuring different exchange processes in human brain. *Neuroimage* 219, 117039. doi:10.1016/j.neuroimage.2020.117039.
- Bai, R., Springer, C.S., Plenz, D., Basser, P.J., 2019. Brain active transmembrane water cycling measured by MR is associated with neuronal activity. *Magn. Reson. Med.* 81, 1280–1295. doi:10.1002/mrm.27473.
- Bai, R., Springer, C.S., Plenz, D., Basser, P.J., 2018. Fast, Na⁺/K⁺ pump driven, steady-state transcytlemmal water exchange in neuronal tissue: a study of rat brain cortical cultures. *Magn. Reson. Med.* 79, 3207–3217. doi:10.1002/mrm.26980.

- Barakovic, M., Tax, C.M.W., Rudrapatna, U., Chamberland, M., Rafael-Patino, J., Granziera, C., Thiran, J.P., Daducci, A., Canales-Rodríguez, E.J., Jones, D.K., 2021. Resolving bundle-specific intra-axonal T2 values within a voxel using diffusion-relaxation tract-based estimation. *Neuroimage* 227, 117617. doi:10.1016/j.neuroimage.2020.117617.
- Baylis, P.H., 1988. *Water Movement Through Lipid Bilayers, Pores and Plasma Membranes: Theory and Reality*, 228. John Wiley and Sons Ltd, New York (Distinguished lecture series of the society of general physiologists, volume 4). Alan Finkelsteinpages, £38.45 (1987). *Cell biochemistry and function* 6, 223–223. doi:10.1002/cbf.290060323.
- Benga, G., Chapman, B.E., Kuchel, P.W., 2009. Comparative NMR studies of diffusional water permeability of red blood cells from different species: XV. Agile wallaby (*Macropus agilis*), red-necked wallaby (*Macropus rufogriseus*) and Goodfellow's tree kangaroo (*Dendrolagus goodfellowi*). *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* 154, 105–109. doi:10.1016/j.cbpa.2009.05.008.
- Benga, G., Kuchel, P.W., Chapman, B.E., Cox, G.C., Ghiran, I., Gallagher, C.H., 2000. Comparative cell shape and diffusional water permeability of red blood cells from indian elephant (*elephas maximus*) and man (*homo sapiens*). *Comp. Haematol. Int.* 10, 1–8. doi:10.1007/s005800070020.
- Bondareff, W., Pysh, J.J., 1968. Distribution of the extracellular space during postnatal maturation of rat cerebral cortex. *Anat. Rec.* 160, 773–780. doi:10.1002/ar.1091600412.
- Boss, D., Kühn, J., Jourdain, P., Depeursinge, C.D., Magistretti, P.J., Marquet, P.P., 2013. Measurement of absolute cell volume, osmotic membrane water permeability, and refractive index of transmembrane water and solute flux by digital holographic microscopy. *JBO* 18, 036007. doi:10.1117/1.JBO.18.3.036007.
- Brusini, L., Menegaz, G., Nilsson, M., 2019. Monte carlo simulations of water exchange through myelin wraps: implications for diffusion MRI. *IEEE Trans. Med. Imaging* 38, 1438–1445. doi:10.1109/TMI.2019.2894398.
- Burcaw, L.M., Fieremans, E., Novikov, D.S., 2015. Mesoscopic structure of neuronal tracts from time-dependent diffusion. *Neuroimage* 114, 18–37. doi:10.1016/j.neuroimage.2015.03.061.
- Callaghan, P.T., Jolley, K.W., Lelievre, J., 1979. Diffusion of water in the endosperm tissue of wheat grains as studied by pulsed field gradient nuclear magnetic resonance. *Biophys. J.* 28, 133–141. doi:10.1016/s0006-3495(79)85164-4.
- Callaghan, R., Alexander, D.C., Palombo, M., Zhang, H., 2020. Config: contextual fibre growth to generate realistic axonal packing for diffusion MRI simulation. *Neuroimage* 220, 117107. doi:10.1016/j.neuroimage.2020.117107.
- Chakwizira, A., Westin, C.F., Brabec, J., Lasič, S., Knutsson, L., Szczepankiewicz, F., Nilsson, M., 2021. Probing restricted diffusion and exchange using free gradient waveforms: validation by numerical simulations. arXiv:2111.11094 [physics]. arXiv preprint arXiv:2111.11094.
- Coelho, S., Fieremans, E., Novikov, D.S., 2021. How do we know we measure tissue parameters, not the prior? *Proc. Int. Soc. Magn. Reson. Med.* 30, 397.
- Coelho, S., Pozo, J.M., Jespersen, S.N., Jones, D.K., Frangi, A.F., 2019. Resolving degeneracy in diffusion MRI biophysical model parameter estimation using double diffusion encoding. *Magn. Reson. Med.* 82, 395–410. doi:10.1002/mrm.27714.
- Daducci, A., Canales-Rodríguez, E.J., Zhang, H., Dyrby, T.B., Alexander, D.C., Thiran, J.-P., 2015. Accelerated microstructure imaging via convex optimization (AMICO) from diffusion MRI data. *Neuroimage* 105, 32–44. doi:10.1016/j.neuroimage.2014.10.026.
- Dhital, B., Kellner, E., Kiselev, V.G., Reisert, M., 2018. The absence of restricted water pool in brain white matter. *NeuroImage Microstruct. Imaging* 182, 398–406. doi:10.1016/j.neuroimage.2017.10.051.
- Dhital, B., Reisert, M., Kellner, E., Kiselev, V.G., 2019. Intra-axonal diffusivity in brain white matter. *Neuroimage* 189, 543–550. doi:10.1016/j.neuroimage.2019.01.015.
- Diao, Y., Jelescu, I.O., 2022. Parameter estimation for WMTI-Watson model of white matter using encoder-decoder recurrent neural network. arXiv doi:10.48550/arXiv.2203.00595.
- Does, M.D., Parsons, E.C., Gore, J.C., 2003. Oscillating gradient measurements of water diffusion in normal and globally ischemic rat brain. *Magn. Reson. Med. Off. J. Soc. Magn. Reson. Med. Soc. Magn. Reson. Med.* 49, 206–215. doi:10.1002/mrm.10385.
- Duong, T.Q., Ackerman, J.J., Ying, H.S., Neil, J.J., 1998. Evaluation of extra- and intra-cellular apparent diffusion in normal and globally ischemic rat brain via 19F NMR. *Magn. Reson. Med. Off. J. Soc. Magn. Reson. Med. Soc. Magn. Reson. Med.* 40, 1–13.
- Efron, B., Hastie, T., Johnstone, I., Tibshirani, R., 2004. Least angle regression. *Ann. Stat.* 32, 407–499. doi:10.1214/009053604000000067.
- Fieremans, E., Burcaw, L.M., Lee, H.H., Lemberskiy, G., Veraart, J., Novikov, D.S., 2016. *In vivo* observation and biophysical interpretation of time-dependent diffusion in human white matter. *Neuroimage* 129, 414–427. doi:10.1016/j.neuroimage.2016.01.018.
- Fieremans, E., Jensen, J.H., Helpert, J.A., 2011. White matter characterization with diffusional kurtosis imaging. *Neuroimage* 58, 177–188. doi:10.1016/j.neuroimage.2011.06.006.
- Fieremans, E., Novikov, D.S., Jensen, J.H., Helpert, J.A., 2010. Monte Carlo study of a two-compartment exchange model of diffusion. *NMR Biomed.* 23, 711–724. doi:10.1002/nbm.1577.
- Ginsburger, K., Matuschke, F., Poupon, F., Mangin, J.-F., Axer, M., Poupon, C., 2019. MEDUSA: a GPU-based tool to create realistic phantoms of the brain microstructure using tiny spheres. *Neuroimage* 193, 10–24. doi:10.1016/j.neuroimage.2019.02.055.
- Gyori, N.G., Clark, C.A., Alexander, D.C., Kaden, E., 2021a. On the potential for mapping apparent neural soma density via a clinically viable diffusion MRI protocol. *Neuroimage* 239, 118303. doi:10.1016/j.neuroimage.2021.118303.
- Gyori, N.G., Palombo, M., Clark, C.A., Zhang, H., Alexander, D.C., 2021b. Training data distribution significantly impacts the estimation of tissue microstructure with machine learning. *Magn Reson Med.* 87: 932–947. doi:10.1002/mrm.29014.

- Harkins, K.D., Galons, J.P., Secomb, T.W., Trouard, T.P., 2009. Assessment of the effects of cellular tissue properties on ADC measurements by numerical simulation of water diffusion. *Magn. Reson. Med.* 62, 1414–1422. doi:10.1002/mrm.22155.
- Hill, I., Palombo, M., Santin, M., Branzoli, F., Philippe, A.C., Wassermann, D., Aigrot, M.S., Stankoff, B., Baron-Van Evercooren, A., Felfli, M., Langui, D., Zhang, H., Lehericy, S., Petiet, A., Alexander, D.C., Ciccarelli, O., Drobnjak, I., 2021. Machine learning based white matter models with permeability: an experimental study in cuprizone treated *in-vivo* mouse model of axonal demyelination. *Neuroimage* 224, 117425. doi:10.1016/j.neuroimage.2020.117425.
- Howard, A.F., Lange, F.J., Mollink, J., Cottaar, M., Drakesmith, M., Rudrapatna, S.U., Jones, D.K., Miller, K.L., Jbabdi, S., 2020. Estimating intra-axonal axial diffusivity in the presence of fibre orientation dispersion. *bioRxiv* doi:10.1101/2020.10.09.332700.
- Hutter, J., Slator, P.J., Christiaens, D., Teixeira, R.P.A.G., Roberts, T., Jackson, L., Price, A.N., Malik, S., Hajnal, J.V., 2018. Integrated and efficient diffusion-relaxometry using ZEBRA. *Sci. Rep.* 8, 15138. doi:10.1038/s41598-018-33463-2.
- Ianus, A., Alexander, D.C., Zhang, H., Palombo, M., 2020. Mapping complex cell morphology in the grey matter with double diffusion encoding MR: A simulation study. *Neuroimage* 241, 118424. doi:10.1016/j.neuroimage.2021.118424.
- Jelescu, I.O., Palombo, M., Bagnato, F., Schilling, K.G., 2020. Challenges for biophysical modeling of microstructure. *J. Neurosci. Methods* 344, 108861. doi:10.1016/j.jneumeth.2020.108861.
- Jelescu, I.O., Uhl, Q., 2022. *Ex vivo* gray matter is complex: exchange and disorder exponents. *Proc. Int. Soc. Magn. Reson. Med.* 31, 253 London, UK.
- Jelescu, I.O., Veraart, J., Fieremans, E., Novikov, D.S., 2016. Degeneracy in model parameter estimation for multi-compartmental diffusion in neuronal tissue. *NMR Biomed.* 29, 33–47. doi:10.1002/nbm.3450.
- Jenkinson, M., Bannister, P., Brady, M., Smith, S., 2002. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* 17, 825–841.
- Jensen, J.H., Helpert, J.A., Ramani, A., Lu, H., Kaczynski, K., 2005. Diffusional kurtosis imaging: the quantification of non-gaussian water diffusion by means of magnetic resonance imaging. *Magn. Reson. Med. Off. J. Soc. Magn. Reson. Med. Soc. Magn. Reson. Med.* 53, 1432–1440. doi:10.1002/mrm.20508.
- Jespersen, S.N., Bjarkam, C.R., Nyengaard, J.R., Chakravarty, M.M., Hansen, B., Vosegaard, T., Ostergaard, L., Yablonskiy, D., Nielsen, N.C., Vestergaard-Poulsen, P., 2010. Neurite density from magnetic resonance diffusion measurements at ultrahigh field: comparison with light microscopy and electron microscopy. *Neuroimage* 49, 205–216. doi:10.1016/j.neuroimage.2009.08.053.
- Jespersen, S.N., Kroenke, C.D., Ostergaard, L., Ackerman, J.J., Yablonskiy, D.A., 2007. Modeling dendrite density from magnetic resonance diffusion measurements. *Neuroimage* 34, 1473–1486. doi:10.1016/j.neuroimage.2006.10.037.
- Jespersen, S.N., Olesen, J.L., Hansen, B., Shemesh, N., 2018. Diffusion time dependence of microstructural parameters in fixed spinal cord. *Neuroimage* 182, 329–342. doi:10.1016/j.neuroimage.2017.08.039.
- Kärger, J., 1985. NMR self-diffusion studies in heterogeneous systems. *Adv. Colloid Interface Sci.* 23, 129–148. doi:10.1016/0001-8686(85)80018-X.
- Kellner, E., Dhital, B., Kiselev, V.G., Reiser, M., 2016. Gibbs-ringing artifact removal based on local subvoxel-shifts. *Magn. Reson. Med.* 76, 1574–1581. doi:10.1002/mrm.26054.
- Kiselev, V.G., Il'yasov, K.A., 2007. Is the “biexponential diffusion” biexponential? *Magn. Reson. Med.* 57, 464–469. doi:10.1002/mrm.21164.
- Kroenke, C.D., Ackerman, J.J., Yablonskiy, D.A., 2004. On the nature of the NAA diffusion attenuated MR signal in the central nervous system. *Magn. Reson. Med. Off. J. Soc. Magn. Reson. Med. Soc. Magn. Reson. Med.* 52, 1052–1059. doi:10.1002/mrm.20260.
- Kunz, N., da Silva, A.R., Jelescu, I.O., 2018. Intra- and extra-axonal axial diffusivities in the white matter: which one is faster? *Neuroimage* 181, 314–322. doi:10.1016/j.neuroimage.2018.07.020.
- Lampinen, B., Szczepankiewicz, F., van Westen, D., Englund, E., C Sundgren, P., Lätt, J., Ståhlberg, F., Nilsson, M., 2017. Optimal experimental design for filter exchange imaging: apparent exchange rate measurements in the healthy brain and in intracranial tumors. *Magn. Reson. Med.* 77, 1104–1114. doi:10.1002/mrm.26195.
- Latour, L.L., Svoboda, K., Mitra, P.P., Sotak, C.H., 1994. Time-dependent diffusion of water in a biological model system. *Proc. Natl. Acad. Sci. USA* 91, 1229–1233.
- Lazari, A., Lipp, I., 2021. Can MRI measure myelin? Systematic review, qualitative assessment, and meta-analysis of studies validating microstructural imaging with myelin histology. *Neuroimage* 230, 117744. doi:10.1016/j.neuroimage.2021.117744.
- Lee, H.H., Fieremans, E., Huang, S.Y., Qiyuan, T., Novikov, D.S., 2021. Localization regime of diffusion in human gray matter on a high-gradient MR system: sensitivity to soma size. *Proc. Int. Soc. Mag. Reson. Med.* 30, 639.
- Lee, H.H., Fieremans, E., Novikov, D.S., 2018. LEMONADE(t): exact relation of time-dependent diffusion signal moments to neuronal microstructure. *Proc. Int. Soc. Magn. Reson. Med.* 27, 884.
- Lee, H.H., Papaioannou, A., Kim, S.L., Novikov, D.S., Fieremans, E., 2020a. A time-dependent diffusion MRI signature of axon caliber variations and beading. *Commun. Biol.* 3, 1–13. doi:10.1038/s42003-020-1050-x.
- Lee, H.H., Papaioannou, A., Novikov, D.S., Fieremans, E., 2020b. *In vivo* observation and biophysical interpretation of time-dependent diffusion in human cortical gray matter. *Neuroimage* 222, 117054. doi:10.1016/j.neuroimage.2020.117054.
- Li, J.G., Stanis, G.J., Henkelman, R.M., 1998. Integrated analysis of diffusion and relaxation of water in blood. *Magn. Reson. Med.* 40, 79–88. doi:10.1002/mrm.1910400112.
- Mancini, M., Karakuzu, A., Cohen-Adad, J., Mercignani, M., Nichols, T.E., Stikov, N., 2020. An interactive meta-analysis of MRI biomarkers of myelin. *Life* 9, e61523. doi:10.7554/eLife.61523.
- Martins, J.P.D.A., Nilsson, M., Lampinen, B., Palombo, M., While, P.T., Westin, C.F., Szczepankiewicz, F., 2021. Neural networks for parameter estimation in microstructural MRI: a study with a high-dimensional diffusion-relaxation model of white matter microstructure. *Neuroimage* 244, 118601. doi:10.1016/j.neuroimage.2021.118601.
- McKinnon, E.T., Jensen, J.H., Glenn, G.R., Helpert, J.A., 2017. Dependence on *b*-value of the direction-averaged diffusion-weighted imaging signal in brain. *Magn. Reson. Imaging* 36, 121–127. doi:10.1016/j.mri.2016.10.026.
- Meier, C., Dreher, W., Leibfritz, D., 2003. Diffusion in compartmental systems. I. a comparison of an analytical model with simulations. *Magn. Reson. Med.* 50, 500–509. doi:10.1002/mrm.10557.
- Motta, A., Berning, M., Boergens, K.M., Staffler, B., Beining, M., Loomba, S., Hennig, P., Wissler, H., Helmstaedter, M., 2019. Dense connectomic reconstruction in layer 4 of the somatosensory cortex. *Science* 366. doi:10.1126/science.aay3134.
- Moutal, N., Nilsson, M., Topgaard, D., Grebenkov, D., 2018. The Kärger vs bi-exponential model: theoretical insights and experimental validations. *J. Magn. Reson.* 296, 72–78. doi:10.1016/j.jmr.2018.08.015.
- Nedjati-Gilani, G.L., Schneider, T., Hall, M.G., Cawley, N., Hill, I., Ciccarelli, O., Drobnjak, I., Wheeler-Kingshott, C.A.M.G., Alexander, D.C., 2017. Machine learning based compartment models with permeability for white matter microstructure imaging. *Neuroimage* 150, 119–135. doi:10.1016/j.neuroimage.2017.02.013.
- Nilsson, M., Latt, J., van Westen, D., Brockstedt, S., Lasic, S., Stahlberg, F., Topgaard, D., 2013a. Noninvasive mapping of water diffusional exchange in the human brain using filter-exchange imaging. *Magn. Reson. Med.* 69, 1573–1581. doi:10.1002/mrm.24395.
- Nilsson, M., van Westen, D., Ståhlberg, F., Sundgren, P.C., Lätt, J., 2013b. The role of tissue microstructure and water exchange in biophysical modelling of diffusion in white matter. *Magn. Reson. Mater. Phys.* 26, 345–370. doi:10.1007/s10334-013-0371-x.
- Novikov, D.S., 2021. The present and the future of microstructure MRI: from a paradigm shift to normal science. *J. Neurosci. Methods* 351, 108947. doi:10.1016/j.jneumeth.2020.108947.
- Novikov, D.S., Fieremans, E., Jespersen, S.N., Kiselev, V.G., 2019. Quantifying brain microstructure with diffusion MRI: theory and parameter estimation. *NMR Biomed.* 32, e3998. doi:10.1002/nbm.3998.
- Novikov, D.S., Jensen, J.H., Helpert, J.A., Fieremans, E., 2014. Revealing mesoscopic structural universality with diffusion. *Proc. Natl. Acad. Sci. USA* 111, 5088–5093. doi:10.1073/pnas.1316944111.
- Novikov, D.S., Kiselev, V.G., Jespersen, S.N., 2018a. On modeling. *Magn. Reson. Med.* 79, 3172–3193. doi:10.1002/mrm.27101.
- Novikov, D.S., Veraart, J., Jelescu, I.O., Fieremans, E., 2018b. Rotationally-invariant mapping of scalar and orientational metrics of neuronal microstructure with diffusion MRI. *Neuroimage* 174, 518–538. doi:10.1016/j.neuroimage.2018.03.006.
- Olesen, J.L., Ostergaard, L., Shemesh, N., Jespersen, S.N., 2022. Diffusion time dependence, power-law scaling, and exchange in gray matter. *Neuroimage* 251, 118976. doi:10.1016/j.neuroimage.2022.118976.
- Olesen, J.L., Ostergaard, L., Shemesh, N., Jespersen, S.N., 2021. Beyond the diffusion standard model in fixed rat spinal cord with combined linear and planar encoding. *Neuroimage* 231, 117849. doi:10.1016/j.neuroimage.2021.117849.
- Pacheco, C.R., Morales, C.N., Ramirez, A.E., Muñoz, F.J., Gallegos, S.S., Caviedes, P.A., Aguayo, L.G., Opazo, C.M., 2015. Extracellular α -synuclein alters synaptic transmission in brain neurons by perforating the neuronal plasma membrane. *J. Neurochem.* 132, 731–741. doi:10.1111/jnc.13060.
- Palombo, M., Alexander, D.C., Zhang, H., 2021. Large-scale analysis of brain cell morphology informs microstructure modelling of gray matter. *Proc. Int. Soc. Magn. Reson. Med.* 30, 642.
- Palombo, M., Alexander, D.C., Zhang, H., 2019. A generative model of realistic brain cells with application to numerical simulation of the diffusion-weighted MR signal. *Neuroimage* 188, 391–402. doi:10.1016/j.neuroimage.2018.12.025.
- Palombo, M., Ianus, A., Guerrero, M., Nunes, D., Alexander, D.C., Shemesh, N., Zhang, H., 2020. SANDI: a compartment-based model for non-invasive apparent soma and neurite imaging by diffusion MRI. *Neuroimage* 215, 116835. doi:10.1016/j.neuroimage.2020.116835.
- Pyatigorskaya, N., Le Bihan, D., Reynaud, O., Ciobanu, L., 2014. Relationship between the diffusion time and the diffusion MRI signal observed at 17.2 Tesla in the healthy rat brain cortex. *Magn. Reson. Med.* 72, 492–500. doi:10.1002/mrm.24921.
- Quirk, J.D., Bretthorst, G.L., Duong, T.Q., Snyder, A.Z., Springer, C.S., Ackerman, J.J.H., Neil, J.J., 2003. Equilibrium water exchange between the intra- and extracellular spaces of mammalian brain. *Magn. Reson. Med.* 50, 493–499. doi:10.1002/mrm.10565.
- Reiser, M., Kellner, E., Dhital, B., Hennig, J., Kiselev, V.G., 2017. Disentangling micro from mesostructure by diffusion MRI: a Bayesian approach. *Neuroimage* 147, 964–975. doi:10.1016/j.neuroimage.2016.09.058.
- Reiser, M., Kiselev, V.G., Dhital, B., 2019. A unique analytical solution of the white matter standard model using linear and planar encodings. *Magn. Reson. Med.* 81, 3819–3825. doi:10.1002/mrm.27685.
- Sehy, J.V., Ackerman, J.J.H., Neil, J.J., 2002. Evidence that both fast and slow water ADC components arise from intracellular space. *Magn. Reson. Med.* 48, 765–770. doi:10.1002/mrm.10301.
- Shapson-Coe, A., Januszewski, M., Berger, D.R., Pope, A., Wu, Y., Blakely, T., Schalek, R.L., Li, P., Wang, S., Maitin-Shepard, J., Karlupia, N., Dorkenwald, S., Sjostedt, E., Leavitt, L., Lee, D., Bailey, L., Fitzmaurice, A., Kar, R., Field, B., Wu, H., Wagner-Carena, J., Aley, D., Lau, J., Lin, Z., Wei, D., Pfister, H., Peleg, A., Jain, V., Lichtman, J.W., 2021. A connectomic study of a petascale fragment of human cerebral cortex. *bioRxiv* doi:10.1101/2021.05.29.446289.
- Shepherd, T.M., Thelwall, P.E., Stanis, G.J., Blackband, S.J., 2009. Aldehyde fixative solutions alter the water relaxation and diffusion properties of nervous tissue. *Magn. Reson. Med.* 62, 26–34. doi:10.1002/mrm.21977.

- Spocter, M.A., Hopkins, W.D., Barks, S.K., Bianchi, S., Hehmeyer, A.E., Anderson, S.M., Stimpson, C.D., Fobbs, A.J., Hof, P.R., Sherwood, C.C., 2012. Neuropil distribution in the cerebral cortex differs between humans and chimpanzees. *J. Comp. Neurol.* 520, 2917–2929. doi:10.1002/cne.23074.
- Stanisz, G.J., Szafer, A., Wright, G.A., Henkelman, R.M., 1997. An analytical model of restricted diffusion in bovine optic nerve. *Magn. Reson. Med.* 37, 103–111 official journal of the society of magnetic resonance in medicine /society of magnetic resonance in medicine.
- Szczepankiewicz, F., van Westen, D., Englund, E., Westin, C.-F., Ståhlberg, F., Lätt, J., Sundgren, P.C., Nilsson, M., 2016. The link between diffusion MRI and tumor heterogeneity: mapping cell eccentricity and density by diffusional variance decomposition (DIVIDE). *Neuroimage* 142, 522–532. doi:10.1016/j.neuroimage.2016.07.038.
- Tax, C.M.W., Kleban, E., Chamberland, M., Baraković, M., Rudrapatna, U., Jones, D.K., 2021. Measuring compartmental T2-orientational dependence in human brain white matter using a tiltable RF coil and diffusion-T2 correlation MRI. *Neuroimage* 236, 117967. doi:10.1016/j.neuroimage.2021.117967.
- Thelwall, P.E., Shepherd, T.M., Stanisz, G.J., Blackband, S.J., 2006. Effects of temperature and aldehyde fixation on tissue water diffusion properties, studied in an erythrocyte ghost tissue model. *Magn. Reson. Med.* 56, 282–289. doi:10.1002/mrm.20962.
- Veraart, J., Fieremans, E., Novikov, D.S., 2019. On the scaling behavior of water diffusion in human brain white matter. *Neuroimage* 185, 379–387. doi:10.1016/j.neuroimage.2018.09.075.
- Veraart, J., Fieremans, E., Novikov, D.S., 2016a. Diffusion MRI noise mapping using random matrix theory. *Magn. Reson. Med.* 76, 1582–1593. doi:10.1002/mrm.26059.
- Veraart, J., Fieremans, E., Rudrapatna, U., Jones, D.K., Novikov, D.S., 2018a. Biophysical modeling of the gray matter: does the “stick” model hold? *Proc. Int. Soc. Magn. Reson. Med.* 27, 1094.
- Veraart, J., Novikov, D.S., Christiaens, D., Ades-aron, B., Sijbers, J., Fieremans, E., 2016b. Denoising of diffusion MRI using random matrix theory. *Neuroimage* 142, 394–406. doi:10.1016/j.neuroimage.2016.08.016.
- Veraart, J., Novikov, D.S., Fieremans, E., 2018b. TE dependent diffusion imaging (TEdDI) distinguishes between compartmental T2 relaxation times. *Neuroimage* 182, 360–369. doi:10.1016/j.neuroimage.2017.09.030.
- Veraart, J., Nunes, D., Rudrapatna, U., Fieremans, E., Jones, D.K., Novikov, D.S., Shemesh, N., 2020. Noninvasive quantification of axon radii using diffusion MRI. *Elife* 9, e49855. doi:10.7554/eLife.49855.
- Veraart, J., Sijbers, J., Sunaert, S., Leemans, A., Jeurissen, B., 2013. Weighted linear least squares estimation of diffusion MRI parameters: strengths, limitations, and pitfalls. *Neuroimage* 81, 335–346. doi:10.1016/j.neuroimage.2013.05.028.
- Vestergaard-Poulsen, P., Hansen, B., Østergaard, L., Jakobsen, R., 2007. Microstructural changes in ischemic cortical gray matter predicted by a model of diffusion-weighted MRI. *J. Magn. Reson. Imaging* 26, 529–540. doi:10.1002/jmri.21030.
- Wang, S.S.H., Shultz, J.R., Burish, M.J., Harrison, K.H., Hof, P.R., Towns, L.C., Wagers, M.W., Wyatt, K.D., 2008. Functional trade-offs in white matter axonal scaling. *J. Neurosci.* 28, 4047–4056. doi:10.1523/JNEUROSCI.5559-05.2008.
- Wang, Y., Wang, Q., Haldar, J.P., Yeh, F.C., Xie, M., Sun, P., Tu, T.W., Trinkaus, K., Klein, R.S., Cross, A.H., Song, S.K., 2011. Quantification of increased cellularity during inflammatory demyelination. *Brain J. Neurol.* 134, 3590–3601. doi:10.1093/brain/awr307.
- Williamson, N.H., Ravin, R., Benjamini, D., Merkle, H., Falgairolle, M., O'Donovan, M.J., Blivis, D., Ide, D., Cai, T.X., Ghorashi, N.S., Bai, R., Basser, P.J., 2019. Magnetic resonance measurements of cellular and sub-cellular membrane structures in live and fixed neural tissue. *Elife* 8, e51101. doi:10.7554/eLife.51101.
- Yang, D.M., Huettner, J.E., Bretthorst, G.L., Neil, J.J., Garbow, J.R., Ackerman, J.J.H., 2018. Intracellular water preexchange lifetime in neurons and astrocytes. *Magn. Reson. Med.* 79, 1616–1627. doi:10.1002/mrm.26781.
- Zhang, H., Schneider, T., Wheeler-Kingshott, C.A., Alexander, D.C., 2012. NODDI: practical *in vivo* neurite orientation dispersion and density imaging of the human brain. *Neuroimage* 61, 1000–1016. doi:10.1016/j.neuroimage.2012.03.072.