

# Berman's Inequality under Random Scaling

Enkelejd Hashorva and Zhichao Weng

University of Lausanne, Faculty of Business and Economics (HEC Lausanne), Lausanne 1015, Switzerland

April 23, 2014

**Abstract:** Berman's inequality is the key for establishing asymptotic properties of maxima of Gaussian random sequences and supremum of Gaussian random fields. This contribution shows that, asymptotically an extended version of Berman's inequality can be established for randomly scaled Gaussian random vectors. Two applications presented in this paper demonstrate the use of Berman's inequality under random scaling.

**Key words:** Berman's inequality; Limit distribution; Extremal index; Random scaling; Hüsler-Reiss distribution.

**AMS Classification:** Primary 60G15; Secondary 60G70

## 1 Introduction

In the analysis of extreme values of Gaussian processes and Gaussian random fields Berman's inequality plays a crucial role. Essentially, for given two Gaussian distribution functions in  $\mathbb{R}^d$  it bounds their difference by comparing the covariances. The key result that motivated this comparison method is Plackett's partial differential equation given in [28]. As explained in [20], it was then developed by Slepian [29], Berman [1, 2], Cramér [4], Piterbarg [26, 27] and then by Li and Shao [22]. Specifically, the developed results are summarised by Berman's inequality which we formulate below in the most general form derived in [22]. Let therefore  $\mathbf{X} = (X_1, \dots, X_n)$  and  $\mathbf{Y} = (Y_1, \dots, Y_n)$  be two Gaussian random vectors with  $N(0, 1)$  components and covariance matrices  $\Lambda_1 = (\lambda_{ij}^{(1)})$  and  $\Lambda_2 = (\lambda_{ij}^{(2)})$ , respectively. For arbitrary constants  $u_i, i \leq n$ , [22] obtained

$$\mathbb{P}(X_i \leq u_i, 1 \leq i \leq n) - \mathbb{P}(Y_i \leq u_i, 1 \leq i \leq n) \leq \frac{1}{2\pi} \sum_{1 \leq i < j \leq n} A_{ij} \exp\left(-\frac{u_i^2 + u_j^2}{2(1 + \rho_{ij})}\right),$$

where

$$\rho_{ij} := \max(|\lambda_{ij}^{(1)}|, |\lambda_{ij}^{(2)}|), \quad A_{ij} = |\arcsin(\lambda_{ij}^{(1)}) - \arcsin(\lambda_{ij}^{(2)})|. \quad (1.1)$$

Clearly, for arbitrary constants  $v_i, u_i, i \leq n$ , set  $w := \min_{1 \leq i \leq n} \min(|u_i|, |v_i|)$

$$\mathbb{P}(-v_i < X_i \leq u_i, 1 \leq i \leq n) - \mathbb{P}(-v_i < Y_i \leq u_i, 1 \leq i \leq n) \leq \frac{2}{\pi} \sum_{1 \leq i < j \leq n} A_{ij} \exp\left(-\frac{w^2}{1 + \rho_{ij}}\right), \quad (1.2)$$

for a detailed proof see [21], see also [23] for recent extensions.

Berman's inequality can be applied also to non-Gaussian random vectors. For instance, consider two random vectors  $\tilde{\mathbf{X}} = (S_1 X_1, \dots, S_n X_n)$  and  $\tilde{\mathbf{Y}} = (S_1 Y_1, \dots, S_n Y_n)$  with  $S, S_i, i \leq n$  some positive independent random variables with common distribution function  $G$  being further independent from  $\mathbf{X}$  and  $\mathbf{Y}$ . In the special case  $G$  is the uniform distribution on  $(0, 1)$ , the upper bound in (1.2) implies

$$\begin{aligned} \Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v}) &:= \mathbb{P}(-v_i < S_i X_i \leq u_i, 1 \leq i \leq n) - \mathbb{P}(-v_i < S_i Y_i \leq u_i, 1 \leq i \leq n) \\ &\leq \frac{2}{\pi} \sum_{1 \leq i < j \leq n} A_{ij} \int_0^1 \int_0^1 \exp\left(-\frac{(w/s_i)^2 + (w/s_j)^2}{2(1 + \rho_{ij})}\right) ds_i ds_j \\ &\leq \frac{2}{\pi} \sum_{1 \leq i < j \leq n} A_{ij} \exp\left(-\frac{w^2}{1 + \rho_{ij}}\right). \end{aligned} \quad (1.3)$$

Another tractable case is when  $G(x) = 1 - \exp(-x), x > 0$  is the exponential distribution. Indeed, by (1.2) for all  $0 < a, b < 1$  we have

$$\Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v}) \leq \frac{2}{\pi} \sum_{1 \leq i < j \leq n} A_{ij} \int_0^\infty \int_0^\infty \exp\left(-\frac{(w/s_i)^2 + (w/s_j)^2}{2(1 + \rho_{ij})} - s_i - s_j\right) ds_i ds_j$$

$$\begin{aligned}
&= \frac{2}{\pi} \sum_{1 \leq i < j \leq n} A_{ij} \int_0^\infty \int_0^\infty \exp\left(-\frac{(w/s_i)^2 + (w/s_j)^2}{2(1 + \rho_{ij})} - as_i - bs_j\right) \exp(-(1-a)s_i - (1-b)s_j) ds_i ds_j \\
&\leq \frac{2}{\pi(1-a)(1-b)} \sum_{1 \leq i < j \leq n} A_{ij} \exp\left(-\frac{3}{2}(a^{\frac{2}{3}} + b^{\frac{2}{3}})(1 + \rho_{ij})^{-\frac{1}{3}} w^{\frac{2}{3}}\right).
\end{aligned} \tag{1.4}$$

Clearly, if we do not know the distribution function of  $S$  it is not possible to obtain an explicit upper bound for  $\Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v})$ . Since for the analysis of extremes of Gaussian random sequences or processes Berman's inequality is applied for large values of the  $u_i$ 's and  $v_i$ 's (see e.g., [27]), in this paper we are concerned with the derivation of Berman's inequality for some general scaling random variable  $S$  and all  $u_i$ 's and  $v_i$ 's sufficiently large. We shall consider two particular cases for the random vector  $\mathbf{S} = (S_1, \dots, S_n)$ , namely it has independent components, and it is comonotonic with  $\mathbf{S} = (S, \dots, S) =: \mathbf{S}\mathbf{1}$ . From the proofs it can be seen that the joint dependence of  $(S_i, S_j)$  for any pair  $(i, j)$  is crucial; our results can be in fact extended for certain tractable dependence models. We shall deal for simplicity only with these two cases.

Since random scaling is a natural phenomena related to the time-value of money in finance, measurement errors in experimental data, or physical constrains, the extension of Berman's inequality for inflated/deflated Gaussian random vectors is of certain interest also for statistical applications.

Of course, Berman's inequality alone is not enough for extending [17] to randomly scaled Gaussian triangular arrays; some additional results (see [15, 16]) which show that for some tractable tail assumptions on  $S$  the scaled random vector  $\widetilde{\mathbf{X}}$  behaves similarly to  $\mathbf{X}$  are also important. Specifically, we shall deal with two large classes of random scaling: a)  $S$  is a bounded random variable with a tractable tail behaviour at the right endpoint of its distribution function, including in particular the case that its survival function is regularly varying, and b)  $S$  is a Weibull-type random variable.

In view of our findings, several known results for Gaussian random sequences and processes can be extended to the scaled Gaussian framework; we shall demonstrate this with two representative applications.

Organisation of the rest of the paper: Section 2 presents Berman's inequality for scaled Gaussian random vectors. In Section 3 we display two applications, while the proofs are relegated to Section 4.

## 2 Main Results

We consider first the case that  $S$  is non-negative with distribution function  $G$  which has right endpoint equal 1. Intuitively, large values of  $S$  do not influence significantly large values of the product say  $SX$  if  $X$  is a Gaussian random variable being independent of  $S$ . It turns out that the following asymptotic upper bound

$$\mathbb{P}(S > 1 - 1/u) \leq c_A u^{-\tau} \tag{2.1}$$

valid for all  $u$  large and some  $c_A > 0, \tau \geq 0$  is sufficient for the derivation of a useful upper bound for  $\Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v})$  defined above.

A canonical example of such  $S$  is the beta random variable, which is a special case of a power-tail random variables  $S$ , namely

$$\mathbb{P}(S > 1 - 1/u) = (1 + o(1))cu^{-\tau}, \quad u \rightarrow \infty \tag{2.2}$$

holds for some  $c > 0, \tau \geq 0$ . Hereafter we set  $w = \min_{1 \leq i \leq n} \min(|u_i|, |v_i|)$  and write  $\Delta_{\mathbf{S}\mathbf{1}}(\mathbf{u}, \mathbf{v})$  instead of  $\Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v})$  if  $\mathbf{S} = (S, \dots, S)$ . Further write  $\Delta_{\mathbf{S}}(u\mathbf{1})$  and  $\Delta_{\mathbf{S}\mathbf{1}}(u\mathbf{1})$  instead of  $\Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v})$  if all components of  $\mathbf{v}$  equal  $-\infty$ ,  $\mathbf{u} = (u, \dots, u) =: u\mathbf{1}$  and the covariance matrix  $\Lambda_2$  of  $\mathbf{Y}$  is identity matrix.

**Theorem 2.1.** Let  $\mathbf{X}, \widetilde{\mathbf{X}}, \mathbf{Y}, \widetilde{\mathbf{Y}}, S, S_i, i \leq n$  be as above. If (2.1) holds, then for all  $u_i, v_i, 1 \leq i \leq n$  large and  $\epsilon > 0$  we have

$$\Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v}) \leq (\mathbb{K}_A + \epsilon)w^{-4\tau} \sum_{1 \leq i < j \leq n} A_{ij}(1 + \rho_{ij})^{2\tau} \exp\left(-\frac{w^2}{1 + \rho_{ij}}\right) \quad (2.3)$$

and

$$\Delta_{S1}(\mathbf{u}, \mathbf{v}) \leq (\mathbb{K}_A^* + \epsilon)w^{-2\tau} \sum_{1 \leq i < j \leq n} A_{ij}(1 + \rho_{ij})^\tau \exp\left(-\frac{w^2}{1 + \rho_{ij}}\right), \quad (2.4)$$

where  $\mathbb{K}_A = \frac{2}{\pi}c_A^2(\Gamma(\tau + 1))^2$  and  $\mathbb{K}_A^* = \frac{2^{1-\tau}}{\pi}c_A\Gamma(\tau + 1)$ .

**Corollary 2.2.** Under the conditions of Theorem 2.1, for all  $u$  large and some positive constants  $\mathcal{Q}$  we have

$$\Delta_{\mathbf{S}}(u\mathbf{1}) \leq \mathcal{Q}u^{-4\tau} \sum_{1 \leq i < j \leq n} |\lambda_{ij}^{(1)}| \exp\left(-\frac{u^2}{1 + |\lambda_{ij}^{(1)}|}\right) \quad (2.5)$$

and

$$\Delta_{S1}(u\mathbf{1}) \leq \mathcal{Q}u^{-2\tau} \sum_{1 \leq i < j \leq n} |\lambda_{ij}^{(1)}| \exp\left(-\frac{u^2}{1 + |\lambda_{ij}^{(1)}|}\right). \quad (2.6)$$

We shall investigate below the more difficult case that the scaling random variable  $S$  has distribution function with an infinite right endpoint. Motivated by the example of the exponential distribution in Introduction, we shall assume that  $S$  has tail behaviour similar to a Weibull distribution. Specifically, for given constants  $\alpha \in \mathbb{R}, c_B, L, p \in (0, \infty)$  suppose that

$$\mathbb{P}(S > u) = (1 + o(1))c_B u^\alpha \exp(-Lu^p), \quad u \rightarrow \infty \quad (2.7)$$

is valid. The class of distribution functions satisfying (2.7) is quite large. More importantly, under (2.7)  $SX$  has also a Weibull tail behaviour if  $X$  is a  $N(0, 1)$  random variable independent of  $S$ , see e.g., [16]. We state next our second result for Weibull-type random scaling.

**Theorem 2.3.** Let  $\mathbf{X}, \widetilde{\mathbf{X}}, \mathbf{Y}, \widetilde{\mathbf{Y}}, S, S_i, i \leq n$  be as above. If (2.7) holds, then for all  $u_i, v_i, 1 \leq i \leq n$  large and  $\epsilon > 0$  we have

$$\Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v}) \leq (\mathbb{K}_B + \epsilon)w^{\frac{4\alpha+2p}{2+p}} \sum_{1 \leq i < j \leq n} A_{ij}(1 + \rho_{ij})^{-\frac{2\alpha-p}{p+2}} \exp\left(-2(1 + \rho_{ij})^{-\frac{p}{2+p}} T w^{\frac{2p}{2+p}}\right) \quad (2.8)$$

and

$$\Delta_{S1}(\mathbf{u}, \mathbf{v}) \leq (\mathbb{K}_B^* + \epsilon)w^{\frac{2\alpha+p}{2+p}} \sum_{1 \leq i < j \leq n} A_{ij}(1 + \rho_{ij})^{-\frac{2\alpha-p}{2(p+2)}} \exp\left(-2(1 + \rho_{ij})^{-1} \frac{p}{2+p} T w^{\frac{2p}{2+p}}\right), \quad (2.9)$$

where  $T = L^{\frac{2}{p+2}} p^{-\frac{p}{p+2}} + (Lp)^{\frac{2}{p+2}} 2^{-1}$ ,  $\mathbb{K}_B = 4c_B^2 (Lp)^{\frac{2(1-\alpha)}{p+2}} (p+2)^{-1}$  and  $\mathbb{K}_B^* = 2^{\frac{3+2p+\alpha}{2+p}} \pi^{-\frac{1}{2}} c_B (Lp)^{\frac{1-\alpha}{p+2}} (p+2)^{-\frac{1}{2}}$ .

**Corollary 2.4.** Under the conditions of Theorem 2.3, for all  $u$  large and some positive constants  $\mathcal{Q}$  we have

$$\Delta_{\mathbf{S}}(u\mathbf{1}) \leq \mathcal{Q}u^{\frac{4\alpha+2p}{2+p}} \sum_{1 \leq i < j \leq n} |\lambda_{ij}^{(1)}| \exp\left(-2(1 + |\lambda_{ij}^{(1)}|)^{-\frac{p}{2+p}} T u^{\frac{2p}{2+p}}\right) \quad (2.10)$$

and

$$\Delta_{S1}(u\mathbf{1}) \leq \mathcal{Q}u^{\frac{2\alpha+p}{2+p}} \sum_{1 \leq i < j \leq n} |\lambda_{ij}^{(1)}| \exp\left(-2(1 + |\lambda_{ij}^{(1)}|)^{-1} \frac{p}{2+p} T u^{\frac{2p}{2+p}}\right). \quad (2.11)$$

**Remarks:** a) Clearly, when  $S$  is uniformly distributed on  $(0, 1)$  then condition (2.1) holds with  $c_A = \tau = 1$ . For this case we have two results, the one derived in the Introduction and that given by (2.3). We see that the bound obtained by (2.3) with  $c_A = \tau = 1$  is better due to the term  $w^{-4\tau}$ .

b) Also for the case  $S$  is a unit exponential random variables we have two bounds, one which holds for all values of  $u_i, v_i, i \leq n$  and the asymptotic one given in Theorem 2.3. The bound implied by (2.8) with  $c_B = 1, \alpha = 0, p = 1, L = 1$  is asymptotically better than that implied by (1.4).

### 3 Applications

An important contribution in extreme value theory concerned with maxima of triangular arrays of Gaussian random variables is [17]. Motivated by the findings of Hüsler and Reiss in 1989 (see [18]) the aforementioned contribution considered a triangular array of  $N(0, 1)$  random variables  $\{X_{n,i}, i, n \geq 1\}$  such that for each  $n$ ,  $\{X_{n,i}, i \geq 1\}$  is a stationary Gaussian random sequence. Assume that  $\varrho_{n,j} = \mathbb{E}(X_{n,i}X_{n,i+j})$  satisfies for any  $j \geq 1$

$$\lim_{n \rightarrow \infty} (1 - \varrho_{n,j}) \ln n = \delta_j \in (0, \infty), \quad \delta_0 := 0 \quad (3.1)$$

and for each  $n$ ,  $\varrho_{n,j}$  decays fast enough as  $j$  increases. Under some additional conditions (see Theorem 3.1 below) the deep contribution [17] shows that for the maxima  $M_n = \max_{1 \leq i \leq n} X_{n,i}$

$$\lim_{n \rightarrow \infty} \mathbb{P}(M_n \leq a_n x + b_n) = \exp(-\vartheta \exp(-x)), \quad x \in \mathbb{R}, \quad (3.2)$$

where

$$a_n = (2 \ln n)^{-\frac{1}{2}}, \quad b_n = (2 \ln n)^{\frac{1}{2}} - \frac{1}{2}(2 \ln n)^{-\frac{1}{2}}(\ln \ln n + \ln 4\pi) \quad (3.3)$$

and

$$\vartheta = \mathbb{P}\left(E/2 + \sqrt{\delta_{k-1}}W_k \leq \delta_{k-1}, \text{ for all } k \geq 2\right),$$

with  $E$  a unit exponential random variable independent of  $W_k$  and  $\{W_k, k \geq 2\}$  being jointly Gaussian with zero means and covariances

$$\mathbb{E}(W_i W_j) = \frac{\delta_{i-1} + \delta_{j-1} - \delta_{|i-j|}}{2\sqrt{\delta_{i-1}\delta_{j-1}}}.$$

The proof of (3.2) strongly relies on Berman's inequality. Hence, our first application extends the result of [17] to triangular arrays of randomly scaled Gaussian random variables. In the following we investigate the effect of a comonotonic random scaling considering a bounded  $S$  and thus  $\mathbf{S} = \mathbf{S1}$ .

**Theorem 3.1.** *Let  $\{X_{n,i}, i, n \geq 1\}$  be a triangular array of standard Gaussian random variables defined as above satisfying (3.1), being further independent of the iid non-negative random variables  $\{S_n, n \geq 1\}$  where  $S_1$  satisfies (2.2). If there exist positive integers  $r_n, l_n$  such that*

$$\lim_{n \rightarrow \infty} \frac{l_n}{r_n} = 0, \quad \lim_{n \rightarrow \infty} \frac{r_n}{n} = 0, \quad (3.4)$$

$$\lim_{n \rightarrow \infty} \frac{n^2}{r_n} c_n^{-\tau} \sum_{j=l_n}^n \frac{|\varrho_{n,j}|(1 + |\varrho_{n,j}|)^\tau}{\sqrt{1 - \varrho_{n,j}^2}} \exp\left(-\frac{c_n}{1 + |\varrho_{n,j}|}\right) = 0, \quad c_n := 2 \ln n - (2\tau + 1) \ln \ln n \quad (3.5)$$

and further

$$\lim_{m \rightarrow \infty} \limsup_{n \rightarrow \infty} \sum_{j=m}^{r_n} n^{-\frac{1 - \varrho_{n,j}}{1 + \varrho_{n,j}}} \frac{(\ln n)^{\frac{\tau(1 - \varrho_{n,j}) - \varrho_{n,j}}{1 + \varrho_{n,j}}}}{\sqrt{1 - \varrho_{n,j}^2}} = 0, \quad (3.6)$$

then for the maxima  $M_n = \max_{1 \leq i \leq n} S_n X_{n,i}$  the result in (3.2) holds with  $\vartheta$  defined as above and

$$a_n = (2 \ln n)^{-1/2}, \quad b_n = (2 \ln n)^{1/2} + (2 \ln n)^{-1/2} \left( \ln(c(2\pi)^{-1/2} \Gamma(1 + \tau)) - \frac{2\tau + 1}{2} (\ln \ln n + \ln 2) \right). \quad (3.7)$$

**Remark:** Using similar arguments as in the proof of Theorem 3.1, the findings of the recent contribution [6] can also be extended by considering randomly scaled Gaussian field on a lattice.

In our second application we consider scaled Gaussian random vectors where the scaling vector  $\mathbf{S}$  has independent components. Let  $\{\mathbf{X}_{n,k} = (X_{n,k}^{(1)}, X_{n,k}^{(2)}), 1 \leq k \leq n, n \geq 1\}$  be a triangular array of bivariate centered stationary Gaussian random vectors with unit-variance and correlation given by

$$\text{corr}\left(X_{n,k}^{(1)}, X_{n,k}^{(2)}\right) = \lambda_0(n), \quad \text{corr}\left(X_{n,k}^{(i)}, X_{n,l}^{(j)}\right) = \lambda_{ij}(|k - l|, n),$$

where  $1 \leq k \neq l \leq n$  and  $i, j \in \{1, 2\}$ . Further, let  $\{\hat{\mathbf{X}}_{n,k} = (\hat{X}_{n,k}^{(1)}, \hat{X}_{n,k}^{(2)}), 1 \leq k \leq n, n \geq 1\}$  denote the associated iid triangular array of  $\{\mathbf{X}_{n,k}\}$ , i.e.,  $\text{corr}(\hat{X}_{n,k}^{(1)}, \hat{X}_{n,k}^{(2)}) = \lambda_0(n)$  and  $\text{corr}(\hat{X}_{n,k}^{(i)}, \hat{X}_{n,l}^{(j)}) = 0$ , for  $1 \leq k \neq l \leq n$  and  $i, j \in \{1, 2\}$ . Let  $\{S_{n,k}, 1 \leq k \leq n, n \geq 1\}$  be iid random variables being independent of  $\{\mathbf{X}_{n,k}, 1 \leq k \leq n, n \geq 1\}$  and  $\{\hat{\mathbf{X}}_{n,k}, 1 \leq k \leq n, n \geq 1\}$ , respectively. Assume that the correlation  $\lambda_0(n)$  satisfies

$$\lim_{n \rightarrow \infty} \frac{b_n}{a_n} (1 - \lambda_0(n)) = 2\lambda^2 \quad \text{with } \lambda \in [0, \infty], \quad (3.8)$$

where

$$a_n = \frac{1}{1 - F(b_n)} \int_{b_n}^{\infty} (1 - F(x)) dx, \quad b_n = F^{-1}\left(1 - \frac{1}{n}\right),$$

with  $F^{-1}$  the inverse of the df  $F$  of  $S_{1,1}\hat{X}_{1,1}^{(1)}$ . It is well-known (see e.g., [10]) that

$$\lim_{n \rightarrow \infty} \sup_{x, y \in \mathbb{R}} \left| \mathbb{P}\left(\max_{1 \leq k \leq n} S_{n,k} \hat{X}_{n,k}^{(1)} \leq u_n(x), \max_{1 \leq k \leq n} S_{n,k} \hat{X}_{n,k}^{(2)} \leq u_n(y)\right) - H_\lambda(x, y) \right| = 0,$$

where  $u_n(z) = a_n z + b_n$ ,  $z \in \mathbb{R}$  and the Hüsler-Reiss distribution function  $H_\lambda$  is given by

$$H_\lambda(x, y) = \exp\left(-e^{-x} \Phi\left(\lambda + \frac{y-x}{2\lambda}\right) - e^{-y} \Phi\left(\lambda + \frac{x-y}{2\lambda}\right)\right), \quad (3.9)$$

with  $\Phi$  the distribution function of an  $N(0, 1)$  random variable.

In the following we are interested in the case that only a fraction of random vectors is observed. Assume therefore that  $\varepsilon_{n,k}$  is an indicator random variable of the event that the random vector  $\mathbf{X}_{n,k}$  is observed. Then  $\Xi_n = \sum_{k=1}^n \varepsilon_{n,k}$  is the number of observed random vectors from the set  $\{\mathbf{X}_{n,1}, \dots, \mathbf{X}_{n,n}\}$ .

We shall impose the following condition:

**Condition E.** The indicator random variables  $\varepsilon_{n,k}$  are independent of  $\mathbf{X}_{n,k}$  and  $S_{n,k}$ . Further, the convergence in probability

$$\frac{\Xi_n}{n} \xrightarrow{P} \eta, \quad n \rightarrow \infty$$

holds with  $\eta$  some random variable taking values in  $(0, 1]$  almost surely.

For notational simplicity we set

$$\mathbf{M}_n(\varepsilon_n) := \begin{cases} \max\{S_{n,k} \mathbf{X}_{n,k}, 1 \leq k \leq n, \varepsilon_{n,k} = 1\}, & \text{if } \sum_{k=1}^n \varepsilon_{n,k} \geq 1, \\ \inf\{\mathbf{x} | \mathbb{P}(S_{n,k} \mathbf{X}_{n,k} \leq \mathbf{x}) > \mathbf{0}\}, & \text{otherwise,} \end{cases}$$

$$\mathbf{m}_n(\varepsilon_n) := \begin{cases} \min\{S_{n,k} \mathbf{X}_{n,k}, 1 \leq k \leq n, \varepsilon_{n,k} = 1\}, & \text{if } \sum_{k=1}^n \varepsilon_{n,k} \geq 1, \\ \inf\{\mathbf{x} | \mathbb{P}(S_{n,k} \mathbf{X}_{n,k} \leq \mathbf{x}) > \mathbf{0}\}, & \text{otherwise} \end{cases}$$

and  $\mathbf{M}_n = \max\{S_{n,k} \mathbf{X}_{n,k}, 1 \leq k \leq n\}$ ,  $\mathbf{m}_n = \min\{S_{n,k} \mathbf{X}_{n,k}, 1 \leq k \leq n\}$ .

For  $S_{n,k} = 1, 1 \leq k \leq n$  almost surely, according to [12], under Condition **E** we have

$$\lim_{n \rightarrow \infty} \sup_{\substack{x_1, y_1 \in \mathbb{R} \\ x_1 \leq y_1}} \left| \mathbb{P}\left(M_n^{(1)}(\varepsilon_n) \leq u_n(x_1), M_n^{(1)} \leq u_n(y_1)\right) - \mathbb{E}\left(\Lambda^\eta(x_1) \Lambda^{1-\eta}(y_1)\right) \right| = 0,$$

where  $u_n(x) = a_n x + b_n$  with  $a_n$  and  $b_n$  defined in (3.3) and  $\Lambda(x) = \exp(-e^{-x})$ ,  $x \in \mathbb{R}$ , provided that  $\lim_{n \rightarrow \infty} \max_{l_n < k < n} \lambda_{11}(k, n) \ln n = 0$  with  $l_n = [n^{\hat{\beta}}]$ ,  $0 < \hat{\beta} < (1 - \hat{\sigma}) / (1 + \hat{\sigma})$  and  $\hat{\sigma} = \max_{1 \leq k < n, n \geq 1} |\lambda_{11}(k, n)|$ . Below we obtain a more general result for our 2-dimensional setup considering Weibull-type random scaling.

**Theorem 3.2.** Let  $\{(X_{n,k}^{(1)}, X_{n,k}^{(2)}), 1 \leq k \leq n, n \geq 1\}$  be a bivariate triangular array of standard Gaussian random vectors defined as above. Let  $\{S_{n,k}, 1 \leq k \leq n, n \geq 1\}$  be iid random variables being independent of  $\{(X_{n,k}^{(1)}, X_{n,k}^{(2)}), 1 \leq k \leq n, n \geq 1\}$ . Suppose that the correlation  $\lambda_0(n)$  satisfy (3.8) with  $\lambda \in (0, \infty)$  and condition **E**

holds. Let  $\beta$  be a constant satisfying  $0 < \beta < 2(1 + \sigma)^{-\frac{p}{2+p}} - 1$  with  $\sigma = \max_{\substack{1 \leq k < n, n \geq 1 \\ i, j \in \{1, 2\}}} |\lambda_{ij}(k, n)| < 1$ , and write  $\iota_n = \lceil n^\beta \rceil$ . If (2.7) holds and the covariance function satisfies

$$\lim_{n \rightarrow \infty} \max_{\substack{\iota_n \leq k < n \\ i, j \in \{1, 2\}}} \lambda_{ij}(k, n) \ln n = 0,$$

then we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup_{\substack{x_i, y_i \in \mathbb{R}, i \leq 4 \\ x_1 \leq x_3, x_2 \leq x_4, y_1 \leq y_3, y_2 \leq y_4}} & \left| \mathbb{P} \left( -u_n(y_1) < m_n^{(1)}(\boldsymbol{\varepsilon}_n) \leq M_n^{(1)}(\boldsymbol{\varepsilon}_n) \leq u_n(x_1), -u_n(y_2) < m_n^{(2)}(\boldsymbol{\varepsilon}_n) \leq M_n^{(2)}(\boldsymbol{\varepsilon}_n) \leq u_n(x_2), \right. \right. \\ & \left. \left. -u_n(y_3) < m_n^{(1)} \leq M_n^{(1)} \leq u_n(x_3), -u_n(y_4) < m_n^{(2)} \leq M_n^{(2)} \leq u_n(x_4) \right) \right. \\ & \left. - \mathbb{E} \left( H_\lambda^\eta(x_1, x_2) H_\lambda^\eta(y_1, y_2) H_\lambda^{1-\eta}(x_3, x_4) H_\lambda^{1-\eta}(y_3, y_4) \right) \right| = 0, \end{aligned}$$

where  $H_\lambda$  is defined in (3.9) and norming constants  $a_n$  and  $b_n$  satisfy

$$a_n = \frac{2+p}{2p} T^{-\frac{2+p}{2p}} (\ln n)^{\frac{2-p}{2p}}, \quad b_n = \left( \frac{\ln n}{T} \right)^{\frac{2+p}{2p}} + \frac{2+p}{2p} T^{-\frac{2+p}{2p}} (\ln n)^{\frac{2-p}{2p}} \left( \frac{\alpha}{p} \ln \ln n - \frac{\alpha}{p} \ln T + \ln \varpi_B \right) \quad (3.10)$$

with  $T = 2^{-1}Q^2 + LQ^{-p}$ ,  $\varpi_B = c_B(2+p)^{-\frac{1}{2}}Q^{-\alpha}$  and  $Q = (pL)^{1/(2+p)}$ .

## 4 Proofs

PROOF OF THEOREM 2.1 By the independence of  $\mathbf{S}$  and  $(\mathbf{X}, \mathbf{Y})$  and the generalised Berman's inequality (see Theorem 2.1 in [22] and Lemma 11.1.2 in [21]), if (2.1) holds, then

$$\begin{aligned} \Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v}) &= \mathbb{P}(-v_i < S_i X_i \leq u_i, 1 \leq i \leq n) - \mathbb{P}(-v_i < S_i Y_i \leq u_i, 1 \leq i \leq n) \\ &= \int_{[0,1]^n} \left( \mathbb{P} \left( -\frac{v_i}{s_i} < X_i \leq \frac{u_i}{s_i}, 1 \leq i \leq n \right) - \mathbb{P} \left( -\frac{v_i}{s_i} < Y_i \leq \frac{u_i}{s_i}, 1 \leq i \leq n \right) \right) dG(s_1) \cdots dG(s_n) \\ &\leq \frac{2}{\pi} \int_{[0,1]^n} \sum_{1 \leq i < j \leq n} A_{ij} \exp \left( -\frac{(w/s_i)^2 + (w/s_j)^2}{2(1 + \rho_{ij})} \right) dG(s_1) \cdots dG(s_n) \\ &= \frac{2}{\pi} \sum_{1 \leq i < j \leq n} A_{ij} \int_0^1 \int_0^1 \exp \left( -\frac{(w/s)^2 + (w/t)^2}{2(1 + \rho_{ij})} \right) dG(s) dG(t), \end{aligned}$$

where  $\rho_{ij}$  and  $A_{ij}$  are defined in (1.1) and  $w = \min_{1 \leq i \leq n} \min(|u_i|, |v_i|)$ . Note that for  $1 \leq i, j \leq n$ ,  $\varepsilon > 0$

$$\begin{aligned} & \int_0^1 \exp \left( -\frac{1}{2(1 + \rho_{ij})} \left( \frac{w}{s} \right)^2 \right) dG(s) \\ & \sim \int_{\frac{1}{\varepsilon+1}}^1 \exp \left( -\frac{1}{2(1 + \rho_{ij})} \left( \frac{w}{s} \right)^2 \right) dG(s) \\ & = \int_w^{w(1+\varepsilon)} \mathbb{P} \left( S > \frac{w}{s} \right) d \left( 1 - \exp \left( -\frac{1}{2(1 + \rho_{ij})} s^2 \right) \right) \\ & = \int_0^{\frac{\varepsilon}{1+\rho_{ij}} w^2} \mathbb{P} \left( S > \frac{w}{w + (1 + \rho_{ij}) w^{-1} t} \right) \left( 1 + \frac{1 + \rho_{ij}}{w^2} t \right) \exp \left( -\frac{1}{2(1 + \rho_{ij})} (w^2 + 2(1 + \rho_{ij}) t + (1 + \rho_{ij})^2 w^{-2} t^2) \right) dt \\ & \sim \int_0^{\frac{\varepsilon}{1+\rho_{ij}} w^2} \mathbb{P} \left( S > 1 - \frac{1 + \rho_{ij}}{w^2} t \right) \exp \left( -t - \frac{w^2}{2(1 + \rho_{ij})} \right) dt \\ & \leq c_A (1 + \rho_{ij})^\tau w^{-2\tau} \exp \left( -\frac{w^2}{2(1 + \rho_{ij})} \right) \int_0^{\frac{\varepsilon}{1+\rho_{ij}} w^2} t^\tau \exp(-t) dt \\ & \sim c_A \Gamma(\tau + 1) (1 + \rho_{ij})^\tau w^{-2\tau} \exp \left( -\frac{w^2}{2(1 + \rho_{ij})} \right), \quad w \rightarrow \infty. \end{aligned}$$

Consequently, for any  $\epsilon > 0$  and all large  $u_i, v_i, i \leq n$

$$\Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v}) \leq \frac{2}{\pi} (\Gamma(\tau + 1))^2 (c_A^2 + \epsilon) w^{-4\tau} \sum_{1 \leq i < j \leq n} A_{ij} (1 + \rho_{ij})^{2\tau} \exp\left(-\frac{w^2}{1 + \rho_{ij}}\right).$$

With similar arguments as above we have

$$\begin{aligned} \Delta_{S1}(\mathbf{u}, \mathbf{v}) &= \int_0^1 \left( \mathbb{P}\left(-\frac{v_i}{s} < X_i \leq \frac{u_i}{s}, 1 \leq i \leq n\right) - \mathbb{P}\left(-\frac{v_i}{s} < Y_i \leq \frac{u_i}{s}, 1 \leq i \leq n\right) \right) dG(s) \\ &\leq \frac{2}{\pi} \sum_{1 \leq i < j \leq n} A_{ij} \int_0^1 \exp\left(-\frac{(w/s)^2}{1 + \rho_{ij}}\right) dG(s) \\ &\leq \frac{2^{1-\tau}}{\pi} \Gamma(\tau + 1) (c_A + \epsilon) w^{-2\tau} \sum_{1 \leq i < j \leq n} A_{ij} (1 + \rho_{ij})^\tau \exp\left(-\frac{w^2}{1 + \rho_{ij}}\right), \end{aligned}$$

hence the claim follows.  $\square$

PROOF OF THEOREM 2.3 According to the independence of the scaling factors with the Gaussian random variables and the generalised Berman's inequality (see Theorem 2.1 in [22] and Lemma 11.1.2 in [21]) again if (2.7) holds, then we have

$$\begin{aligned} \Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v}) &= \int_{[0, \infty]^n} \left( \mathbb{P}\left(-\frac{v_i}{s_i} < X_i \leq \frac{u_i}{s_i}, 1 \leq i \leq n\right) - \mathbb{P}\left(-\frac{v_i}{s_i} < Y_i \leq \frac{u_i}{s_i}, 1 \leq i \leq n\right) \right) dG(s_1) \cdots dG(s_n) \\ &\leq \frac{2}{\pi} \int_{[0, \infty]^n} \sum_{1 \leq i < j \leq n} A_{ij} \exp\left(-\frac{(w/s_i)^2 + (w/s_j)^2}{2(1 + \rho_{ij})}\right) dG(s_1) \cdots dG(s_n) \\ &= \frac{2}{\pi} \sum_{1 \leq i < j \leq n} A_{ij} \int_0^\infty \int_0^\infty \exp\left(-\frac{(w/s)^2 + (w/t)^2}{2(1 + \rho_{ij})}\right) dG(s) dG(t), \end{aligned}$$

where  $\rho_{ij}$  and  $A_{ij}$  are defined in (1.1). Note that for  $1 \leq i, j \leq n$  and some positive constants  $c_1, c_2$ , using similar arguments as in the proof of Theorem 2.1 in [16], we have

$$\begin{aligned} &\int_0^\infty \exp\left(-\frac{1}{2(1 + \rho_{ij})} \left(\frac{w}{s}\right)^2\right) dG(s) \\ &\sim \int_{c_1 w^{\frac{2}{p+2}}}^{c_2 w^{\frac{2}{p+2}}} \exp\left(-\frac{1}{2(1 + \rho_{ij})} \left(\frac{w}{s}\right)^2\right) dG(s) \\ &\sim c_B L p \int_{c_1 w^{\frac{2}{p+2}}}^{c_2 w^{\frac{2}{p+2}}} s^{\alpha+p-1} \exp\left(-L s^p - \frac{1}{2(1 + \rho_{ij})} \left(\frac{w}{s}\right)^2\right) ds \\ &= c_B L p \left(\frac{w^2}{L p (1 + \rho_{ij})}\right)^{\frac{\alpha+p}{p+2}} \int_{c_1 (L p (1 + \rho_{ij}))^{\frac{1}{p+2}}}^{c_2 (L p (1 + \rho_{ij}))^{\frac{1}{p+2}}} t^{\alpha+p-1} \exp\left(-L p \left(\frac{w^2}{L p (1 + \rho_{ij})}\right)^{\frac{p}{p+2}} (p^{-1} t^p + 2^{-1} t^{-2})\right) dt \\ &\sim \sqrt{\frac{2\pi}{p+2}} c_B (L p)^{\frac{1-\alpha}{p+2}} (1 + \rho_{ij})^{\frac{-2\alpha-p}{2(p+2)}} w^{\frac{2\alpha+p}{p+2}} \exp\left(- (1 + \rho_{ij})^{-\frac{p}{p+2}} (L^{\frac{2}{p+2}} p^{-\frac{p}{p+2}} + (L p)^{\frac{2}{p+2}} 2^{-1}) w^{\frac{2p}{p+2}}\right), \quad w \rightarrow \infty. \end{aligned}$$

Hence for  $\epsilon > 0$  we have

$$\Delta_{\mathbf{S}}(\mathbf{u}, \mathbf{v}) \leq \frac{4(c_B^2 + \epsilon) (L p)^{\frac{2(1-\alpha)}{p+2}}}{p+2} w^{\frac{4\alpha+2p}{2+p}} \sum_{1 \leq i < j \leq n} A_{ij} (1 + \rho_{ij})^{\frac{-2\alpha-p}{p+2}} \exp\left(-2(1 + \rho_{ij})^{-\frac{p}{2+p}} T w^{\frac{2p}{2+p}}\right),$$

where  $T = L^{\frac{2}{p+2}} p^{-\frac{p}{p+2}} + (L p)^{\frac{2}{p+2}} 2^{-1}$ . Proceeding as above

$$\begin{aligned} \Delta_{S1}(\mathbf{u}, \mathbf{v}) &= \int_0^\infty \left( \mathbb{P}\left(-\frac{v_i}{s} < X_i \leq \frac{u_i}{s}, 1 \leq i \leq n\right) - \mathbb{P}\left(-\frac{v_i}{s} < Y_i \leq \frac{u_i}{s}, 1 \leq i \leq n\right) \right) dG(s) \\ &\leq \frac{2}{\pi} \sum_{1 \leq i < j \leq n} A_{ij} \int_0^\infty \exp\left(-\frac{(w/s)^2}{1 + \rho_{ij}}\right) dG(s) \\ &\leq 2^{\frac{3+2p+\alpha}{2+p}} \pi^{-\frac{1}{2}} (c_B + \epsilon) (L p)^{\frac{1-\alpha}{p+2}} (p+2)^{-\frac{1}{2}} w^{\frac{2\alpha+p}{2+p}} \end{aligned}$$

$$\times \sum_{1 \leq i < j \leq n} A_{ij} (1 + \rho_{ij})^{\frac{-2\alpha - p}{2(p+2)}} \exp\left(- (2(1 + \rho_{ij})^{-1})^{\frac{p}{2+p}} T w^{\frac{2p}{2+p}}\right),$$

hence the proof is complete.  $\square$

**Lemma 4.1.** *Under the conditions of Theorem 3.1, for any bounded set  $K \subset \{2, 3, \dots\}$  we have*

$$\lim_{n \rightarrow \infty} \mathbb{P}(S_n X_{n,k} \leq u_n, k \in K | S_n X_{n,1} > u_n) = \mathbb{P}\left(E/2 + \sqrt{\delta_{k-1}} W_k \leq \delta_{k-1}, k \in K\right),$$

where  $E$  is a standard exponential random variable independent of  $\{W_k, k \in K\}$  and the  $W_k$  have a jointly Gaussian distribution with mean zero and

$$\mathbb{E}(W_i W_j) = \frac{\delta_{i-1} + \delta_{j-1} - \delta_{|i-j|}}{2\sqrt{\delta_{i-1}\delta_{j-1}}}, \quad i, j \in K.$$

PROOF OF LEMMA 4.1 A centered Gaussian random vector  $\mathbf{X}_n = (X_{n,k}, k \in K \cup \{1\})^\top$  with covariance matrix  $B_n^\top B_n = (\varrho_{n,|i-j|})_{i,j \in K \cup \{1\}}$  has stochastic representation

$$(X_{n,k}, k \in K \cup \{1\})^\top \stackrel{d}{=} R B_n^\top \mathbf{U}_{m+1},$$

where  $m$  is the cardinality of set  $K$ ,  $R$  is a positive random variable such that  $R^2$  is chi-squared distributed with  $m + 1$  degrees of freedom and independent of  $\mathbf{U}_{m+1}$  which is a random vector uniformly distributed on the unit sphere of  $\mathbb{R}^{m+1}$ . Since  $S_n$  is independent of  $X_{n,k}$  using Corollary 5 in [3] we have (set  $t_n(y) := u_n + y/u_n$ )

$$(S_n X_{n,k}, k \in K | S_n X_{n,1} = t_n(y))^\top \stackrel{d}{=} R_{m,y} \hat{B}_n^\top \mathbf{U}_m + t_n(y) \Sigma_{12},$$

where  $\Sigma_{12} = (\varrho_{n,k-1}, k \in K)^\top$ ,  $\hat{B}_n^\top \hat{B}_n = (\varrho_{n,|i-j|} - \varrho_{n,i-1}\varrho_{n,j-1})_{i,j \in K}$  and  $R_{m,y}$  is a positive random variable independent of  $\mathbf{U}_m$  with distribution function  $F_{m,y}$  defined by

$$F_{m,y}(x) = \frac{\int_{t_n(y)}^{((t_n(y))^2 + x^2)^{1/2}} (s^2 - (t_n(y))^2)^{\frac{m}{2}-1} s^{1-m} dF_1(s)}{\int_{t_n(y)}^{\infty} (s^2 - (t_n(y))^2)^{\frac{m}{2}-1} s^{1-m} dF_1(s)}, \quad x > 0,$$

with  $F_1$  the distribution function of  $S_n R$ . According to Theorem 3.1 in [11]  $F_1$  in the Gumbel max-domain of attraction and

$$\lim_{n \rightarrow \infty} \frac{\mathbb{P}(S_n X_{n,1} > t_n(y))}{\mathbb{P}(S_n X_{n,1} > u_n)} = e^{-y}, \quad \forall y \in \mathbb{R}. \quad (4.1)$$

Hence, by Theorem 3.1 in [8]

$$\begin{aligned} p_{n,y} &:= \mathbb{P}(S_n X_{n,k} \leq u_n, k \in K | S_n X_{n,1} = t_n(y)) \\ &= \mathbb{P}\left(\frac{u_n(1 - \varrho_{n,k-1}^2)^{1/2}}{2} Z_{n,k} + \frac{\varrho_{n,k-1}}{2} y \leq \frac{u_n^2(1 - \varrho_{n,k-1})}{2}, k \in K\right) \\ &\rightarrow \mathbb{P}\left(\sqrt{\delta_{k-1}} W_k + \frac{y}{2} \leq \delta_{k-1}, k \in K\right), \quad n \rightarrow \infty \end{aligned} \quad (4.2)$$

uniformly on compact sets of  $y$ , where

$$(Z_{n,k}, k \in K)^\top \stackrel{d}{=} R_{m,y} \tilde{B}_n^\top \mathbf{U}_m, \quad \text{with } \tilde{B}_n^\top \tilde{B}_n = \left( \frac{\varrho_{n,|i-j|} - \varrho_{n,i-1}\varrho_{n,j-1}}{\sqrt{(1 - \varrho_{n,i-1}^2)(1 - \varrho_{n,j-1}^2)}} \right)_{i,j \in K}$$

and  $\{W_k, k \in K\}$  being jointly Gaussian with zero means and covariances

$$\mathbb{E}(W_i W_j) = \frac{\delta_{i-1} + \delta_{j-1} - \delta_{|i-j|}}{2\sqrt{\delta_{i-1}\delta_{j-1}}}, \quad i, j \in K.$$

Since further

$$\mathbb{P}(S_n X_{n,k} \leq u_n, k \in K | S_n X_{n,1} > u_n) = \int_0^\infty p_{n,y} d \frac{\mathbb{P}(S_n X_{n,1} \leq t_n(y))}{\mathbb{P}(S_n X_{n,1} > u_n)}$$

the proof is established by applying Lemma 4.4 in [8] (recall (4.1) and (4.2)).  $\square$

PROOF OF THEOREM 3.1 According to (2.4), if  $1 \leq k_1 < \dots < k_s \leq n$  and  $k = \min_{1 \leq i < s} (k_{i+1} - k_i)$  then the joint distribution function  $F_{k_1, \dots, k_s}$  of  $S_n X_{n,k_1}, \dots, S_n X_{n,k_s}$  satisfies

$$\left| F_{k_1, \dots, k_s}(u_n) - \prod_{i=1}^s \mathbb{P}(S_n X_{n,k_i} \leq u_n) \right| \leq \mathcal{Q} u_n^{-2\tau} n \sum_{i=k}^n \frac{|\varrho_{n,i}| (1 + |\varrho_{n,i}|)^\tau}{\sqrt{1 - \varrho_{n,i}^2}} \exp\left(-\frac{u_n^2}{1 + |\varrho_{n,i}|}\right).$$

Suppose now that  $1 \leq i_1 < \dots < i_p < j_1 < \dots < j_{p'} \leq n$  and  $j_1 - i_p \geq l_n$ . Identifying  $\{k_1, \dots, k_s\}$  in turn with  $\{i_1, \dots, i_p, j_1, \dots, j_{p'}\}$ ,  $\{i_1, \dots, i_p\}$  and  $\{j_1, \dots, j_{p'}\}$ , we thus have

$$|F_{i_1, \dots, i_p, j_1, \dots, j_{p'}}(u_n) - F_{i_1, \dots, i_p}(u_n) F_{j_1, \dots, j_{p'}}(u_n)| \leq 3\mathcal{Q} u_n^{-2\tau} n \sum_{i=l_n}^n \frac{|\varrho_{n,i}| (1 + |\varrho_{n,i}|)^\tau}{\sqrt{1 - \varrho_{n,i}^2}} \exp\left(-\frac{u_n^2}{1 + |\varrho_{n,i}|}\right).$$

By Example 1 in [9] and Table 3.4.4 in [5] we have

$$\lim_{n \rightarrow \infty} n \mathbb{P}(S_n X_{n,1} \geq u_n(x)) = e^{-x}, \quad \forall x \in \mathbb{R},$$

where  $u_n(x) = a_n x + b_n$  with  $a_n$  and  $b_n$  defined in (3.7). Consequently, as  $n \rightarrow \infty$

$$u_n^2(x) = 2 \ln n - (2\tau + 1) \ln \ln n + O(1).$$

Hence, in view of (3.4) and (3.5), Theorem 2.1 in [24] implies

$$\lim_{n \rightarrow \infty} \left[ \mathbb{P}\left(\max_{1 \leq i \leq n} S_n X_{n,i} \leq u_n(x)\right) - \exp\left(-n \mathbb{P}(S_n X_{n,1} > u_n(x)) \mathbb{P}\left(\bigcap_{i=2}^{r_n} \{S_n X_{n,i} \leq u_n(x)\} | S_n X_{n,1} > u_n(x)\right)\right)\right] = 0.$$

Note that for  $m \leq j \leq r_n$  we have

$$\mathbb{P}\left(W > u_n \sqrt{\frac{1 - \varrho_{n,j}}{1 + \varrho_{n,j}}} - \frac{y}{u_n} \frac{\varrho_{n,j}}{\sqrt{1 - \varrho_{n,j}^2}}\right) \leq \mathcal{Q} n^{-\frac{1 - \varrho_{n,j}}{1 + \varrho_{n,j}}} \frac{(\ln n)^{\frac{\tau(1 - \varrho_{n,j}) - \varrho_{n,j}}{1 + \varrho_{n,j}}}}{\sqrt{1 - \varrho_{n,j}^2}},$$

where  $W$  is a  $N(0, 1)$  random variable. The claim can then be established by using similar arguments as in the proof of Theorem 2.1 in [17] making further use of (3.6) and Lemma 4.1.  $\square$

Next, for some index sets  $I_n \subset N$  we define

$$\widehat{\mathbf{M}}(I_n, \varepsilon_n) := \begin{cases} \max\{S_{n,k} \widehat{\mathbf{X}}_{n,k}, k \in I_n, \varepsilon_{n,k} = 1\}, & \text{if } \sum_{k \in I_n} \varepsilon_{n,k} \geq 1; \\ \inf\{\mathbf{x} | \mathbb{P}(S_{n,k} \widehat{\mathbf{X}}_{n,k} \leq \mathbf{x}) > \mathbf{0}\}, & \text{otherwise,} \end{cases}$$

$$\widehat{\mathbf{m}}(I_n, \varepsilon_n) := \begin{cases} \min\{S_{n,k} \widehat{\mathbf{X}}_{n,k}, k \in I_n, \varepsilon_{n,k} = 1\}, & \text{if } \sum_{k \in I_n} \varepsilon_{n,k} \geq 1; \\ \inf\{\mathbf{x} | \mathbb{P}(S_{n,k} \widehat{\mathbf{X}}_{n,k} \leq \mathbf{x}) > \mathbf{0}\}, & \text{otherwise.} \end{cases}$$

For simplicity, we write  $\widehat{\mathbf{M}}_n(\varepsilon_n) = \widehat{\mathbf{M}}(\{1, 2, \dots, n\}, \varepsilon_n)$ ,  $\widehat{\mathbf{M}}(I_n) = \max\{S_{n,k} \widehat{\mathbf{X}}_{n,k}, k \in I_n\}$ ,  $\widehat{\mathbf{M}}_n = \max\{S_{n,k} \widehat{\mathbf{X}}_{n,k}, 1 \leq k \leq n\}$ . Similarly we also define  $\widehat{\mathbf{m}}_n(\varepsilon_n)$ ,  $\widehat{\mathbf{m}}(I_n)$ ,  $\widehat{\mathbf{m}}_n$ .

**Lemma 4.2.** *Let  $\{(\widehat{X}_{n,i}^{(1)}, \widehat{X}_{n,i}^{(2)}), 1 \leq i \leq n, n \geq 1\}$  be a triangular array of centered stationary Gaussian random vectors defined as above with the correlation  $\lambda_0(n)$  satisfying (3.8) with  $\lambda \in (0, \infty)$ . Further let  $\{S_{n,k}, 1 \leq k \leq n, n \geq 1\}$  be iid random variables being independent of  $\{(\widehat{X}_{n,i}^{(1)}, \widehat{X}_{n,i}^{(2)}), 1 \leq i \leq n, n \geq 1\}$  and satisfying (2.7). Then we have*

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(-u_n(y_1) < \widehat{m}_n^{(1)} \leq \widehat{M}_n^{(1)} \leq u_n(x_1), -u_n(y_2) < \widehat{m}_n^{(2)} \leq \widehat{M}_n^{(2)} \leq u_n(x_2)\right) = H_\lambda(x_1, x_2) H_\lambda(y_1, y_2).$$

PROOF OF LEMMA 4.2 Our proof is similar to that of Theorem 2.1 in [14]. For any integer  $n$  we may write

$$n(1 - P(n, x_1, x_2, y_1, y_2)) = nP_1(n, x_1, x_2) + nP_2(n, y_1, y_2) - nP_3(n, x_1, y_2) - nP_4(n, y_1, x_2),$$

where

$$\begin{aligned} P(n, x_1, x_2, y_1, y_2) &:= \mathbb{P}\left(-u_n(y_1) < S_{n,1}\hat{X}_{n,1}^{(1)} \leq u_n(x_1), -u_n(y_2) < S_{n,1}\hat{X}_{n,1}^{(2)} \leq u_n(x_2)\right), \\ P_1(n, x_1, x_2) &:= \mathbb{P}\left(S_{n,1}\hat{X}_{n,1}^{(1)} > u_n(x_1)\right) + \mathbb{P}\left(S_{n,1}\hat{X}_{n,1}^{(2)} > u_n(x_2)\right) - \mathbb{P}\left(S_{n,1}\hat{X}_{n,1}^{(1)} > u_n(x_1), S_{n,1}\hat{X}_{n,1}^{(2)} > u_n(x_2)\right), \\ P_2(n, y_1, y_2) &:= \mathbb{P}\left(S_{n,1}\hat{X}_{n,1}^{(1)} \leq -u_n(y_1)\right) + \mathbb{P}\left(S_{n,1}\hat{X}_{n,1}^{(2)} \leq -u_n(y_2)\right) - \mathbb{P}\left(S_{n,1}\hat{X}_{n,1}^{(1)} \leq -u_n(y_1), S_{n,1}\hat{X}_{n,1}^{(2)} \leq -u_n(y_2)\right), \\ P_3(n, x_1, y_2) &:= \mathbb{P}\left(S_{n,1}\hat{X}_{n,1}^{(1)} > u_n(x_1), S_{n,1}\hat{X}_{n,1}^{(2)} \leq -u_n(y_2)\right), \\ P_4(n, y_1, x_2) &:= \mathbb{P}\left(S_{n,1}\hat{X}_{n,1}^{(1)} \leq -u_n(y_1), S_{n,1}\hat{X}_{n,1}^{(2)} > u_n(x_2)\right). \end{aligned}$$

The random vector  $(\hat{X}_{n,1}^{(1)}, \hat{X}_{n,1}^{(2)})$  has the following stochastic representation

$$(\hat{X}_{n,1}^{(1)}, \hat{X}_{n,1}^{(2)}) \stackrel{d}{=} (R \cos \theta, R \cos(\theta - \psi_n)),$$

where  $R$  is a positive random variable being independent of the random variable  $\theta$  which is uniformly distributed in  $(-\pi, \pi)$  and  $\psi_n = \arccos(\lambda_0(n))$ . If  $S_{n,1}$  satisfy (2.7) and is independent of  $(\hat{X}_{n,1}^{(1)}, \hat{X}_{n,1}^{(2)})$ , using Laplace approximation (see e.g., [16]) we have that the distribution function of  $S_{n,1}R$  is in the max-domain of attraction of the Gumbel distribution. Hence, according to Remark 2.2 in [13] we have

$$\lim_{n \rightarrow \infty} n\mathbb{P}\left(S_{n,1}\hat{X}_{n,1}^{(1)} > u_n(x)\right) = e^{-x}, \quad x \in \mathbb{R}, \quad (4.3)$$

where  $u_n(x) = a_n x + b_n$  with  $a_n$  and  $b_n$  defined in (3.10). Moreover, by Theorem 2.1 in [7]

$$\lim_{n \rightarrow \infty} nP_1(n, x_1, x_2) = \Phi\left(\lambda + \frac{x_1 - x_2}{2\lambda}\right) \exp(-x_2) + \Phi\left(\lambda + \frac{x_2 - x_1}{2\lambda}\right) \exp(-x_1) =: D(x_1, x_2)$$

and since  $(-S_{n,1}\hat{X}_{n,1}^{(1)}, -S_{n,1}\hat{X}_{n,1}^{(2)}) \stackrel{d}{=} (S_{n,1}\hat{X}_{n,1}^{(1)}, S_{n,1}\hat{X}_{n,1}^{(2)})$

$$\lim_{n \rightarrow \infty} nP_2(n, y_1, y_2) = D(y_1, y_2).$$

Since  $\lim_{n \rightarrow \infty} \lambda_0(n) = 1$ ,  $\lim_{n \rightarrow \infty} \psi_n = 0$  implying

$$\begin{aligned} &\lim_{n \rightarrow \infty} nP_3(n, x_1, y_2) \\ &= \lim_{n \rightarrow \infty} n\mathbb{P}\left(S_{n,1}R \cos(\theta) > u_n(x_1), S_{n,1}R \cos(\theta - \psi_n) \leq -u_n(y_1)\right) \\ &= \lim_{n \rightarrow \infty} n\mathbb{P}\left(S_{n,1}R \cos(\theta) > u_n(x_1), S_{n,1}R \cos(\theta - \psi_n) \leq -u_n(y_1), \cos(\theta) > 0, \cos(\theta - \psi_n) < 0\right) \\ &= \lim_{n \rightarrow \infty} n\mathbb{P}\left(S_{n,1}R \cos(\theta) > u_n(x_1), S_{n,1}R \cos(\theta - \psi_n) \leq -u_n(y_1), \max\left(-\frac{\pi}{2}, -\pi + \psi_n\right) < \theta < -\frac{\pi}{2} + \psi_n\right) \\ &= 0. \end{aligned}$$

Similarly, we have  $\lim_{n \rightarrow \infty} nP_4(n, y_1, x_2) = 0$ . Hence for all  $x_1, x_2, y_1, y_2 \in \mathbb{R}$

$$\begin{aligned} &\lim_{n \rightarrow \infty} \mathbb{P}\left(-u_n(y_1) < \hat{m}_n^{(1)} \leq \widehat{M}_n^{(1)} \leq u_n(x_1), -u_n(y_2) < \hat{m}_n^{(2)} \leq \widehat{M}_n^{(2)} \leq u_n(x_2)\right) \\ &= \lim_{n \rightarrow \infty} (P(n, x_1, x_2, y_1, y_2))^n \\ &= \lim_{n \rightarrow \infty} (1 - (1 - P(n, x_1, x_2, y_1, y_2)))^n \\ &= \lim_{n \rightarrow \infty} \left(1 - \frac{D(x_1, x_2) + D(y_1, y_2)}{n} + o\left(\frac{1}{n}\right)\right)^n \\ &= \exp(-D(x_1, x_2) - D(y_1, y_2)) = H_\lambda(x_1, x_2)H_\lambda(y_1, y_2), \end{aligned}$$

hence the proof is complete.  $\square$

**Lemma 4.3.** *Under the conditions of Lemma 4.2, if the indicator random variables  $\varepsilon_n = \{\varepsilon_{n,i}, 1 \leq i \leq n\}$  are independent of both  $\{(\hat{X}_{n,i}^{(1)}, \hat{X}_{n,i}^{(2)}), 1 \leq i \leq n\}$  and  $\{S_{n,i}, 1 \leq i \leq n\}$  and satisfying condition **E**, then*

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup_{\substack{x_i, y_i \in \mathbb{R}, i = \{1, 2, 3, 4\} \\ x_1 \leq x_3, x_2 \leq x_4, y_1 \leq y_3, y_2 \leq y_4}} & \left| \mathbb{P} \left( -u_n(y_1) < \hat{m}_n^{(1)}(\varepsilon_n) \leq \widehat{M}_n^{(1)}(\varepsilon_n) \leq u_n(x_1), -u_n(y_2) < \hat{m}_n^{(2)}(\varepsilon_n) \leq \widehat{M}_n^{(2)}(\varepsilon_n) \leq u_n(x_2), \right. \right. \\ & \left. \left. -u_n(y_3) < \hat{m}_n^{(1)} \leq \widehat{M}_n^{(1)} \leq u_n(x_3), -u_n(y_4) < \hat{m}_n^{(2)} \leq \widehat{M}_n^{(2)} \leq u_n(x_4) \right) \right. \\ & \left. - \mathbb{E} \left( H_\lambda^\eta(x_1, x_2) H_\lambda^\eta(y_1, y_2) H_\lambda^{1-\eta}(x_3, x_4) H_\lambda^{1-\eta}(y_3, y_4) \right) \right| = 0. \end{aligned}$$

PROOF OF LEMMA 4.3 Using similar arguments as for the derivation of [19], let  $K_s = \{j : (s-1)\nu + 1 \leq j \leq s\nu\}, 1 \leq s \leq l, \nu = \lfloor \frac{n}{l} \rfloor, \mathbf{x} = (x_1, x_2, x_3, x_4), \mathbf{y} = (y_1, y_2, y_3, y_4)$  and  $\beta_n = \{\beta_{n,k}, 1 \leq k \leq n\}$  be a nonrandom triangular array consisting of 0's and 1's. For some random variable  $\eta$  such that  $0 \leq \eta \leq 1$  a.s., write

$$B_{\mu,l} = \left\{ \omega : \eta(\omega) \in \begin{cases} [0, \frac{1}{2^l}], & \mu = 0, \\ (\frac{\mu}{2^l}, \frac{\mu+1}{2^l}], & 0 < \mu \leq 2^l - 1 \end{cases} \right\},$$

$$B_{\mu,l,\beta_n} = \{\omega : \varepsilon_{n,k}(\omega) = \beta_{n,k}, 1 \leq k \leq n\} \cap B_{\mu,l}.$$

Set

$$\begin{aligned} & P(K_s, \beta_n, \mathbf{x}, \mathbf{y}) \\ = & \mathbb{P} \left( -u_n(y_1) < \hat{m}^{(1)}(K_s, \beta_n) \leq \widehat{M}^{(1)}(K_s, \beta_n) \leq u_n(x_1), -u_n(y_2) < \hat{m}^{(2)}(K_s, \beta_n) \leq \widehat{M}^{(2)}(K_s, \beta_n) \leq u_n(x_2), \right. \\ & \left. -u_n(y_3) < \hat{m}^{(1)}(K_s) \leq \widehat{M}^{(1)}(K_s) \leq u_n(x_3), -u_n(y_4) < \hat{m}^{(2)}(K_s) \leq \widehat{M}^{(2)}(K_s) \leq u_n(x_4) \right) \end{aligned}$$

and

$$\begin{aligned} & P(n, \beta_n, \mathbf{x}, \mathbf{y}) \\ = & \mathbb{P} \left( -u_n(y_1) < \hat{m}_n^{(1)}(\beta_n) \leq \widehat{M}_n^{(1)}(\beta_n) \leq u_n(x_1), -u_n(y_2) < \hat{m}_n^{(2)}(\beta_n) \leq \widehat{M}_n^{(2)}(\beta_n) \leq u_n(x_2), \right. \\ & \left. -u_n(y_3) < \hat{m}_n^{(1)} \leq \widehat{M}_n^{(1)} \leq u_n(x_3), -u_n(y_4) < \hat{m}_n^{(2)} \leq \widehat{M}_n^{(2)} \leq u_n(x_4) \right). \end{aligned}$$

Using similar arguments as in the proof of Lemma 3.3 in [25] for  $n$  large we can choose a positive integer  $\tilde{\nu}_n$  such that  $l < \tilde{\nu}_n < \nu$  and  $\tilde{\nu}_n = o(n)$ , by (4.3) we have

$$\begin{aligned} & \left| P(n, \beta_n, \mathbf{x}, \mathbf{y}) - \prod_{s=1}^l P(K_s, \beta_n, \mathbf{x}, \mathbf{y}) \right| \\ \leq & (4l+2)\tilde{\nu}_n \left( \mathbb{P} \left( S_{n,1} \hat{X}_{n,1}^{(1)} \leq -u_n(y_1) \right) + \mathbb{P} \left( S_{n,1} \hat{X}_{n,1}^{(1)} > u_n(x_1) \right) \right. \\ & \left. + \mathbb{P} \left( S_{n,1} \hat{X}_{n,1}^{(2)} \leq -u_n(y_2) \right) + \mathbb{P} \left( S_{n,1} \hat{X}_{n,1}^{(2)} > u_n(x_2) \right) \right) \\ \rightarrow & 0, \quad n \rightarrow \infty. \end{aligned} \tag{4.4}$$

Note that

$$\begin{aligned} & 1 - \frac{\nu\mu}{2^l} \Sigma_1 - \nu \left( 1 - \frac{\mu}{2^l} \right) \Sigma_2 + \left( \frac{\sum_{j \in K_s} \beta_{nj}}{\nu} - \frac{\mu}{2^l} \right) \nu (\Sigma_2 - \Sigma_1) \\ \leq & P(K_s, \beta_n, \mathbf{x}, \mathbf{y}) \\ \leq & 1 - \frac{\nu\mu}{2^l} \Sigma_1 - \nu \left( 1 - \frac{\mu}{2^l} \right) \Sigma_2 + \left( \frac{\sum_{j \in K_s} \beta_{nj}}{\nu} - \frac{\mu}{2^l} \right) \nu (\Sigma_2 - \Sigma_1) + \nu \Sigma_3, \end{aligned}$$

where

$$\begin{aligned} \Sigma_1 &= P_1(n, x_1, x_2) + P_2(n, y_1, y_2) - P_3(n, x_1, y_2) - P_4(n, y_1, x_2), \\ \Sigma_2 &= P_1(n, x_3, x_4) + P_2(n, y_3, y_4) - P_3(n, x_3, y_4) - P_4(n, y_3, x_4) \end{aligned}$$

with  $P_i(n, z_1, z_2)$ 's defined in the proof of Lemma 4.2 and

$$\begin{aligned} \Sigma_3 &= \sum_{i,j=\{1,2\}} \sum_{t=2}^{\nu} \left( \mathbb{P} \left( S_{n,1} \hat{X}_{n,(s-1)\nu+1}^{(i)} > u_n(x_i), S_{n,1} \hat{X}_{n,(s-1)\nu+t}^{(j)} > u_n(x_j) \right) \right. \\ &\quad + \mathbb{P} \left( S_{n,1} \hat{X}_{n,(s-1)\nu+1}^{(i)} > u_n(x_i), S_{n,1} \hat{X}_{n,(s-1)\nu+t}^{(j)} \leq -u_n(y_j) \right) \\ &\quad + \mathbb{P} \left( S_{n,1} \hat{X}_{n,(s-1)\nu+1}^{(i)} \leq -u_n(y_i), S_{n,1} \hat{X}_{n,(s-1)\nu+t}^{(j)} > u_n(x_j) \right) \\ &\quad \left. + \mathbb{P} \left( S_{n,1} \hat{X}_{n,(s-1)\nu+1}^{(i)} \leq -u_n(y_i), S_{n,1} \hat{X}_{n,(s-1)\nu+t}^{(j)} \leq -u_n(y_j) \right) \right). \end{aligned}$$

Since  $0 \leq 1 - \frac{\nu\mu}{2^l} \Sigma_1 - \nu \left( 1 - \frac{\mu}{2^l} \right) \Sigma_2 \leq 1$  applying Lemma 3 in [19] we obtain

$$\begin{aligned} &\sum_{\mu=0}^{2^l-1} \sum_{\beta_n \in \{0,1\}^n} \mathbb{E} \left( \left| \prod_{s=1}^l P(K_s, \beta_n, \mathbf{x}, \mathbf{y}) - \prod_{s=1}^l \left[ 1 - \frac{\frac{\mu}{2^l} n \Sigma_1 - \left( 1 - \frac{\mu}{2^l} \right) n \Sigma_2}{l} \right] \right| \mathbb{I}(B_{\mu,l, \beta_n}) \right) \\ &\leq \sum_{\mu=0}^{2^l-1} \sum_{\beta_n \in \{0,1\}^n} \mathbb{E} \left( \sum_{s=1}^l \left| P(K_s, \beta_n, \mathbf{x}, \mathbf{y}) - \left[ 1 - \frac{\frac{\mu}{2^l} n \Sigma_1 - \left( 1 - \frac{\mu}{2^l} \right) n \Sigma_2}{l} \right] \right| \mathbb{I}(B_{\mu,l, \beta_n}) \right) \\ &\leq \sum_{\mu=0}^{2^l-1} \sum_{s=1}^l \mathbb{E} \left( \left| \frac{\sum_{j \in K_s} \varepsilon_{n,j}}{\nu} - \frac{\mu}{2^l} \right| \mathbb{I}(B_{\mu,l}) \right) n(\Sigma_1 - \Sigma_2) + n \Sigma_3 \\ &\leq \sum_{s=1}^l \left[ 2(2s-1) \left( d \left( \frac{\Xi_{\nu s}}{\nu s}, \eta \right) + d \left( \frac{\Xi_{\nu(s-1)}}{\nu(s-1)}, \eta \right) \right) + \frac{1}{2^l} \right] \frac{n(\Sigma_1 - \Sigma_2)}{l} + n \Sigma_3, \end{aligned} \quad (4.5)$$

where  $d(X, Y)$  stands for Ky Fan metric, i.e.,  $d(X, Y) = \inf\{\varepsilon, \mathbb{P}(|X - Y| > \varepsilon) < \varepsilon\}$ . Furthermore,

$$\begin{aligned} &\sum_{\mu=0}^{2^l-1} \sum_{\beta_n \in \{0,1\}^n} \mathbb{E} \left( \left| \prod_{s=1}^l \left[ 1 - \frac{\frac{\mu}{2^l} n \Sigma_1 - \left( 1 - \frac{\mu}{2^l} \right) n \Sigma_2}{l} \right] - \prod_{s=1}^l \left[ 1 - \frac{\eta n \Sigma_1 - \left( 1 - \eta \right) n \Sigma_2}{l} \right] \right| \mathbb{I}(B_{\mu,l, \beta_n}) \right) \\ &\leq \sum_{\mu=0}^{2^l-1} \sum_{s=1}^l \mathbb{E} \left( \left| \frac{\mu}{2^l} - \eta \right| \mathbb{I}(B_{\mu,l}) \right) \frac{n}{l} (\Sigma_1 + \Sigma_2) \\ &\leq \frac{n(\Sigma_1 + \Sigma_2)}{2^l}. \end{aligned} \quad (4.6)$$

By the fact that  $\lim_{\nu \rightarrow \infty} d \left( \frac{\Xi_{\nu s}}{\nu s}, \eta \right) = 0$  and utilising (4.3)-(4.6), by passing to limit for  $n \rightarrow \infty$  and then letting  $\nu \rightarrow \infty$  we obtain

$$\begin{aligned} &\left| P(n, \varepsilon_n, \mathbf{x}, \mathbf{y}) - \mathbb{E} \left( 1 - \frac{\eta(D(x_1, x_2) + D(y_1, y_2)) + (1 - \eta)(D(x_3, x_4) + D(y_3, y_4))}{l} \right) \right| \\ &\leq \frac{D(x_1, x_2) + D(y_1, y_2)}{2^{l-1}} + \frac{1}{l} (e^{-x_1} + e^{-y_1} + e^{-x_2} + e^{-y_2})^2. \end{aligned}$$

Next, letting  $l \rightarrow \infty$  implies

$$\lim_{n \rightarrow \infty} \sup_{\substack{x_i, y_i \in \mathbb{R}, i=\{1,2,3,4\} \\ x_1 \leq x_3, x_2 \leq x_4, y_1 \leq y_3, y_2 \leq y_4}} \left| P(n, \varepsilon_n, \mathbf{x}, \mathbf{y}) - \mathbb{E} \left( H_\lambda^\eta(x_1, x_2) H_\lambda^\eta(y_1, y_2) H_\lambda^{1-\eta}(x_3, x_4) H_\lambda^{1-\eta}(y_3, y_4) \right) \right| = 0,$$

hence the claim follows.  $\square$

PROOF OF THEOREM 3.2 If (2.7) holds, by (2.8) for some positive constant  $\mathcal{Q}$  we have

$$\begin{aligned} &\left| \mathbb{P} \left( -u_n(y_1) < m_n^{(1)}(\varepsilon_n) \leq M_n^{(1)}(\varepsilon_n) \leq u_n(x_1), -u_n(y_2) < m_n^{(2)}(\varepsilon_n) \leq M_n^{(2)}(\varepsilon_n) \leq u_n(x_2), \right. \right. \\ &\quad \left. \left. -u_n(y_3) < m_n^{(1)} \leq M_n^{(1)} \leq u_n(x_3), -u_n(y_4) < m_n^{(2)} \leq M_n^{(2)} \leq u_n(x_4) \right) \right. \\ &\quad \left. - \mathbb{P} \left( -u_n(y_1) < \widehat{m}_n^{(1)}(\varepsilon_n) \leq \widehat{M}_n^{(1)}(\varepsilon_n) \leq u_n(x_1), -u_n(y_2) < \widehat{m}_n^{(2)}(\varepsilon_n) \leq \widehat{M}_n^{(2)}(\varepsilon_n) \leq u_n(x_2), \right. \right. \\ &\quad \left. \left. -u_n(y_3) < \widehat{m}_n^{(1)} \leq \widehat{M}_n^{(1)} \leq u_n(x_3), -u_n(y_4) < \widehat{m}_n^{(2)} \leq \widehat{M}_n^{(2)} \leq u_n(x_4) \right) \right| \end{aligned}$$

$$\leq Qnw^{\frac{4\alpha+2p}{2+p}} \sum_{i,j=1,2} \sum_{k=1}^n |\lambda_{ij}(k, n)| \exp\left(-2(1 + |\lambda_{ij}(k, n)|)^{-\frac{p}{2+p}} Tw^{\frac{2p}{2+p}}\right),$$

where  $w = \min(|u_n(x_i)|, |u_n(y_i)|, 1 \leq i \leq 4)$ . In view of Lemma 3.3 in [13], the sum of the right side of the inequality tends to 0. Thus the claim follows by Lemma 4.3.  $\square$

**Acknowledgments.** We would like to thank the referees for several suggestions which improved our manuscript. The authors have been partially support from the Swiss National Science Foundation grants 200021-140633/1, 200021-134785 and the project RARE -318984 (an FP7 Marie Curie IRSES).

## References

- [1] S.M. Berman. Limit theorems for the maximum term in stationary sequences. *Ann. Math. Statist.*, 35:502–516, 1964.
- [2] S.M. Berman. *Sojourns and Extremes of Stochastic Processes*. The Wadsworth & Brooks/Cole Statistics/Probability Series. Wadsworth & Brooks/Cole Advanced Books & Software, Pacific Grove, CA, 1992.
- [3] S. Cambanis, S. Huang, and G. Simons. On the theory of elliptically contoured distributions. *J. Multivariate Anal.*, 11(3):368–385, 1981.
- [4] H. Cramér and M.R. Leadbetter. *Stationary and related stochastic processes. Sample function properties and their applications*. John Wiley & Sons Inc., New York, 1967.
- [5] P. Embrechts, C. Klüppelberg, and T. Mikosch. *Modelling Extremal Events*. Springer-Verlag, Berlin, 1997.
- [6] J.P. French and R.A. Davis. The asymptotic distribution of the maxima of a Gaussian random field on a lattice. *Extremes*, 16(1):1–26, 2013.
- [7] E. Hashorva. Elliptical triangular arrays in the max-domain of attraction of Hüsler-Reiss distribution. *Statist. Probab. Lett.*, 72(2):125–135, 2005.
- [8] E. Hashorva. Gaussian approximation of conditional elliptic random vectors. *Stoch. Models*, 22(3):441–457, 2006.
- [9] E. Hashorva. Exact tail asymptotics in bivariate scale mixture models. *Extremes*, 15(1):109–128, 2012.
- [10] E. Hashorva. Minima and maxima of elliptical arrays and spherical processes. *Bernoulli*, 19(3):886–904, 2013.
- [11] E. Hashorva, A. G. Pakes, and Q. Tang. Asymptotics of random contractions. *Insurance Math. Econom.*, 47(3):405–414, 2010.
- [12] E. Hashorva, Z. Peng, and Z. Weng. On Piterbarg theorem for maxima of stationary Gaussian sequences. *Lithuanian Mathematical Journal*, 53(3):280–292, 2013.
- [13] E. Hashorva, Z. Peng, and Z. Weng. Limit distributions of exceedances point processes of scaled stationary Gaussian sequences. *Probability and Mathematical Statistics, in press*, 2014.
- [14] E. Hashorva and Z. Weng. Limit laws for extremes of dependent stationary Gaussian arrays. *Statist. Probab. Lett.*, 83(1):320–330, 2013.
- [15] E. Hashorva and Z. Weng. Limit laws for maxima of contracted stationary Gaussian sequences. *Communications in Statistics-Theory and Methods, in press*, 2014.

- [16] E. Hashorva and Z. Weng. Tail asymptotic of Weibull-type risks. *Statistics, in press*, 2014.
- [17] T. Hsing, J. Hüsler, and R.-D. Reiss. The extremes of a triangular array of normal random variables. *Ann. Appl. Probab.*, 6(2):671–686, 1996.
- [18] J. Hüsler and R.-D. Reiss. Maxima of normal random vectors: between independence and complete dependence. *Statist. Probab. Lett.*, 7(4):283–286, 1989.
- [19] T. Krajka. The asymptotic behaviour of maxima of complete and incomplete samples from stationary sequences. *Stochastic Process. Appl.*, 121(8):1705–1719, 2011.
- [20] M.F. Kratz. Level crossings and other level functionals of stationary Gaussian processes. *Probab. Surv.*, 3:230–288, 2006.
- [21] M.R. Leadbetter, G. Lindgren, and H. Rootzén. *Extremes and related properties of random sequences and processes*. Springer Verlag, 1983.
- [22] W.V. Li and Q.-M. Shao. A normal comparison inequality and its applications. *Probab. Theory Related Fields*, 122(4):494–508, 2002.
- [23] D. Lu and X. Wang. Some new normal comparison inequalities related to Gordon’s inequality. *Statist. Probab. Lett.*, 88, 2014.
- [24] G. L. O’Brien. Extreme values for stationary and Markov sequences. *Ann. Probab.*, 15(1):281–291, 1987.
- [25] Z. Peng, Z. Weng and S. Nadarajah. Joint limiting distributions of maxima and minima for complete and incomplete samples from weakly dependent stationary sequences. *J. Comput. Anal. Appl.*, 13(5):875–880, 2011.
- [26] V.I. Piterbarg. On the paper by J. Pickands ”upcrossing probabilities for stationary gaussian processes”. *Vestnik Moscow Univ. Ser. I Mat. Mekh.* 27, 25-30. *English transl. in Moscow Univ. Math. Bull.* 1972,27, 27:25–30, 1972.
- [27] V.I. Piterbarg. *Asymptotic methods in the theory of Gaussian processes and fields*, volume 148 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1996.
- [28] R.L. Plackett. A reduction formula for normal multivariate integrals. *Biometrika*, 41:351–360, 1954.
- [29] D. Slepian. The one-sided barrier problem for Gaussian noise. *Bell System Tech. J.*, 41:463–501, 1962.