

Serveur Académique Lausannois SERVAL [serval.unil.ch](http://serval.unil.ch)

## Author Manuscript

Faculty of Biology and Medicine Publication

**This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.**

Published in final edited form as:

**Title:** Acute and long-lasting effects of oxytocin in cortico-limbic circuits: consequences for fear recall and extinction.

**Authors:** Triana-Del Río R, van den Burg E, Stoop R, Hegoburu C

**Journal:** Psychopharmacology

**Year:** 2019

**Issue:** 236

**Volume:** 1

**Pages:** 339-354

**DOI:** [10.1007/s00213-018-5030-5](https://doi.org/10.1007/s00213-018-5030-5)

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

# Acute and long-lasting effects of oxytocin in cortico-limbic circuits: consequences for fear recall and extinction

Rodrigo Triana-Del Río<sup>1</sup> & Erwin van den Burg<sup>1</sup> & Ron Stoop<sup>1</sup> & Chloé Hegoburu<sup>1</sup>

## Abstract

The extinction of conditioned fear responses entrains the formation of safe new memories to decrease those behavioral responses. The knowledge in neuronal mechanisms of extinction is fundamental in the treatment of anxiety and fear disorders. Interestingly, the use of pharmacological compounds that reduce anxiety and fear has been shown as a potent co-adjuvant in extinction therapy. However, the efficiency and mechanisms by which pharmacological compounds promote extinction of fear memories remains still largely unknown and would benefit from a validation based on functional neuronal circuits, and the neurotransmitters that modulate them. From this perspective, oxytocin receptor signaling, which has been shown in cortical and limbic areas to modulate numerous functions (Eliava et al. *Neuron* 89(6):1291-1304, 2016), among them fear and anxiety circuits, and to enhance the salience of social stimuli (Stoop *Neuron* 76(1):142-59, 2012), may offer an interesting perspective. Experiments in animals and humans suggest that oxytocin could be a promising pharmacological agent at adjusting memory consolidation to boost fear extinction. Additionally, it is possible that long-term changes in endogenous oxytocin signaling can also play a role in reducing expression of fear at different brain targets. In this review, we summarize the effects reported for oxytocin in cortico-limbic circuits and on fear behavior that are of relevance for the modulation and potential extinction of fear memories.

**Keywords** Fear extinction · Fear retrieval · Oxytocin · Central amygdala · Prefrontal cortex

## Abbreviations

ACC Anterior cingulate cortex  
 AVP1a-R Vasopressin receptor  
 BA Basal amygdala  
 BLA Basolateral amygdala  
 CBT Cognitive behavioral therapy  
 CeA Central amygdala  
 CeL Centro-lateral amygdala  
 CeM Centro-medial amygdala  
 CS Conditioned stimulus  
 CSF Cerebrospinal fluid  
 dmPFC Dorso-medial prefrontal cortex  
 GABA Gamma amino butyric acid  
 IL Infralimbic cortex

ITCd Intercalated cell masses (dorsal)  
 ITCv Intercalated cell masses (ventral)  
 LA Lateral amygdala  
 NMDAR N-methyl-d-aspartate receptor  
 Nacc Nucleus accumbens  
 OT Oxytocin  
 OTR Oxytocin receptor  
 PKC $\delta$  Protein kinase delta  
 PL Prelimbic cortex  
 PTSD Posttraumatic stress disorder  
 PVN Paraventricular nucleus of the hypothalamus  
 US Unconditioned stimulus  
 vmPFC Ventromedial prefrontal cortex

\* Chloé Hegoburu  
 chloe.hegoburu@gmail.com

<sup>1</sup> Laboratory of Neurobiology of emotions, Center of Psychiatric Neuroscience, CHUV, Prilly, Lausanne, Switzerland

## Introduction

Anxiety-related disorders are at present a worldwide psychiatric, pharmacological, and economical issue, and according to the World Health Organization (WHO 2016), since 1990, the number of people suffering from anxiety increased by nearly 50%. In fact, close to 10% of the world's population

---

is affected by mental disorders and this accounts for 30% of the global non-fatal disease burden (World Health Organization 2016). Thus, reducing anxiety and fear-related neuropathologies represents a priority in the current health system.

Fear memories are caused by concrete threats. During fear-conditioned learning, an aversive stimulus or unconditioned stimulus (US), like a mild electric foot shock in rodents or gunfire in humans, is associated with a neutral sensory stimulus or conditioned stimulus (CS), like a tone, a smell or a visual cue, resulting in the creation of a conditioned fear memory. When the individual is re-exposed to the CS, the fear memory is retrieved or recalled, generating a fear response, for example, freezing in rats and mice (LeDoux 2000; Alberini 2013), or fear-potentiated startle (often measured in humans) accompanied by the activation of a whole series of physiological responses (Debiec and LeDoux 2006; Quirk and Mueller 2008; Do-Monte et al. 2015). Meanwhile, anxiety states can be induced by circumstantial, potential, imagined, or anticipated threats, which are more related with the unspecific generalization of fear. Anxiety states and fear memories are brought about in part by the same neuronal circuits, involving the different subnuclei of the amygdala and the prefrontal cortex (LeDoux 2000, 2014; Davis et al. 2010; Tye and Deisseroth 2012; Anderson and Adolphs 2014; Tovote et al. 2015), so that treatment options of conditioned fear and anxiety can overlap. Furthermore, anxiety and fear memories are encoded in an engram, which has been represented by a specific population of neurons activated during specific learning events (Tonegawa et al. 2015). This notion provides a cellular substrate that can be subjected to modulation during the recall of the original memory, but also during its extinction. Efficient therapies at modulating engrams to promote fear extinction are therefore needed, and extinction learning is at the basis of this. Indeed, extinction of conditioned fear responses, which encompasses the formation of new memories encoding safety, and thereby inhibiting fear, represents an important mechanism in the treatment of anxiety disorders, such as posttraumatic stress disorder (PTSD), social phobia, panic disorders, and fear generalization. In psychiatry, extinction training is commonly used in cognitive behavioral therapies (CBT) (Cisler et al. 2016; Klumpp et al. 2017; Abramowitz 2013). CBT focuses on identifying and modifying maladaptive patterns of conditioned behavior and is one of the most established methods to reduce fear (Stewart and Chambless 2009). In addition, it targets the neuronal processes that inhibit fear retrieval (Straub et al. 2017; Carpenter et al. 2018).

Extinction training involves repeated presentation of the CS in the absence of the US, and results in a diminished fear response towards the CS (Hofmann et al. 2015; Pavlov 1927; Alberini 2013). This is presumed to be the result of a cascade of genetic, molecular, and cellular changes (Myers and Davis

2007; Orsini and Maren 2012) that alter synaptic efficacy and neuronal network interactions in the pathways for retrieval and extinction of fear memories (Pape and Paré 2010; Tonegawa et al. 2015). The strength of extinction memory (extinction consolidation) is evaluated after the training, in so-called extinction retrieval sessions, in which the conditioned fear is decreased, and therefore, also the potential anxiety that accompanies the fear memory is diminished (Quirk et al. 2006; Schiller et al. 2013). Extinction training, although effective in decreasing the fear response in the short term, does not erase the original fear memory trace, but forms a new, distinct memory (Milad and Quirk 2002; Quirk et al. 2006; Schiller et al. 2013; Kroes et al. 2016; Rashid et al. 2016). As a result, the original conditioned fear response can reoccur with time through spontaneous recovery, reinstatement, and renewal (Rescorla 2004; Kim and Richardson 2010). Remarkably, the extinction memory does not generalize as easily to new contexts, as the original conditioned fear memory does during anxiety states (Woods and Bouton 2006). Because extinction learning does not generalize to situations outside the extinction context, it is therefore not the best clinical tool for long-lasting fear reduction.

A different method to reduce the fear response is the use of pharmacological agents that target precisely the synaptic plasticity of neuronal circuits of extinction memories, whose downstream signaling pathways can allow fear extinction and strengthen extinction memory unremittingly in preclinical and clinical models (for extensive review follow: Bukalo et al. 2014; Singewald et al. 2015). Indeed, pharmacological agents applied after retrieval of the fear memory might facilitate fear extinction, as evidenced by the apparent amnesic effects of propranolol, a beta-adrenergic antagonist, in animals and humans (Kroes et al. 2016; Giustino et al. 2016). Although propranolol has been accepted as a clinical option for general anxiety, its precise mechanism lies in the amnesic effect on retrieved fear memories, and are not fully validated in the circuits for fear extinction (Debiec and Ledoux 2004; Steenen et al. 2016). In the same line, the antibiotics anisomycin (a blocker of protein synthesis) and d-cycloserine (partial NMDA receptor agonist that works as a cognitive enhancer) seem to support fear extinction as well (Nader et al. 2000; Ledgerwood et al. 2005; Anderson and Insel 2006; Hofmann et al. 2015; Bowers and Ressler 2015a, 2015b). While the use of anisomycin is limited by its high toxicity, as it blocks all protein synthesis, d-cycloserine is already being used in combination with CBT to enhance fear extinction. On one hand, it is thought to suppress the initial fear memory, and on the other hand to promote extinction learning through activation of NMDA receptors (Davis et al. 2006; Bowers and Ressler 2015b; Mataix-Cols et al. 2017; Goodman et al. 2018).

Although these examples show the beneficial effects of pharmacological agents, as co-adjuncts of extinction

therapies, some other reports suggest that co-adjuvant use of anxiolytics provides only modest gains in fear extinction or can even impair extinction learning working by acting as potential anxiogenics (Singewald et al. 2015). An example of this can be found in positive allosteric modulation of GABA<sub>A</sub> receptor activity via the benzodiazepine-binding site (Clark et al. 2004; Knoflach et al. 2016; Duke et al. 2018). Because of the sedating and calming effects of benzodiazepines, they might in fact hinder extinction via reduced arousal and decreased release of noradrenaline and glucocorticoids, neuromodulators that support extinction (Otto et al. 2005; Bentz et al. 2010). Similar concerns have been raised for the use of serotonin selective reuptake inhibitors (SSRI). Chronic treatment with some SSRIs (e.g., fluoxetine) but not all (e.g., citalopram) enhances extinction in rodents (Burghardt and Bauer 2013). However, acute SSRI treatment could on the other hand induce anxiogenic effects (Campbell and Merchant 2003; Salchner and Singewald 2006). Finally, another interesting compound is venlafaxine, a combined 5-HT and noradrenaline reuptake inhibitor with weaker affinity for the noradrenaline transporter (Owens et al. 1997). This compound improves extinction retrieval and protects against fear reinstatement in rodents (Yang et al. 2012). In summary, some pharmacological compounds could target the brain networks responsible for fear extinction but in a rather unspecific manner. A more effective and innocuous pharmacological approach should therefore be adopted to specifically target the consolidation and reconsolidation of extinction memories (Alberini 2013) (Fig. 1).

Here, we discuss the neuropeptide oxytocin (OT) as a potential endogenous regulator of the neuronal plasticity behind fear expression and extinction-related behaviors, and which can easily be applied intranasally to improve fear extinction learning in humans (Eckstein et al. 2015, 2017). Oxytocin is a well-established anxiolytic when exogenously applied or endogenously released in several brain regions, including the PVN where it is synthesized (Blume et al. 2008; Jurek et al. 2012). In this area, its anxiolytic activity is brought about by the recruitment of an oxytocin receptor (OTR)–TRPV2 channels (and subsequent Ca<sup>2+</sup> influx)–MEK1/2 pathway (van den Burg et al. 2015). Underlying mechanisms of oxytocin and fear extinction in other brain regions, like the PFC and amygdala, are beginning to be explored, motivated by the discovery of OT release and OTR expression in those areas (Knobloch et al. 2012; Mitre et al. 2016; Li et al. 2016; Rogers et al. 2018).

### **Cortico-limbic circuits for fear retrieval and extinction memories**

As the modern standpoint in psychiatry pursues a more precise circuit-based therapy, a pertinent pharmacological

methodology must target the neuronal populations, where extinction memories are formed and consolidated (Fitzgerald et al. 2014; Bukalo et al. 2014). The circuits between the prefrontal cortex and the amygdala represent the physiological substrate for the recall or retrieval of fear memories, but also extinction memories (Fig. 1). Moreover, these brain areas express the receptor for oxytocin (OTR). Thus, the physiological interactions between the brain areas involved, as well as the neuromodulators such as OT that regulate their functions, represent a clear anatomical substrate to facilitate strengthening of fear extinction memories.

### **Amygdala**

In the classical circuit model of fear conditioning, the lateral nucleus of the amygdala (LA) is the primary site where associations between the CS and the US are formed and stored (Davis 1992; LeDoux 2000; Maren 2001). LA pyramidal neurons project to the central nucleus of the amygdala (CeA), which then mediates the physiological and behavioral expression of conditioned fear responses. CeA output neurons, most of which are located in its medial subdivision (CeM), project to downstream targets in the brainstem and hypothalamus that orchestrate conditioned autonomic and motor responses (Krettek and Price 1978; Veening et al. 1984; LeDoux et al. 1988). Nevertheless, circuits within the CeA have also been shown to be important for acquisition as well as expression of fear (Ciocchi et al. 2010; Haubensak et al. 2010; Duvarci et al. 2011; Amano et al. 2012; Li et al. 2013a).

In these different studies, distinct cell types have been identified based on functional and genetic criteria. In the lateral subdivision of CeA (CeL), PKC $\delta^+$  mRNA-containing neurons exhibit inhibitory CS responses, also called CeL<sub>OFF</sub> cells. These PKC $\delta^+$  neurons receive local inhibitory inputs from CS-activated PKC $\delta^-$  neurons (CeL<sub>ON</sub>), and in turn make inhibitory synaptic contacts with PAG-projecting CeA output neurons, thereby gating acute CeA output through disinhibition. CeM output cells are under tight inhibitory control of this GABAergic microcircuit located in the CeL. This microcircuit can be activated by OT, resulting in inhibition of neuronal activity in the CeM (Cassell 1999; Huber et al. 2005; Ehrlich et al. 2009; Viviani et al. 2011; Knobloch et al. 2012).

Abbreviations: ACC, anterior cingulate cortex; PL, prelimbic cortex; IL, infralimbic cortex; LA, lateral amygdala; BA, basal amygdala; ITCd, intercalated cells dorsal; ITCv, intercalated cells ventral; CeL, centro-lateral amygdala; CeM, centro-medial amygdala; PAG, periaqueducal gray

While this anatomical arrangement is at the basis of the physiological and behavioral expression of fear, recent studies have highlighted that the microcircuitry in the CeA is important for fear memory as well. For example, during fear retrieval and accompanying freezing behavior, excitatory synapses onto somatostatin-positive neurons (CeL<sub>ON</sub>) in the CeL




Figure 1 Hypothetical circuits involved in recall and extinction of fear memories. In this hypothesis, two parallel mechanisms are involved to induce the CS responsiveness of CeM output neurons after conditioning. The balance between excitatory inputs from BA and the inhibitory inputs from CeL and ITCv inputs to CeM triggered its CS responsiveness. This balance can be modulated by the ACC and prefrontal cortex (PL or IL) activities. Recall pathways (upper part of the panel): The CS-activated LA neurons projecting to BA neuron subtype (fear neurons, ▲ pink triangle) to excite CeM cells, whereas another type of BA neurons (extinction neurons, ▲ green triangle) are inhibited. The response from LA and BA fear neurons can be modulated through excitatory interactions with each other and/or with prelimbic (PL) cells and anterior cingulate cortex, would reinforce the activity in LA during CS presentation. Conversely, the activation of LA neurons also leads to the recruitment of ITCd inhibitory cells and of a subgroup of CeL interneurons named CeL<sub>Off</sub>/PKCδ<sup>-</sup>. ITCd neurons would then inhibit ITCv cells, disinhibiting CeM neurons. In addition, ITCd cells would inhibit another subgroups of CeL interneurons named CeL<sub>Off</sub>/PKCδ<sup>+</sup>. These activations cause a further inhibition

of PKCδ<sup>+</sup> neurons and disinhibition of CeM cells leading to freezing response. These mechanisms take place only when the threat is inescapable, otherwise the neurons from BA lead to escape behavior. Extinction pathways (lower part of the panel): These balance between two parallel mechanisms leads to the decrease of fear response to the CS presentation and the activation of CeM output neurons during extinction. During this new phase of learning, the CS diminished rapidly the recruitment of BA fear neurons leading to a disinhibition of BA extinction neurons (green). Moreover, the projections from Infralimbic to BA (and reciprocally) might also enhance the excitability of BA extinction cells. The disinhibition of extinction cells induced the activation of a subgroup of interneurons in BA inhibiting fear neurons. In parallel, the reduction of activity in LA causes a disaffiliation of ITCd neurons leading to an activation by disinhibition of ITCv neurons, and simultaneous activation by BA 'Extinction' glutamatergic neurons. Moreover, the subsets of CeL cells PKCδ<sup>+</sup> can be disinhibited leading to the inactivation of CeM neurons and less fear responses.

experience a robust synaptic potentiation (Li et al. 2013a, b). Additionally, optogenetic stimulation of basolateral amygdala (BLA) terminals in the CeA results in anxiolysis during the retrieval of fear context (Tye et al. 2011), possibly brought about by activation of CeL<sub>OFF</sub> cells expressing the OTR (Haubensak et al. 2010; Viviani et al. 2011). Thus, the observed changes in plasticity (the cellular correlate of learning) and bidirectional modulation of fear-related behavior through CeL<sub>ON</sub> and CeL<sub>OFF</sub> cell activity establish the CeA as an important node in the expression of fear, and a locus for fear memory.

Another relevant mechanism for extinction involves the activation of the intercalated cell masses (ITC) of the amygdala (Royer et al. 1999; Quirk et al. 2003; Paré et al. 2004; Amano et al. 2010; Amir et al. 2011; Bukalo et al. 2014). The ITCs include clusters of tightly packed, GABAergic neurons connected with the fiber bundles that lie between the BA/LA and the CeL/CeM nuclei. Further reports have highlighted the heterogeneity of diverse ITCs. The dorsal and ventral ITC nuclei are heavily interconnected, exhibit miscellaneous electrophysiological and molecular profiles, and are differentially activated by fear and fear extinction, while dorsal ITC activity emerge during fear recall, ventral ITC activity is enhanced during fear extinction (Geracitano et al. 2007; Kaoru et al. 2010; Busti et al. 2011; Bukalo et al. 2014). In fact, general excitatory synaptic signaling onto the ventral ITC population appears to be necessary for fear extinction, as a mouse model lacking the serine protease inhibitor protease-nexin-1 (PN-1), that modulates NMDAR-mediated synaptic currents in the vITC, displays impaired fear extinction (Meins et al. 2010).

### Prelimbic versus infralimbic cortices

The fundamental role of the prefrontal cortex to retrieve, consolidate, and extinguish fear memories has been well-demonstrated in a plethora of pharmacological, optogenetic, and electrophysiological studies. These roles seem to be conserved between rodents and humans, and homologous subnuclei in the prefrontal cortex have been found in both species (Fig. 2). In rodents, the subnuclei are named, from dorsal to ventral: anterior cingulate cortex (ACC), prelimbic cortex (PL), and infralimbic cortex (IL). They correspond respectively to the human anterior cingulate cortex (ACC), dorso-medial prefrontal cortex (dmPFC), and ventro-medial prefrontal cortex (vmPFC) (Milad et al. 2006; Quirk and Milad 2010; Likhtik et al. 2014; Duvarci and Pare 2014; Giustino and Maren 2015).

The PL and IL play opposing roles in encoding fear memories, triggering fear retrieval, and fear extinction, respectively (Vidal-Gonzalez et al. 2006; Herry et al. 2008; Quirk and Milad 2010; Milad and Quirk 2012; Little and Carter 2013; Courtin et al. 2014; Senn et al. 2014; Dejean et al. 2016; Karalis et al. 2016; McGarry and Carter 2017). This difference is due to the differential functional connectivity patterns of PL and IL with

unique populations of pyramidal cells in the BLA, and is amplified by mutual inhibition driven by interneuron populations in both prefrontal subnuclei (Vogel et al. 2016; Bukalo et al. 2015; Vertes 2004; Vidal-Gonzalez et al. 2006; Cho et al. 2013; Courtin et al. 2014; Saffari et al. 2016; Karalis et al. 2016). The BLA pyramidal neurons project back to both the PL and IL sub regions, creating a feedback loop (Gabbott et al. 2006; Hoover and Vertes 2007; Senn et al. 2014; Burgos-Robles et al. 2017; McGarry and Carter 2017). These BLA inputs to the PL and IL modulate the excitatory/inhibitory balance on cortical pyramidal cells, possibly via engagement of local interneurons (Ishikawa and Nakamura 2003; Sun and Laviolette 2012; Sotres-Bayon et al. 2012; Dilgen et al. 2013). Then, the bidirectional projections between PL/IL and the BLA encode the experience-dependent retrieval versus extinction processing of fear memories by balancing PL vs. IL activity.

### Anterior cingulate cortex

Besides the PL and IL, the anterior cingulate cortex (ACC) also sends projections to the BLA, and modulates the expression and retrieval of fear (Vertes 2004; Sierra et al. 2017; Webb et al. 2017; Yin et al. 2018). In rodents, this area is responsible for the enhancement of contextual fear generalization after fear conditioning, and for the transfer of socially derived fear through its connections with the amygdala (Cullen et al. 2015; Pisansky et al. 2017; Allsop et al. 2018). In humans, local GABAergic signaling in the ACC promotes fear extinction, which suggests that inhibition of the dorsal ACC may initially inhibit the retrieval of fear memories, and thus facilitates extinction learning (Levar et al. 2017a, 2017b).


Additionally, some structural and functional deficits in this area have been found in fMRI studies in neuropsychiatric disorders, such as generalized anxiety disorder (Andreescu et al. 2017) and PTSD (Shvil et al. 2014), which both affect severely the extinction of fear memories. In the case of PTSD, interestingly, the activation of ACC, together with the vmPFC and hippocampus, is sexually dimorphic: men with PTSD exhibited increased activation in the left rostral dACC during extinction recall compared to women with PTSD (Shvil et al. 2014; Yoon et al. 2017). Further, animal and human studies (Cullen et al. 2015; Levar et al. 2017a, 2017b) have positioned the ACC as an important hub to trigger fear retrieval alongside the PL, and its inhibition might be a way to boost fear extinction.

### Modulation of fear circuits by oxytocin in preclinical animal models

#### Rodents

There is quite a lot of evidence that OT acts in different brain areas to decrease anxiety responses and conditioned fear

Fig. 2 Anatomical locations of the anterior cingulate cortex (ACC), prelimbic (PL), and infralimbic (IL) subregions of the medial PFC, showing equivalence between the human and rat brains. The rodent PL region is largely associated to Brodmann area 32, while the IL is analogous to Brodmann area 25. The ACC is mainly associated with the Brodmann area 24. Usually these observations are based on the different type of thalamic inputs to these regions



responses during retrieval in rodents (Fig. 3). Indeed, OTRs and OT-binding sites have been found in those circuits, especially in the ACC, PL, and CeL (Huber et al. 2005; Nakajima et al. 2014; Burkett et al. 2016; Li et al. 2016; Sabihi et al. 2017) (Jiménez et al. 2015), that together constitute the fear retrieval network. The OTRs are expressed specifically by GABAergic interneurons in all of the brain regions involved in fear retrieval (Huber et al. 2005; Owen et al. 2013; Nakajima et al. 2014; Marlin et al. 2015; Sabihi et al. 2017). Importantly, OT-positive fibers originating from the PVN project to these areas (Knobloch et al. 2012), and, at least in the CeL, OT is released from these fibers as observed following optogenetic stimulation and application of an OTR antagonist. Furthermore, OT release in the CeL is sufficient to decrease conditioned fear responses, clearly demonstrating the anxiolytic potential of OT in a pathway that mediates fear retrieval when the animals are re-exposed to the conditioned fear context. It has further been shown that the OTR mRNA is indeed present in part of the CeL<sub>OFF</sub> cells in the CeL (Haubensak et al. 2010), and we found that activation of these receptors inhibits the CeM and its output to downstream effector regions (Huber et al. 2005; Viviani et al. 2011; Knobloch et al. 2012). Furthermore, the OTR has been found in the basolateral amygdala (BLA), another hub for extinction memories, by means of immunohistochemistry and autoradiography (Stamatakis et al. 2016; Mitre et al. 2016). In fact, it has been suggested that the extinction of conditioned fear is facilitated

by local infusions of OT or TGOT (specific agonist of the OTR) in the BLA, but not in the CeA (Campbell-Smith et al. 2015).

By using an OTR antagonist, it has further been shown that endogenous OT has similar anxiolytic effects in the PL, as it has in the CeL (Nakajima et al. 2014; Li et al. 2016; Sabihi et al. 2017). In addition, optogenetic and pharmacological activation of the OTR in the PL reduces anxiety-like behavior, in a gender-dependent manner (Li et al. 2016), further characterizing the differential role of different prefrontal-amygdala circuits during the modulation of fear memories in female and male rodents (Shansky et al. 2010; Gruene et al. 2015). Moreover, Mitre et al. (2016) found a higher density of the OTR expression within the PL of lactating females compared to virgins or males rodents. Consistent with this, Sabihi et al. (Sabihi et al. 2014) showed that blocking the OTR in this area enhances anxiety in lactating but not virgin females. In addition, in the ACC, the OTR is also expressed (Mitre et al. 2016), and an OTR antagonist infused into this region abolished the partner-directed consolation behavior during fear transfer in prairie voles (Burkett et al. 2016).

While OT can exert acute anxiolytic effects by reducing the activity of the fear retrieval network, especially in the central amygdala (Fig. 3), it might also modulate plasticity of the fear extinction network to bring about the reduction of fear responses. This would require the presence of OTRs for instance in the IL, or inhibitory actions in the PL or ACC (as in the CeA). Indeed, administration of OT in the IL of fear-




Fig. 3 Modulation by oxytocin signaling of the balance between fear recall and fear extinction. In this hypothesis, oxytocin would allow a switch from fear recall to fear extinction, based on the hypothetical model of circuitry presented in Fig. 1. This model revealed a complex mechanism involving always a balance between inhibition and excitation leading to increase or decrease the CS responsiveness from CeM neurons. In oxytocin model of switching fear to extinction, the presence of oxytocin would activate the receptors localized only on the interneurons of ACC, PL, and CeL. Interestingly, in CeL the OT-receptors are only on cell PKC $\delta$ +. During the recall, the CS activated LA neurons projecting to BA neuron subtype (bfear neurons, ^ pink triangle) to excite CeM cells, whereas another type of BA neurons (bextinction neurons, ^ green triangle) are inhibited. Oxytocin receptors are localized on interneurons of regions (ACC, PL, CeL) involved in recall, thus the activation of these receptors by oxytocin leads to inhibit the bfear neurons and to recruit more bextinction neurons in BA. Oxytocin would diminish rapidly the

capacity from the CS to recruit bfear neurons in BA leading to a disinhibition of BA bextinction neurons (green). Moreover, the projections from infralimbic to BA (and reciprocally) might also enhance the excitability of BA bextinction cells. The disinhibition of extinction cells induced the activation of a subgroup of interneurons in BA inhibiting bfear neurons. In parallel, the reduction of activity in LA causes a disaffiliation of ITCd neurons leading to an activation by disinhibition of ITCv neurons, and simultaneous activation by BA bextinction glutamatergic neurons. Moreover, the subsets of CeL cells PKC $\delta$ + can be disinhibited leading to the inactivation of CeM neurons and less fear responses. Abbreviations: ACC, anterior cingulate cortex; PL, prelimbic cortex; IL, infralimbic cortex; LA, lateral amygdala; BA, basal amygdala; ITCd, intercalated cells dorsal; ITCv, intercalated cells ventral; CeL, centro-lateral amygdala; CeM, centro-medial amygdala; PAG, periaqueductal gray

conditioned rats resulted in facilitation of subsequent extinction (Lahoud and Maroun 2013). Furthermore, social interaction during extinction training facilitates the consolidation of extinction, and this effect is mediated by the pharmacological manipulation of OTR signaling in IL (Brill-Maoz and Maroun 2016). However, interpretation of these observation is not as straight forward as one might think, as OTRs and OT-releasing fibers are absent in this region (Knobloch et al. 2012; Mitre et al. 2016). Nonetheless, a different study showed the presence of OTR-binding sites in IL of rats using autoradiography (Hansson et al. 2018).

While OTR expression has clearly been demonstrated in rodent brain regions involved in fear extinction and fear recall, this

### Non-human primates

---

seems to be different in non-human primates, since the identification of these receptors has certain technical limitations. Here, OTRs appear rather expressed in brain areas related to the integration of visual stimuli (Freeman et al. 2014; Freeman and Young 2016) that are upstream of the fear-controlling regions. For example, the nucleus basalis of Meynert (NBM) projects to the amygdala and it is the primary source of cholinergic input to the BLA (Nagai et al. 1982; Woolf and Butcher 1982). This cholinergic input is required for memory consolidation (Power et al. 2003), possibly promoting the oxytocin signaling-dependent encoding of memory during sustained attention to visual stimuli (Freeman and Young 2016), with relevance for social- and fear-related behaviors.

In chimpanzees, OT-immunoreactive fibers have been found in the ACC, but not in that of macaque monkeys (Rogers et al. 2018). In addition, OTRs seem to be completely

absent in these animals, and then OT is taken over by the closely related vasopressin receptor 1. Exogenous OT and AVP can bind to AVP receptors in the ACC of male macaques, with behavioral consequences (Jiang and Platt 2018), suggesting that exogenous OT may shape social and fear retrieval memories in primates via binding to AVP receptors (Terashima et al. 1999; Chini et al. 2008; Loyens et al. 2012). This finding supports the direct effect of OT in the CeA of different non-human primates, where the AVP receptors are densely expressed, but not the OTRs (Young and Flanagan-Cato 2012; Freeman et al. 2014).

### **Modulation of fear circuits by oxytocin in humans**

In humans, OTR-binding sites have been found in specific relays of the fear retrieval pathway in postmortem tissue. For example, Hansson et al. (2018) observed OTR expression by autoradiography in the ACC and Uhrig et al. (2016) also showed *OTR* mRNA and OT-binding sites in Brodmann area 10, which is part of the rostromedial prefrontal cortex, and hence, of the fear retrieval pathway. Furthermore, as in the brain of chimpanzees, OT-positive fibers terminate in the ACC (Rogers et al. 2018), indicating a possible divergent evolution of the OT and AVP systems in more evolved primates. Taken together, it seems that OT signaling modulates the activity of fear retrieval and recall pathways across many species, including humans.

In humans, although the intranasal route of delivery does not allow for a precise targeting of brain regions (Leng and Ludwig 2016), it has been indeed the method of choice to deliver OT into the brain. Various key reports have even demonstrated that OT levels increase in the CSF after intranasal administration (Chang et al. 2012; Neumann et al. 2013; Striepens et al. 2013; Dal Monte et al. 2014; Modi et al. 2014). In fact, fMRI studies detail widespread changes in BOLD signals in the majority of brain regions expressing OTRs after this application, among which the amygdala subnuclei (Gamer et al. 2010; Paloyelis et al. 2016). Some of these fMRI studies highlight the anxiolytic effects, with a locus in brain areas and subnuclei that belong to the fear retrieval circuit after intranasal administration of OT (Gamer et al. 2010). As in rodents, OT has general anxiolytic acute properties in humans, decreasing the reactivity of the amygdala to fearful faces (Kirsch et al. 2005; Domes et al. 2007; Labuschagne et al. 2010). Noteworthy, intranasal administration of OT can promote differential behavioral outcomes in men and women, in terms of emotional salience (Domes et al. 2010; Rilling et al. 2014; Luo et al. 2017). Interestingly, in the ACC, activation of the dorsal part in response to fearful faces is diminished following intranasal OT application in men, but increased in women (Luo et al. 2017), indicating a sex-

dependent outcome of OT modulation in fear retrieval. This sexual dimorphism in cortical oxytocin signaling resembles that observed in the PL of mice (Li et al. 2016). Similarly, different reports show that while OT decreases amygdala reactivity to threatening faces in men, it rather enhances the same amygdala reactivity to similar threatening faces in women, when they are in the luteal phase of the menstrual cycle (Domes et al. 2010; Lischke et al. 2012). This observation suggests a sex-specific difference in OT-dependent threat processing. It is potentially due to the interaction of OTR with gonadal steroids, such as progesterone and estradiol, which have a more pronounced impact in the female brain. Adding to this idea, intranasal OT shows sex-specific functional roles via the amygdala to enhance the salience and attractiveness of positive social attributes in women but enhancing the salience of the negative ones in men (Gao et al. 2016). This is a behavioral and functional proof of how this molecule's network could have evolved in a different way in men and women, in order to display social relations and threat responses.

These studies support the idea that OTR signaling can be sex-dependent in the modulation of fear circuits, and that its role to enhance the sensitivity to salient social stimuli can be affected by sex-dependent networks, like the gonadal steroid hormones. Importantly, steroid hormones, such as estradiol and progesterone have been found to modulate the OTR: estradiol enhances OTR affinity for OT, while progesterone decreases receptor binding (Gimpl et al. 2002; Choleris et al. 2008). These hormones regulate synaptic plasticity, NMDAR function and LTP, fundamental effectors of learning and memory (Milad et al. 2009; Gillies and McArthur 2010; Shansky et al. 2010; Zeidan et al. 2011; Milad and Quirk 2012). Such gender difference might also be quite relevant for psychiatric conditions related with disrupted fear extinction, like PTSD, that are more prevalent in women than in men (Kessler et al. 2017; Ramikie and Ressler 2018). Therefore, when applying OT in humans, it is important to consider these gender differences, and this also applies to preclinical research in animal models, where these differences have not either been consistently addressed.

While research in rodents has clarified many aspects of acute anxiolytic actions of OT, and of the inhibition of fear retrieval by OT signaling, research in humans has been much more ambivalent. Besides potential differences in the way the induced behavioral and emotional changes are interpreted, this may also be due to the different methods of OT administration (Leng and Ludwig 2016; Walum et al. 2016). Indeed, human studies typically administrate oxytocin itself by intranasal application route. As a result, in these studies, OT may also reach vasopressin receptors through which it is known to often exert opposite effects, e.g., in the amygdala (Huber et al. 2005). Animal studies, in contrast, often apply the much more specific OTR agonist Thr4Gly7-OT (TGOT), or the specific OTR

---

agonist (OTA) and target these compounds through local injection to the brain regions under study.

Nonetheless, in fear extinction procedures, intranasal OT after Pavlovian fear conditioning of healthy subjects increases electrodermal responses and vmPFC signals to conditioned fear in the early phase, and decreases amygdala activity in the late phase of extinction (Eckstein et al. 2015). Similarly, intranasal OT promotes recall of fear extinction up to 24 h after fear extinction training (Acheson et al. 2013). To promote the consolidation of fear extinction, OT administered in men inhibits amygdala responses to negative stimuli, and facilitates left insula responses for subsequent memory retrieval (Striepens et al. 2012), to form a memory of aversive social stimuli (Striepens et al. 2012). This is a very fascinating discovery that has not been found before in animals and postulates that OT could have long-lasting effects in memory consolidation for socially aversive cues. In addition, Eckstein and colleagues showed that exogenous OT increases the functional connectivity of the BLA with the dmPFC, and decreases that of the CeM, output region of the amygdala for negative emotional salience (Eckstein et al. 2017). Together, these examples show that OT could modulate memory consolidation to boost fear extinction in a long-lasting manner. This makes OT a promising pharmacological agent for the treatment of fear and anxiety disorders.

However, surprisingly, exogenous administration of OT in humans can also have contrasting effects (MacDonald et al. 2013), for example, facilitating Pavlovian fear learning, correlated with heightened activity in ACC (Eckstein et al. 2016). Moreover, in experimental designs using unpredictable threats, OT promotes anxiogenic effects by enhancing the potentiated startle reflex (Grillon et al. 2013). This could be explained from the functional role of OT to bind and stimulate the BNST (Moaddab and Dabrowska 2017; Dabrowska et al. 2011; Knobloch et al. 2012; Martinon and Dabrowska 2018; Duque-Wilckens et al. 2018), a structure that underlies sustained anxiety-like behaviors to uncertainty by directly activating corticotrophin releasing factor neurons. A better understanding of the consequences of OT administration in different brain areas, and in diverse social and non-social contexts, as well as a fully characterization of the dose-response function in clinical studies is warranted. Moreover, although current data supports the inhibition of cortico-amygdaloid retrieval circuits by exogenous OT applications, more knowledge about the specific cell types or synaptic modulations in those networks are required, for example, in a clinical setting, through a PET scan that could screen OT tonic activity in the brain, as it exists for the dopaminergic signaling (Volkow et al. 2005). Indeed, there have been recent efforts to develop radioligands of OT that can be used for monitoring OT receptor distribution in humans (Smith et al. 2016a).

## Oxytocin treatment: mode of action

In addition to acting as an acute anxiolytic to diminish fear responses, mainly in the CeA and other areas (Litvin et al. 2016; Viviani et al. 2011; de la Mora et al. 2016; Sabihi et al. 2017; Grund et al. 2017; Menon et al. 2018), OT could also act on fear learning, and fear extinction learning in a long-lasting manner. In other words, it could work on neuronal plasticity in the fear-related neural networks. Unfortunately, studies describing the mechanism of OT on synaptic plasticity in the ACC, PL, IL, BLA, and CeL are still scarce. However, a recent report demonstrated that, while intranasal acute administration of OT increases cellular activity in ACC and correlated with the transfer of socially derived fear, as in Burkett et al. 2016, chronic administration of OT had similar behavioral effects that correlated with decreased OTR mRNA in the CeA (Pisansky et al. 2017). Considering that intranasal OT could also affect VP receptors, these findings should be interesting to consider in light of the opposite effects both receptors exert in the CeA on expression of fear (Huber et al. 2005).

Furthermore, OT signaling in other brain areas brings about plastic synaptic changes that rely on the OTR (Fang et al. 2008). For example, in the nucleus accumbens, OT acts pre-synaptically to induce long-term depression (LTD), which has been associated with social reward (Dölen et al. 2013; Gur et al. 2014; Marlin et al. 2015; Mitre et al. 2016). Likewise, in the medial amygdala, OT induces LTD, which is fundamental for social recognition (Gur et al. 2014). Finally, OT has been shown to promote long-term potentiation in the auditory cortex in response to vocal calls of the pups (Marlin et al. 2015; Mitre et al. 2016) and in the hippocampus of lactating mothers through activation of MEK1/2 and CREB to improve spatial learning (Tomizawa et al. 2003). Taken the results of these studies together, it appears that OT can induce plastic changes in social contexts, in line with its proposed pro-social dynamic function (Fang et al. 2008; Marlin et al. 2015; Stoop et al. 2015; Modi et al. 2015; Coria-Avila et al. 2014; Triana-Del Rio et al. 2015). Therefore, we propose that putative contributions of OT to neuronal plasticity in the fear retrieval and extinction circuits can be best studied in relation to social interactions. In agreement with this, OT is released in the CeA in lactating mother rats in the presence of their pups (i.e., a social stimulus) to reduce freezing and increase active defensive behavior to protect the pups (Rickenbacher et al. 2017). The proof of whether OT release under socially relevant conditions induces plastic changes in the circuits of fear extinction is thus an attractive question waiting to be answered.

Finally, at the clinical level, OT has successfully been used to treat PTSD, having an acute anxiolytic effect, but only in those patients with less severe symptoms (Sack et al. 2017). Anxiolysis might be explained here by restoration of the

pathologically low OT endogenous brain concentration in those patients (Donadon et al. 2018). At the circuit level, it thus appears that the increased release of endogenous OT from the PVN can activate different inhibitory neuronal populations in the brain expressing the OTR, that collaborate all together to diminish conditioned fear responses and anxiety during retrieval and, simultaneously, allow for extinction of fear memories in a social context-dependent modulatory manner.

#### Conclusion and perspectives

Recent research has not only established OT as an acute anxiolytic, but also as a modulator of long-lasting synaptic plasticity, especially in social contexts. This makes possible to combine OT with cognitive behavioral therapy for the treatment of anxiety disorders and trauma. However, the precise contribution of OT to fear extinction learning is hard to distinguish from its anxiolytic effects. This is partially due to almost a unique behavioral readout to evaluate fear responses in animal studies (freezing), and the experimental need to look for other outcomes (Blanchard et al. 1991; Myers and Davis 2007; Hegoburu et al. 2011; Wöhr and Schwarting 2013). For instance, we have found that, although OT does decrease freezing when released in the CeA, it is without effect on other parameters of fear, especially heart rate variability (Viviani et al. 2011). Instead, OT might induce a switch from passive to active defensive behavior as demonstrated in lactating mothers (Bosch et al. 2005; Rickenbacher et al. 2017), in the presence of social stimuli. Thus, there is still a lot to be discovered about the role of OT in anxiety-like behavior, not in the least about plastic long-lasting changes in the cortico- limbic fear memory circuits.

Altogether, taking in consideration its region-specific and sex-specific effects, OT presents itself as a candidate molecule to facilitate fear extinction and anxiolysis, with the advantages that it is easy to apply, is endogenous, and its potential use is based on well-characterized neural circuitry controlling fear and anxiety responses, in animal models as well as in humans.

Funding information RT is supported by the Synapsis Foundation, CH by a Marie-Heim Vögtlin grant from the Swiss National Science Foundation, and EvdB by a Swiss Federal grant from the Commission of Technology and Innovation.

## References

Abramowitz JS (2013) The practice of exposure therapy: relevance of cognitive-behavioral theory and extinction theory. *Behav Ther* 44: 548-558. <https://doi.org/10.1016/j.beth.2013.03.003>

Acheson D, Feifel D, de Wilde S et al (2013) The effect of intranasal oxytocin treatment on conditioned fear extinction and recall in a healthy human sample. *Psychopharmacology* 229:199-208. <https://doi.org/10.1007/s00213-013-3099-4>

Alberini CM (ed) (2013) *Memory reconsolidation*, 1 edition. Academic press, London

Allsop SA, Wichmann R, Mills F et al (2018) Corticoamygdala transfer of socially derived information gates observational learning. *Cell* 173:1329-1342.e18. <https://doi.org/10.1016/j.cell.2018.04.004>

Amano T, Unal CT, Paré D (2010) Synaptic correlates of fear extinction in the amygdala. *Nat Neurosci* 13:489-494. <https://doi.org/10.1038/nn.2499>

Amano T, Amir A, Goswami S, Paré D (2012) Morphology, PKC $\delta$  expression, and synaptic responsiveness of different types of rat central lateral amygdala neurons. *J Neurophysiol* 108:3196-3205. <https://doi.org/10.1152/jn.00514.2012>

Amir A, Amano T, Paré D (2011) Physiological identification and infralimbic responsiveness of rat intercalated amygdala neurons. *J Neurophysiol* 105:3054-3066. <https://doi.org/10.1152/jn.00136.2011>

Anderson DJ, Adolphs R (2014) A framework for studying emotions across species. *Cell* 157:187-200. <https://doi.org/10.1016/j.cell.2014.03.003>

Anderson KC, Insel TR (2006) The promise of extinction research for the prevention and treatment of anxiety disorders. *Biol Psychiatry* 60: 319-321. <https://doi.org/10.1016/j.biopsych.2006.06.022>

Andreescu C, Tudorascu D, Sheu LK et al (2017) Brain structural changes in late-life generalized anxiety disorder. *Psychiatry Res* 268:15-21. <https://doi.org/10.1016/j.psychres.2017.08.004>

Bentz D, Michael T, de Quervain DJ-F, Wilhelm FH (2010) Enhancing exposure therapy for anxiety disorders with glucocorticoids: from basic mechanisms of emotional learning to clinical applications. *J Anxiety Disord* 24:223-230. <https://doi.org/10.1016/j.janxdis.2009.10.011>

Blanchard DC, Weatherspoon A, Shepherd J et al (1991) Paradoxical effects of morphine on antipredator defense reactions in wild and laboratory rats. *Pharmacol Biochem Behav* 40:819-828

Blume A, Bosch OJ, Miklos S et al (2008) Oxytocin reduces anxiety via ERK1/2 activation: local effect within the rat hypothalamic paraventricular nucleus. *Eur J Neurosci* 27:1947-1956. <https://doi.org/10.1111/j.1460-9568.2008.06184.x>

Bosch OJ, Meddle SL, Beiderbeck DI et al (2005) Brain oxytocin correlates with maternal aggression: link to anxiety. *J Neurosci* 25:6807-6815. <https://doi.org/10.1523/JNEUROSCI.1342-05.2005>

Bowers ME, Ressler KJ (2015a) Interaction between the cholecystokinin and endogenous cannabinoid systems in cued fear expression and extinction retention. *Neuropsychopharmacology* 40:688-700. <https://doi.org/10.1038/npp.2014.225>

Bowers ME, Ressler KJ (2015b) An overview of translationally informed treatments for posttraumatic stress disorder: animal models of pavlovian fear conditioning to human clinical trials. *Biol Psychiatry* 78: E15-E27. <https://doi.org/10.1016/j.biopsych.2015.06.008>

Brill-Maoz N, Maroun M (2016) Extinction of fear is facilitated by social presence: synergism with prefrontal oxytocin. *Psychoneuroendocrinology* 66:75-81. <https://doi.org/10.1016/j.psyneuen.2016.01.003>

Bukalo O, Pinard CR, Holmes A (2014) Mechanisms to medicines: elucidating neural and molecular substrates of fear extinction to identify novel treatments for anxiety disorders. *Br J Pharmacol* 171:4690-4718. <https://doi.org/10.1111/bph.12779>

Bukalo O, Pinard CR, Silverstein S, et al (2015) Prefrontal inputs to the amygdala instruct fear extinction memory formation. *Sci Adv* 1(6): pii: e1500251

Burghardt NS, Bauer EP (2013) Acute and chronic effects of selective serotonin reuptake inhibitor treatment on fear conditioning: implications for underlying fear circuits. *Neuroscience* 247:253-272. <https://doi.org/10.1016/j.neuroscience.2013.05.050>

Burgos-Robles A, Kimchi EY, Izadmehr EM et al (2017) Amygdala inputs to prefrontal cortex guide behavior amid conflicting cues of reward and punishment. *Nat Neurosci* 20:824-835. <https://doi.org/10.1038/nn.4553>

- Burkett JP, Andari E, Johnson ZV et al (2016) Oxytocin-dependent consolation behavior in rodents. *Science* 351:375-378. <https://doi.org/10.1126/science.aac4785>
- Busti D, Geracitano R, Whittle N et al (2011) Different fear states engage distinct networks within the intercalated cell clusters of the amygdala. *J Neurosci* 31:5131-5144. <https://doi.org/10.1523/JNEUROSCI.6100-10.2011>
- Campbell BM, Merchant KM (2003) Serotonin 2C receptors within the basolateral amygdala induce acute fear-like responses in an open-field environment. *Brain Res* 993:1-9
- Campbell-Smith EJ, Holmes NM, Lingawi NW et al (2015) Oxytocin signaling in basolateral and central amygdala nuclei differentially regulates the acquisition, expression, and extinction of context-conditioned fear in rats. *Learn Mem* 22:247-257. <https://doi.org/10.1101/lm.036962.114>
- Carpenter JK, Andrews LA, Witcraft SM et al (2018) Cognitive behavioral therapy for anxiety and related disorders: a meta-analysis of randomized placebo-controlled trials. *Depress Anxiety* 35(6):502-514. <https://doi.org/10.1002/da.22728>
- Cassell EJ (1999) Diagnosing suffering: a perspective. *Ann Intern Med* 131:531-534
- Chang SWC, Barter JW, Ebitz RB et al (2012) Inhaled oxytocin amplifies both vicarious reinforcement and self reinforcement in rhesus macaques (*Macaca mulatta*). *Proc Natl Acad Sci U S A* 109:959-964. <https://doi.org/10.1073/pnas.1114621109>
- Chini B, Manning M, Guillon G (2008) Affinity and efficacy of selective agonists and antagonists for vasopressin and oxytocin receptors: an easy guide to receptor pharmacology. *Prog Brain Res* 170:513-517. [https://doi.org/10.1016/S0079-6123\(08\)00438-X](https://doi.org/10.1016/S0079-6123(08)00438-X)
- Cho J-H, Deisseroth K, Bolshakov VY (2013) Synaptic encoding of fear extinction in mPFC-amygdala circuits. *Neuron* 80:1491-1507. <https://doi.org/10.1016/j.neuron.2013.09.025>
- Choleris E, Devidze N, Kavaliers M, Pfaff DW (2008) Steroidal/neuropeptide interactions in hypothalamus and amygdala related to social anxiety. *Prog Brain Res* 170:291-303. [https://doi.org/10.1016/S0079-6123\(08\)00424-X](https://doi.org/10.1016/S0079-6123(08)00424-X)
- Ciocchi S, Herry C, Grenier F et al (2010) Encoding of conditioned fear in central amygdala inhibitory circuits. *Nature* 468:277-282. <https://doi.org/10.1038/nature09559>
- Cisler JM, Sigel BA, Steele JS, Smitherman S, et al (2016) Changes in functional connectivity of the amygdala during cognitive reappraisal predict symptom reduction during trauma-focused cognitive-behavioral therapy among adolescent girls with post-traumatic stress disorder. *Psychol Med* 46(14):3013-3023
- Clark RE, Xie H, Brunette MF (2004) Benzodiazepine prescription practices and substance abuse in persons with severe mental illness. *J Clin Psychiatry* 65:151-155
- Coria-Avila GA, Manzo J, Garcia LI et al (2014) Neurobiology of social attachments. *Neurosci Biobehav Rev* 43:173-182. <https://doi.org/10.1016/j.neubiorev>
- Courtin J, Chaudun F, Rozeske RR et al (2014) Prefrontal parvalbumin interneurons shape neuronal activity to drive fear expression. *Nature* 505:92-96. <https://doi.org/10.1038/nature12755>
- Cullen PK, Gilman TL, Winiecki P et al (2015) Activity of the anterior cingulate cortex and ventral hippocampus underlie increases in contextual fear generalization. *Neurobiol Learn Mem* 124:19-27. <https://doi.org/10.1016/j.nlm.2015.07.001>
- Dabrowska J, Hazra R, Ahern TH et al (2011) Neuroanatomical evidence for reciprocal regulation of the corticotrophin-releasing factor and oxytocin systems in the hypothalamus and the bed nucleus of the stria terminalis of the rat: implications for balancing stress and affect. *Psychoneuroendocrinology* 36:1312-1326. <https://doi.org/10.1016/j.psyneuen.2011.03.003>
- Dal Monte O, Noble PL, Turchi J et al (2014) CSF and blood oxytocin concentration changes following intranasal delivery in macaque. *PLoS One* 9:e103677. <https://doi.org/10.1371/journal.pone.0103677>
- Davis M (1992) The role of the amygdala in fear and anxiety. *Annu Rev Neurosci* 15:353-375. <https://doi.org/10.1146/annurev.ne.15.030192.002033>
- Davis M, Myers KM, Chhatwal J, Ressler KJ (2006) Pharmacological treatments that facilitate extinction of fear: relevance to psychotherapy. *NeuroRx* 3:82-96. <https://doi.org/10.1016/j.nurx.2005.12.008>
- Davis S, Renaudineau S, Poirier R et al (2010) The formation and stability of recognition memory: what happens upon recall? *Front Behav Neurosci* 4:177. <https://doi.org/10.3389/fnbeh.2010.00177>
- de la Mora MP, Pérez-Carrera D, Crespo-Ramírez M et al (2016) Signaling in dopamine D2 receptor- oxytocin receptor heterocomplexes and its relevance for the anxiolytic effects of dopamine and oxytocin interactions in the amygdala of the rat. *Biochim Biophys Acta* 1862:2075-2085. <https://doi.org/10.1016/j.bbadis.2016.07.004>
- Debiec J, LeDoux JE (2004) Disruption of reconsolidation but not consolidation of auditory fear conditioning by noradrenergic blockade in the amygdala. *Neuroscience* 129:267-272. <https://doi.org/10.1016/j.neuroscience.2004.08.018>
- Debiec J, LeDoux JE (2006) Noradrenergic signaling in the amygdala contributes to the reconsolidation of fear memory: treatment implications for PTSD. *Ann NY Acad Sci* 1071:521-524. <https://doi.org/10.1196/annals.1364.056>
- Dejean C, Courtin J, Karalis N et al (2016) Prefrontal neuronal assemblies temporally control fear behaviour. *Nature* 535:420-424. <https://doi.org/10.1038/nature18630>
- Dilgen J, Tejada HA, O'Donnell P (2013) Amygdala inputs drive feedforward inhibition in the medial prefrontal cortex. *J Neurophysiol* 110:221-229. <https://doi.org/10.1152/jn.00531.2012>
- Dölen G, Darvishzadeh A, Huang KW, Malenka RC (2013) Social reward requires coordinated activity of nucleus accumbens oxytocin and serotonin. *Nature* 501:179-184. <https://doi.org/10.1038/nature12518>
- Domes G, Heinrichs M, Gläscher J et al (2007) Oxytocin attenuates amygdala responses to emotional faces regardless of valence. *Biol Psychiatry* 62:1187-1190. <https://doi.org/10.1016/j.biopsych.2007.03.025>
- Domes G, Lischke A, Berger C et al (2010) Effects of intranasal oxytocin on emotional face processing in women. *Psychoneuroendocrinology* 35:83-93. <https://doi.org/10.1016/j.psyneuen.2009.06.016>
- Do-Monte FH, Manzano-Nieves G, Quiñones-Laracuente K et al (2015) Revisiting the role of infralimbic cortex in fear extinction with optogenetics. *J Neurosci* 35:3607-3615. <https://doi.org/10.1523/JNEUROSCI.3137-14.2015>
- Donadon MF, Martin-Santos R, Osório F de L (2018) The associations between oxytocin and trauma in humans: a systematic review. *Front Pharmacol* 9:154. <https://doi.org/10.3389/fphar.2018.00154>
- Duke AN, Meng Z, Platt DM et al (2018) Evidence that sedative effects of benzodiazepines involve unexpected GABA<sub>A</sub> receptor subtypes: quantitative observation studies in rhesus monkeys. *J Pharmacol Exp Ther* 366:145-157. <https://doi.org/10.1124/jpet.118.249250>
- Duque-Wilckens N, Steinman MQ, Busnelli M et al (2018) Oxytocin receptors in the anteromedial bed nucleus of the stria terminalis promote stress-induced social avoidance in female California mice. *Biol Psychiatry* 83:203-213. <https://doi.org/10.1016/j.biopsych.2017.08.024>
- Duvarci S, Pare D (2014) Amygdala microcircuits controlling learned fear. *Neuron* 82:966-980. <https://doi.org/10.1016/j.neuron.2014.04.042>
- Duvarci S, Popa D, Paré D (2011) Central amygdala activity during fear conditioning. *J Neurosci* 31:289-294. <https://doi.org/10.1523/JNEUROSCI.4985-10.2011>

- Eckstein M, Becker B, Scheele D et al (2015) Oxytocin facilitates the extinction of conditioned fear in humans. *Biol Psychiatry* 78:194-202. <https://doi.org/10.1016/j.biopsych.2014.10.015>
- Eckstein M, Scheele D, Patin A et al (2016) Oxytocin facilitates Pavlovian fear learning in males. *Neuropsychopharmacology* 41(4):932-939. <https://doi.org/10.1038/npp.2015.245>
- Eckstein M, Markett S, Kendrick KM et al (2017) Oxytocin differentially alters resting state functional connectivity between amygdala subregions and emotional control networks: inverse correlation with depressive traits. *Neuroimage* 149:458-467. <https://doi.org/10.1016/j.neuroimage.2017.01.078>
- Ehrlich I, Humeau Y, Grenier F et al (2009) Amygdala inhibitory circuits and the control of fear memory. *Neuron* 62:757-771. <https://doi.org/10.1016/j.neuron.2009.05.026>
- Eliava M, Melchior M, Knobloch-Bollmann HS, Wahis J et al (2016) A new population of parvocellular oxytocin neurons controlling magnocellular neuron activity and inflammatory pain processing. *Neuron* 89(6):1291-1304. <https://doi.org/10.1016/j.neuron>
- Fang L-Y, Quan R-D, Kaba H (2008) Oxytocin facilitates the induction of long-term potentiation in the accessory olfactory bulb. *Neurosci Lett* 438:133-137. <https://doi.org/10.1016/j.neulet.2007.12.070>
- Fitzgerald PJ, Seemann JR, Maren S (2014) Can fear extinction be enhanced? A review of pharmacological and behavioral findings. *Brain Res Bull* 105:46-60. <https://doi.org/10.1016/j.brainresbull.2013.12.007>
- Freeman SM, Young LJ (2016) Comparative perspectives on oxytocin and vasopressin receptor research in rodents and primates: translational implications. *J Neuroendocrinol* 28. <https://doi.org/10.1111/jne.12382>
- Freeman SM, Inoue K, Smith AL et al (2014) The neuroanatomical distribution of oxytocin receptor binding and mRNA in the male rhesus macaque (*Macaca mulatta*). *Psychoneuroendocrinology* 45:128-141. <https://doi.org/10.1016/j.psyneuen.2014.03.023>
- Gabbott PLA, Warner TA, Busby SJ (2006) Amygdala input monosynaptically innervates parvalbumin immunoreactive local circuit neurons in rat medial prefrontal cortex. *Neuroscience* 139:1039-1048. <https://doi.org/10.1016/j.neuroscience.2006.01.026>
- Gamer M, Zurowski B, Büchel C (2010) Different amygdala subregions mediate valence-related and attentional effects of oxytocin in humans. *Proc Natl Acad Sci U S A* 107:9400-9405. <https://doi.org/10.1073/pnas.1000985107>
- Gao S, Becker B, Luo L et al (2016) Oxytocin, the peptide that bonds the sexes also divides them. *Proc Natl Acad Sci U S A* 113:7650-7654. <https://doi.org/10.1073/pnas.1602620113>
- Geracitano R, Kaufmann WA, Szabo G et al (2007) Synaptic heterogeneity between mouse paracapsular intercalated neurons of the amygdala. *J Physiol Lond* 585:117-134. <https://doi.org/10.1113/jphysiol.2007.142570>
- Gillies GE, McArthur S (2010) Estrogen actions in the brain and the basis for differential action in men and women: a case for sex-specific medicines. *Pharmacol Rev* 62:155-198. <https://doi.org/10.1124/pr.109.002071>
- Gimpl G, Wiegand V, Burger K, Fahrenholz F (2002) Cholesterol and steroid hormones: modulators of oxytocin receptor function. *Prog Brain Res* 139:43-55
- Giustino TF, Maren S (2015) The role of the medial prefrontal cortex in the conditioning and extinction of fear. *Front Behav Neurosci* 9:298. <https://doi.org/10.3389/fnbeh.2015.00298>
- Giustino TF, Fitzgerald PJ, Maren S (2016) Revisiting propranolol and PTSD: memory erasure or extinction enhancement? *Neurobiol Learn Mem* 130:26-33. <https://doi.org/10.1016/j.nlm.2016.01.009>
- Goodman AM, Harnett NG, Knight DC (2018) Pavlovian conditioned diminution of the neurobehavioral response to threat. *Neurosci Biobehav Rev* 84:218-224. <https://doi.org/10.1016/j.neubiorev.2017.11.021>
- Grillon C, Krinsky M, Charney DR et al (2013) Oxytocin increases anxiety to unpredictable threat. *Mol Psychiatry* 18:958-960. <https://doi.org/10.1038/mp.2012.156>
- Gruene TM, Flick K, Stefano A, et al (2015) Sexually divergent expression of active and passive conditioned fear responses in rats. *Elife* 4. doi: <https://doi.org/10.7554/eLife.11352>
- Grund T, Goyon S, Li Y et al (2017) Neuropeptide S activates paraventricular oxytocin neurons to induce anxiolysis. *J Neurosci* 37:12214-12225. <https://doi.org/10.1523/JNEUROSCI.2161-17.2017>
- Gur R, Tendler A, Wagner S (2014) Long-term social recognition memory is mediated by oxytocin-dependent synaptic plasticity in the medial amygdala. *Biol Psychiatry* 76:377-386. <https://doi.org/10.1016/j.biopsych.2014.03.022>
- Hansson AC, Koopmann A, Uhrig S et al (2018) Oxytocin reduces alcohol cue-reactivity in alcohol-dependent rats and humans. *Neuropsychopharmacology* 43:1235-1246. <https://doi.org/10.1038/npp.2017.257>
- Haubensak W, Kunwar PS, Cai H et al (2010) Genetic dissection of an amygdala microcircuit that gates conditioned fear. *Nature* 468:270-276. <https://doi.org/10.1038/nature09553>
- Hegoburu C, Shionoya K, Garcia S et al (2011) The RUB cage: respiration-ultrasonic vocalizations-behavior acquisition setup for assessing emotional memory in rats. *Front Behav Neurosci* 5:25. <https://doi.org/10.3389/fnbeh.2011.00025>
- Herry C, Ciochi S, Senn V et al (2008) Switching on and off fear by distinct neuronal circuits. *Nature* 454:600-606. <https://doi.org/10.1038/nature07166>
- Hofmann SG, Otto MW, Pollack MH, Smits JA (2015) D-cycloserine augmentation of cognitive behavioral therapy for anxiety disorders: an update. *Curr Psychiatry Rep* 17:532. <https://doi.org/10.1007/s11920-014-0532-2>
- Hoover WB, Vertes RP (2007) Anatomical analysis of afferent projections to the medial prefrontal cortex in the rat. *Brain Struct Funct* 212:149-179. <https://doi.org/10.1007/s00429-007-0150-4>
- Huber D, Veinante P, Stoop R (2005) Vasopressin and oxytocin excite distinct neuronal populations in the central amygdala. *Science* 308:245-248. <https://doi.org/10.1126/science.1105636>
- Ishikawa A, Nakamura S (2003) Convergence and interaction of hippocampal and amygdalar projections within the prefrontal cortex in the rat. *J Neurosci* 23:9987-9995
- Jiang Y, Platt ML (2018) Oxytocin and vasopressin flatten dominance hierarchy and enhance behavioral synchrony in part via anterior cingulate cortex. *Sci Rep* 8:8201. <https://doi.org/10.1038/s41598-018-25607-1>
- Jiménez A, Young LJ, Triana-Del Río R, LaPrairie JL et al (2015) Neuroanatomical distribution of oxytocin receptor binding in the female rabbit forebrain: variations across the reproductive cycle. *Brain Res* 1629:329-339. <https://doi.org/10.1016/j.brainres>
- Jurek B, Slaterry DA, Maloumy R et al (2012) Differential contribution of hypothalamic MAPK activity to anxiety-like behaviour in virgin and lactating rats. *PLoS One* 7:e37060. <https://doi.org/10.1371/journal.pone.0037060>
- Kaoru T, Liu F-C, Ishida M et al (2010) Molecular characterization of the intercalated cell masses of the amygdala: implications for the relationship with the striatum. *Neuroscience* 166:220-230. <https://doi.org/10.1016/j.neuroscience.2009.12.004>
- Karalis N, Dejean C, Chaudun F et al (2016) 4-Hz oscillations synchronize prefrontal-amygdala circuits during fear behavior. *Nat Neurosci* 19:605-612. <https://doi.org/10.1038/nn.4251>
- Kessler RC, Aguilar-Gaxiola S, Alonso J et al (2017) The associations of earlier trauma exposures and history of mental disorders with PTSD after subsequent traumas. *Mol Psychiatry*. <https://doi.org/10.1038/mp.2017.194>
- Kim JH, Richardson R (2010) New findings on extinction of conditioned fear early in development: theoretical and clinical implications. *Biol*

- Psychiatry 67:297-303. <https://doi.org/10.1016/j.biopsych.2009.09.003>
- Kirsch P, Esslinger C, Chen Q et al (2005) Oxytocin modulates neural circuitry for social cognition and fear in humans. *J Neurosci* 25: 11489-11493. <https://doi.org/10.1523/JNEUROSCI.3984-05.2005>
- Klump H, Fitzgerald JM, Kinney KL et al (2017) Predicting cognitive behavioral therapy response in social anxiety disorder with anterior cingulate cortex and amygdala during emotion regulation. *Neuroimage Clin* 15:25-34. <https://doi.org/10.1016/j.nicl.2017.04.006>
- Knobloch HS, Charlet A, Hoffmann LC et al (2012) Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron* 73:553-566. <https://doi.org/10.1016/j.neuron.2011.11.030>
- Knoflach F, Hernandez M-C, Bertrand D (2016) GABAA receptor-mediated neurotransmission: not so simple after all. *Biochem Pharmacol* 115:10-17. <https://doi.org/10.1016/j.bcp.2016.03.014>
- Krettek JE, Price JL (1978) Amygdaloid projections to subcortical structures within the basal forebrain and brainstem in the rat and cat. *J Comp Neurol* 178:225-254. <https://doi.org/10.1002/cne.901780204>
- Kroes MCW, Schiller D, LeDoux JE, Phelps EA (2016) Translational approaches targeting reconsolidation. *Curr Top Behav Neurosci* 28:197-230. [https://doi.org/10.1007/7854\\_2015\\_5008](https://doi.org/10.1007/7854_2015_5008)
- Labuschagne I, Phan KL, Wood A et al (2010) Oxytocin attenuates amygdala reactivity to fear in generalized social anxiety disorder. *Neuropsychopharmacology* 35:2403-2413. <https://doi.org/10.1038/npp.2010.123>
- Lahoud N, Maroun M (2013) Oxytocinergic manipulations in corticolimbic circuit differentially affect fear acquisition and extinction. *Psychoneuroendocrinology* 38:2184-2195. <https://doi.org/10.1016/j.psyneuen.2013.04.006>
- Ledgerwood L, Richardson R, Cranney J (2005) D-cycloserine facilitates extinction of learned fear: effects on reacquisition and generalized extinction. *Biol Psychiatry* 57:841-847. <https://doi.org/10.1016/j.biopsych.2005.01.023>
- LeDoux JE (2000) Emotion circuits in the brain. *Annu Rev Neurosci* 23: 155-184. <https://doi.org/10.1146/annurev.neuro.23.1.155>
- LeDoux JE (2014) Coming to terms with fear. *Proc Natl Acad Sci U S A* 111:2871-2878. <https://doi.org/10.1073/pnas.1400335111>
- LeDoux JE, Iwata J, Cicchetti P, Reis DJ (1988) Different projections of the central amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear. *J Neurosci* 8:2517-2529
- Leng G, Ludwig M (2016) Intranasal oxytocin: myths and delusions. *Biol Psychiatry* 79:243-250. <https://doi.org/10.1016/j.biopsych.2015.05.003>
- Levar N, van Leeuwen JMC, Denys D, van Wingen GA (2017a) Divergent influences of anterior cingulate cortex GABA concentrations on the emotion circuitry. *Neuroimage* 158:136-144. <https://doi.org/10.1016/j.neuroimage.2017.06.055>
- Levar N, van Leeuwen JMC, Puts NAJ et al (2017b) GABA concentrations in the anterior cingulate cortex are associated with fear network function and fear recovery in humans. *Front Hum Neurosci* 11:202. <https://doi.org/10.3389/fnhum.2017.00202>
- Li H, Penzo MA, Taniguchi H, Kopec CD, Huang ZJ, Li B (2013a) Experience-dependent modification of a central amygdala fear circuit. *Nat Neurosci* 16(3):332-339. <https://doi.org/10.1038/nn.3322>
- Li Y, Meloni EG, Carlezon WA et al (2013b) Learning and reconsolidation implicate different synaptic mechanisms. *Proc Natl Acad Sci U S A* 110:4798-4803. <https://doi.org/10.1073/pnas.1217878110>
- Li K, Nakajima M, Ibañez-Tallon I, Heintz N (2016) A cortical circuit for sexually dimorphic oxytocin-dependent anxiety behaviors. *Cell* 167:60-72.e11. <https://doi.org/10.1016/j.cell.2016.08.067>
- Likhtik E, Stujenske JM, Topiwala MA et al (2014) Prefrontal entrainment of amygdala activity signals safety in learned fear and innate anxiety. *Nat Neurosci* 17:106-113. <https://doi.org/10.1038/nn.3582>
- Lischke A, Berger C, Prehn K et al (2012) Intranasal oxytocin enhances emotion recognition from dynamic facial expressions and leaves eye-gaze unaffected. *Psychoneuroendocrinology* 37:475-481. <https://doi.org/10.1016/j.psyneuen.2011.07.015>
- Little JP, Carter AG (2013) Synaptic mechanisms underlying strong reciprocal connectivity between the medial prefrontal cortex and basolateral amygdala. *J Neurosci* 33:15333-15342. <https://doi.org/10.1523/JNEUROSCI.2385-13.2013>
- Litvin Y, Turner CA, Rios MB et al (2016) Fibroblast growth factor 2 alters the oxytocin receptor in a developmental model of anxiety-like behavior in male rat pups. *Horm Behav* 86:64-70. <https://doi.org/10.1016/j.yhbeh.2011.11.059>
- Loyens E, Vermoesen K, Schallier A et al (2012) Proconvulsive effects of oxytocin in the generalized pentylenetetrazol mouse model are mediated by vasopressin 1a receptors. *Brain Res* 1436:43-50. <https://doi.org/10.1016/j.brainres.2011.11.059>
- Luo L, Becker B, Geng Y et al (2017) Sex-dependent neural effect of oxytocin during subliminal processing of negative emotion faces. *Neuroimage* 162:127-137. <https://doi.org/10.1016/j.neuroimage.2017.08.079>
- MacDonald K, MacDonald TM, Brüne M et al (2013) Oxytocin and psychotherapy: a pilot study of its physiological, behavioral and subjective effects in males with depression. *Psychoneuroendocrinology* 38:2831-2843. <https://doi.org/10.1016/j.psyneuen.2013.05.014>
- Maren S (2001) Neurobiology of Pavlovian fear conditioning. *Annu Rev Neurosci* 24:897-931. <https://doi.org/10.1146/annurev.neuro.24.1.897>
- Marlin BJ, Mitre M, D'Amour JA et al (2015) Oxytocin enables maternal behavior by balancing cortical inhibition. *Nature* 520:499-504. <https://doi.org/10.1038/nature14402>
- Martinon D, Dabrowska J (2018) Corticotropin-releasing factor receptors modulate oxytocin release in the dorsolateral bed nucleus of the stria terminalis (BNST) in male rats. *Front Neurosci* 12:183. <https://doi.org/10.3389/fnins.2018.00183>
- Mataix-Cols D, Fernández de la Cruz L, Monzani B et al (2017) D-Cycloserine augmentation of exposure-based cognitive behavior therapy for anxiety, obsessive-compulsive, and posttraumatic stress disorders: a systematic review and meta-analysis of individual participant data. *JAMA Psychiatry* 74:501-510. <https://doi.org/10.1001/jamapsychiatry.2016.3955>
- McGarry LM, Carter AG (2017) Prefrontal cortex drives distinct projection neurons in the basolateral amygdala. *Cell Rep* 21:1426-1433. <https://doi.org/10.1016/j.celrep.2017.10.046>
- Meins M, Herry C, Müller C et al (2010) Impaired fear extinction in mice lacking protease nexin-1. *Eur J Neurosci* 31:2033-2042. <https://doi.org/10.1111/j.1460-9568.2010.07221.x>
- Menon R, Grund T, Zoicas I et al (2018) Oxytocin signaling in the lateral septum prevents social fear during lactation. *Curr Biol* 28:1066-1078.e6. <https://doi.org/10.1016/j.cub.2018.02.044>
- Milad MR, Quirk GJ (2002) Neurons in medial prefrontal cortex signal memory for fear extinction. *Nature* 420:70-74. <https://doi.org/10.1038/nature01138>
- Milad MR, Quirk GJ (2012) Fear extinction as a model for translational neuroscience: ten years of progress. *Annu Rev Psychol* 63:129-151. <https://doi.org/10.1146/annurev.psych.121208.131631>
- Milad MR, Rauch SL, Pitman RK, Quirk GJ (2006) Fear extinction in rats: implications for human brain imaging and anxiety disorders. *Biol Psychol* 73:61-71. <https://doi.org/10.1016/j.biopsycho.2006.01.008>
- Milad MR, Pitman RK, Ellis CB et al (2009) Neurobiological basis of failure to recall extinction memory in posttraumatic stress disorder. *Biol Psychiatry* 66:1075-1082. <https://doi.org/10.1016/j.biopsych.2009.06.026>



- Mitre M, Marlin BJ, Schiavo JK et al (2016) A distributed network for social cognition enriched for oxytocin receptors. *J Neurosci* 36: 2517-2535. <https://doi.org/10.1523/JNEUROSCI.2409-15.2016>
- Moaddab M, Dabrowska J (2017) Oxytocin receptor neurotransmission in the dorsolateral bed nucleus of the stria terminalis facilitates the acquisition of cued fear in the fear-potentiated startle paradigm in rats. *Neuropharmacology* 121:130-139. <https://doi.org/10.1016/j.neuropharm>
- Modi ME, Connor-Stroud F, Landgraf R et al (2014) Aerosolized oxytocin increases cerebrospinal fluid oxytocin in rhesus macaques. *Psychoneuroendocrinology* 45:49-57. <https://doi.org/10.1016/j.psyneuen.2014.02.011>
- Modi ME, Inoue K, Barrett CE, Kittelberger KA et al (2015) Melanocortin receptor agonists facilitate oxytocin-dependent partner preference formation in the prairie vole. *Neuropsychopharmacology* 40(8):1856-1865. <https://doi.org/10.1038/npp>
- Myers KM, Davis M (2007) Mechanisms of fear extinction. *Mol Psychiatry* 12:120-150. <https://doi.org/10.1038/sj.mp.4001939>
- Nader K, Schafe GE, Le Doux JE (2000) Fear memories require protein synthesis in the amygdala for reconsolidation after retrieval. *Nature* 406:722-726. <https://doi.org/10.1038/35021052>
- Nagai T, Kimura H, Maeda T et al (1982) Cholinergic projections from the basal forebrain of rat to the amygdala. *J Neurosci* 2:513-520
- Nakajima M, Görlich A, Heintz N (2014) Oxytocin modulates female sociosexual behavior through a specific class of prefrontal cortical interneurons. *Cell* 159:295-305. <https://doi.org/10.1016/j.cell.2014.09.020>
- Neumann ID, Maloumy R, Beiderbeck DI et al (2013) Increased brain and plasma oxytocin after nasal and peripheral administration in rats and mice. *Psychoneuroendocrinology* 38:1985-1993. <https://doi.org/10.1016/j.psyneuen.2013.03.003>
- Orsini CA, Maren S (2012) Neural and cellular mechanisms of fear and extinction memory formation. *Neurosci Biobehav Rev* 36:1773-1802. <https://doi.org/10.1016/j.neubiorev.2011.12.014>
- Otto MW, Bruce SE, Deckersbach T (2005) Benzodiazepine use, cognitive impairment, and cognitive-behavioral therapy for anxiety disorders: issues in the treatment of a patient in need. *J Clin Psychiatry* 66(Suppl 2):34-38
- Owen SF, Tuncdemir SN, Bader PL et al (2013) Oxytocin enhances hippocampal spike transmission by modulating fast-spiking interneurons. *Nature* 500:458-462. <https://doi.org/10.1038/nature12330>
- Owens MJ, Morgan WN, Plott SJ, Nemeroff CB (1997) Neurotransmitter receptor and transporter binding profile of antidepressants and their metabolites. *J Pharmacol Exp Ther* 283:1305-1322
- Paloyelis Y, Krahé C, Maltezos S, et al (2016) The analgesic effect of oxytocin in humans: a double-blind, placebo-controlled cross-over study using laser-evoked potentials. *J Neuroendocrinol* 28. doi: <https://doi.org/10.1111/jne.12347>
- Pape H-C, Paré D (2010) Plastic synaptic networks of the amygdala for the acquisition, expression, and extinction of conditioned fear. *Physiol Rev* 90:419-463. <https://doi.org/10.1152/physrev.00037.2009>
- Paré D, Quirk GJ, Ledoux JE (2004) New vistas on amygdala networks in conditioned fear. *J Neurophysiol* 92:1-9. <https://doi.org/10.1152/jn.00153.2004>
- Pavlov IP (1927) *Conditioned reflexes*. Oxford University Press
- Pisansky MT, Hanson LR, Gottesman II, Gewirtz JC (2017) Oxytocin enhances observational fear in mice. *Nat Commun* 8:2102. <https://doi.org/10.1038/s41467-017-02279-5>
- Power AE, Vazdarjanova A, McGaugh JL (2003) Muscarinic cholinergic influences in memory consolidation. *Neurobiol Learn Mem* 80:178-193
- Quirk GJ, Milad MR (2010) Neuroscience: editing out fear. *Nature* 463: 36-37. <https://doi.org/10.1038/463036a>
- Quirk GJ, Mueller D (2008) Neural mechanisms of extinction learning and retrieval. *Neuropharmacology* 33:56-72. <https://doi.org/10.1038/sj.npp.1301555>
- Quirk GJ, Likhtik E, Pelletier JG, Paré D (2003) Stimulation of medial prefrontal cortex decreases the responsiveness of central amygdala output neurons. *J Neurosci* 23:8800-8807
- Quirk GJ, Garcia R, González-Lima F (2006) Prefrontal mechanisms in extinction of conditioned fear. *Biol Psychiatry* 60:337-343. <https://doi.org/10.1016/j.biopsych.2006.03.010>
- Ramikie TS, Ressler KJ (2018) Mechanisms of sex differences in fear and posttraumatic stress disorder. *Biol Psychiatry* 83:876-885. <https://doi.org/10.1016/j.biopsych.2017.11.016>
- Rashid AJ, Yan C, Mercaldo Vet al (2016) Competition between engrams influences fear memory formation and recall. *Science* 353:383-387. <https://doi.org/10.1126/science.aaf0594>
- Rescorla RA (2004) Spontaneous recovery varies inversely with the training-extinction interval. *Learn Behav* 32:401-408
- Rickenbacher E, Perry RE, Sullivan RM, Moita MA (2017) Freezing suppression by oxytocin in central amygdala allows alternate defensive behaviours and mother-pup interactions. *Elife* 6. doi: <https://doi.org/10.7554/eLife.24080>
- Rilling JK, Demarco AC, Hackett PD et al (2014) Sex differences in the neural and behavioral response to intranasal oxytocin and vasopressin during human social interaction. *Psychoneuroendocrinology* 39: 237-248. <https://doi.org/10.1016/j.psyneuen.2013.09.022>
- Rogers CN, Ross AP, Sahu SP, et al (2018) Oxytocin- and arginine vasopressin-containing fibers in the cortex of humans, chimpanzees, and rhesus macaques. *Am J Primatol* e22875. doi: <https://doi.org/10.1002/ajp.22875>
- Royer S, Martina M, Paré D (1999) An inhibitory interface gates impulse traffic between the input and output stations of the amygdala. *J Neurosci* 19:10575-10583
- Sabihi S, Dong SM, Durosko NE, Leuner B (2014) Oxytocin in the medial prefrontal cortex regulates maternal care, maternal aggression and anxiety during the postpartum period. *Front Behav Neurosci* 8:258. <https://doi.org/10.3389/fnbeh.2014.00258>
- Sabihi S, Dong SM, Maurer SD et al (2017) Oxytocin in the medial prefrontal cortex attenuates anxiety: anatomical and receptor specificity and mechanism of action. *Neuropharmacology* 125:1-12. <https://doi.org/10.1016/j.neuropharm.2017.06.024>
- Sack M, Spieler D, Wizelman L et al (2017) Intranasal oxytocin reduces provoked symptoms in female patients with posttraumatic stress disorder despite exerting sympathomimetic and positive chronotropic effects in a randomized controlled trial. *BMC Med* 15:40. <https://doi.org/10.1186/s12916-017-0801-0>
- Saffari R, Teng Z, Zhang M et al (2016) NPY<sup>+</sup>, but not PV<sup>+</sup>- GABAergic neurons mediated long-range inhibition from infra-to prefrontal cortex. *Transl Psychiatry* 6:e736. <https://doi.org/10.1038/tp.2016.7>
- Salchner P, Singewald N (2006) 5-HT receptor subtypes involved in the anxiogenic-like action and associated Fos response of acute fluoxetine treatment in rats. *Psychopharmacology* 185:282-288. <https://doi.org/10.1007/s00213-005-0247-5>
- Schiller D, Kanen JW, LeDoux JE et al (2013) Extinction during reconsolidation of threat memory diminishes prefrontal cortex involvement. *Proc Natl Acad Sci U S A* 110:20040-20045. <https://doi.org/10.1073/pnas.1320322110>
- Senn V, Wolff SBE, Herry C et al (2014) Long-range connectivity defines behavioral specificity of amygdala neurons. *Neuron* 81:428-437. <https://doi.org/10.1016/j.neuron.2013.11.006>
- Shansky RM, Hamo C, Hof PR et al (2010) Estrogen promotes stress sensitivity in a prefrontal cortex-amygdala pathway. *Cereb Cortex* 20:2560-2567. <https://doi.org/10.1093/cercor/bhq003>
- Shvil E, Sullivan GM, Schafer S et al (2014) Sex differences in extinction recall in posttraumatic stress disorder: a pilot fMRI study. *Neurobiol Learn Mem* 113:101-108. <https://doi.org/10.1016/j.nlm.2014.02.003>

- Sierra RO, Pedraza LK, Zanona QK et al (2017) Reconsolidation-induced rescue of a remote fear memory blocked by an early cortical inhibition: involvement of the anterior cingulate cortex and the mediation by the thalamic nucleus reuniens. *Hippocampus* 27:596-607. <https://doi.org/10.1002/hipo.22715>
- Singewald N, Schmuckermair C, Whittle N et al (2015) Pharmacology of cognitive enhancers for exposure-based therapy of fear, anxiety and trauma-related disorders. *Pharmacol Ther* 149:150-190. <https://doi.org/10.1016/j.pharmthera.2014.12.004>
- Smith AL, Freeman SM, Barnhart TE, Abbott DH, Ahlers EO, Kukis DL, Bales KL, Goodman MM, Young LJ (2016a) Initial investigation of three selective and potent small molecule oxytocin receptor PET ligands in New World monkeys. *Bioorg Med Chem Lett* 26(14):3370-3375. <https://doi.org/10.1016/j.bmcl>
- Smith AS, Tabbaa M, Lei K et al (2016b) Local oxytocin tempers anxiety by activating GABAA receptors in the hypothalamic paraventricular nucleus. *Psychoneuroendocrinology* 63:50-58. <https://doi.org/10.1016/j.psycheneu.2015.09.017>
- Sotres-Bayon F, Sierra-Mercado D, Pardilla-Delgado E, Quirk GJ (2012) Gating of fear in prelimbic cortex by hippocampal and amygdala inputs. *Neuron* 76:804-812. <https://doi.org/10.1016/j.neuron.2012.09.028>
- Stamatakis A, Manatos V, Kalpachidou T, Stylianopoulou F (2016) Exposure to a mildly aversive early life experience leads to prefrontal cortex deficits in the rat. *Brain Struct Funct* 221:4141-4157. <https://doi.org/10.1007/s00429-015-1154-0>
- Steenen SA, van Wijk AJ, van der Heijden GJM et al (2016) Propranolol for the treatment of anxiety disorders: systematic review and meta-analysis. *J Psychopharmacol (Oxford)* 30:128-139. <https://doi.org/10.1177/0269881115612236>
- Stewart RE, Chambless DL (2009) Cognitive-behavioral therapy for adult anxiety disorders in clinical practice: a meta-analysis of effectiveness studies. *J Consult Clin Psychol* 77:595-606. <https://doi.org/10.1037/a0016032>
- Stoop R (2012) Neuromodulation by oxytocin and vasopressin. *Neuron* 76(1):142-159. <https://doi.org/10.1016/j.neuron.2012.09.025>
- Stoop R, Hegoburu C, van den Burg E (2015) New opportunities in vasopressin and oxytocin research: a perspective from the amygdala. *Annu Rev Neurosci* 38:369-388. <https://doi.org/10.1146/annurev-neuro-071714-033904>
- Straub J, Metzger CD, Plener PL et al (2017) Successful group psychotherapy of depression in adolescents alters fronto-limbic resting-state connectivity. *J Affect Disord* 209:135-139. <https://doi.org/10.1016/j.jad.2016.12.019>
- Striepens N, Scheele D, Kendrick KM et al (2012) Oxytocin facilitates protective responses to aversive social stimuli in males. *Proc Natl Acad Sci U S A* 109:18144-18149. <https://doi.org/10.1073/pnas.1208852109>
- Striepens N, Kendrick KM, Hanking V et al (2013) Elevated cerebrospinal fluid and blood concentrations of oxytocin following its intranasal administration in humans. *Sci Rep* 3:3440. <https://doi.org/10.1038/srep03440>
- Sun N, Laviolette SR (2012) Inactivation of the basolateral amygdala during opiate reward learning disinhibits prelimbic cortical neurons and modulates associative memory extinction. *Psychopharmacology* 222:645-661. <https://doi.org/10.1007/s00213-012-2665-5>
- Terashima Y, Kondo K, Oiso Y (1999) Administration of oxytocin affects vasopressin V2 receptor and aquaporin-2 gene expression in the rat. *Life Sci* 64:1447-1453
- Tomizawa K, Iga N, Lu Y-F et al (2003) Oxytocin improves long-lasting spatial memory during motherhood through MAP kinase cascade. *Nat Neurosci* 6:384-390. <https://doi.org/10.1038/nn1023>
- Tonegawa S, Pignatelli M, Roy DS, Ryan TJ (2015) Memory engram storage and retrieval. *Curr Opin Neurobiol* 35:101-109. <https://doi.org/10.1016/j.conb.2015.07.009>
- Tovote P, Fadok JP, Lüthi A (2015) Neuronal circuits for fear and anxiety. *Nat Rev Neurosci* 16:317-331. <https://doi.org/10.1038/nrn3945>
- Triana-Del Rio R, Tecamachaltzi-Silvarán MB, Díaz-Estrada VX et al (2015) Conditioned same-sex partner preference in male rats is facilitated by oxytocin and dopamine: effect on sexually dimorphic brain nuclei. *Behav Brain Res* 283:69-77. <https://doi.org/10.1016/j.bbr.2015.04.019>
- Tye KM, Deisseroth K (2012) Optogenetic investigation of neural circuits underlying brain disease in animal models. *Nat Rev Neurosci* 13:251-266. <https://doi.org/10.1038/nrn3171>
- Tye KM, Prakash R, Kim S-Y et al (2011) Amygdala circuitry mediating reversible and bidirectional control of anxiety. *Nature* 471:358-362. <https://doi.org/10.1038/nature09820>
- Uhrig S, Hirth N, Broccoli L et al (2016) Reduced oxytocin receptor gene expression and binding sites in different brain regions in schizophrenia: a post-mortem study. *Schizophr Res* 177:59-66. <https://doi.org/10.1016/j.schres.2016.04.019>
- Van den Burg EH, Stindl J, Grund T et al (2015) Oxytocin stimulates extracellular Ca<sup>2+</sup> influx through TRPV2 channels in hypothalamic neurons to exert its anxiolytic effects. *Neuropsychopharmacology* 40:2938-2947. <https://doi.org/10.1038/npp.2015.147>
- Veening JG, Swanson LW, Sawchenko PE (1984) The organization of projections from the central nucleus of the amygdala to brainstem sites involved in central autonomic regulation: a combined retrograde transport-immunohistochemical study. *Brain Res* 303:337-357
- Vertes RP (2004) Differential projections of the infralimbic and prelimbic cortex in the rat. *Synapse* 51:32-58. <https://doi.org/10.1002/syn.10279>
- Vidal-Gonzalez I, Vidal-Gonzalez B, Rauch SL, Quirk GJ (2006) Microstimulation reveals opposing influences of prelimbic and infralimbic cortex on the expression of conditioned fear. *Learn Mem* 13:728-733. <https://doi.org/10.1101/lm.306106>
- Viviani D, Charlet A, van den Burg E et al (2011) Oxytocin selectively gates fear responses through distinct outputs from the central amygdala. *Science* 333:104-107. <https://doi.org/10.1126/science.1201043>
- Vogel E, Krabbe S, Gründemann J, Wamsteeker Cusulin JI, Lüthi A (2016) Projection-specific dynamic regulation of inhibition in amygdala micro-circuits. *Neuron* 91(3):644-651. <https://doi.org/10.1016/j.neuron.2016.06.016>
- Volkow ND, Wang G-J, Fowler JS, Ding Y-S (2005) Imaging the effects of methylphenidate on brain dopamine: new model on its therapeutic actions for attention-deficit/hyperactivity disorder. *Biol Psychiatry* 57:1410-1415. <https://doi.org/10.1016/j.biopsych.2004.11.006>
- Walum H, Waldman ID, Young LJ (2016) Statistical and methodological considerations for the interpretation of intranasal oxytocin studies. *Biol Psychiatry* 79:251-257. <https://doi.org/10.1016/j.biopsych.2015.06.016>
- Webb WM, Sanchez RG, Perez G et al (2017) Dynamic association of epigenetic H3K4me3 and DNA 5hmC marks in the dorsal hippocampus and anterior cingulate cortex following reactivation of a fear memory. *Neurobiol Learn Mem* 142:66-78. <https://doi.org/10.1016/j.nlm.2017.02.010>
- WHO (2016) Investing in treatment for depression and anxiety leads to fourfold return. In: WHO. <http://www.who.int/mediacentre/news/releases/2016/depression-anxiety-treatment/en/>. Accessed 5 Apr 2018
- Wöhr M, Schwarting RKW (2013) Affective communication in rodents: ultrasonic vocalizations as a tool for research on emotion and motivation. *Cell Tissue Res* 354:81-97. <https://doi.org/10.1007/s00441-013-1607-9>
- Woods AM, Bouton ME (2006) D-cycloserine facilitates extinction but does not eliminate renewal of the conditioned emotional response. *Behav Neurosci* 120:1159-1162. <https://doi.org/10.1037/0735-7044.120.5.1159>

- 
- Woolf NJ, Butcher LL (1982) Cholinergic projections to the basolateral amygdala: a combined Evans Blue and acetylcholinesterase analysis. *Brain Res Bull* 8:751-763
- Yang C-H, Shi H-S, Zhu W-L et al (2012) Venlafaxine facilitates between-session extinction and prevents reinstatement of auditory-cue conditioned fear. *Behav Brain Res* 230:268-273. <https://doi.org/10.1016/j.bbr.2012.02.023>
- Yin S, Liu Y, Petro NM, et al (2018) Amygdala adaptation and temporal dynamics of the salience network in conditioned fear: a single-trial fMRI study. *eNeuro* 5. doi: <https://doi.org/10.1523/ENEURO.0445-17.2018>
- Yoon S, Kim JE, Hwang J et al (2017) Recovery from posttraumatic stress requires dynamic and sequential shifts in amygdalar connectivities. *Neuropsychopharmacology* 42:454-461. <https://doi.org/10.1038/npp.2016.136>
- Young LJ, Flanagan-Cato LM (2012) Editorial comment: oxytocin, vasopressin and social behavior. *Horm Behav* 61:227-229. <https://doi.org/10.1016/j.yhbeh.2012.02.019>
- Zeidan MA, Igoe SA, Linnman C et al (2011) Estradiol modulates medial prefrontal cortex and amygdala activity during fear extinction in women and female rats. *Biol Psychiatry* 70:920-927. <https://doi.org/10.1016/j.biopsych.2011.05.016>