

**Serveur Académique Lausannois SERVAL [serval.unil.ch](http://serval.unil.ch)**

## **Author Manuscript**

### **Faculty of Biology and Medicine Publication**

**This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.**

Published in final edited form as:

**Title:** Seventy-five genetic loci influencing the human red blood cell.

**Authors:** van der Harst P, Zhang W, Mateo Leach I, Rendon A, Verweij N, Sehmi J, Paul DS, Elling U, Allayee H, Li X, Radhakrishnan A, Tan ST, Voss K, Weichenberger CX, Albers CA, Al-Hussani A, Asselbergs FW, Ciullo M, Danjou F, Dina C, Esko T, Evans DM, Franke L, Gögele M, Hartiala J, Hersch M, Holm H, Hottenga JJ, Kanoni S, Kleber ME, Lagou V, Langenberg C, Lopez LM, Lytikäinen LP, Melander O, Murgia F, Nolte IM, O'Reilly PF, Padmanabhan S, Parsa A, Pirastu N, Porcu E, Portas L, Prokopenko I, Ried JS, Shin SY, Tang CS, Teumer A, Traglia M, Ulivi S, Westra HJ, Yang J, Zhao JH, Anni F, Abdellaoui A, Attwood A, Balkau B, Bandinelli S, Bastardot F, Benyamin B, Boehm BO, Cookson WO, Das D, de Bakker PI, de Boer RA, de Geus EJ, de Moor MH, Dimitriou M, Domingues FS, Döring A, Engström G, Eyjolfsson GI, Ferrucci L, Fischer K, Galanello R, Garner SF, Genser B, Gibson QD, Girotto G, Gudbjartsson DF, Harris SE, Hartikainen AL, Hastie CE,

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

Published in final edited form as:

*Nature*. 2012 December 20; 492(7429): 369–375. doi:10.1038/nature11677.

## Seventy-five genetic loci influencing the human red blood cell

Pim van der Harst<sup>1,2,\*</sup>, Weihua Zhang<sup>3,4,\*</sup>, Irene Mateo Leach<sup>1,\*</sup>, Augusto Rendon<sup>5,6,7,8,\*</sup>, Niek Verweij<sup>1,\*</sup>, Joban Sehmi<sup>4,9,\*</sup>, Dirk S. Paul<sup>10,\*</sup>, Ulrich Elling<sup>11,\*</sup>, Hooman Allayee<sup>12</sup>, Xinzhong Li<sup>13,14</sup>, Aparna Radhakrishnan<sup>5,6,8,10</sup>, Sian-Tsung Tan<sup>4,9</sup>, Katrin Voss<sup>5,6,8</sup>, Christian X. Weichenberger<sup>15</sup>, Cornelis A. Albers<sup>5,6,10</sup>, Abtehale Al-Hussani<sup>3</sup>, Folkert W. Asselbergs<sup>16,17,18</sup>, Marina Ciullo<sup>19</sup>, Fabrice Danjou<sup>20</sup>, Christian Dina<sup>21,22,23</sup>, Tõnu Esko<sup>24,25</sup>, David M. Evans<sup>26</sup>, Lude Franke<sup>2</sup>, Martin Gögele<sup>15</sup>, Jaana Hartiala<sup>12</sup>, Micha Hersch<sup>27,28</sup>, Hilma Holm<sup>29</sup>, Jouke-Jan Hottenga<sup>30</sup>, Stavroula Kanoni<sup>10</sup>, Marcus E. Kleber<sup>31,32</sup>, Vasiliki Lagou<sup>33,34</sup>, Claudia Langenberg<sup>35</sup>, Lorna M. Lopez<sup>36,37</sup>, Leo-Pekka Lyytikäinen<sup>38,39</sup>, Olle Melander<sup>40</sup>, Federico Murgia<sup>41</sup>, Ilja M. Nolte<sup>42</sup>, Paul F. O'Reilly<sup>3</sup>, Sandosh Padmanabhan<sup>43</sup>, Afshin Parsa<sup>44</sup>, Nicola Pirastu<sup>45</sup>, Eleonora Porcu<sup>46</sup>, Laura Portas<sup>41</sup>, Inga Prokopenko<sup>33,34</sup>, Janina S. Ried<sup>47</sup>, So-Youn Shin<sup>10</sup>, Clara S. Tang<sup>48</sup>, Alexander Teumer<sup>49</sup>, Michela Traglia<sup>50</sup>, Sheila Ulivi<sup>51</sup>, Harm-Jan Westra<sup>2</sup>, Jian Yang<sup>52</sup>, Jing Hua Zhao<sup>35</sup>, Franco Anni<sup>20</sup>, Abdel Abdellaoui<sup>30</sup>, Antony Attwood<sup>5,6,8,10</sup>, Beverley Balkau<sup>53,54</sup>, Stefania Bandinelli<sup>55</sup>, François Bastardot<sup>56,57</sup>, Beben Benyamin<sup>48,58</sup>, Bernhard O. Boehm<sup>59</sup>, William O. Cookson<sup>9</sup>, Debashish Das<sup>60</sup>, Paul I. W. de Bakker<sup>17,18,61,62</sup>, Rudolf A. de Boer<sup>1</sup>, Eco J. C. de Geus<sup>30</sup>, Marleen H. de Moor<sup>30</sup>, Maria Dimitriou<sup>63</sup>, Francisco S.

© 2012 Macmillan Publishers Limited. All rights reserved

Correspondence and requests for materials should be addressed to J.C.C. (john.chambers@ic.ac.uk), C.G. (christian.gieger@helmholtz-muenchen.de), P.v.d.H. (p.van.der.harst@umcg.nl), J.S.K. (j.kooner@ic.ac.uk), W.H.O. (who1000@cam.ac.uk) and N.S. (ns6@sanger.ac.uk).

\*These authors contributed equally to this work.

Supplementary Information is available in the online version of the paper.

**Author Contributions** Study organisation: J.C.C., C.G., P.v.d.H., J.S.K., W.H.O. and N.S. Manuscript preparation: H.A., J.S.B., J.C.C., G.V.D., P.D., C.G., P.v.d.H., A.A. Hicks, J.S.K., I.M.-L., W.H.O., A. Radhakrishnan, A. Rendon, S.S., J. Sehmi, N.S., D.S.P., M.U., N.V. and W.Z. All authors reviewed and had the opportunity to comment on the manuscript. Data collection and analysis in the participating genome-wide association, replication and phenotype cohorts: **ALSPAC**: D.M.E., J.P.K., S.M.R., G.D.S.; **AMISH**: Q.D.G., B.D.M., A. Parsa, A.R.S.; **Beta-thalassaemia**: F.A., F.D., P. Fortina, R.G., L. Perseu, A. Piga, S.S., M.U.; **CBR**: A. Attwood, J.D., S.F.G., H.L.-J., C. Moore, W.H.O., J. Sambrook; **CoLAUS**: F.B., J.S.B., M.H., P.V.; **DeCODE**: G.I.E., D.F.G., H.H., I.O., P.T.O., K.S., P.S., U.T.; **DESIR**: B. Balkau, C.D., P. Froguel, R. Sladek; **EGCUT**: T.E., K.F., A.M., E.M., A.S.; **EPIC**: K.-T.K., C.L., R.J.F.L., N.J.W., J.-H.Z.; **Genebank**: H.A., J.H., S.L.H., W.H.W.T.; **INGI CARL**: P.G., G.G., N.P.; **INGI CILENTO**: M.C., T.N., D.R., R. Sorice.; **INGI FVG**: A.P.d.A., A. Robino, S.U.; **INGI Val Borbera**: G.P., C.S., D.T., M.T.; **KORA**: A.D., C.G., T.I., C. Meisinger, J.S.R.; **LBC**: I.J.D., S.E.H., L.M.L., J.M.S.; **LIFELINES**: R.A.d.B., I.P.K., I.M.-L., G.N., P.v.d.H., L.J.v.P., N.V., B.H.R.W.; **LOLIPOP**: A. Al-Hussani, J.C.C., D.D., P.E., J.S.K., X.L., K.M., J. Scott, J. Sehmi, S.-T.T., W.Z.; **LURIC**: B.G., B.O.B., M.E.K., W.M., B.R.W.; **MDC**: A.F.D., G.E., B.H., C.E.H., O.M., S.P., J.G.S.; **MICROS**: M.G., A. AHicks, A.S.-P., P.P.P.; **NESDA**: I.M.N., B.W.P., J.H.S., H. Snieder; **NFBC1966**: A.-L.H., M.-R.J., P.F.O., A. Pouta, A. Ruokonen.; **NTR**: A. Abdellaoui, D.I.B., E.J.C.d.G., J.-J.H., M.H.d.M., G. Willemsen; **OGP**: F.M., D.P., L. Portas, M.P.; **PREVEND**: R.A.d.B., I.M.-L., G.N., P.v.d.H., W.H.v.G., D.J.v.V., N.V.; **QIMR**: B. Benyamin, M.A.F., N.G.M., S.E.M., G.W.M., C.S.T., P.M.V., J.B.W.; **Sardinia**: F.C., E.P., S.S., M.U.; **SHIP**: A.G., M. Nauck, C.O.S., A. Teumer, U.V.; **SMART**: A. Algra, F.W.A., P.I.W.d.B., V.T.; **SORBS**: V.L., I.P., M.S., A. Tönjes.; **TwinsUK**: Y.M., S.-Y.S., N.S., T.D.S.; **UKBS**: J.J., W.H.O., N.S., J. Stephens; **Young Finns**: M.K., T.L., L.-P.L., O.R. Functional studies: *Drosophila*, U.E., F.S.D., A.A. Hicks, M. Novatchkova, J.M.P., U.P., C.X.W., G. Wirsberger; expression profiling, W.O.C., L. Franke, L.L., M.F.M., A. Rendon, E.S., H.-J.W.; **FAIRE**, C.A.A., P.D., W.H.O., D.S.P., A. Rendon, N.S. Data analysis and bioinformatics: A. Al-Hussani, S.B., J.C.C., M.D., L. Ferrucci, P.v.d.H., S.K., X.L., I.M.-L., K.M., S.M., A. Radhakrishnan, A. Rendon, R.R.-S., H. Schepers, J. Sehmi, N.S., H.H.W.S., S.T., T.T., N.V., K.V., P.V., J.Y., W.Z.

Summary statistics from the genome-wide association study are available from the European Genome-Phenome Archive (EGA, <http://www.ebi.ac.uk/ega>) under accession number EGAS00000000132.

Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints).

The authors declare no competing financial interests.

Readers are welcome to comment on the online version of the paper.

**Full Methods** and any associated references are available in the online version of the paper.

Domingues<sup>15</sup>, Angela Döring<sup>64</sup>, Gunnar Engström<sup>40</sup>, Gudmundur Ingi Eyjolfsson<sup>65</sup>, Luigi Ferrucci<sup>66</sup>, Krista Fischer<sup>24</sup>, Renzo Galanello<sup>20</sup>, Stephen F. Garner<sup>5,6,8</sup>, Bernd Genser<sup>31</sup>, Quince D. Gibson<sup>44,67</sup>, Giorgia Giotto<sup>45</sup>, Daniel Fannar Gudbjartsson<sup>29</sup>, Sarah E. Harris<sup>37,68</sup>, Anna-Liisa Hartikainen<sup>69</sup>, Claire E. Hastie<sup>43</sup>, Bo Hedblad<sup>40</sup>, Thomas Illig<sup>70,71</sup>, Jennifer Jolley<sup>5,6,8</sup>, Mika Kähönen<sup>72,73</sup>, Ido P. Kema<sup>74</sup>, John P. Kemp<sup>26</sup>, Liming Liang<sup>75</sup>, Heather Lloyd-Jones<sup>5,6,8</sup>, Ruth J. F. Loos<sup>35</sup>, Stuart Meacham<sup>5,6,8,10</sup>, Sarah E. Medland<sup>48</sup>, Christa Meisinger<sup>76</sup>, Yasin Memari<sup>10,77</sup>, Evelin Mihailov<sup>19</sup>, Kathy Miller<sup>4</sup>, Miriam F. Moffatt<sup>9</sup>, Matthias Nauck<sup>78</sup>, Maria Novatchkova<sup>11</sup>, Teresa Nutile<sup>19</sup>, Isleifur Olafsson<sup>79</sup>, Pall T. Onundarson<sup>80,81</sup>, Debora Parracciani<sup>82</sup>, Brenda W. Penninx<sup>83,84,85</sup>, Lucia Perseu<sup>46</sup>, Antonio Piga<sup>86</sup>, Giorgio Pistis<sup>50</sup>, Anneli Pouta<sup>87,88</sup>, Ursula Puc<sup>11</sup>, Olli Raitakari<sup>89,90</sup>, Susan M. Ring<sup>91</sup>, Antonietta Robino<sup>45</sup>, Daniela Ruggiero<sup>19</sup>, Aimo Ruukonen<sup>92</sup>, Aude Saint-Pierre<sup>15</sup>, Cinzia Sala<sup>50</sup>, Andres Salumets<sup>93,94</sup>, Jennifer Sambrook<sup>5,6,8</sup>, Hein Schepers<sup>95,96</sup>, Carsten Oliver Schmidt<sup>97</sup>, Herman H. W. Silljé<sup>1</sup>, Rob Sladek<sup>98</sup>, Johannes H. Smit<sup>83</sup>, John M. Starr<sup>37,99</sup>, Jonathan Stephens<sup>5,6,8</sup>, Patrick Sulem<sup>29</sup>, Toshiko Tanaka<sup>66</sup>, Unnur Thorsteinsdottir<sup>29,100</sup>, Vinicius Tragante<sup>16</sup>, Wiek H. van Gilst<sup>1</sup>, L. Joost van Pelt<sup>74</sup>, Dirk J. van Veldhuisen<sup>1</sup>, Uwe Völker<sup>49</sup>, John B. Whitfield<sup>48</sup>, Gonneke Willemssen<sup>30</sup>, Bernhard R. Winkelmann<sup>101</sup>, Gerald Wirnsberger<sup>11</sup>, Ale Algra<sup>17,102</sup>, Francesco Cucca<sup>46,103</sup>, Adamo Pio d'Adamo<sup>45</sup>, John Danesh<sup>104</sup>, Ian J. Deary<sup>36,37</sup>, Anna F. Dominiczak<sup>43</sup>, Paul Elliott<sup>3,105</sup>, Paolo Fortina<sup>106,107</sup>, Philippe Froguel<sup>108,109</sup>, Paolo Gasparini<sup>45</sup>, Andreas Greinacher<sup>110</sup>, Stanley L. Hazen<sup>111</sup>, Marjo-Riitta Jarvelin<sup>3,87,105,112,113</sup>, Kay Tee Khaw<sup>114</sup>, Terho Lehtimäki<sup>38,39</sup>, Winfried Maerz<sup>31,115</sup>, Nicholas G. Martin<sup>48</sup>, Andres Metspalu<sup>24,25</sup>, Braxton D. Mitchell<sup>44</sup>, Grant W. Montgomery<sup>48</sup>, Carmel Moore<sup>104</sup>, Gerjan Navis<sup>116</sup>, Mario Pirastu<sup>41</sup>, Peter P. Pramstaller<sup>15,117,118</sup>, Ramiro Ramirez-Solis<sup>10</sup>, Eric Schadt<sup>119</sup>, James Scott<sup>9</sup>, Alan R. Shuldiner<sup>44,120</sup>, George Davey Smith<sup>26</sup>, J. Gustav Smith<sup>40,121</sup>, Harold Snieder<sup>42</sup>, Rossella Sorice<sup>19</sup>, Tim D. Spector<sup>122</sup>, Kari Stefansson<sup>29,100</sup>, Michael Stumvoll<sup>123,124</sup>, W. H. Wilson Tang<sup>111</sup>, Daniela Toniolo<sup>50,125</sup>, Anke Tönjes<sup>123,124</sup>, Peter M. Visscher<sup>37,48,52,58</sup>, Peter Vollenweider<sup>56,57</sup>, Nicholas J. Wareham<sup>35</sup>, Bruce H. R. Wolffenbuttel<sup>126</sup>, Dorret I. Boomsma<sup>30</sup>, Jacques S. Beckmann<sup>27,127</sup>, George V. Dedoussis<sup>63</sup>, Panos Deloukas<sup>10</sup>, Manuel A. Ferreira<sup>48</sup>, Serena Sanna<sup>46</sup>, Manuela Uda<sup>46</sup>, Andrew A. Hicks<sup>15,\*</sup>, Josef Martin Penninger<sup>11,\*</sup>, Christian Gieger<sup>47,\*</sup>, Jaspal S. Kooner<sup>4,9,128,\*</sup>, Willem H. Ouwehand<sup>5,6,8,10,\*</sup>, Nicole Soranzo<sup>10,\*</sup>, and John C Chambers<sup>3,4,14,128,\*</sup>

<sup>1</sup>Department of Cardiology, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, The Netherlands <sup>2</sup>Department of Genetics, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, The Netherlands <sup>3</sup>Department of Epidemiology and Biostatistics, Imperial College London, London W2 1PG, UK <sup>4</sup>Ealing Hospital NHS Trust, Middlesex UB1 3HW, UK <sup>5</sup>Department of Haematology, University of Cambridge, Cambridge CB2 0XY, UK <sup>6</sup>NHS Blood and Transplant, Cambridge CB2 0PT, UK <sup>7</sup>MRC Biostatistics Unit, Institute of Public Health, Cambridge CB2 2SR, UK <sup>8</sup>NIHR Cambridge Biomedical Research Centre, Cambridge CB2 0QQ, UK <sup>9</sup>National Heart and Lung Institute, Imperial College London, London W12 0NN, UK <sup>10</sup>Wellcome Trust Sanger Institute, Hinxton, Cambridge CB10 1SA, UK <sup>11</sup>Institute of Molecular Biotechnology of the Austrian Academy of Sciences, 1030 Vienna, Austria <sup>12</sup>Department of Preventive Medicine, University of Southern California Keck School of Medicine, Los Angeles, California 90033, USA <sup>13</sup>Institute of Clinical Sciences, Imperial College London, London W12 0NN, UK <sup>14</sup>NIHR Cardiovascular Biomedical Research Unit, Royal Brompton and Harefield NHS Foundation Trust and Imperial College London, London SW3 6NP, UK <sup>15</sup>Center for Biomedicine, European Academy Bozen/Bolzano (EURAC), 39100 Bolzano, Italy <sup>16</sup>Department of Cardiology, Division Heart and Lungs, University Medical Center Utrecht, 3508 Utrecht, The Netherlands <sup>17</sup>Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, 3508 Utrecht, The Netherlands <sup>18</sup>Department of Medical Genetics, Biomedical Genetics, University Medical Center Utrecht, 3508 Utrecht, The Netherlands <sup>19</sup>Institute of Genetics and Biophysics "Adriano Buzzati-Traverso"-CNR, 80131 Naples, Italy

<sup>20</sup>Clinica Pediatrica 2a, Dipartimento di Scienze Biomediche e Biotecnologie - Università di Cagliari, Ospedale Regionale Microcitemie ASL8, 09121 Cagliari, Italy <sup>21</sup>Institut National de la Santé et de la Recherche Médicale (INSERM) Unité Mixte de Recherche (UMR) 1087, BP 70721 44007 Nantes cedex, 1 Nantes, France <sup>22</sup>Centre National de la Recherche Scientifique (CNRS) UMR 6291, BP 70721 44007 Nantes cedex 1, Nantes, France <sup>23</sup>School of Medicine, Nantes University, 44000 Nantes, France <sup>24</sup>Estonian Genome Center of University of Tartu, 51010 Tartu, Estonia <sup>25</sup>Institute of Molecular and Cell Biology, University of Tartu, 51010 Tartu, Estonia <sup>26</sup>MRC Centre for Causal Analyses in Translational Epidemiology, School of Social and Community Medicine, University of Bristol, Bristol BS8 2BN, UK <sup>27</sup>Department of Medical Genetics, University of Lausanne, CH-1005 Lausanne, Switzerland <sup>28</sup>Swiss Institute of Bioinformatics, CH-1015 Lausanne, Switzerland <sup>29</sup>deCODE genetics, 101 Reykjavik, Iceland <sup>30</sup>Department of Biological Psychology, VU University, 1081 BT Amsterdam, The Netherlands <sup>31</sup>Mannheim Institute of Public Health, Social and Preventive Medicine, Medical Faculty of Mannheim, University of Heidelberg, D-68167 Mannheim, Germany <sup>32</sup>LURIC Study nonprofit LLC, D-79098 Freiburg, Germany <sup>33</sup>Oxford Centre for Diabetes, Endocrinology and Metabolism, Radcliffe Department of Medicine, University of Oxford OX3 7LJ, UK <sup>34</sup>Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford OX3 7BN, UK <sup>35</sup>MRC Epidemiology Unit, Institute of Metabolic Science, Addenbrooke's Hospital, Cambridge CB2 0QQ, UK <sup>36</sup>Department of Psychology, The University of Edinburgh, Edinburgh EH8 9JZ, UK <sup>37</sup>Centre for Cognitive Ageing and Cognitive Epidemiology, The University of Edinburgh, Edinburgh EH8 9JZ, UK <sup>38</sup>Department of Clinical Chemistry, Fimlab Laboratories, Tampere University Hospital, FIN-33521 Tampere, Finland <sup>39</sup>Department of Clinical Chemistry, University of Tampere School of Medicine, FIN-33521 Tampere, Finland <sup>40</sup>Department of Clinical Sciences, Lund University, SE-205 02Malmö, Sweden <sup>41</sup>Institute of Population Genetics, National Research Council of Italy, 07100 Sassari, Italy <sup>42</sup>Department of Epidemiology, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, The Netherlands <sup>43</sup>Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow G12 8QQ, United Kingdom <sup>44</sup>University of Maryland School of Medicine, Baltimore, Maryland 21201, USA <sup>45</sup>Institute for Maternal and Child Health-IRCCS "Burlo Garofolo"-Trieste, University of Trieste, 34137 Trieste, Italy <sup>46</sup>Istituto di Ricerca Genetica e Biomedica, Consiglio Nazionale delle Ricerche, c/o Cittadella Universitaria di Monserrato, Monserrato, Cagliari 09042, Italy <sup>47</sup>Institute of Genetic Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, D-85764 Neuherberg, Germany <sup>48</sup>Queensland Institute of Medical Research, Brisbane, Queensland 4006, Australia <sup>49</sup>Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald, D-17487 Greifswald, Germany <sup>50</sup>Division of Genetics and Cell Biology, San Raffaele Scientific Institute, 20132 Milano, Italy <sup>51</sup>Institute for Maternal and Child Health - IRCCS "Burlo Garofolo"- Trieste, 34137 Trieste, Italy <sup>52</sup>University of Queensland Diamantina Institute, The University of Queensland, Princess Alexandra Hospital, Brisbane, Queensland 4102, Australia <sup>53</sup>Inserm, CESP Centre for research in Epidemiology and Population Health, U1018, Villejuif F-94807, France <sup>54</sup>University Paris Sud 11, UMRS 1018, Villejuif F-94807, France <sup>55</sup>Geriatric Unit, Azienda Sanitaria Firenze, 50125 Florence, Italy <sup>56</sup>Centre Hospitalier Universitaire Vaudois, CH-1011 Lausanne, Switzerland <sup>57</sup>Department of Internal Medicine, University of Lausanne, CH-1011 Lausanne, Switzerland <sup>58</sup>Queensland Brain Institute, The University of Queensland, Brisbane, Queensland 4072, Australia <sup>59</sup>Division of Endocrinology and Diabetes, Department of Medicine, University Hospital, Ulm D-89075, Germany <sup>60</sup>The Hatter Cardiovascular Institute, University College London, London WC1E 6HX, UK <sup>61</sup>Division of Genetics, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115, USA <sup>62</sup>Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, Massachusetts 02142, USA <sup>63</sup>Nutrition and Dietetics, Harokopio University, Kallithea 17671, Athens, Greece <sup>64</sup>Institute of Epidemiology I and Institute of Epidemiology II, Helmholtz Zentrum München, German Research Center for Environmental

Health, D-85764 Neuherberg, Germany <sup>65</sup>The Laboratory in Mjodd, 108 Reykjavik, Iceland  
<sup>66</sup>Clinical Research Branch, National Institute on Aging, Baltimore, Maryland 21250, USA  
<sup>67</sup>Instituto de Saúde Coletiva, Federal University of Bahia, Salvador, Bahia 40110-040, Brazil  
<sup>68</sup>Medical Genetics Section, The University of Edinburgh, Edinburgh EH4 2XU, UK <sup>69</sup>Institute of  
Clinical Sciences, Obstetrics and Gynecology, University of Oulu FIN-90220 Oulu, Finland  
<sup>70</sup>Research Unit of Molecular Epidemiology, Helmholtz Zentrum München, German Research  
Center for Environmental Health, D-85764 Neuherberg, Germany <sup>71</sup>Hannover Unified Biobank,  
Hannover Medical School, D-30625 Hannover, Germany <sup>72</sup>Department of Clinical Physiology,  
Tampere University Hospital, FIN-33521 Tampere, Finland <sup>73</sup>Department of Clinical Physiology,  
University of Tampere School of Medicine, FIN-33521 Tampere, Finland <sup>74</sup>Department of  
Laboratory Medicine, University of Groningen, University Medical Center Groningen, 9700 RB  
Groningen, The Netherlands <sup>75</sup>Department of Epidemiology, Department of Biostatistics, Harvard  
School of Public Health, Cambridge, Massachusetts 02115, USA <sup>76</sup>Institute of Epidemiology II,  
Helmholtz Zentrum München, German Research Center for Environmental Health, D-85764  
Neuherberg, Germany <sup>77</sup>Department of Twin Research and Genetic Epidemiology, Kings College  
London, London SE1 7EH, UK <sup>78</sup>Institute of Clinical Chemistry and Laboratory Medicine,  
University Medicine Greifswald, D-17475 Greifswald, Germany <sup>79</sup>Department of Clinical  
Biochemistry, Landspítali University Hospital, 101 Reykjavik, Iceland <sup>80</sup>Faculty of Medicine,  
University of Iceland, 101 Reykjavik, Iceland <sup>81</sup>Laboratory of Hematology and Coagulation  
Disorder Center, Landspítali University Hospital, 101 Reykjavik, Iceland <sup>82</sup>Genetic Park of  
Ogliastra, Perdasdefogu, Sardinia, Italy <sup>83</sup>Department of Psychiatry/EMGO Institute/Neuroscience  
Campus, VU University Medical Centre, 1081 BT Amsterdam, The Netherlands <sup>84</sup>Department of  
Psychiatry, University of Groningen, University Medical Center Groningen, 9700 RB Groningen,  
The Netherlands <sup>85</sup>Department of Psychiatry, Leiden University Medical Centre, 2333 Leiden,  
The Netherlands <sup>86</sup>Division of Pediatrics and Thalassemia Centre, Department of Clinical and  
Biological Sciences, University of Torino, 10043 Orbassano, Turin, Italy <sup>87</sup>Institute of Health  
Sciences, University of Oulu, FIN-90220 Oulu, Finland <sup>88</sup>National Institute of Health and Welfare,  
Aapistie 1, P.O. Box 310, FIN-90101 Oulu, Finland <sup>89</sup>Department of Clinical Physiology and  
Nuclear Medicine, Turku University Hospital, FIN-20521 Turku, Finland <sup>90</sup>Research Centre of  
Applied and Preventive Cardiovascular Medicine, University of Turku, FIN-20521 Turku, Finland  
<sup>91</sup>The School of Social and Community Medicine, University of Bristol, Bristol BS82PS, UK  
<sup>92</sup>Institute of Diagnostics, University of Oulu, FIN-90014 Oulu, Finland <sup>93</sup>Competence Centre on  
Reproductive Medicine and Biology, 50410 Tartu, Estonia <sup>94</sup>Institute of General and Molecular  
Pathology, University of Tartu, 51014 Tartu, Estonia <sup>95</sup>Department of Experimental Hematology,  
University of Groningen, University Medical Center Groningen, 9700 RB Groningen, The  
Netherlands <sup>96</sup>Department of Stem Cell Biology, University of Groningen, University Medical  
Center Groningen, 9700 RB Groningen, The Netherlands <sup>97</sup>Institute for Community Medicine,  
University Medicine Greifswald, D-17475 Greifswald, Germany <sup>98</sup>Departments of Human  
Genetics and Medicine, Faculty of Medicine, McGill University, Montreal, Quebec H3A 1B1,  
Canada <sup>99</sup>Geriatric Medicine Unit, The University of Edinburgh, Western General Hospital,  
Edinburgh EH4 2XU, UK <sup>100</sup>Faculty of Medicine, University of Iceland, 101 Reykjavik, Iceland  
<sup>101</sup>Clin Phenomics Study Center, D-60594 Frankfurt, Germany <sup>102</sup>Utrecht Stroke Center,  
Department of Neurology and Neurosurgery, Rudolf Magnus Institute of Neuroscience, University  
Medical Center Utrecht, 3584 CG Utrecht, The Netherlands <sup>103</sup>Dipartimento di Scienze  
Biomediche, Università di Sassari, 07100 Sassari, Italy <sup>104</sup>Department of Public Health and  
Primary Care, University of Cambridge, Cambridge CB1 8RN, UK <sup>105</sup>MRC-HPA Centre for  
Environment and Health, Imperial College London, London W2 1PG, UK <sup>106</sup>Department of  
Cancer Biology, Thomas Jefferson University Jefferson Medical College, Philadelphia,  
Pennsylvania 19107, USA <sup>107</sup>Dipartimento di Medicina Molecolare, Università La Sapienza,  
00161 Roma, Italy <sup>108</sup>Centre National de la Recherche Scientifique (CNRS)-Unité mixte de  
recherche (UMR) 8199, Lille Pasteur Institute, Lille 59100, France <sup>109</sup>Department of Genomics of

Common Disease, School of Public Health, Imperial College London, London W2 1PG, UK  
<sup>110</sup>Institute for Immunology and Transfusion Medicine, University Medicine Greifswald, D-17487 Greifswald, Germany <sup>111</sup>Center for Cardiovascular Diagnostics and Prevention, Department of Cell Biology, Lerner Research Institute, Cleveland Clinic, Cleveland, Ohio 44195, USA  
<sup>112</sup>Department of Lifecourse and Service, National Institute for Health and Welfare, FIN-90101 Oulu, Finland <sup>113</sup>Biocenter Oulu, University of Oulu, FIN-90220 Oulu, Finland <sup>114</sup>Clinical Gerontology Unit, Box 251, Addenbrooke's Hospital, Hills Road, Cambridge CB2 2QQ, UK  
<sup>115</sup>Synlab Academy, D-68165 Mannheim, Germany <sup>116</sup>Department of Internal Medicine, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, The Netherlands <sup>117</sup>Department of Neurology, General Central Hospital, 39100 Bolzano, Italy  
<sup>118</sup>Department of Neurology, University of Lübeck, D-23538 Lübeck, Germany <sup>119</sup>Institute for Genomics and Multiscale Biology, Mount Sinai School of Medicine, New York 10029-6574, USA  
<sup>120</sup>Geriatric Research and Education Clinical Center, Veterans Administration Medical Center, Baltimore, Maryland 21201, USA <sup>121</sup>Department of Cardiology, Lund University, 22185 Lund, Sweden <sup>122</sup>Department of Twin Research and Genetic Epidemiology, Kings College London, London SE1 7EH, UK <sup>123</sup>Department of Medicine, University of Leipzig, Liebigstr. 18, D-04103 Leipzig, Germany <sup>124</sup>University of Leipzig, IFB Adiposity Diseases, D-04103 Leipzig, Germany  
<sup>125</sup>Institute of Molecular Genetics, CNR, 27100 Pavia, Italy <sup>126</sup>Department of Endocrinology, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, The Netherlands <sup>127</sup>Service of Medical Genetics, Centre Hospitalier Universitaire Vaudois, 1011 Lausanne, Switzerland <sup>128</sup>Imperial College Healthcare NHS Trust, London W12 0HS, UK

## Abstract

Anaemia is a chief determinant of global health, contributing to cognitive impairment, growth retardation and impaired physical capacity. To understand further the genetic factors influencing red blood cells, we carried out a genome-wide association study of haemoglobin concentration and related parameters in up to 135,367 individuals. Here we identify 75 independent genetic loci associated with one or more red blood cell phenotypes at  $P < 10^{-8}$ , which together explain 4–9% of the phenotypic variance per trait. Using expression quantitative trait loci and bioinformatic strategies, we identify 121 candidate genes enriched in functions relevant to red blood cell biology. The candidate genes are expressed preferentially in red blood cell precursors, and 43 have haematopoietic phenotypes in *Mus musculus* or *Drosophila melanogaster*. Through open-chromatin and coding-variant analyses we identify potential causal genetic variants at 41 loci. Our findings provide extensive new insights into genetic mechanisms and biological pathways controlling red blood cell formation and function.

---

Haemoglobin, an iron-containing metalloprotein found in the red blood cells of all vertebrates, provides the primary mechanism for oxygen transport in the circulation. Haemoglobin levels and related red blood cell phenotypes are tightly regulated, including an important genetic component<sup>1–5</sup>. To refine our understanding of the genetic factors influencing red blood cell formation and function, we carried out a meta-analysis of genome-wide association studies (GWAS) and staged follow-up genotyping of six red blood cell phenotypes: haemoglobin, mean cell haemoglobin (MCH), mean cell haemoglobin concentration (MCHC), mean cell volume (MCV), packed cell volume (PCV) and red blood cell count (RBC).

Our study design is summarized in Supplementary Fig. 1. In brief, we combined genome-wide association data from 71,861 individuals of European or South Asian ancestry, with up to 2,644,161 autosomal single-nucleotide polymorphisms (SNPs) and 67,645 X-chromosome SNPs. Characteristics of participants, genotyping arrays and imputation are

summarized in Supplementary Tables 1–3. Meta-analysis was carried out among Europeans and South Asians separately, followed by a final combined analysis of results from the two populations. We performed replication testing of 22 loci showing suggestive association ( $10^{-8} < P < 10^{-7}$ ) in a further 63,506 individuals using a combination of *in silico* data and direct genotyping (Supplementary Tables 1, 2 and Supplementary Note). Genome-wide significance was set at  $P < 1 \times 10^{-8}$ , allowing a Bonferroni correction both for the  $\sim 10^6$  independent SNPs tested<sup>6</sup>, as well as for the six inter-related red blood cell phenotypes (Supplementary Note)<sup>7</sup>.

Seventy-five independent genetic loci reached genome-wide significance for association with one or more red blood cell phenotypes (Table 1 and Supplementary Fig. 2), 43 of which are novel. For descriptive and downstream purposes, we identified a single ‘sentinel’ SNP for each of the 75 loci, defined as the SNP with the lowest *P* value against any phenotype at each locus; regional plots for the 75 loci are shown in Supplementary Fig. 3. Full lists of the SNPs associated with phenotype at  $P < 10^{-6}$  and of the sentinel SNPs are provided (Supplementary Tables 4 and 5). Of the 38 loci previously reported to be associated with red blood cell traits<sup>1–5</sup>, we replicate 32 loci ( $P < 10^{-8}$ ) and find three to be nominally associated ( $P < 0.05$ ; Supplementary Table 6). The remaining three loci, initially reported in an East Asian GWAS<sup>4</sup>, were not associated with red blood cell phenotypes in our sample (Supplementary Fig. 4 and Supplementary Note).

Among the 75 genomic loci identified, we found that 31 were associated with one red blood cell phenotype, and 44 with two or more phenotypes, at  $P < 10^{-8}$ . The total number of locus–phenotype associations identified at  $P < 10^{-8}$  was 156, of which 92 are novel (Supplementary Fig. 5 and Supplementary Table 7). In addition, at 8 of the 75 loci we found evidence for multiple SNPs independently associated with red blood cell phenotype at  $P < 10^{-8}$  in conditional analyses<sup>8</sup>, suggesting the presence of possible secondary genetic mechanisms at these loci (Supplementary Table 8).

## Identification of candidate genes

There are >3,000 protein-coding genes within 1 megabase (Mb) of the sentinel SNPs from the 75 genetic loci associated with red blood cell phenotypes. We prioritized genes as probable candidates underlying the observed genetic associations using the following criteria: (1) gene nearest to the sentinel SNP, and any other gene within 10 kilo-bases (kb) (97 genes; Table 1); (2) gene containing a non-synonymous SNP in high linkage disequilibrium ( $r^2 > 0.8$ ) with the sentinel SNP (24 genes; Supplementary Table 9); (3) gene with expression quantitative trait loci (eQTL) associated with sentinel SNP in peripheral blood lymphocytes (27 genes; Supplementary Table 10); and (4) gene relationships among implicated loci (GRAIL) literature analysis<sup>9</sup> (9 genes; Supplementary Table 11). This strategy identified 121 candidate genes (Table 1 and Supplementary Fig. 6).

Pathway analysis revealed that the list of candidate genes is strongly enriched for genes known to be involved in haematological development and function ( $P = 10^{-63}$ ), as well as in cellular proliferation, development and death, and immunological processes (Supplementary Tables 12 and 13). Current knowledge of gene function for all 121 candidates is summarized in Supplementary Table 14. Of note, some of the genes within these regions are known to underlie the Mendelian red blood cell disorders of elliptocytosis, ovalocytosis and spherocytosis (*ANK1*, *SLC4A1*, *SPTA1*)<sup>10</sup>, haemolytic anaemia (*HK1*)<sup>11</sup> and iron deficiency or overload (*TMPRSS6*, *HFE*, *TFR2*)<sup>12</sup>. Furthermore, somatic mutations of *IKZF1*, *KIT*, *SH2B3*, *SH3GL1* and *TAL1* (also known as *SCL*) underlie several haematologic proliferative disorders (Supplementary Table 14).

## Gene expression during haematopoiesis

We next explored expression of the 121 candidate genes using an atlas of 38 different haematopoietic cell types (Supplementary Table 15)<sup>13</sup>. Ninety-seven genes could be reliably assigned a probe on the Affymetrix HG\_U133AAofAv2 array (Fig. 1a); these transcripts were, on average, expressed at higher levels in late erythroblasts (or the precursors of red blood cells, EB3–EB5) compared to other transcripts in the same cell type ( $P < 0.01$  after Bonferroni correction; Fig. 1b). Furthermore, expression was more likely to be upregulated in EB3–5 relative to other cell types ( $P = 1.2 \times 10^{-6}$ , rank-sum test).

To further investigate lineage-specific effects, we assessed temporal patterns of gene expression during *in vitro* differentiation of haematopoietic stem cells to erythroblasts<sup>14</sup>. On average, candidate genes have increasing expression over time along the erythroid lineage ( $P = 0.006$ , rank-sum test; Fig. 1c). These data support the view that the gene set identified here is enriched for genes relevant to red blood cell biology, including a number of candidate genes differentially regulated to increase their expression in late erythropoiesis.

## Coding and regulatory sequence variants

To better capture common sequence variation at the 75 loci, we searched the 1000 Genomes Project data set ([www.1000genomes.org](http://www.1000genomes.org)) and identified 39 non-synonymous SNPs that are in high linkage disequilibrium ( $r^2 > 0.8$ ) with sentinel SNPs at the red blood cell loci (Supplementary Table 9). This represents a ~sixfold enrichment compared to the expectation under the null hypothesis ( $P = 0.01$ ; Supplementary Note). Although re-sequencing will be needed to obtain a complete assessment of variants at these loci, these non-synonymous sites represent an initial set of candidates for genetic variants underlying the observed associations with red blood cell phenotypes, potentially mediated through changes in protein function.

We next searched for sequence variants at the red blood cell loci that might influence gene regulation. We used formaldehyde-assisted isolation of regulatory elements followed by next-generation sequencing (FAIRE-seq) to identify nucleosome-depleted regions (NDRs) that may represent active regulatory elements<sup>15</sup>. We studied three haematologic cell types, and found 103,308 unique NDRs, of which 38,014 were present in erythroblasts, 50,372 in megakaryocytes and 34,833 in monocytes. We then searched the 1000 Genomes Project data set and found 60 SNPs located within one of these NDRs that are either: (1) one of the 75 sentinel SNPs from the red blood cell GWAS, or (2) in high linkage disequilibrium ( $r^2 > 0.8$ ) and located within 1 Mb of a sentinel SNP (Supplementary Table 16). The NDRs overlapping these 60 SNPs were more likely to be erythroblast specific than expected by chance (1.8-fold enrichment compared to background distribution of NDRs;  $P = 0.007$ , Bonferroni-adjusted binomial test); by contrast, there were fewer megakaryocyte-specific NDRs coinciding with red blood cell SNPs (0.4-fold enrichment;  $P = 0.007$ ; Fig. 1d). This pattern of erythroblast enrichment and megakaryocyte depletion was robust to the stringency of NDR peak-calling (Supplementary Table 17). Our results indicate that regulatory variation within the erythroid lineage may underlie the associations observed at several of the loci identified in our red blood cell GWAS. The 19 genes closest to the 25 erythroblast-specific NDRs were more likely to be upregulated during erythropoiesis compared to all other expressed transcripts ( $P = 6.3 \times 10^{-6}$ , rank-sum test; Supplementary Table 18), lending further support to the view that the NDRs identified have a role in the regulation of genes involved in erythropoiesis<sup>16,17</sup>. Interestingly, the SNPs associated with MCH at 16p11 overlap an erythroblast-specific NDR that coincides with the NPRL3 regulatory element in the locus control region of the downstream haemoglobin- $\alpha$  locus<sup>18,19</sup>.



Together our coding- and regulatory-variant analyses thus identify a set of ~100 SNPs across 41 regions that are candidates for functional genomic elements influencing red blood cell formation and function, and which constitutes a priority set for future experimental evaluation.

## Insights from mouse models

A systematic search of the Mouse Genome Informatics database reveals haematologic phenotypes for 29 of the 100 candidate genes that have mouse homologues (Supplementary Fig. 6 and Supplementary Tables 14, 19), including genes involved in cell cycle regulation: *CCNA2* (4q27), *CCND2* (12p13) and *CCND3* (6p21); genes coding for transcription factors and their interacting proteins: *BCL11A* (2p16), *CITED2* (6q24), *IKZF1* (7p13) and *TAL1* (1p32); and genes involved in growth factor or cytokine signalling: *KIT* (4q11), *KITLG* (12q22), *SH2B3* (12q24) and *PTPRCAP* (11q13). Among the gene products encoded at the newly identified loci, KITLG, also known as stem cell factor, is the cognate ligand for the KIT tyrosine kinase receptor<sup>20</sup>. KIT signalling is involved in the perinatal transition from fetal to adult haemoglobin, in addition to maintenance, proliferation and differentiation of haematopoietic stem cells<sup>21</sup>. *Kitlg*<sup>-/-</sup> and *Kir*<sup>-/-</sup> mice have low red blood cell concentrations, anaemia and other haematological abnormalities. CCNA2, CCND2 and CCND3 are cyclin-dependent kinases that contribute to initiation and progression of cell division<sup>22</sup>. Knock-out models of these genes have a number of haematological abnormalities, including reduced stem cell and red blood cell concentrations, and anaemia<sup>22</sup>. Of the 29 candidate genes with a blood phenotype in mouse, 25 were identified as the genes nearest to the sentinel SNP, and 15 through the eQTL ( $n=2$ ), coding-variant ( $n=6$ ) or GRAIL ( $n=8$ ) analyses (Supplementary Table 19).

## RNAi silencing in *D. melanogaster*

We used haemocyte-specific RNA interference (RNAi) silencing in *D. melanogaster* to further evaluate the candidate genes for their role in blood cell formation. We first carried out permutation testing in a genome-wide *D. melanogaster* RNAi silencer screen (Supplementary Note). Results confirmed that the 121 candidates are enriched for genes with a blood cell phenotype in *D. melanogaster*, supporting the view that our GWAS identifies a set of genes conserved across phyla and involved in blood cell formation or survival.

We next created haemocyte-specific RNAi knockdowns for 96 *D. melanogaster* genes that are orthologues for 74 of the 121 candidate genes, and assessed blood cell formation (crystal cells and plasmotocytes) in early- and late-stage L3 larvae<sup>23</sup>. We found 19 out of the 74 candidate genes with orthologues in *D. melanogaster* to have a blood cell phenotype, of which 5 also have a haematological phenotypes in mouse models: *KIT*, *HK1*, *CCNA2*, *AP3D1* and *PSMB10* (Supplementary Tables 19 and 20). Among the genes highlighted, RNAi silencing of *KIT* and *CCNA2* orthologues was associated with a profound reduction in plasmocyte formation (Fig. 2), consistent with their established role in cytokinesis<sup>20,22</sup>. *AP3D1* is involved in vesicular trafficking and dense granule formation in platelets<sup>24</sup>, whereas *PSMB10* is a component of a widely distributed proteasome linked to inflammation and ubiquitin signalling<sup>25</sup>. *UBE2L3* is also involved in ubiquitin signalling and immune regulation<sup>26</sup>, and genetic variants in *UBE2L3* are strongly associated with several autoimmune diseases known to influence blood cell counts<sup>27,28</sup>. *EIF5* (14q32) is involved in activation of the ribosomal initiation complex<sup>29</sup>, whereas *RPS6KB2* (22q11) is a key component of growth factor and other signalling cascades that regulate ribosomal function, cellular proliferation and survival<sup>30</sup>. For most of the genes identified, the mechanisms

underlying their potential relationship to red cell biology remain to be elucidated; our gene set thus provides a rich resource for future experimental evaluation and discovery.

## Contribution to clinical phenotype

The 75 sentinel SNPs together account for between 3.9% (PCV) and 8.9% (MCV) of population variation in red blood cell phenotypes (Supplementary Table 21). Individuals in the highest quartile of genetic risk score (GRS; on the basis of weighted effect of the 75 sentinel SNPs) are 3–5-fold more likely to be in the highest quartile for population distribution of MCH, MCV and RBC (Fig. 3). GRS is associated with haemoglobin concentrations across the physiological range, including at haemoglobin levels that predict adverse outcomes in pregnancy, cardiovascular and neurologic disease, in addition to mortality in the elderly<sup>31–34</sup>.

We next investigated the association of the 75 sentinel SNPs with red blood cell phenotypes in thalassaemia, a group of genetic disorders characterized by defects in haemoglobin synthesis and anaemia. We confirmed association of several of the sentinel SNPs with respective blood cell trait, and found that GRS predicts phenotype similarly, among 460  $\beta$ -thalassaemia heterozygotes (Supplementary Table 22 and Supplementary Note). In separate experiments, GRS predicts time to first blood transfusion among 495 patients with thalassaemia major ( $P=6.9 \times 10^{-4}$ ); however, this effect was fully accounted for by the *MYB-HBS1L* locus, which modifies the severity of thalassaemia major through its effect on fetal haemoglobin levels (Supplementary Note)<sup>35</sup>. Together, our findings demonstrate that the common genetic variants identified contribute to phenotypic variation in the general population, and suggest that they may also act as genetic modifiers in clinically relevant red blood cell abnormalities.

## Conclusions

Our genome-wide association and replication study in 135,367 individuals identifies 75 genetic loci influencing red blood cell phenotypes, and 156 locus–phenotype associations; most of these discoveries are novel. Through open-chromatin and coding-variant studies, we identify a first set of SNPs as potential causal variants. In parallel, our bioinformatic strategies identify a core set of genes, differentially regulated in haematologic precursor cells, which are candidates for mediating the effects on red blood cell phenotypes. However, despite our extensive GWAS, bioinformatic and experimental data, the precise identities of the causal variants, regulatory regions and genes remain to be determined; definitive identification will require further detailed experimental evaluation. Our results thus provide new insights into the genes and gene variants that may influence haemoglobin levels and related red blood cell indices, and will underpin a deeper knowledge of the biological mechanisms involved in haematopoiesis and red blood cell function.

## METHODS

### Genome-wide association

Genome-wide association was carried out in 62,553 people of European ancestry and 9,308 people of South Asian ancestry, using up to 2,644,161 autosomal and 67,645 X-chromosome SNPs. Imputation was done using haplotypes from HapMap Phase 2. Characteristics of participants, genotyping arrays and imputation are summarized in Supplementary Tables 1 and 2. Participants with extreme measurements ( $> \pm 3$  s.d. from mean) were excluded on a per-phenotype basis. Each population cohort was approved by a research ethics committee, and all participants gave informed consent.

SNP associations with each phenotype were tested by linear regression using an additive genetic model. Associations were tested separately in men and women in each cohort, with principal components and other study-specific factors as covariates to account of population substructure as described in Supplementary Table 2. Test statistics from each cohort were then corrected for their respective genomic-control inflation factor to adjust for residual population sub-structure; genomic-control inflation factors are summarized in Supplementary Table 3. We then carried out a meta-analysis of results from the individual cohorts using  $Z$ -scores weighted by the square root of sample size. The meta-analysis was varied out among Europeans and South Asians separately. There were no South-Asian-specific discoveries, but also little evidence for heterogeneity of effect at known or new genetic loci (Supplementary Table 23); we therefore carried out a final combined analysis of results for the two populations. SNPs with minor allele frequency  $<1\%$  (weighted average across cohorts) were removed, as were SNPs with weight  $<50\%$  of phenotype sample size. There was no evidence for inflation of test statistics at SNPs not known to be associated with red blood cell phenotypes (Supplementary Table 3), and genomic control was not applied to the final meta-analysis results. We used the function ‘clump’ implemented in PLINK to cluster the SNPs into genomic loci using a 2-Mb window; clustering was done separately for each phenotype. Inverse variance meta-analysis was used to quantify effect sizes for SNPs of interest.

Genome-wide significance was inferred at  $P < 1 \times 10^{-8}$ . This choice of statistical threshold was grounded on the guidelines derived from studies of the ENCODE (encyclopedia of DNA elements) regions<sup>6</sup>, combined with results of permutation testing to determine the additional adjustment needed for the six red blood cell phenotypes studied (Supplementary Tables 24, 25 and Supplementary Note). As an alternative strategy, a  $P$ -value threshold of  $P < 3.2 \times 10^{-9}$  would provide correction for the number of SNP–phenotype combinations tested without any adjustment for the correlations between the SNPs or phenotypes tested. We note that 70 of the 75 loci identified would exceed such a highly stringent threshold, including all four of the loci identified through the joint analysis of European and South Asian data.

### Replication testing

We carried out replication testing of 22 SNPs selected on the basis of the following criteria: (1) the lead SNP from each of 17 loci showing suggestive evidence for association with one or more red blood cell phenotypes in Europeans ( $P > 10^{-8}$  and  $P < 10^{-7}$ ), and (2) the lead SNP from each of the loci identified through combined analysis of genome-wide association data for Europeans and South Asians. Replication testing was done using a combination of *in silico* results and direct genotyping among 63,506 people from four population cohorts.

*In silico* data were available for 34,843 people from Iceland participating in the deCODE (diabetes epidemiology: collaborative analysis of diagnostic criteria in Europe) study<sup>37</sup> (Supplementary Table 1). SNPs were directly genotyped with the Illumina HumanHap300 or CNV370 chips or imputed from one or more of four sources: the HapMap2 CEU sample (60 triads), the 1000 Genomes Project data (179 individuals) and Icelandic samples genotyped with the Illumina Human1 M-Duo (123 triads) or the HumanOmni1-Quad chips (505 individuals), as previously described in ref. 37. The 22 SNPs were tested for association against their respective discovery phenotypes, under an additive genetic model; results were combined with the genome-wide association data by weighted- $Z$ -score meta-analysis.

We found that for 7 of the 22 SNPs carried forward for replication, their associations with phenotype remained inconclusive after *in silico* testing ( $P > 10^{-8}$  but  $P < 10^{-7}$ ). For these SNPs we carried out additional direct genotyping using Sequenom assays, among up to

20,066 people from three population cohorts (Supplementary Table 1). Associations were tested in each cohort separately, and results combined across the replication cohorts, and then with the genome-wide association data, by weighted-*Z*-score meta-analysis (Supplementary Table 26).

### Conditional analysis

We performed conditional-association analysis using the summary statistics from the meta-analysis to test for the association of each SNP while conditioning on the top SNPs, with correlations between SNPs due to linkage disequilibrium estimated from the imputed genotype data from the atherosclerosis risk in communities (ARIC) cohort<sup>8,38</sup>. Secondary-association signals were selected with conditional-association  $P < 1 \times 10^{-8}$ .

### Identification of candidate genes

We considered the nearest gene, and any other gene located within 10 kb of the sentinel SNP, to be a candidate for mediating the association with red blood cell phenotype. We also used coding variant, eQTL and literature analyses to identify candidate genes. On the basis of analysis of linkage-disequilibrium relations at the 75 genetic loci, we defined genomic region as the 1-Mb interval either side of the sentinel SNP for our functional genomic studies (Supplementary Fig. 7).

### Coding variation

We identified all non-synonymous SNPs that were in linkage disequilibrium with one or more of the sentinel SNPs at  $r^2 > 0.8$  in 1000 Genomes Project data set (released in March 2012). We considered the gene to be a candidate when the non-synonymous and sentinel SNPs were in linkage disequilibrium at  $r^2 > 0.8$  and with no evidence for heterogeneity of effect on phenotype. This strategy identified 39 non-synonymous SNPs distributed between 24 genes (Supplementary Table 9), representing a ~sixfold enrichment compared to the mean number expected under the null hypothesis generated by permutation testing of SNP sets matched for allele frequency ( $\pm 0.05$ ) and number of genes in proximity ( $\pm 10$  kb), but selected otherwise at random ( $P = 0.01$ ; Supplementary Note).

### Expression analyses

To identify the possible genes influencing red blood cell phenotypes at the 75 loci, we examined the association of the sentinel SNPs with eQTL data from two data sets: (1) peripheral blood lymphocytes from 206 families of European descent (830 parents and offspring)<sup>39</sup> and (2) peripheral blood lymphocytes from 1,469 unrelated individuals<sup>40</sup>.

SNPs were tested for association with expression of nearby (1 Mb) genes ( $P < 0.05$  after Bonferroni correction for number of SNP–transcript associations tested). Where eQTLs were identified, we used the whole-genome SNP data available in these data sets (imputed with HapMap Phase 2 genotypes), to identify the SNP at the locus most closely associated with transcript level (the transcript SNP). We then tested whether the sentinel SNP and the transcript SNP were coincident, defined as  $r^2 > 0.8$  with no evidence for heterogeneity of effect on phenotype or transcript level ( $P > 0.05$ ). This strategy identified eQTLs involving 28 genes from 18 loci (Supplementary Table 10).

### GRAIL analyses

We carried out a literature analysis using the GRAIL algorithm<sup>9</sup>, a statistical tool that uses text mining of PubMed abstracts to annotate candidate genes from loci associated with phenotypic traits. We carried out the analysis using the 2006 data set to avoid confounding by subsequent GWAS discoveries; results identified candidate genes at nine loci ( $P < 0.05$ ;

Supplementary Table 11). Results are also shown for a GRAIL analysis using the 2011 PubMed data set, although these were not used for the final analysis.

### Gene expression in haematopoietic precursors

Cord-blood-derived CD34<sup>+</sup> haematopoietic stem cells were differentiated *in vitro* along the erythroid lineage in the presence of 6 U ml<sup>-1</sup> erythropoietin (R&D Systems), 10 ng ml<sup>-1</sup> inter-leukin (IL)-3 (Miltenyi Biotec) and 100 ng ml<sup>-1</sup> stem cell factor (R&D Systems). Cells were collected at days 3, 5, 7, 9 and 10 in three biological replicates and gene expression was assayed using Illumina human WGv3.0 microarrays<sup>41</sup>. For each gene, we determined the relationship of gene expression with time using linear regression, and calculated the *t*-statistic for the difference in  $\beta$  from zero. We then classified gene-expression patterns as increasing, decreasing or unchanged on the basis of the 2.5% and 97.5% quartiles of the *t* distribution with 4 degrees of freedom. To test whether a gene set was enriched for differentially regulated genes, a Wilcoxon signed-rank test of the *t* scores in the gene set relative to all other genes that were expressed in at least one time point was calculated.

### FAIRE-seq

We generated maps of chromatin accessibility ('open chromatin') in primary human erythroblasts and megakaryocytes, and in peripheral blood monocytes using FAIRE-seq. Cord-blood-derived CD34<sup>+</sup> haematopoietic progenitor cells from two unrelated individuals were differentiated *in vitro* into either erythroblasts (in the presence of erythropoietin, IL-3 and stem cell factor) or megakaryocytes (in the presence of thrombopoietin and IL-1 $\beta$ ). Monocytes were purified from leukocyte cones of apheresis collections from another two individuals.

FAIRE experiments were performed as previously described in ref. 42. FAIRE DNA was processed following the Illumina paired-end library-generation protocol. Genomic libraries derived from erythroblast and megakaryocyte cultures were sequenced with 54-bp paired-end reads on Illumina Genome Analyzer II. Libraries derived from monocyte extractions were sequenced with 50-bp paired-end reads on Illumina HiSeq. Raw sequence reads were aligned to the human reference sequence (NCBI build 37) using the read mapper Stampy<sup>43</sup>. Reads were realigned around known insertions and deletions, followed by base-quality recalibration using the Genome Analysis Toolkit (GATK)<sup>44</sup>. Duplicates were flagged using the software Picard (<http://picard.sourceforge.net/>) and excluded from subsequent analyses. For each cell type, we merged all read fragments into one data set. NDRs were identified as regions of sequencing enrichment (peaks) using the software F-Seq<sup>36</sup>. We applied a feature length of  $L = 600$  bp and a s.d. threshold of  $T = 8.0$  over the mean across a local background. In order to reduce false-positive peak calls, we removed regions of collapsed repeats as recently described, applying a threshold of 0.1%<sup>45</sup>. For each associated locus, candidate functional SNPs were selected by identifying all biallelic SNPs with an  $r^2 > 0.8$  and within 1 Mb of the sentinel SNP in the European samples of the 1000 Genomes Project (data released June 2011).

### D. melanogaster gene-silencing models

We used haemocyte-specific RNAi silencing to investigate whether the 121 candidate genes identified in the red blood cell GWAS influenced blood cell formation in *D. melanogaster*. We identified *D. melanogaster* genes predicted to be orthologues of human genes using the Ensembl v65 Compara pipeline, an established phylogenetic-tree-based approach for orthology prediction<sup>46</sup>; this revealed 96 *D. melanogaster* orthologues for 74 of the 121 human candidate genes (Supplementary Table 27). We evaluated each of the 96 orthologues for a blood cell phenotype in *D. melanogaster*. We obtained all 225 available *D. melanogaster* lines carrying inducible siRNA constructs from the Vienna *Drosophila* RNAi

Center (VDRC)<sup>23</sup>. To achieve haemocyte-specific knockdowns, flies were crossed to the blood-specific *Hml-Gal4* line driving Gal4 expression under the control of a hemolymph promoter<sup>47</sup>. Flies were crossed at 29 °C, and early and late L3 larvae analysed 7 days after mating. Upstream activating sequence–green fluorescent protein enabled microscopic visualization of plasmotocytes and evaluation of cell size and cell number (L3 larvae only). Early- and late-stage larvae were incubated at 60 °C for 15 min, a process that turns the crystal cells black and allows quantification of crystal cells microscopically. For each orthologue, all available RNAi silencer constructs were investigated, and in addition, each construct was assayed in duplicate, blind to initial result. Cell counts were quantified visually (0–3, decreased or increased) and the mean of the duplicate measurements calculated.

We separately carried out permutation testing in a genome-wide screen of 5,658 *D. melanogaster* genes to simulate expectations under the null hypothesis (Supplementary Fig. 8 and Supplementary Note); results confirmed that the 121 candidate genes were enriched for blood cell phenotype in *D. melanogaster* orthologues ( $P < 0.05$ ), and showed that this was robust to threshold for calling.

### Contribution of the genetic loci identified to population variation in red blood cell phenotypes

This was investigated in participants from the Estonian Genome Center of University of Tartu (EGCUT), LIFELINES, Ludwigshafen Risk and Cardiovascular Health Study (LURIC) and Young Finns cohorts using samples that were not included in the discovery experiment (Supplementary Table 1). The contribution of the SNPs to population variation in red blood cell phenotypes was quantified using two models: model 1, limited to SNPs associated with respective phenotype at  $P < 1 \times 10^{-8}$ ; and model 2, comprising all of the 75 sentinel SNPs identified. Estimates of population variance explained were made in each study separately, and average values calculated weighted by sample size (Supplementary Table 21).

We then investigated whether the 75 sentinel SNPs influenced the probability of being in the highest versus the lowest quartile for population distribution of phenotype. Two SNP scores were calculated for each phenotype: score 1, limited to SNPs associated with respective phenotype at  $P < 1 \times 10^{-8}$ , and score 2, containing all 75 sentinel SNPs identified. For both, SNP score was calculated as the sum of number of effect (trait raising) alleles present, weighted according to effect size. We then calculated the odds ratio for being in the highest versus the lowest quartile of phenotype, associated with SNP scores in the second, third and fourth quartiles, compared to first quartile of SNP score. Odds ratios were calculated in each study separately, and then combined by inverse variance meta-analysis (Fig. 3).

### Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

### Acknowledgments

A detailed list of acknowledgements is provided in the Supplementary Material.

### References


1. Chambers JC, et al. Genome-wide association study identifies variants in *TM6SS6* associated with hemoglobin levels. *Nature Genet.* 2009; 41:1170–1172. [PubMed: 19820698]

2. Ganesh SK, et al. Multiple loci influence erythrocyte phenotypes in the CHARGE Consortium. *Nature Genet.* 2009; 41:1191–1198. [PubMed: 19862010]
3. Soranzo N, et al. A genome-wide meta-analysis identifies 22 loci associated with eight hematological parameters in the HaemGen consortium. *Nature Genet.* 2009; 41:1182–1190. [PubMed: 19820697]
4. Kamatani Y, et al. Genome-wide association study of hematological and biochemical traits in a Japanese population. *Nature Genet.* 2010; 42:210–215. [PubMed: 20139978]
5. Ding K, et al. Genetic loci implicated in erythroid differentiation and cell cycle regulation are associated with red blood cell traits. *Mayo Clin Proc.* 2012; 87:461–474. [PubMed: 22560525]
6. Pe'er I, Yelensky R, Altshuler D, Daly MJ. Estimation of the multiple testing burden for genomewide association studies of nearly all common variants. *Genet Epidemiol.* 2008; 32:381–385. [PubMed: 18348202]
7. Nyholt DR. A simple correction for multiple testing for single-nucleotide polymorphisms in linkage disequilibrium with each other. *Am J Hum Genet.* 2004; 74:765–769. [PubMed: 14997420]
8. Yang J, et al. Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nature Genet.* 2012; 44:369–375. [PubMed: 22426310]
9. Raychaudhuri S, et al. Identifying relationships among genomic disease regions: predicting genes at pathogenic SNP associations and rare deletions. *PLoS Genet.* 2009; 5:e1000534. [PubMed: 19557189]
10. An X, Mohandas N. Disorders of red cell membrane. *Br J Haematol.* 2008; 141:367–375. [PubMed: 18341630]
11. van Wijk R, Rijksen G, Huizinga EG, Nieuwenhuis HK, van Solinge WW. HK Utrecht: missense mutation in the active site of human hexokinase associated with hexokinase deficiency and severe nonspherocytic hemolytic anemia. *Blood.* 2003; 101:345–347. [PubMed: 12393545]
12. Camaschella C, Poggiali E. Inherited disorders of iron metabolism. *Curr Opin Pediatr.* 2011; 23:14–20. [PubMed: 21150441]
13. Novershtern N, et al. Densely interconnected transcriptional circuits control cell states in human hematopoiesis. *Cell.* 2011; 144:296–309. [PubMed: 21241896]
14. Gieger C, et al. New gene functions in megakaryopoiesis and platelet formation. *Nature.* 2011; 480:201–208. [PubMed: 22139419]
15. Paul DS, et al. Maps of open chromatin guide the functional follow-up of genome-wide association signals: application to hematological traits. *PLoS Genet.* 2011; 7:e1002139. [PubMed: 21738486]
16. Forrester WC, Thompson C, Elder JT, Groudine M. A developmentally stable chromatin structure in the human beta-globin gene cluster. *Proc Natl Acad Sci USA.* 1986; 83:1359–1363. [PubMed: 3456593]
17. Tuan D, Solomon W, Li Q, London IM. The “beta-like-globin” gene domain in human erythroid cells. *Proc Natl Acad Sci USA.* 1985; 82:6384–6388. [PubMed: 3879975]
18. Kowalczyk MS, et al. Intragenic enhancers act as alternative promoters. *Mol Cell.* 2012; 45:447–458. [PubMed: 22264824]
19. Baù D, et al. The three-dimensional folding of the  $\alpha$ -globin gene domain reveals formation of chromatin globules. *Nature Struct Mol Biol.* 2011; 18:107–114. [PubMed: 21131981]
20. Zsebo KM, et al. Stem cell factor is encoded at the *Sf* locus of the mouse and is the ligand for the *c-kit* tyrosine kinase receptor. *Cell.* 1990; 63:213–224. [PubMed: 1698556]
21. Heissig B, et al. Recruitment of stem and progenitor cells from the bone marrow niche requires MMP-9 mediated release of kit-ligand. *Cell.* 2002; 109:625–637. [PubMed: 12062105]
22. Kozar K, et al. Mouse development and cell proliferation in the absence of D-cyclins. *Cell.* 2004; 118:477–491. [PubMed: 15315760]
23. Dietzl G, et al. A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. *Nature.* 2007; 448:151–156. [PubMed: 17625558]
24. Clark RH, et al. Adaptor protein 3-dependent microtubule-mediated movement of lytic granules to the immunological synapse. *Nature Immunol.* 2003; 4:1111–1120. [PubMed: 14566336]

25. Berhane S, et al. Adenovirus E1A interacts directly with, and regulates the level of expression of, the immunoproteasome component MECL1. *Virology*. 2011; 421:149–158. [PubMed: 22018786]
26. Tiwari S, Weissman AM. Endoplasmic reticulum (ER)-associated degradation of T cell receptor subunits. Involvement of ER-associated ubiquitin-conjugating enzymes (E2s). *J Biol Chem*. 2001; 276:16193–16200. [PubMed: 11278356]
27. Fransen K, et al. Analysis of SNPs with an effect on gene expression identifies UBE2L3 and BCL3 as potential new risk genes for Crohn's disease. *Hum Mol Genet*. 2010; 19:3482–3488. [PubMed: 20601676]
28. Zhernakova A, et al. Meta-analysis of genome-wide association studies in celiac disease and rheumatoid arthritis identifies fourteen non-HLA shared loci. *PLoS Genet*. 2011; 7:e1002004. [PubMed: 21383967]
29. Das S, Ghosh R, Maitra U. Eukaryotic translation initiation factor 5 functions as a GTPase-activating protein. *J Biol Chem*. 2001; 276:6720–6726. [PubMed: 11092890]
30. Fenton TR, Gout IT. Functions and regulation of the 70 kDa ribosomal S6 kinases. *Int J Biochem Cell Biol*. 2011; 43:47–59. [PubMed: 20932932]
31. Scanlon KS, Yip R, Schieve LA, Cogswell ME. High and low hemoglobin levels during pregnancy: differential risks for preterm birth and small for gestational age. *Obstet Gynecol*. 2000; 96:741–748. [PubMed: 11042311]
32. Shah RC, Buchman AS, Wilson RS, Leurgans SE, Bennett DA. Hemoglobin level in older persons and incident Alzheimer disease: prospective cohort analysis. *Neurology*. 2011; 77:219–226. [PubMed: 21753176]
33. Sabatine MS, et al. Association of hemoglobin levels with clinical outcomes in acute coronary syndromes. *Circulation*. 2005; 111:2042–2049. [PubMed: 15824203]
34. Zakai NA, et al. A prospective study of anemia status, hemoglobin concentration, and mortality in an elderly cohort: the Cardiovascular Health Study. *Arch Intern Med*. 2005; 165:2214–2220. [PubMed: 16246985]
35. Galanello R, et al. Amelioration of Sardinian  $\beta^0$  thalassemia by genetic modifiers. *Blood*. 2009; 114:3935–3937. [PubMed: 19696200]
36. Boyle AP, Guinney J, Crawford GE, Furey TS. F-Seq: a feature density estimator for high-throughput sequence tags. *Bioinformatics*. 2008; 24:2537–2538. [PubMed: 18784119]
37. Holm H, et al. A rare variant in MYH6 is associated with high risk of sick sinus syndrome. *Nature Genet*. 2011; 43:316–320. [PubMed: 21378987]
38. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet*. 2011; 88:76–82. [PubMed: 21167468]
39. Dixon AL, et al. A genome-wide association study of global gene expression. *Nature Genet*. 2007; 39:1202–1207. [PubMed: 17873877]
40. Dubois PC, et al. Multiple common variants for celiac disease influencing immune gene expression. *Nature Genet*. 2010; 42:295–302. [PubMed: 20190752]
41. Anderson RJ, et al. Reduced dependency on arteriography for penetrating extremity trauma: influence of wound location and noninvasive vascular studies. *J Trauma*. 1990; 30:1059–1063. [PubMed: 2213941]
42. Giresi PG, Lieb JD. Isolation of active regulatory elements from eukaryotic chromatin using FAIRE (Formaldehyde Assisted Isolation of Regulatory Elements). *Methods*. 2009; 48:233–239. [PubMed: 19303047]
43. Lunter G, Goodson M. Stampy: a statistical algorithm for sensitive and fast mapping of Illumina sequence reads. *Genome Res*. 2011; 21:936–939. [PubMed: 20980556]
44. McKenna A, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res*. 2010; 20:1297–1303. [PubMed: 20644199]
45. Pickrell JK, Gaffney DJ, Gilad Y, Pritchard JK. False positive peaks in ChIP-seq and other sequencing-based functional assays caused by unannotated high copy number regions. *Bioinformatics*. 2011; 27:2144–2146. [PubMed: 21690102]
46. Vilella AJ, et al. EnsemblCompara GeneTrees: complete, duplication-aware phylogenetic trees in vertebrates. *Genome Res*. 2009; 19:327–335. [PubMed: 19029536]




47. Goto A, et al. A *Drosophila* haemocyte-specific protein, hemolectin, similar to human von Willebrand factor. *Biochem J.* 2001; 359:99–108. [PubMed: 11563973]




**Figure 1. Gene-expression patterns for 121 putative candidate genes, and tissue distribution of NDRs**

**a.** Heat-map of candidate genes in the Differentiation Map of Hematology<sup>13</sup>. Cell acronyms refer to original source (summarized in Supplementary Table 15). Expression above a  $\log_2$  signal intensity (SI) of 6 is consistently above background. **b.**  $-\log_{10} P$  of the signed-rank test for candidate genes being more highly expressed in each cell type than non-candidate genes. **c.** Time-course of differentiation of cord-blood haematopoietic stem cells cultured along the erythroid lineage. Putative candidate genes are shown as upregulated (red), downregulated (blue) or with the slope not being significantly different from zero (grey). **d.** Tissue distribution of NDRs containing a potential causal variant. NDRs were ranked by peak score (proportional to their peak height in FAIRE-seq). The rankings were then used to divide the NDRs into cumulative tranches to explore the effect of calling-thresholds on results (left bar, tranche containing the 5,000 top-ranked NDRs of each cell type; penultimate bar, tranche containing the 50,000 top-ranked NDRs of each cell type). The solid line indicates the number of SNPs overlapping the tranche-specific NDRs that are potential causal variants (defined as a sentinel SNP from the red blood cell GWAS, or a SNP in high linkage disequilibrium ( $r^2 > 0.8$ ) and located within 1 Mb of a sentinel SNP; right-hand y axis); the bar summarizes the tissue distribution of these SNPs (as a percentage of tranche-specific total). The right-hand bar represents the expected tissue distribution for the SNPs under the null hypothesis. Results show that the potential causal variants are most commonly found in erythroblast-specific NDRs, and that this is true across the spectrum of peak-calling thresholds.



**Figure 2. RNAi silencing in *D. melanogaster***

**a**, Plasmatocytes imaged by green fluorescent protein expression (light green spots on posterior dorsal end of L3 larvae) from wild-type (WT) cells and cells with RNAi silencing of orthologues of the following human genes: *CRHR1* (106381, increased cell counts (CC)), *KIT* (13502, decreased CC) and *CCNA2* (32421, increased CC). Numbers represent the unique Flybase IDs corresponding to the *D. melanogaster* orthologues. Scale bar, 0.5 mm. Bottom right, plasmatocyte size is also increased in *CCNA2* compared to wild type. Scale bars, 0.1 mm. **b**, Crystal cells (black spots visualized by heating larvae to 60 °C) in wild-type larvae, and in RNAi silencing of *ATP50* (12794, increased CC), *UBE2L3* (110767, decreased CC) or *ATP2B4* (101743, aggregated). Scale bars, 0.5 mm.



**Figure 3. Association of SNP score with red blood cell phenotypes**

Results presented as odds ratio (95% confidence interval) for participants in each SNP score quartile (Q) having phenotype level in the top quartile versus the lowest quartile of the respective population distribution, compared to people in the lowest quartile of SNP score (Q1, reference group). HB, haemoglobin; *n*, number of participants in the respective comparison of SNP score quartiles.

Table 1

Genomic loci associated with red blood cell phenotypes

Region	Sentinel SNP	Position (B36)	Alleles (EA/OA)	EAF	Phenotype	Effect (SE)	P	Candidate genes
1p36	rs1175550	3,681,388	G/A	0.22	MCHC	0.008 (0.013)	$8.6 \times 10^{-15}$	<i>CCDC27<sup>n</sup>, LRRCA8<sup>n</sup></i>
1p34 <sup>†</sup>	rs3916164	39,842,526	G/A	0.71	MCH	0.008 (0.004)	$3.1 \times 10^{-10}$	<i>HEYL<sup>n</sup></i>
1p32	rs741959	47,448,820	G/A	0.57	MCV	0.157 (0.025)	$6.0 \times 10^{-10}$	<i>TAL1<sup>n</sup></i>
1q23 <sup>†</sup>	rs857684	156,842,353	C/T	0.74	MCHC	-0.006 (0.011)	$3.5 \times 10^{-16}$	<i>OR6Y1<sup>n</sup>, OR10Z1<sup>n</sup>, SPTA1<sup>n</sup></i>
1q32 <sup>†</sup>	rs7529925	197,273,831	C/T	0.28	RBC	0.014 (0.002)	$8.3 \times 10^{-9}$	<i>MIR181A<sup>n</sup></i>
1q32	rs7551442	201,921,744	A/G	0.09	MCHC	-0.023 (0.017)	$9.7 \times 10^{-12}$	<i>ATP2B4<sup>n</sup></i>
1q32	rs9660992	203,516,073	G/A	0.42	MCH	0.007 (0.004)	$7.1 \times 10^{-10}$	<i>TMCC2<sup>n</sup></i>
1q44 <sup>†</sup>	rs3811444	246,106,074	T/C	0.35	RBC	0.018 (0.003)	$4.5 \times 10^{-10}$	<i>TRIM58<sup>n</sup></i>
2p21 <sup>†</sup>	rs4953318	46,208,555	A/C	0.62	PCV	0.152 (0.018)	$3.1 \times 10^{-19}$	<i>PRKCE<sup>n</sup></i>
2p16 <sup>†</sup>	rs243070	60,473,790	T/A	0.72	MCV	-0.181 (0.027)	$4.4 \times 10^{-13}$	<i>BCL11A<sup>n</sup></i>
2q13	rs10207392	111,566,130	G/A	0.44	MCV	-0.132 (0.025)	$4.4 \times 10^{-11}$ *	<i>ACOXL<sup>n</sup></i>
3p24 <sup>†</sup>	rs9310736	24,325,815	A/G	0.35	MCV	-0.210 (0.026)	$6.1 \times 10^{-16}$	<i>THRB<sup>n</sup></i>
3q22	rs6776003	142,749,183	A/G	0.44	MCV	-0.138 (0.026)	$3.7 \times 10^{-11}$ *	<i>RASA2<sup>n</sup></i>
3q23	rs13061823	143,603,476	T/C	0.56	MCV	-0.168 (0.025)	$4.7 \times 10^{-13}$	<i>XRN1<sup>n</sup></i>
3q29 <sup>†</sup>	rs11717368	197,318,754	C/G	0.52	MCH	0.008 (0.004)	$6.6 \times 10^{-19}$	<i>TFRC1<sup>n</sup></i>
4q11 <sup>†</sup>	rs218238	55,089,781	A/T	0.78	RBC	0.033 (0.003)	$2.8 \times 10^{-39}$	<i>KIT<sup>n</sup></i>
4q27	rs13152701	122,970,511	A/G	0.37	MCV	0.150 (0.026)	$9.0 \times 10^{-10}$	<i>BBS7<sup>n</sup>, CCNA2<sup>n</sup></i>
6p23	rs6914805	16,389,166	C/T	0.75	MCH	0.012 (0.004)	$1.2 \times 10^{-19}$	<i>GMPR1<sup>n</sup></i>
6p21 <sup>†</sup>	rs1408272	25,950,930	G/T	0.07	MCH	0.033 (0.009)	$4.8 \times 10^{-67}$	<i>HFE<sup>n</sup>, SLC17A3<sup>n</sup></i>
6p22	rs13219787	27,969,649	A/G	0.09	MCH	0.023 (0.007)	$5.9 \times 10^{-17}$	<i>HIST1H2AM<sup>n</sup>, HIST1H2BO<sup>n</sup>, HIST1H3<sup>n</sup></i>
6p22	rs2097775	30,462,282	A/T	0.15	HB	0.055 (0.008)	$1.3 \times 10^{-10}$	<i>TRIM39-RPP2<sup>n</sup></i>
6p21	rs9272219	32,710,247	G/T	0.72	RBC	0.015 (0.002)	$4.3 \times 10^{-10}$	<i>HLA-DQA1<sup>n</sup>, HLA-DQA2<sup>n</sup></i>
6p21 <sup>†</sup>	rs9349204	42,022,356	G/A	0.27	MCV	-0.367 (0.028)	$2.4 \times 10^{-40}$	<i>CCND3<sup>n</sup></i>
6p12	rs9369427	43,919,408	A/C	0.68	HB	0.042 (0.006)	$5.6 \times 10^{-12}$	<i>VEGFA<sup>n</sup></i>
6q21 <sup>†</sup>	rs1008084	109,733,658	G/A	0.56	MCH	-0.010 (0.003)	$6.4 \times 10^{-26}$	<i>CCDC162<sup>n</sup></i>

Region	Sentinel SNP	Position (B36)	Alleles (EA/OA)	EAF	Phenotype	Effect (SE)	P	Candidate genes
6q23.1	rs9389269	135,468,852	T/C	0.72	MCV	-0.600 (0.028)	$2.6 \times 10^{-19}$	<i>HBSIL</i> <sup>n</sup>
6q24.1	rs590856	139,886,122	G/A	0.43	MCV	0.313 (0.026)	$5.0 \times 10^{-36}$	<i>CITED2</i> <sup>n</sup>
6q26	rs736661	164,402,826	A/G	0.62	MCH	0.007 (0.004)	$1.6 \times 10^{-11}$	<i>QKP</i> <sup>n</sup>
7p13.1	rs12718598	50,395,939	T/C	0.51	MCV	-0.204 (0.030)	$1.6 \times 10^{-13}$	<i>IKZF1</i> <sup>n</sup>
7q22.1	rs2075672	100,078,232	A/G	0.39	RBC	0.022 (0.003)	$1.9 \times 10^{-20}$	<i>ACTL6B</i> <sup>n</sup> , <i>TFR2</i> <sup>ng</sup>
7q36.1	rs10480300	151,036,938	C/T	0.72	HB	0.052 (0.007)	$7.8 \times 10^{-15}$	<i>PRKAG2</i> <sup>ng</sup>
8p11	rs4737009	41,749,562	G/A	0.74	MCHC	-0.014 (0.013)	$4.9 \times 10^{-11}$	<i>ANK</i> <sup>ng</sup>
8p11	rs6987853	42,576,607	C/T	0.62	MCHC	-0.002 (0.010)	$6.1 \times 10^{-11}$	<i>C8orf40</i> <sup>ng</sup>
9p24.1	rs2236496	4,834,265	C/T	0.22	MCV	-0.279 (0.031)	$1.4 \times 10^{-19}$	<i>RCL1</i> <sup>n</sup>
9q34.1	rs579459	135,143,989	T/C	0.8	RBC	0.021 (0.003)	$9.3 \times 10^{-18}$	<i>ABO</i> <sup>n</sup>
10q11.1	rs901683	45,286,428	A/G	0.08	MCV	0.364 (0.050)	$1.5 \times 10^{-16}$	<i>MARCH8</i> <sup>ng</sup>
10q22.1	rs10159477	70,769,894	A/G	0.16	HB	0.087 (0.010)	$4.4 \times 10^{-20}$	<i>HK1</i> <sup>ng</sup>
10q24	rs11190134	101,272,190	G/A	0.6	MCH	-0.011 (0.004)	$1.3 \times 10^{-10}$ *	<i>NKX2-3</i> <sup>n</sup>
11p15	rs11042125	8,894,625	A/T	0.6	HB	0.032 (0.006)	$1.5 \times 10^{-9}$	<i>AKIP1</i> <sup>ng</sup> , <i>C11orf10</i> <sup>ng</sup> , <i>NRIP3</i> <sup>n</sup> , <i>ST5</i> <sup>n</sup>
11p15	rs7936461	9,997,462	C/T	0.75	PCV	0.121 (0.021)	$1.0 \times 10^{-9}$	<i>SBF2</i> <sup>n</sup>
11q13	rs2302264	66,964,002	G/A	0.58	MCV	0.140 (0.025)	$1.3 \times 10^{-10}$	<i>CORO1B</i> <sup>ng</sup> , <i>PTPRCAP</i> <sup>ng</sup> , <i>RPS6KB2</i> <sup>ng</sup>
11q13	rs7125949	72,686,732	A/G	0.11	HB	0.053 (0.010)	$2.1 \times 10^{-9}$	<i>ARHGGEF17</i> <sup>ng</sup> , <i>P2RY6</i> <sup>n</sup>
12p13	rs7312105	2,393,616	G/A	0.36	PCV	0.104 (0.019)	$3.2 \times 10^{-9}$ *	<i>CACNA1C</i> <sup>n</sup>
12p13	rs10849023	4,202,739	C/T	0.79	MCH	-0.008 (0.005)	$7.5 \times 10^{-12}$	<i>CCND2</i> <sup>ng</sup>
12q22	rs11104870	87,353,425	C/T	0.3	RBC	0.013 (0.002)	$6.2 \times 10^{-11}$ *	<i>KITLG</i> <sup>n</sup>
12q24.1	rs3184504	110,368,991	T/C	0.48	HB	0.051 (0.006)	$4.3 \times 10^{-19}$	<i>ATXN2</i> <sup>n</sup> , <i>SH2B3</i> <sup>ng</sup>
12q24	rs3829290	119,610,821	C/T	0.44	MCV	-0.153 (0.026)	$2.1 \times 10^{-9}$	<i>ACADS</i> <sup>n</sup> , <i>MLEC</i> <sup>n</sup>
14q23.1	rs7155454	64,571,992	A/G	0.51	MCH	0.002 (0.004)	$1.8 \times 10^{-12}$	<i>FNTB</i> <sup>n</sup> , <i>MAX</i> <sup>n</sup>
14q24	rs11627546	69,435,677	C/A	0.84	MCV	0.162 (0.032)	$1.1 \times 10^{-9}$ *	<i>SMOC1</i> <sup>n</sup>
14q32.1	rs17616316	102,892,515	G/C	0.07	MCH	0.014 (0.009)	$8.2 \times 10^{-11}$ *	<i>EIF5</i> <sup>n</sup>
15q21.1	rs1532085	56,470,658	G/A	0.59	HB	0.034 (0.006)	$6.7 \times 10^{-11}$ *	<i>LIPC</i> <sup>n</sup>
15q22.1	rs2572207	63,857,747	C/T	0.74	MCV	0.153 (0.029)	$3.4 \times 10^{-9}$	<i>DENND4A</i> <sup>n</sup> , <i>PTPLAD1</i> <sup>n</sup>

Region	Sentinel SNP	Position (B36)	Alleles (EA/OA)	EAF	Phenotype	Effect (SE)	P	Candidate genes
15q24	rs8028632	73,108,315	T/C	0.8	MCV	0.188 (0.032)	$6.9 \times 10^{-10}$	<i>PPCDC<sup>h</sup>, SCAMP3<sup>h</sup></i>
15q24	rs11072566	74,081,026	A/G	0.48	HB	0.028 (0.006)	$3.0 \times 10^{-10}$ *	<i>NRG4<sup>h</sup></i>
15q25	rs2867932	76,378,092	G/A	0.61	MCHC	-0.021 (0.010)	$3.3 \times 10^{-9}$	<i>DNAJA4<sup>h</sup>, WDR6<sup>h</sup></i>
16p11 <sup>†</sup>	rs11248850	103,598	G/A	0.5	MCH	0.007 (0.004)	$6.3 \times 10^{-23}$	<i>NPRL3<sup>h</sup></i>
16q22	rs2271294	66,459,827	T/A	0.15	RBC	0.017 (0.003)	$1.1 \times 10^{-9}$	<i>CTRL<sup>h</sup>, DUS2L<sup>h</sup>, EDC4<sup>h</sup>, NUTTF2<sup>h</sup>, PSMB10<sup>h</sup></i>
16q24 <sup>†</sup>	rs10445033	87,367,963	G/A	0.37	MCHC	0.020 (0.012)	$1.5 \times 10^{-22}$	<i>PIEZO1<sup>h</sup></i>
17p11	rs888424	19,926,019	A/G	0.43	MCH	0.006 (0.004)	$5.4 \times 10^{-20}$	<i>SPECC1<sup>h</sup></i>
17q11	rs2070265	24,099,550	T/C	0.2	MCH	0.013 (0.004)	$5.1 \times 10^{-14}$	<i>C17orf63<sup>h</sup>, ERALF<sup>h</sup>, NEK8<sup>h</sup>, TRAF4<sup>h</sup></i>
17q12	rs8182252	34,981,476	C/T	0.18	RBC	0.016 (0.003)	$5.9 \times 10^{-9}$	<i>CDK12<sup>h</sup>, NEUROD2<sup>h</sup></i>
17q21	rs2269906	39,649,863	C/A	0.36	MCHC	0.027 (0.010)	$2.0 \times 10^{-11}$	<i>SLC44A1B<sup>h</sup>, UBT1<sup>h</sup></i>
17q21	rs12150672	41,182,408	A/G	0.23	RBC	0.017 (0.003)	$4.7 \times 10^{-12}$	<i>ARHGAP27<sup>h</sup>, ARL17B<sup>h</sup>, C17orf69<sup>h</sup>, CRHR1<sup>h</sup>, SPPL2C<sup>h</sup>, KANSL1<sup>h</sup>, MAPT<sup>h</sup>, STHF</i>
17q25	rs4969184	73,905,008	G/A	0.53	HB	0.031 (0.006)	$7.0 \times 10^{-9}$	<i>PGS1<sup>h</sup></i>
18q21	rs4890633	42,087,276	G/A	0.27	MCH	0.005 (0.004)	$1.9 \times 10^{-23}$	<i>C18orf25<sup>h</sup></i>
19p13	rs2159213	2,087,102	C/T	0.5	HB	0.032 (0.006)	$1.9 \times 10^{-9}$	<i>AP3D1<sup>h</sup></i>
19p13	rs732716	4,317,219	A/G	0.71	MCV	0.201 (0.028)	$1.5 \times 10^{-14}$	<i>MPND1, SH3GL1<sup>h</sup>, UBXYN6<sup>h</sup></i>
19p13 <sup>†</sup>	rs741702	12,885,250	A/C	0.35	MCH	0.006 (0.004)	$8.2 \times 10^{-20}$	<i>CALR<sup>h</sup>, FARS4<sup>h</sup>, SYCE2<sup>h</sup></i>
19q13	rs3892630	37,873,324	T/C	0.18	MCV	0.176 (0.034)	$1.0 \times 10^{-10}$ *	<i>NUDT19<sup>h</sup></i>
20q13 <sup>†</sup>	rs737092	55,423,811	C/T	0.49	MCV	0.216 (0.033)	$4.0 \times 10^{-13}$	<i>RBM38<sup>h</sup></i>
21q22 <sup>†</sup>	rs2032314	34,276,393	T/C	0.08	PCV	0.154 (0.034)	$7.5 \times 10^{-10}$ *	<i>ATP5O1<sup>h</sup></i>
22q11 <sup>†</sup>	rs5754217	20,269,675	G/T	0.83	MCV	0.194 (0.031)	$8.6 \times 10^{-10}$	<i>UBE2L3<sup>h</sup>, YD1C<sup>h</sup></i>
22q12 <sup>†</sup>	rs5749446	31,210,585	T/C	0.62	MCH	0.007 (0.004)	$3.3 \times 10^{-13}$	<i>FBXO7<sup>h</sup></i>
22q12 <sup>†</sup>	rs855791	35,792,882	G/A	0.57	MCH	0.012 (0.004)	$1.0 \times 10^{-69}$	<i>KCTD17<sup>h</sup>, TMPRSS6<sup>h</sup></i>
22q13 <sup>†</sup>	rs140522	49,318,132	C/T	0.67	MCV	0.287 (0.030)	$4.5 \times 10^{-23}$	<i>TYMP<sup>h</sup>, NCAPH2<sup>h</sup>, ODF3B<sup>h</sup>, SCO2<sup>h</sup></i>

Candidate gene superscripts indicate the method of identification.

<sup>h</sup> Replication testing performed.

<sup>†</sup> Previously reported.

<sup>‡</sup> Discovered from combined analysis of European and South Asian genome-wide association data.

c, coding variant; e, eQTL; EA, effect allele; EAF, effect allele frequency; g, GRAIL; HB, haemoglobin; z, nearby; OA, other allele; SE, standard error.