



**Table 1 Clinical model for assessment of deep vein thrombosis**

Clinical finding	Points
Active cancer (treatment ongoing or within previous 6 months or palliative)	1
Paralysis, paresis or recent plaster immobilization of the lower extremities	1
Recent immobilization > 3 days or major surgery within 12 weeks requiring general or regional anesthesia	1
Localized tenderness along the distribution of the deep venous system	1
Entire leg swollen	1
Calf swelling 3 cm > asymptomatic side (measured 10 cm below tibial tuberosity)	1
Pitting edema confined to the symptomatic leg	1
Collateral superficial veins (nonvaricose)	1
Alternative diagnosis as likely or greater than that of DVT	-2

Adapted from [8]. Clinical probability for DVT calculated as follows: high (3); moderate (1 or 2); low (0). DVT, deep vein thrombosis.

different parameters on postoperative D-dimer levels was studied using univariate and multivariate analyses. The study was approved by the Ethics Committee of the institution and was registered at [clinicaltrials.gov](http://clinicaltrials.gov) (NCT00450528).

### Study population

Between January 2007 and July 2007, 154 patients undergoing abdominal surgery at our institution were included in the study, 108 in the study cohort and 46 in the validation cohort. Informed consent was obtained from all participants. Exclusion criteria were age less than 18 years, pregnancy, oral anticoagulation, history of former VTE and preoperative concomitant inflammatory diseases (e.g. sepsis, pneumonia). Additionally, patients were only allowed to participate in one single scientific study. Patients, who received fresh frozen plasma or fibrinogen during or after surgery or who had to be resubmitted to the operating theatre because of a complication, were secondarily excluded from the study. All patients received low molecular weight heparin (LMWH) once daily (Fragmin, dalteparin sodium, Pharmacia and Upjohn, Dubendorf, Switzerland; 5000 units subcutaneous) during their hospital stay, starting the evening before surgery. Additionally, graded compression stockings were used for all patients during the entire hospital stay. All patients were clinically evaluated at discharge to exclude VTE.

### Determination of D-dimers

In the first phase, plasma D-dimers levels were determined preoperatively (day 0) and at various time points after surgery (day 1, 3, 5, 7, 10, 14, and weekly thereafter until D-dimer values returned to normal). After hospital discharge, blood samples were taken in the outpatient clinic. A value of less than 500 ng/ml or equal was considered as normal. D-dimers were analyzed using a quantitative enzyme-linked immunosorbent assay (ELISA) (VIDAS D-dimer; bioMérieux, Geneva, Switzerland) as previously described [13].

### Stratification of surgical procedures, patients' risk and surgical morbidity

Surgical operations were stratified based on the degree of severity (type I: superficial surgery not opening the abdominal cavity such as open hernia repairs; type II: intraabdominal procedures such as colon resections; type III: retroperitoneal surgery such as pancreatic or esophageal resections and liver surgery) as previously defined [14]. Preoperative patients' risk was estimated by means of the grading systems of the American Society of Anesthesiologists (ASA) [15], the Nutrition Risk Score [16], and the Charlson Risk Index (CRI) [17]. The CRI is a comorbidity measure that contains a list of 19 different comorbidities. A numerical value from 1 to 6 is assigned to each of the comorbidities and the sum of these values reflects the CRI. In patients with a CRI of at least 3, mortality was shown to be doubled after surgery [18].

### Statistical analyses

Continuous data are presented as mean  $\pm$  standard deviation or median (range) and are compared between groups using the Mann-Whitney test or Kruskal-Wallis test in cases appropriate. Spearman rank correlations were used to analyze correlations between continuous variables. Categorical data are presented as number with percentage and are compared between groups using the  $\chi^2$  test [two degrees of freedom (DF) with the exception of type of surgery and ASA] or Fisher's exact test in cases appropriate. A logarithmic transformation of D-dimer values was used to approach a normal distribution. Analysis of covariance for repeated measures was applied to transformed data to analyze degradation over time. Paired *t*-tests were performed to compare values between days, and unpaired *t*-tests and analysis of variation (ANOVA) were used to compare values between groups. Box plots are given with medians and interquartile ranges (IQR). Outliers are indicated by open dots and asterisks: 'Extreme' outliers or those which lay more than three times the IQR (3 IQR) to the left and right from the first and third quartiles, respectively, are indicated by the presence of an asterisk. 'Mild' outliers – that is, those observations that lay more than 1.5 times the IQR from the first and third quartile, but are not extreme outliers, are indicated by the presence of a open dot. Single observations are indicated by a bar. SPSS 13.0 (SPSS, Inc., Chicago, Illinois, USA) was used for statistical analyses. Two-sided *P* values less or equal to 0.05 are considered statistically significant.

## Results

### Patient demographics

Informed consent has been obtained from 172 patients; 18 patients had to be excluded after surgery due to reoperation because of a complication (13 patients) or due to perioperative transfusions of fresh frozen plasma (five patients). A total of 154 patients were finally included in the study (108 as study cohort and 46 as

Table 2 Patient characteristics

	Overall <i>n</i> = 154	Study cohort <i>n</i> = 108	Validation cohort <i>n</i> = 46	<i>P</i>
Age; years (SD)	55.1 (±14.5)	55.5 (±14.4)	54.4 (±14.9)	0.98
Female patients, <i>n</i> (%)	70 (45.5)	49 (45.4)	21 (45.7)	0.91
D-dimer preoperative > 500 ng/ml; <i>n</i> (%)	36 (23.4)	22 (20.4)	14 (30.4)	0.12
Length of surgery; min (range)	180 (25–480)	180 (25–480)	240 (40–450)	0.04
Blood loss, ml (range)	100 (0–2000)	100 (0–2000)	150 (0–2000)	0.10
Length of hospitalization; days (range)	8 (1–90)	8 (2–90)	8 (1–64)	0.80
Laparoscopic procedures; <i>n</i> (%)	63 (40.9)	49 (45.4)	14 (30.4)	0.06
Cancer; <i>n</i> (%)	59 (38.3)	35 (32.4)	24 (52.2)	0.03
Type of surgery				0.78
Type I; <i>n</i> (%)	20 (13.0)	15 (13.9)	5 (10.9)	NA
Type II; <i>n</i> (%)	66 (42.9)	46 (42.6)	20 (43.5)	NA
Type III; <i>n</i> (%)	68 (44.1)	47 (43.5)	21 (45.7)	NA
ASA				0.93
ASA I; <i>n</i> (%)	30 (19.5)	22 (20.4)	8 (17.4)	NA
ASA II; <i>n</i> (%)	82 (53.2)	57 (52.8)	25 (54.3)	NA
ASA III; <i>n</i> (%)	42 (27.3)	29 (26.8)	13 (38.3)	NA

Types of surgery: I, without opening of abdominal cavity; II, abdominal procedures w/o retroperitoneal/liver surgery; III: retroperitoneal/liver surgery. ASA, American Society of Anesthesiologists; NA, not available.

validation cohort); Table 2 shows the patients characteristics. Mean age was 55.1 years (±14.5) with 45.5% female participants in the entire patient cohort. Cancer was present in 38.3% of the patients. Patients (19.5%) were estimated to be grade I according to the ASA, 53.2% of grade II and 27.3% of grade III. Few of the surgical procedures (40.9%) were performed laparoscopically. Thirteen percent of the procedures comprised type I surgery, 42.9% were type II and 44.1% type III procedures. Median duration of surgery was 180 min (25–480), median blood loss was 100 ml (0–2000) and median length of hospitalization was 8 days (1–90). Surgical morbidity rate was 22.7% (35/154). Of note, none of the patients presented with clinical signs or symptoms of VTE during hospitalization or during follow-up, and none of them was readmitted to the hospital with symptomatic VTE.

There was no difference between the study cohort and the validation cohort regarding age ( $P=0.98$ ), blood loss ( $P=0.10$ ), preoperative D-dimer levels ( $P=0.12$ ) and length of hospitalization ( $P=0.80$ ). However, length of surgery was significantly longer in the patients used for validation ( $P=0.04$ ) with more patients operated on for cancer ( $P=0.03$ ).

#### Kinetics of D-dimer

D-dimer level before surgery was 290 ng/ml (100–3800). In 34 out of 154 (22.1%) patients, D-dimer levels were found to be above normal values already before surgery. Malignant disease and age were significantly associated with an elevated D-dimer before surgery ( $P=0.01$  and 0.02, respectively). After surgery, D-dimer levels increased postoperatively reaching a peak on day 7 ( $P\leq 0.001$  vs. all other days except day 5). After type I surgery, peak D-dimer levels did not exceed normal range (300 ng/ml, 100–500). Peak D-dimer level after type II procedures was 1500 ng/ml (200–7800) and

returned to normal values after a period of 25 days (±14). After type III surgery, peak D-dimer level was 4000 ng/ml (500–14 400) and normalized within 38 days (±11). Forty-three out of 46 patients with a type II procedure and 39 of 47 patients undergoing a type III procedure, respectively, had D-dimer values within normal range after 25 days. After 38 days, D-dimer values were normal in all of the patients who underwent type II procedures and in 44 of 47 patients with a type III procedure.


Degradation of D-dimer was exponential after having reached the peak with a clearance of 6.0% per day [95% confidence interval (CI) 4.8–7.1%] in the study cohort. By this clearance, D-dimer values could be adequately predicted in the patients of the validation cohort after determination of the peak D-dimer level ( $r^2=0.72$  on a log scale). Clearance in the validation cohort was similar with 6.7% per day (95% CI 2.4–10.4%;  $P=0.65$ ). Daily clearance of the entire patient cohort was 6.1% (95% CI 4.9–7.2%). Hence, half-life time of postoperative D-dimers was 11 days. The kinetics of D-dimers after surgery is shown for the entire patient cohort for all procedures (Fig. 1) and stratified by surgical type (Fig. 2).

#### Determinants of peak D-dimer levels

In univariate analyses, peak D-dimer was significantly determined by age ( $P<0.001$ ), duration of surgery ( $P<0.001$ ), blood loss ( $P<0.001$ ), peak levels of C-reactive protein ( $P<0.001$ ), preoperative D-dimer levels (greater than normal;  $P<0.001$ ), length of hospitalization ( $P<0.001$ ), type of surgery ( $P<0.001$ ), the Nutrition Risk Score ( $P<0.001$ ), the CRI ( $P<0.001$ ), malignant disease ( $P<0.001$ ), surgical technique (open vs. laparoscopic;  $P<0.001$ ) and the ASA scoring ( $P=0.02$ ).

By multivariate regression analysis of univariate significant parameters, duration and type of surgery ( $P<0.001$ ,

Fig. 1



D-dimer values before (day 0) and after general surgery (n = 154).

$\beta = 0.233$ ,  $r^2 = 0.44$  and  $P < 0.001$ ,  $\beta = 0.338$ ,  $r^2 = 0.38$ ) and preoperative D-dimer levels (greater than normal;  $P < 0.001$ ,  $\beta = 0.278$ ,  $r^2 = 0.16$ ) were identified as independent determinants of peak D-dimer levels. By taking all independent risk factors into account, the prediction of the D-dimer levels could be markedly improved


( $r^2 = 0.63$ ; Fig. 3) compared with the predictive capacity of each factor alone.

### Discussion

Kinetics of D-dimer is largely unknown limiting the use of D-dimer testing for VTE exclusion after surgery. This is the first study showing the postoperative kinetics of D-dimers after abdominal surgery. D-dimer levels peak on the 7th postoperative day determined by the duration and the type of surgery as well as elevated preoperative D-dimer levels. After having reached the peak, D-dimers decrease with a rate of 6% per day, irrespective of the surgical type. On the basis of these findings, the duration of D-dimer elevation after surgery might be estimated for each patient.


D-dimer testing in combination with clinical judgment has been shown to be cost effective in the diagnostic algorithm of VTE in the outpatient setting due to its high negative predictive value. The lack of specificity [7], however, prevents its use in the diagnosis of VTE particularly for patients after surgery. Despite prophylaxis with LMWH, compression stockings and physical therapy, VTE remains a challenging problem after surgery. Incidence of symptomatic VTE is reported to be 3% in cancer patients undergoing abdominal surgery [3] with most of the VTE occurring after patients' discharge [5,6]. Activation of the coagulation/ fibrinolysis system is well described in surgical patients [9–12]

Fig. 2



D-dimer values before (day 0) and after surgery stratified by surgical type.

Fig. 3



Correlation between predicted and measured peak D-dimer levels (multivariate regression model).

leading to an increased risk of VTE [19]. Alterations in the coagulation induced by the surgical trauma have been shown to persist for 4 weeks after hip replacement [20] and for more than 7 weeks in patients after coronary bypass surgery [21]. In abdominal surgery, D-dimer has been shown to rise after open [11] and laparoscopic cholecystectomy [9,11], after open and laparoscopic gastric bypass surgery [12] and after a selection of abdominal procedures [22] with a maximal observation time of 2 weeks [9,22]. However, no comprehensive data are available so far regarding the kinetics of D-dimers in postoperative patients without symptomatic VTE. We demonstrate that D-dimers peak at day 7 after surgery and, thereafter, are cleared irrespective of the type of surgery as well as the occurrence of complications not requiring intervention. The calculated clearance of the D-dimer (6% per day in the study cohort) could be validated by the second cohort of patients (validation cohort). D-dimers normalized within 25 and 38 days after type II and type III procedures, respectively. Only 3% of the patients showed a D-dimer level above the normal range 5 weeks after those procedures.

Our study reveals that D-dimers after minor surgery (type I procedures) do not increase. However, opening of the peritoneum leads to an increase of D-dimer levels depending on the duration of surgery, the invasiveness of the operation (type II vs. type III surgery) and elevated D-dimer levels. Interestingly, parameters such as age or malignancy did not influence the D-dimer levels after surgery – a finding, which seems to be in contrast to the literature [23,24]. This discrepancy might be explained by the fact that most of these studies focus on nonsurgical patients and by the lack of multivariate analyses to

determine independent predictive factors of D-dimer elevation. However, malignant disease and age were associated in our study with increased preoperative D-dimer levels.

Beside its role as part of the diagnostic algorithm, D-dimer recently emerged as possible predictive factor for VTE [25,26]. There is evidence that D-dimer levels after surgery might be used to estimate the patients' risk for the development of postoperative VTE [26]. Hence, determination of postoperative D-dimer levels might be used to identify the patients who may benefit from extended thromboprophylaxis. In general surgery, four clinical trials have investigated the use of extended prophylaxis beyond the period of hospitalization following general surgery demonstrating a significant reduction of VTE with prolonged thromboprophylaxis [27–30]. On the basis of these studies, posthospital discharge prophylaxis with LMWH is suggested for high-risk general surgery patients, especially for patients after major cancer surgery [31]. However, high-risk patients are poorly defined, and the duration of the hypercoagulable state after general surgery remains elusive. Our study evaluated the duration of the activated coagulation/fibrinolysis system after abdominal surgery by elevated D-dimer levels, which might help to define the reasonable duration of postoperative thromboprophylaxis. We demonstrate that D-dimer levels are elevated for up to 6 weeks after major abdominal surgery. This finding may well correspond to the data from orthopedic surgery showing that extending thromboprophylaxis with LMWH for 4–5 weeks significantly reduces the incidence of VTE in patients undergoing total knee or hip replacement [32,33]. However, data from orthopedic surgery may not be directly transferred to general surgery, and the correlation between postoperative D-dimer levels and the risk of VTE has not yet been established.

Some considerations have to be made while interpreting our results. First, we did not perform routine duplex sonography or venography to exclude asymptomatic VTE after surgery. However, the focus of the study was to analyze D-dimer kinetics in patients without symptomatic VTE in order to apply the conclusions of this study to the vast majority of surgical patients that do not suffer from symptomatic VTE. Of note, none of the patients suffered from symptomatic VTE during follow-up. Second, only one assay was used for the measurement of D-dimers. Substantial variation has been described between different analytical methods to determine D-dimer levels [34]. However, quantitative rapid ELISA (VIDAS), as used in this study, has been identified as one of the most sensitive assay available [34].

In conclusion, this is the first report demonstrating the kinetics of D-dimers after abdominal surgery in an average surgical population. It allows estimating at which

point in time D-dimers reach normal values in asymptomatic patients after surgery. This might have an important clinical implication: it indicates when D-dimers may be used again in the diagnostic algorithm for VTE exclusion after surgery in patients with low or moderate clinical probability. After type I operations, D-dimer testing can be utilized after surgery in cases of suspected VTE without a lag phase whereas it appears futile after intraabdominal procedures (type II surgery) for a time span of 4 weeks and for 5–6 weeks after major abdominal surgery (type III surgery).

## References

- Agnelli G, Bolis G, Capussotti L, Scarpa RM, Tonelli F, Bonizzoni E, *et al.* A clinical outcome-based prospective study on venous thromboembolism after cancer surgery: the @RISTOS project. *Ann Surg* 2006; **243**:89–95.
- Nguyen NT, Hinojosa MW, Fayad C, Varela E, Konyalian V, Stamos MJ, *et al.* Laparoscopic surgery is associated with a lower incidence of venous thromboembolism compared with open surgery. *Ann Surg* 2007; **246**:1021–1027.
- White RH, Zhou H, Romano PS. Incidence of symptomatic venous thromboembolism after different elective or urgent surgical procedures. *Thromb Haemost* 2003; **90**:446–455.
- Prandoni P, Piccioli A, Girolami A. Cancer and venous thromboembolism: an overview. *Haematologica* 1999; **84**:437–445.
- Huber O, Bounameaux H, Borst F, Rohner A. Postoperative pulmonary embolism after hospital discharge. An underestimated risk. *Arch Surg* 1992; **127**:310–313.
- White RH, Romano PS, Zhou H, Rodrigo J, Bargar W. Incidence and time course of thromboembolic outcomes following total hip or knee arthroplasty. *Arch Intern Med* 1998; **158**:1525–1531.
- Wells PS, Owen C, Doucette S, Fergusson D, Tran H. Does this patient have deep vein thrombosis? *JAMA* 2006; **295**:199–207.
- Wells PS, Anderson DR, Bormanis J, Guy F, Mitchell M, Gray L. Value of assessment of pretest probability of deep-vein thrombosis in clinical management. *Lancet* 1997; **350**:1795–1798.
- Lippi G, Veraldi GF, Fraccaroli M, Manzato F, Cordiano C, Guidi G. Variation of plasma D-dimer following surgery: implications for prediction of postoperative venous thromboembolism. *Clin Exp Med* 2001; **1**:161–164.
- Parolari A, Mussoni L, Frigerio M, Naliato M, Alamanni F, Polvani GL. Increased prothrombotic state lasting as long as one month after on-pump and off-pump coronary surgery. *J Thorac Cardiovasc Surg* 2005; **130**:303–308.
- Schietroma M, Carlei F, Mownah A, Franchi L, Mazzolta C, Sozio A, *et al.* Changes in the blood coagulation, fibrinolysis, and cytokine profile during laparoscopic and open cholecystectomy. *Surg Endosc* 2004; **18**:1090–1096.
- Nguyen NT, Owings JT, Gosselin R, Pevec WC, Lee SJ, Goldman C, *et al.* Systemic coagulation and fibrinolysis after laparoscopic and open gastric bypass. *Arch Surg* 2001; **136**:909–916.
- Pittet JL, de Moerloose P, Reber G, Durand C, Villard C, Piga N, *et al.* VIDAS D-dimer: fast quantitative ELISA for measuring D-dimer in plasma. *Clin Chem* 1996; **42**:410–415.
- Dindo D, Muller MK, Weber M, Clavien PA. Obesity in general elective surgery. *Lancet* 2003; **361**:2032–2035.
- Anesthesiologists ASo. New classification of physical status. *Anesthesiology* 1963; **24**:111.
- Kondrup J, Rasmussen HH, Hamberg O, Stanga Z. Nutritional risk screening (NRS 2002): a new method based on an analysis of controlled clinical trials. *Clin Nutr* 2003; **22**:321–336.
- Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index. *J Clin Epidemiol* 1994; **47**:1245–1251.
- Rieker RJ, Hammer E, Eisele R, Schmid E, Högel J. The impact of comorbidity on the overall survival and the cause of death in patients after colorectal cancer resection. *Langenbecks Arch Surg* 2002; **387**:72–76.
- Iversen LH, Thorlacius-Ussing O. Relationship of coagulation test abnormalities to tumour burden and postoperative DVT in resected colorectal cancer. *Thromb Haemost* 2002; **87**:402–408.
- Dahl OE, Aspelin T, Arnesen H, Seljeflot I, Kierulf P, Ruyter R, *et al.* Increased activation of coagulation and formation of late deep venous thrombosis following discontinuation of thromboprophylaxis after hip replacement surgery. *Thromb Res* 1995; **80**:299–306.
- Parolari A, Camera M, Alamanni F, Naliato M, Polvani GI, Agrifoglio M, *et al.* Systemic inflammation after on-pump and off-pump coronary bypass surgery: a one-month follow-up. *Ann Thorac Surg* 2007; **84**:823–828.
- Galster H, Kolb G, Kohsytorz A, Seidlmeyer C, Paal V. The pre-, peri-, and postsurgical activation of coagulation and the thromboembolic risk for different risk groups. *Thromb Res* 2000; **100**:381–388.
- Harper PL, Theakston E, Ahmed J, Ockelford P. D-dimer concentration increases with age reducing the clinical value of the D-dimer assay in the elderly. *Intern Med J* 2007; **37**:607–613.
- Dirix LY, Salgado R, Weytjens R, Colpaert C, Benoy I, Huget P. Plasma fibrin D-dimer levels correlate with tumour volume, progression rate and survival in patients with metastatic breast cancer. *Br J Cancer* 2002; **86**:389–395.
- Palareti G, Legnani C, Cosmi B, Guazzaloca G, Pancani C, Coccheri S. Risk of venous thromboembolism recurrence: high negative predictive value of D-dimer performed after oral anticoagulation is stopped. *Thromb Haemost* 2002; **87**:7–12.
- Peetz D, Hafner G, Hansen M, Mayer A, Ripplin G, Rommens PM, *et al.* Dose-adjusted thrombosis prophylaxis in trauma surgery according to levels of D-Dimer. *Thromb Res* 2000; **98**:473–483.
- Lausen I, Jensen R, Jorgensen LN, Rasmussen MS, Lyng KM, Andersen M, *et al.* Incidence and prevention of deep venous thrombosis occurring late after general surgery: randomised controlled study of prolonged thromboprophylaxis. *Eur J Surg* 1998; **164**:657–663.
- Bergqvist D, Agnelli G, Cohen AT, Eldor A, Nilsson PE, Le Moigne-Amrani A, *et al.* Duration of prophylaxis against venous thromboembolism with enoxaparin after surgery for cancer. *N Engl J Med* 2002; **346**:975–980.
- Rasmussen MS. Preventing thromboembolic complications in cancer patients after surgery: a role for prolonged thromboprophylaxis. *Cancer Treat Rev* 2002; **28**:141–144.
- Rasmussen MS, Jorgensen LN, Wille-Jorgensen P, Nielsen JD, Horn A, Mohn AC, *et al.* Prolonged prophylaxis with dalteparin to prevent late thromboembolic complications in patients undergoing major abdominal surgery: a multicenter randomized open-label study. *J Thromb Haemost* 2006; **4**:2384–2390.
- Geerts WH, Pineo GF, Heit JA, Bergqvist D, Lassen MR, Colwell CW, *et al.* Prevention of venous thromboembolism: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004; **126**:338S–400S.
- Hull RD, Pineo GF, Francis C, Bergqvist D, Fellenius C, Soderberg K, *et al.* Low-molecular-weight heparin prophylaxis using dalteparin extended out-of-hospital vs in-hospital warfarin/out-of-hospital placebo in hip arthroplasty patients: a double-blind, randomized comparison. North American Fragmin Trial Investigators. *Arch Intern Med* 2000; **160**:2208–2215.
- Bergqvist D, Benoni G, Bjorgell O, Fredin H, Hedlundh U, Nicolas S, *et al.* Low-molecular-weight heparin (enoxaparin) as prophylaxis against venous thromboembolism after total hip replacement. *N Engl J Med* 1996; **335**:696–700.
- Stein PD, Hull RD, Patel KC, Olson RE, Ghali WA, Brant R, *et al.* D-dimer for the exclusion of acute venous thrombosis and pulmonary embolism: a systematic review. *Ann Intern Med* 2004; **140**:589–602.