

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Male-pattern baldness susceptibility locus at 20p11.

Authors: Richards JB, Yuan X, Geller F, Waterworth D, Bataille V, Glass D, Song K, Waeber G, Vollenweider P, Aben KK, Kiemenev LA, Walters B, Soranzo N, Thorsteinsdottir U, Kong A, Rafnar T, Deloukas P, Sulem P, Stefansson H, Stefansson K, Spector TD, Mooser V

Journal: Nature genetics

Year: 2008 Nov

Volume: 40

Issue: 11

Pages: 1282-4

DOI: 10.1038/ng.255

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

Published in final edited form as:

Nat Genet. 2008 November ; 40(11): 1282–1284. doi:10.1038/ng.255.

Male-pattern baldness susceptibility locus at 20p11

J Brent Richards^{1,2}, Xin Yuan³, Frank Geller⁴, Dawn Waterworth³, Veronique Bataille¹, Daniel Glass¹, Kijoung Song³, Gerard Waeber⁵, Peter Vollenweider⁵, Katja K H Aben^{6,7}, Lambertus A Kiemeny^{8,9}, Bragi Walters⁴, Nicole Soranzo^{1,10}, Unnur Thorsteinsdottir⁴, Augustine Kong⁴, Thorunn Rafnar⁴, Panos Deloukas¹⁰, Patrick Sulem⁴, Hreinn Stefansson⁴, Kari Stefansson⁴, Tim D Spector^{1,11}, and Vincent Mooser^{3,11}

¹Department of Twin Research and Genetic Epidemiology, King's College London, London SE1 7EH, UK ²Department of Medicine, Jewish General Hospital, Faculty of Medicine, McGill University, Montréal, Québec H3T 1E2, Canada ³Genetics Division, GlaxoSmithKline, King of Prussia, Pennsylvania 19406, USA ⁴deCODE Genetics, Reykjavik 101, Iceland ⁵Department of Internal Medicine, CHUV University Hospital, Lausanne 1011, Switzerland ⁶Comprehensive Cancer Center IKO, Radboud University Nijmegen Medical Center, Nijmegen 6501 BG, The Netherlands ⁷Department of Epidemiology and Biostatistics, Radboud University Nijmegen Medical Center, Nijmegen 6500 HB, The Netherlands ⁸Department of Epidemiology and Biostatistics, Radboud University Nijmegen Medical Center, Nijmegen 6500 HB, The Netherlands ⁹Department of Urology, Radboud University Nijmegen Medical Center, Nijmegen 6525 GA, The Netherlands ¹⁰Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton CB10 1SA, UK

Abstract

We conducted a genome-wide association study for androgenic alopecia in 1,125 men and identified a newly associated locus at chromosome 20p11.22, confirmed in three independent cohorts ($n = 1,650$; OR = 1.60, $P = 1.1 \times 10^{-14}$ for rs1160312). The one man in seven who harbors risk alleles at both 20p11.22 and *AR* (encoding the androgen receptor) has a sevenfold-increased odds of androgenic alopecia (OR = 7.12, $P = 3.7 \times 10^{-15}$).

Androgenic alopecia is a common disorder affecting 40% of adult men and women¹. Men and women with hair loss experience negative body-image perceptions². Moreover, the mechanisms involved in androgenic alopecia may influence common medical disorders, such as coronary heart disease and metabolic syndrome³. Underscoring the social

© 2008 Nature Publishing Group

Correspondence should be addressed to T.D.S. (tim.spector@kcl.ac.uk).

¹¹These authors contributed equally to this work.

AUTHOR CONTRIBUTIONS

V.M., P.V., G.W. and D.W. designed and implemented the CoLaus Study and the primary nested male-pattern baldness study. J.B.R., F.G., L.A.K., H.S., K. Stefansson, T.D.S. and V.M. designed the final study with appropriate replication datasets. J.B.R., X.Y., F.G., D.W., K. Song and V.M. analyzed the data. J.B.R., V.B., D.G., G.W., P.V., D.W., K.K.H.A., L.A.K., B.W., U.T., A.K., T.R., P.S., T.D.S. and V.M. contributed to data collection and phenotype definitions. V.B., G.W., T.D.S. and V.M. obtained funding. N.S. and P.D. performed genotyping for the women from the TwinsUK cohort. J.B.R. wrote the first draft of the paper and all authors made important contributions to the final version of the paper.

COMPETING INTERESTS STATEMENT

The authors declare competing financial interests: details accompany the full-text HTML version of the paper at <http://www.nature.com/naturegenetics/>.

Published online at <http://www.nature.com/naturegenetics/>

Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions/>

Note: Supplementary information is available on the Nature Genetics website.

implications of baldness, global annual sales of a medical therapy for male-pattern baldness recently surpassed \$405 million (Merck Financial Disclosures, Fourth Quarter 2007).

Androgenic alopecia is a highly heritable condition, with heritability estimates of over 80%⁴. Genetic variants in, or in proximity to, the *AR* (androgen receptor) gene have been previously associated with male-pattern baldness⁵⁻⁹. However, as the inheritance pattern of this trait seems to be polygenic, we undertook a two-stage genome-wide association (GWA) study for androgenic alopecia.

We carried out an extreme discordant case-control GWA scan for androgenic alopecia nested within the CoLaus population-based study in Lausanne, Switzerland¹⁰. A total of 578 male individuals with early-onset alopecia were compared with 547 male control individuals without alopecia (**Supplementary Methods** online). In the second stage, we sought replication in independent samples from the UK, Iceland and The Netherlands. The UK cohort comprised 453 men (176 affected individuals) from a population-based sample of British twins; 1,308 women (95 affected individuals) from this sample were also assessed for hair loss. The Dutch cohort comprised 463 men (147 affected individuals) previously diagnosed with prostate cancer and also assessed for androgenic alopecia. The Icelandic study included 734 Icelandic men (536 affected individuals) and 878 Icelandic women (397 affected individuals). The overall sample size thus comprised 4,961 individuals.

The GWA study (Supplementary Fig. 1 online) was done using the Affymetrix GeneChip Human Mapping 500K array in the CoLaus cohort. All SNPs from this GWA study with a P value $< 10^{-5}$ were from two loci (Supplementary Fig. 2 online). We confirmed the importance of the previously described *AR* gene⁵⁻⁹ (OR for lead SNP rs6625163[A] = 3.30 (2.31-4.71), $P = 5.0 \times 10^{-11}$). We also located a new susceptibility locus on chromosome 20p11.22 (OR = 1.79 (1.49-2.15), $P = 3.2 \times 10^{-10}$ for rs1160312[A] and OR = 1.80 (1.49-2.16), $P = 3.5 \times 10^{-10}$ for rs913063[T]; the r^2 between rs1160312 and rs913063 was 1.0; Table 1). These SNPs are located over 350 kb distant to the nearest annotated gene, *PAX1*.

This same risk allele at rs1160312, rs1160312[A], was associated with an increased risk of androgenic alopecia in the TwinsUK cohort (OR = 1.68 (1.22-2.33), $P = 2.0 \times 10^{-3}$), hair loss in men in the Icelandic cohort (OR = 1.41 (1.10-1.79), $P = 6.1 \times 10^{-3}$) and androgenic alopecia in the Dutch cohort (OR = 1.40 (1.06-1.85), $P = 0.018$). In a combined analysis of all male cases and controls, the risk allele was strongly associated with hair loss (OR = 1.60 (1.42-1.80), $P = 1.1 \times 10^{-14}$). In addition, the same risk allele was associated with hair loss in women (OR = 1.24 (1.05-1.47), $P = 0.012$) for combined data from Icelandic and TwinsUK women (OR = 1.29 (1.05-1.57), $P = 0.01$ for Icelandic women; OR = 1.13 (0.82-1.56), $P = 0.45$ for women from the TwinsUK cohort).

We found that 14% of men harbored at least one risk allele at both 20p11.22 and the *AR* gene and that this was associated with a markedly increased risk of androgenic alopecia (OR = 7.12 (4.34-11.68), $P = 3.7 \times 10^{-15}$) in the CoLaus cohort. We observed no significant statistical interaction between the two loci. The variance in androgenic alopecia explained by the presence of at least one risk allele at both loci was 13.7%. Using all men from the population-based CoLaus study who did not meet case or control definitions and who were not included in the GWA scan ($n = 940$), we found that the ability of these two risk alleles to exclude the development of androgenic alopecia was high, but lacked specificity (negative predictive value = 96.5%, positive predictive value = 12.2%, sensitivity = 98.2%, specificity = 6.6%).

The rs1160312 SNP lies between *PAX1* and *FOXA2* on chromosome 20. It is not immediately clear how this gene might affect androgenic alopecia. Although rs1160312 is over 350 kb away from both genes, it remains possible that this SNP (or another variant in linkage disequilibrium with it) may influence the expression of either transcript through long-range control, as has been demonstrated for other genes involved in development¹¹. How these variants affect disease outcomes deserves further investigation.

The fact that, despite a relatively small number of subjects, rs1160312 had a genome-wide significant association with androgenic alopecia in the discovery cohort alone and several other SNPs within the LD block bearing rs1160312 were associated with androgenic alopecia (Fig. 1) reflects the strength of the extreme discordant age-based case-control definitions. The phenotype from the Icelandic and the female TwinsUK cohorts was derived from self-reported hair loss, without the use of matching diagrams. Although this phenotype definition is less precise than that derived using diagram-matching techniques, which have been previously validated¹², the association for males was replicated in Iceland and the results for females are consistent between the cohorts. However, it is necessary to investigate the association for female hair loss in larger and carefully phenotyped cohorts.

Even though the definition of hair loss and the method of sampling differed between the four study groups, the fact that the association was observed in all groups underscores the generalizability of our findings to other European populations. We also note that, as in any GWA study, the identified variant is likely to be in linkage disequilibrium with the causal SNP.

These results may have clinical significance. Androgenic alopecia is a disease of considerable social concern and the scalp is one of the few areas that are directly accessible by liposome–DNA mixtures¹³; this method has been demonstrated to result in hair-follicle transfection¹⁴. Therefore, if a variant at 20p11.22 influences expression of a transcript, such as *PAX1* or *FOXA2*, then manipulation of this pathway may lead to a target for gene therapy. However, demonstration of an influence of the chromosome 20p11.22 locus on transcription of either gene requires further investigation. The relatively high prevalence and magnitude of risk attributed to the risk alleles at both *AR* and 20p11.22 loci suggests that these markers may assist in the identification of groups of men at high risk for androgenic alopecia, and the absence of these risk alleles can be helpful in excluding its development.

In summary, we report a replicated GWA study for androgenic alopecia providing evidence from four distinct European populations for a newly associated locus that influences a common disorder of social importance. The risk alleles at this newly identified locus and *AR* are common in Europeans and impart a relatively large risk for androgenic alopecia. Given the feasibility of gene therapy in human follicles, our results may point to an intriguing new potential target for the treatment of hair loss.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We wish to acknowledge the cooperation and dedication of the study participants and the personnel at the recruitment centers and core facilities. We would like to acknowledge the genotyping contributions of D. Smyth and J. Todd. This study was funded in part by GlaxoSmithKline; deCODE Genetics; the Wellcome Trust; NIHR Biomedical Research Centre (grant to Guys' and St. Thomas' Hospitals and King's College London); the Chronic Disease Research Foundation; the Canadian Institutes of Health Research (J.B.R.) and the Sixth Framework Program of the EU contract number 018827 (Polygene).

References

1. Olsen, E. Disorders of hair growth: Diagnosis and Treatment. McGraw-Hill; New York: 1994. p. 257
2. Cash TF, Price VH, Savin RC. J. Am. Acad. Dermatol. 1993; 29:568–575. [PubMed: 8408792]
3. Lo JC, et al. J. Clin. Endocrinol. Metab. 2006; 91:1357–1363. [PubMed: 16434451]
4. Nyholt DR, Gillespie NA, Heath AC, Martin NG. J. Invest. Dermatol. 2003; 121:1561–1564. [PubMed: 14675213]
5. Ellis JA, Stebbing M, Harrap SB. J. Invest. Dermatol. 2001; 116:452–455. [PubMed: 11231320]
6. Hayes VM, et al. Cancer Epidemiol. Biomarkers Prev. 2005; 14:993–996. [PubMed: 15824176]
7. Hillmer AM, et al. Am. J. Hum. Genet. 2005; 77:140–148. [PubMed: 15902657]
8. Levy-Nissenbaum E, Bar-Natan M, Frydman M, Pras E. Eur. J. Dermatol. 2005; 15:339–340. [PubMed: 16172040]
9. Prodi DA, et al. J. Invest. Dermatol. 2008; 128:2268–2270. [PubMed: 18385763]
10. Sandhu MS, et al. Lancet. 2008; 371:483–491. [PubMed: 18262040]
11. Kleinjan DA, van Heyningen V. Am. J. Hum. Genet. 2005; 76:8–32. [PubMed: 15549674]
12. Littman AJ, White E. Ann. Epidemiol. 2005; 15:771–772. [PubMed: 15949955]
13. Li L, Hoffman RM. Nat. Med. 1995; 1:705–706. [PubMed: 7585157]
14. Domashenko A, Gupta S, Cotsarelis G. Nat. Biotechnol. 2000; 18:420–423. [PubMed: 10748523]




Figure 1.

Association between SNPs near chromosome 20p11.22 and androgenic alopecia. **(a)** $-\log(P \text{ value})$ measures for association between SNPs and chromosomal position. **(b)** Recombination hot spots (HapMap CEPH population, NCBI build 36). **(c)** Linkage disequilibrium in GOLD heat map Haploview 4.0 color scheme, CEPH population. The x axis represents genomic position in Mb **(a)** and in kb **(b,c)**. All P values are derived from the CoLaus cohort, except that for the lead SNP, rs11603132, which is derived from the combined P value from the CoLaus, TwinsUK, Icelandic and Dutch cohorts.

Table 1
Summary results for the lead SNP from the 20p11.22 locus for androgenic alopecia in men

Marker	Chr.	Cohort	Sample size		Risk-allele frequency		OR per risk allele (95% CI)	P	P _{combined}
			Controls	Cases	Controls	Cases			
rs1160312	20	CoLaus	547	578	0.43	0.57	1.79 (1.49-2.15)	3.2 × 10 ⁻¹⁰	
A/G		TwinsUK	277	176	0.45	0.58	1.68 (1.22-2.33)	2.0 × 10 ⁻³	
		Icelandic	198	536	0.56	0.64	1.41 (1.10-1.79)	6.1 × 10 ⁻³	
		Dutch	316	147	0.52	0.60	1.40 (1.06-1.85)	0.018	
		All men	1,338	1,437			1.60 (1.42-1.80)	1.1 × 10 ⁻¹⁴	