

Doublesex Evolution Is Correlated with Social Complexity in Ants

Ling-Yi Jia¹, Li Chen², Laurent Keller³, John Wang⁴, Jin-Hua Xiao^{5,*}, and Da-Wei Huang^{1,5,*}

¹Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Beijing, China

²State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing, China

³Department of Ecology and Evolution, Biophore, University of Lausanne, Switzerland

⁴Biodiversity Research Center, Academia Sinica, Taipei, Taiwan

⁵College of Life Sciences, Nankai University, Tianjin, China

*Corresponding authors: E-mails: xiaojinhua2015@163.com; huangdw@ioz.ac.cn.

Accepted: November 9, 2018

Data deposition: Publicly available RNA-sequencing (RNA-seq) data sets used in this study are listed in supplementary table S1, Supplementary Material online. These sequence data are from NCBI's Sequence Read Archive (SRA) (<https://www.ncbi.nlm.nih.gov/sra>). New *Solenopsis invicta* RNA-seq data have been deposited at the SRA database under the accession SRP143788. Sequence alignments used are available by the corresponding authors upon request.

Abstract

The *Dmrt* (*doublesex* and *mab-3-related transcription factor*) genes are transcription factors crucial for sex determination and sexual differentiation. In some social insects, *doublesex* (*dsx*) exhibits widespread caste-specific expression across different tissues and developmental stages and has been suggested as a candidate gene for regulating division of labor in social insects. We therefore conducted a molecular evolution analysis of the *Dmrt* gene family in 20 ants. We found that the insect-specific oligomerization domain of DSX, oligomerization domain 2, was absent in all ants, except for the two phylogenetically basal ant species (Ponerinae), whose social structure and organization resemble the presumed ancestral condition in ants. Phylogenetic reconstruction and selection analysis revealed that *dsx* evolved faster than the other three members of the *Dmrt* family. We found evidence for positive selection for *dsx* in the ant subfamilies with more advanced social organization (Myrmicinae and Formicinae), but not in the Ponerinae. Furthermore, we detected expression of two *Dmrt* genes, *dsx* and *DMRT11E*, in adult ants, and found a clear male-biased expression pattern of *dsx* in most species for which data are available. Interestingly, we did not detect male-biased expression of *dsx* in the two ant species that possess a genetic caste determination system. These results possibly suggest an association between the evolution of *dsx* and social organization as well as reproductive division of labor in ants.

Key words: sex determination, caste determination, eusocial insect, transcription factor, transcriptome, alternative splicing.

Introduction

Ants (Hymenoptera, Formicidae), which are one of the most successful eusocial taxa (Gadau et al. 2012), display large variation in the complexity of social organization, ranging from some "primitive" ponerine species, whose workers have full reproductive potential, to species where workers are totally sterile and exhibit morphologically distinct subcastes (Hölldobler and Wilson 1990, 2009). In the majority of ants, only queens can mate and reproduce, whereas workers are incapable of mating, because they lack functional spermathecae (Bourke 1988).

The genetic pathways regulating the determination of the queen and worker castes include the insulin/insulin-like growth signaling (IIS), target of rapamycin, epidermal growth factor receptor, juvenile hormone (JH), and vitellogenin. External environmental stimuli, such as nutrition and temperature, are important inputs into these integrative pathways. However, the downstream effector pathways connecting these regulators to growth and differentiation remain largely unknown (reviewed in Corona et al. [2016]).

In solitary insects, the sex-differentiation gene *doublesex* (*dsx*) links development to nutrition and sex. For example,

© The Author(s) 2018. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

dsx regulates sex-specific mandible growth via JH signaling in stag beetles (Gotoh et al. 2014). *Drosophila melanogaster dsx* was the first *Dmrt* gene (*doublesex* and *mab-3-related transcription factor*) identified. *dsx* encodes two sex-specific splice variants that regulate transcription of target genes to trigger sexual differentiation (Nagoshi et al. 1988). Both the male- and female-specific protein products, DSX^M and DSX^F, share an N-terminal DM domain, which consists of a zinc finger-related DNA-binding domain and an oligomerization domain (OD1) (Erdman and Burtis 1993; An et al. 1996). *Dmrt* family members are defined by the presence of the DM domain and have been found in almost all eumetazoans (Wexler et al. 2014). There are eight *Dmrt* genes in *Homo sapiens* and 11 in *Caenorhabditis elegans* (Ottolenghi et al. 2002; Volff et al. 2003). In insects, four *Dmrt* genes are known (*DMRT11E*, *DMRT93B*, *DMRT99B*, and *dsx*) (Volff et al. 2003).

The C-termini of DSX^M and DSX^F contain another oligomerization domain, OD2, which consists of both a shared oligomerization sequence and sex-specific sequences. DSX forms dimers to bind DNA and regulate the transcription of target genes. The dimerization of the monomer DSXs is caused by OD1 and OD2 independently (An et al. 1996; Erdman et al. 1996). DSX^F and DSX^M bind to the same dimorphic *cis*-regulatory-element, but the sex-specific OD2 domains may recruit different sets of regulatory proteins, thereby regulating sex-specific expression of target genes (reviewed in Verhulst and van de Zande [2015]). Consequently, OD2 is important to the double-switch function of DSX in regulating sexual differentiation.

As a pleiotropic transcription factor, *dsx* controls a wide range of downstream patterning genes responsible for the development of sex-related traits. Among them, the most ancestral function of *dsx* was likely the determination of gonadal sex (Matson and Zarkower 2012), including regulating the differentiation of genital structures, such as the spermathecae (Sanchez et al. 2001). *dsx* also regulates biosynthesis of pheromone (Shirangi et al. 2009) and sexual behaviors (Rideout et al. 2010). Given the important roles *dsx* plays in multiple aspects of reproduction, *dsx* may be involved in the polyphenism of eusocial insects between and within sexes. In line with this view, several studies have revealed caste-specific expression of *dsx* across different tissues in the honeybee *Apis mellifera* or development stages in the ant *Cardiocondyla obscurior* and hypothesized that *dsx* may regulate polyphenism in social insects (Johnson and Jasper 2016; Klein et al. 2016).

Compared with other eusocial insects, the gradient of social complexity (Hölldobler and Wilson 1990, 2009) along with their rich taxonomic diversity makes ants an ideal model system for investigating the evolution of caste polyphenism. In this study, we annotated *Dmrt* genes in 20 ant species and predicted their conserved domains. Overall, the OD2 domain is present in some phylogenetically basal ants that show weak reproductive division of labor but is lost in several more advanced ant lineages. We also found that the evolutionary rate

of *dsx* is relatively high among the four *Dmrt* genes, especially in ants showing more advanced social organizations. These results suggest that the coding sequences of *dsx* genes are diverging congruently with variation in social organization. We then used transcriptome data to analyze sex- or caste-biased gene expression and alternative splicing of *Dmrt* genes, and found that transcription of *dsx* was higher in males for most examined Hymenopteran species, except for two Myrmicinae ants, which are known for their peculiar caste-determination mechanisms (Fournier et al. 2005; Ohkawara et al. 2006). Furthermore, the exon-usage of *dsx* in these two Myrmicinae species shows significant differences not only between sexes but also among castes. Taken together, the evolutionary and expression analyses provide multiple lines of evidence that the evolutionary characteristics of *dsx* map well to the evolution of the caste determination in ants.

Materials and Methods

Identification of *Dmrt* Genes and Phylogenetic Reconstruction

The annotated genes and genome assemblies of 48 insect species, including 39 Hymenopteran species (20 ants and 19 nonant Hymenoptera) who were the only Hymenopteran species with available genomes at the start of this study, and nine representatives for non-Hymenopteran species whose genomes were also available. Besides, six Crustaceans whose *Dmrt* genes had been reported (Kato et al. 2008, 2011; Toyota et al. 2013; Zhang et al. 2014) were also included in our research as outgroups to the insects. Our analysis was focused on 20 ant species. Four *D. melanogaster Dmrt* genes (*Dmel_DMRT11E*, *DMRT93B*, *DMRT99B*, and *dsx*) were used as queries in the BLASTp search (Altschul 1997), with an *E*-value cutoff of 10^{-5} . We reiterated a tBLASTn search in the genome assemblies using each candidate *Dmrt* sequences as queries, with an *E*-value cutoff of 10^{-4} . Then, a hidden Markov model (HMM)-based tool FGENE SH+ (Buschinger 1968) in the SOFTBERRY package (<http://linux1.softberry.com>; Last Accessed December 7, 2018) was used to improve the exon–intron boundaries. Gene fragments without a full DM domain or with an open reading frame (ORF) length <500 bp were categorized as pseudogenes.

Eighty-two amino acid sequences of *Dmrt* genes from 20 ant species and two outgroup Vespidae species were aligned using MAFFT version 7 (Kato and Standley 2013). The resulting alignment was analyzed using PROTTEST3 to predict the bestfit substitution model of molecular evolution (Darriba et al. 2011). The phylogenetic tree was constructed using the PHYML3.0 program (Crisuolo 2011) with the Jones–Taylor–Thornton model, gamma distribution of rates, optimized equilibrium frequencies and invariant site categories, with 100 bootstrap replications.




FIG. 1.—Domain organization of ant DMRTs. Domains of DMRT proteins for 20 ant species, two outgroup Vespidae species, and *D. melanogaster*. Full-length proteins for each of the four *DMRT* genes are shown as hollow boxes in four distinct colors. DM, DMA, and OD2 domains are illustrated as solid boxes in black, gray, and red, respectively. OD2-like domains found in *H. saltator* and *Di. quadriceps* are shown as hashed red boxes. The branches belonging to the subfamilies of Myrmicinae, Formicinae, and Ponerinae are colored in blue, green, and yellow, respectively. The phylogenetic relationship is based on Moreau and Bell (2013) and Ward et al. (2015). The plus signs indicate the six branches undergone positive selection.

Selection Analysis

PAML analyses were performed based on the species tree shown in figure 1. The ratios of nonsynonymous (d_N) and synonymous (d_S) substitution rates ($\omega = d_N/d_S$) were evaluated using CODEML in PAML version 4.5 (Yang 2007). For each *Dmrt* subfamily, tests using different site models were conducted: the basic model (M0: one ratio) assuming ω is invariable among sites and branches (Goldman and Yang 1994), the site-specific models assuming ω varies among sites but not branches, including M1a (neutral), M2a (selection), M3 (discrete), M7 (beta), M8 (beta & $\omega > 1$), and M8a (beta & $\omega = 1$) (Yang et al. 2000). Pairwise comparisons of M0 versus M3, M1a versus M2a, M7 versus M8, and M8a versus M8 were used to perform likelihood ratio tests (LRTs). For *dsx* genes, subsequently analysis using the branch-site model, allowing ω to vary both among sites and branches (model = 2, NSSites = 2) (Zhang et al. 2005), was tested (df = 1) against a null model (fixed ω of 1). LRTs using the pairwise comparisons of branch-site model versus branch-site null model were performed. Significance tests were performed using the χ^2 distribution. After, if the LRT was significant, a Bayes empirical Bayes (BEB) analysis was conducted to identify positively selected sites ($P > 0.95$).

Gene Expression Analysis

Solenopsis invicta colonies were collected in Guangdong Province, South China, on October 16, 2017. We selected a colony whose monogyne status was verified by using a polymerase chain reaction protocol previously described (Krieger and Ross 2002). After being reared in the lab for 2 weeks, three biological replicated samples were chosen from the same three colonies. Total RNA was isolated from whole-body adults of three castes, (workers, egg-laying queens,

and presumptive haploid males) separately, with three biological replicates for each caste. For the RNA extraction, the TransZol Up Plus RNA Kit (Transgen Biotech, Beijing, China) was used in accordance with the manufacturer's instructions. These mRNA samples were then submitted to the Berry Genomics Company (Beijing, China) for mRNA-seq library construction and sequencing. The sequencing was conducted using an Illumina HiSeq 4000 (Illumina, San Diego, CA) platform, with a paired-end module (PE150 mode). The transcriptome data of *S. invicta* have been deposited to the NCBI's Sequence Read Archive (SRA) database (<http://www.ncbi.nlm.nih.gov/Traces/sra>; Last Accessed December 7, 2018) and given the accession number SRP143788. Fastq files for other species (supplementary table S1, Supplementary Material online) were downloaded from the SRA database. All the RNA-sequencing (RNA-seq) data were then mapped against their assembled genome sequences using Tophat (Trapnell et al. 2009) with default parameters. To estimate the transcript level abundance in each sample, we calculated fragments per kilobase of transcript per million fragments mapped (FPKM) using CUFFLINKS (Trapnell et al. 2012). Genes with FPKM > 10 were used for subsequent analyses. For the *dsx* gene which expressed multiple alternatively spliced transcripts, FPKMs for each of the transcripts were calculated separately by CUFFLINKS and were then summed up to evaluate the expression level of the gene. Only gene isoforms with a DM domain were analyzed. In order to control for the differences caused by the different sequencing protocols, the expression levels of the housekeeping genes *ribosomal protein L18*, *elongation factor 1*, and *NADH dehydrogenase* were also determined for each caste, using the same method.

Evidence of differential exon-usage in the *dsx* and *DMRT11E* genes was tested using the DEXSeq package's (version 1.22.0) "exon-centric" differential splicing analysis

(Li et al. 2015) in R (R Core Team 2000). Prior to analysis, transcript models produced by CUFFLINKS were used as input for the script to prepare the annotation used in DEXSeq. An LRT was performed, comparing the full model (\sim sample + exon + caste:exon) and reduced model (\sim sample + exon), to detect differences in exon-usage.

Results

Gene Annotation

We searched the genome assemblies and the predicted gene sets from 20 ant species and 19 other Hymenopteran species (supplementary table S2, Supplementary Material online) for *Dmrt* homologs. We combined these Hymenopteran *Dmrt* genes with 45 additional *Dmrt*s from nine representatives of other insect orders and six Crustaceans as insect outgroup and then built a phylogeny. Examination of the *Dmrt* tree topology suggested a diversification of the four *Dmrt* subfamilies prior to the diversification of Crustacea and Hexapoda (supplementary fig. S1, Supplementary Material online). We used a nomenclature for insect *Dmrt* genes according to the *Dmrt* genes in *D. melanogaster* (*DMRT11E*, *DMRT93B*, *DMRT99B*, and *dsx*) (Voff et al. 2003).

In Hymenoptera, lineage-specific gene duplication and loss are rare (supplementary table S2, Supplementary Material online). We annotated partial gene fragments on separated scaffolds as pseudogenes, but these are likely genome assembly artifacts. Aside from these artificial pseudogenes, our analysis suggests that all the clear cases of expansion and contraction of *Dmrt* genes took place in the lineages giving rise to *Atta cephalotes*, *Pseudomyrmex gracilis*, and *Wasmannia auropunctata* (bold in supplementary table S2, Supplementary Material online).

Loss of OD2 Domain in Ants

We used the NCBI's Conserved Domain Detector (CDD) (Marchler-Bauer et al. 2015) to identify conserved domains of the *Dmrt* genes. All the examined genes had a DM domain that is characteristic for the *Dmrt* gene family. All predicted orthologs of *DMRT11E*, *DMRT93B*, and *DMRT99B* shared the same domain organization as *D. melanogaster*, including the presence of a DMA domain in *DMRT93B* and *DMRT99B* (Ottolenghi et al. 2002; Voff et al. 2003).

Interestingly, the conserved domain composition of DSX proteins was more variable (fig. 1 and supplementary fig. S1, Supplementary Material online). We did not detect the OD2 domain in any ant DSX, even when we relaxed the *E*-value threshold to 0.01. To search for remote homologs, we further queried the 20 ant genomic data sets via BLASTn/tBLASTn (*E*-value = 1.0) with the OD2 sequence of *A. mellifera*. There was evidence for the presence of a DSX OD2 domain only in *Harpegnathos saltator* and *Dinoponera quadricaps*, both of which are ponerine ants. The alignment

of Hsal_DSX and Dqua_DSX with other Hymenopteran DSXs indicated an exclusive 5-amino acid insertion in the middle of the OD2 domain in these two ponerine ant sequences (supplementary fig. S2, Supplementary Material online). When aligned pairwise with other Hymenopteran DSXs, the average sequence identity was lower for Hsal_DSX and Dqua_DSX (0.23) than other Hymenopteran DSXs (0.42). The high level of sequence divergence was likely the reason why we could not detect the OD2 domain using the CDD database.

To ascertain that the loss of OD2 domain was not an assembly artifact, we queried the short-read sequencing data of the 20 ants with the seed alignments for the OD2 (or OD2-like) domain sequences of *D. melanogaster*, *H. saltator*, and *Di. quadricaps* via SRA-BLAST (Camacho et al. 2009). In addition, an HMM profile (HMMER version 3.1) (Eddy 1998) was created with the same alignment of OD2 domain sequences as used in the SRA-BLAST search, and a profile nHMMER search was carried out against the genome databases of the 20 ants, with a threshold *E*-value of 0.01. Both of these searches failed to detect OD2 domain homologs in the other 18 ant species.

We further examined the Hymenopteran *dsx* genes with and without the OD2 domain for any additional structural differences. We found that the 5'-terminal sequences of *dsx* (DM domain coding sequences) are conserved among all the species. In contrast, the 3'-terminal sequences of *dsx* are quite variable, both for ants (without OD2) and for nonant species (with OD2). We have aligned the OD2 domain of DSX in different Hymenopteran species (with OD2 for nonant species, and OD2-like domain for the two ponerine ants), as shown in figure S2, Supplementary Material online. But it seems impossible to align the 3'-terminal sequences of *dsx* for "higher" ants (without OD2), because the sequence similarity is too low.

Gene Phylogeny

Our phylogenetic analysis (fig. 2) revealed that the Vespidae *DMRT11E*, *DMRT93B*, and *DMRT99B* genes form outgroups to the Formicidae homologs, which is consistent with the taxonomic relationship of these species. However, the pattern was strikingly different for *dsx*, with the two Vespidae *dsx* not forming an ancestral lineage to the Formicidae lineages. After conducting a phylogenetic analysis with only DSX sequences, the two Vespidae *dsx* sequences rooted basal to the Formicidae *dsx* sequences (supplementary fig. S3, Supplementary Material online), suggesting that the ingroup placement of Vespidae *dsx* in figure 2 was caused by long-branch attraction (Felsenstein 1978). The branch lengths separating taxa in the *dsx* clade were long, indicating high evolutionary divergence of *dsx* across the different species included in the analysis. A phylogenetic tree built with only the DM domain showed a similar topological structure (supplementary fig. S4, Supplementary Material online), indicating




FIG. 2.—Maximum-likelihood phylogenetic analysis of the deduced amino acid sequences of *Dmrt* homologs in ants and two vespids using PHYML. Bootstrap values are indicated for each node (of 100 resamplings).

that the evolutionary rate of the DM domain, which is thought to be conserved among *Dmrt* homologs, is extremely high in *dsx*. Taken together, these results indicate that the *dsx* genes have the fastest rate of evolution among the four *Dmrt* genes.

dsx Genes of Ants Have Evolved under Positive Selection

The finding that the tree topology of the *Dmrt* genes (fig. 2) does not mirror that of the species tree (fig. 1) hinted that the *Dmrt* genes might undergo lineage-specific selection in ants. We further estimated the ratios of nonsynonymous (d_N) and synonymous (d_S) substitution rates for the four *Dmrt* genes using the PAML package (Yang 2007). Because of the

extreme divergence of the C-terminus of DSX, it was difficult to determine homologous codon positions outside the DM domain. So only the coding sequence for the N-terminus of the DSX protein, which includes the DM domain, was used in the PAML analysis. We analyzed the four *Dmrt* genes separately under multiple site-specific models, which allow $\omega = d_N/d_S$ to vary among sites (table 1 and supplementary tables S3–S5, Supplementary Material online). Under the basic model (M0), which assumes ω to be invariable among sites and branches (Goldman and Yang 1994), *dsx* showed a higher ω (0.62) than the other three *Dmrt* genes (0.05–0.16). To test whether ω values among sites were variable, we compared M0 with the discrete model (M3) using LRTs. Another two LRTs, which compared M1a (neutral) with M2a

Table 1

Results of Site-Specific Positive Selection Analysis of the *dsx* Gene

| Models | P | In L | Parameters | PS Sites | Significant after LRT Correction |
|--------------------------|----|---------------|---|----------------------------------|----------------------------------|
| M0: One ratio | 40 | -4,240.610423 | $\omega = 0.62265$ | None | — |
| M1a: neutral | 41 | -4,131.232251 | $p0 = 0.33583, p1 = 0.66417$ $\omega0 = 0.16958, \omega1 = 1.00000$ | Not allowed | — |
| M2a: selection | 43 | -4,107.690927 | $p0 = 0.19604, p1 = 0.53092$ ($p2 = 0.27304$), $\omega0 = 0.10031$ $\omega1 = 1.00000, \omega2 = 3.19922$ | BEB $6 > 0.95$ $6 > 0.99$ | M1a–M2a: yes |
| M3: discrete | 44 | -4,097.544406 | $p0 = 0.04743$ ($p1 = 0.52106$) $p2 = 0.43152, \omega0 = 0.00510$ $\omega1 = 0.51352, \omega2 = 1.46193$ | NEB $5 > 0.95$ $20 > 0.99$ | M0–M3: yes |
| M7: beta | 41 | -4,110.962547 | $p = 0.46067, q = 0.25504$ | Not allowed | — |
| M8: beta & $\omega > 1$ | 43 | -4,094.79483 | $p0 = 0.74246, p = 0.49554$ $q = 0.36279$ ($p1 = 0.25754$), $\omega = 2.53805$ | BEB $5 > 0.95$ $5 > 0.99$ | M7–M8: yes |
| M8a: beta & $\omega = 1$ | 42 | -4,110.28144 | $p0 = 0.57543, p = 0.63016$ $q = 1.07704$ ($p1 = 0.42457$), $\omega = 1.00000$ | Not allowed | M8a–M8: yes |

NOTE.—The table gives the number of sites predicted to be under positive selection for each site-specific model. Abbreviations are as follows. Models, site-specific models; P, number of parameters; In L, log likelihood scores; p, proportion of sites under a particular omega value; ω , ratio of nonsynonymous to synonymous substitution rates; PS sites, number of positively selected sites; NEB, naive empirical Bayes analysis.

(selection), and M7 (beta) with M8 (beta & ω), were also conducted. For all four *Dmrt* genes, M3 was a significantly better fit than M0, indicating that ω is different among sites. Using a BEB approach to identify positively selected sites (in the case of models M2a and M8), we found no positively selected sites for *DMRT11E*, *DMRT93B*, and *DMRT99B* (supplementary tables S3–S5, Supplementary Material online). By contrast, multiple amino acid sites of *dsx* (12 sites in M2a and 10 sites in M8) apparently underwent adaptive evolution, and LRTs performed were significant for both of these models (table 1).

The alignment of the DSX N-terminal coding sequences was further analyzed using the branch-site model A in PAML, which allows ω to vary both among branches and sites (table 2). *dsx* genes from each of the ant species were tested as foreground branches. Positively selected sites could be detected in six species, four of which belong to the Myrmicinae subfamily and the other two to the Formicinae subfamily (fig. 1 and table 2). LRT analysis indicated that the unconstrained model (model H_A) fitted the data sets significantly better than the model fixing ω for the foreground branches at 1 (model H₀) in all six cases. Interestingly, no positively selected sites could be found in Ponerinae ants (table 2). This suggests that among the four *Dmrt* genes, positive selection occurred only in *dsx*, and that this occurred only in the two “higher” ant subfamilies, but not in the “primitive” Ponerinae.

Overall Male-Biased Expression of *dsx* in Ants

To test whether *Dmrt* genes show sex- or caste-specific gene expression in ants, we gathered transcriptomic data for five

ant species (*Camponotus floridanus*, *H. saltator*, *S. invicta*, *Vollenhovia emeryi*, and *W. auropunctata*). The expression levels of the housekeeping genes as *ribosomal protein L18*, *elongation factor 1*, and *NADH dehydrogenase*, which have proven to be valid controls in Hymenopteran species (Scharlaken et al. 2008; Cheng et al. 2013; Helmkampf et al. 2015; Livramento et al. 2018), were used to normalize gene expression across different samples. The expression of these reference genes was consistent within species (coefficients of variation ≤ 0.74), making them suitable reference genes to compare *Dmrt* gene expression levels within species (supplementary fig. S5, Supplementary Material online).

There were common features and also interesting species-specific differences in *Dmrt* expression profiles. *dsx* and *DMRT11E* showed the strongest expression in ants, with *DMRT11E* expression levels being generally lower than *dsx*. The expression levels of the other two *Dmrt* genes were too low to be detected in most cases (fig. 3). In contrast, *dsx* and *DMRT99B*, but not *DMRT11E*, were expressed at higher levels than the other two *Dmrt* genes in three other nonant Hymenopterans (supplementary fig. S6, Supplementary Material online). In summary, among the four *Dmrt* genes, *dsx* was widely expressed in all Hymenopteran species included in this study, whereas *DMRT11E* was expressed only in the ant lineage.

We then tested whether *Dmrt* expression was sex- or caste-biased in ants. We detected male-biased expression of *dsx* in three out of the five ant species (fig. 3). The expression of *dsx* was significantly higher in males than in queens and workers in the fire ant *S. invicta* (one-way ANOVA, *P*-value < 0.01, followed by the Tukey’s Honestly Significant Difference [HSD] post hoc test). *dsx* gene expression was similarly male-

Table 2Tests for Positive Selection on *dsx* Homologs in Ants Using the PAML Branch-Site Model A

| Branch | H ₀ : p2a (ω 2 = 1) | H _A : p2a, ω 2 | PS Sites (BEB) | Significant after LRT Correction |
|-------------------------------------|--|----------------------------------|----------------|----------------------------------|
| Myrmicinae | | | | |
| <i>Atta colombica</i> | 0.00 | 0.02, 1.00 | None | No |
| <i>Atta cephalotes</i> | 0.01 | 0.06, 2.61 | None | No |
| <i>Acromyrmex echinatior</i> | 0.01 | 0.10, 999.00 | 4 > 0.95 | Yes |
| <i>Trachymyrmex septentrionalis</i> | 0.07 | 0.07, 1.86 | None | No |
| <i>Trachymyrmex cornetzi</i> | 0.05 | 0.04, 5.37 | None | No |
| <i>Trachymyrmex zeteki</i> | 0.10 | 0.24, 1.00 | 1 > 0.95 | Yes |
| | | | 1 > 0.99 | |
| <i>Cyphomyrmex costatus</i> | 0.00 | 0.03, 17.94 | None | Yes |
| <i>Wasmannia auropunctata</i> | 0.08 | 0.08, 22.14 | 2 > 0.95 | Yes |
| | | | 1 > 0.99 | |
| <i>Vollenhovia emeryi</i> | 0.00 | 0.01, 18.18 | None | No |
| <i>Cardiocondyla obscurior</i> | 0.00 | 0.03, 207.50 | None | No |
| <i>Solenopsis invicta</i> | 0.06 | 0.07, 88.31 | 1 > 0.95 | Yes |
| | | | 1 > 0.99 | |
| <i>Monomorium pharaonicum</i> | 0.01 | 0.03, 13.05 | None | No |
| <i>Pogonomyrmex barbatus</i> | 0.00 | 0.00, 1.00 | None | No |
| Formicinae | | | | |
| <i>Lasius niger</i> | 0.13 | 0.09, 999.00 | 1 > 0.95 | Yes |
| <i>Camponotus floridanus</i> | 0.11 | 0.13, 51.08 | 1 > 0.95 | Yes |
| | | | 2 > 0.99 | |
| Cerapachyinae | | | | |
| <i>Ooceraea biroi</i> | 0.02 | 0.04, 133.61 | None | No |
| Ponerinae | | | | |
| <i>Dinoponera quadriceps</i> | 0.00 | 0.00, 1.00 | None | No |
| <i>Harpegnathos saltator</i> | 0.00 | 0.00, 1.00 | None | No |

biased in *Cam. floridanus* and *H. saltator* as well as the three nonant Hymenopteran species examined, although statistical tests were not possible in these species (fig. 3 and [supplementary fig. S6, Supplementary Material](#) online).

The other two ant species did not display male-biased expression of *dsx*. In *V. emeryi*, *dsx* and *DMRT11E* were expressed in all three castes, resembling the expression profiles of the other four ants (fig. 3). However, the expression levels of *dsx* and *DMRT11E* did not differ significantly among the different castes (one-way ANOVA, *dsx*: P -value = 0.40; *DMRT11E*: P -value = 0.10). No difference in *dsx* expression levels was detected even when queens and workers were pooled as females in comparison to males (t -test, P -value = 0.24). In addition, *DMRT11E* expression levels were significantly higher than *dsx* in queens (t -test, P -value = 0.04) and males (t -test, P -value = 0.01). In *W. auropunctata*, *dsx* expression was significantly higher in queens than both males and workers (one-way ANOVA, P -value < 0.01, followed by the Tukey's HSD post hoc test), and *DMRT11E* expression was significantly higher in workers compared with queens (one-way ANOVA, P -value = 0.02, followed by the Tukey's HSD post hoc test). The queen-biased rather than male-biased expression of *dsx* stands in contrast to the other species (fig. 3) and is distinctive among all other insects included in the analysis.

dsx Has Different Exon-Usage Patterns Compared with *DMRT11E* in Ants

The 3'-sequence of *dsx*, which includes the OD2 domain, is alternatively spliced in a sex-specific manner in other insects (Baker and Wolfner 1988; Cho et al. 2007). We were curious as to whether ant *dsx* genes were alternatively spliced in the condition of losing OD2. We applied DEXSeq to the RNA-seq data to test for differential usage of *dsx* and *DMRT11E* exons caused by the differentiation of castes. Common *dsx* transcripts that are both expressed in females and males have been found in several other insects (Cho et al. 2007; Duan et al. 2013), and there are also some nonsense splice forms that are possibly degraded by nonstop-mediated decay (Duan et al. 2013). So with the existence of the common transcripts and the nonsense transcripts, the expression patterns of *dsx* exons detected by DEXseq were expected to overlap between the two sexes or among castes. But still, the specific spliced exons could be deduced from the significant difference of the exon usage abundance among the three castes. Unexpectedly, we found significant differential usage of *dsx* exons in different castes for all three ant species investigated (*S. invicta*, *V. emeryi*, and *W. auropunctata*; fig. 4 and [supplementary table S6, Supplementary Material](#) online). In contrast,




Fig. 3.—Sex- and caste-specific gene expression levels of *Dmrt* genes in five ant species. Gene expression levels are shown in FPKM. The phylogenetic relationship is based on Moreau and Bell (2013). For species with at least three biological replicates (*W. auropunctata*, *V. emeryi*, and *S. invicta*), error bars represent standard error of the mean (SEM), and asterisks indicate *P*-value from a one-way ANOVA ($P < 0.05$), followed by the Tukey’s HSD post hoc test. Q, queen; M, male; W, worker; MiW, minor worker; MaW, major worker (*Cam. floridanus* has two nonreproductive castes); G, gamergate (workers having the ability to replace the queen when the queen dies [Peeters et al. 2000]). Genes that are not expressed in any caste are not shown in the graph.

there was no evidence of caste-specific alternative splicing for the *DMRT11E* ORFs (supplementary fig. S7, Supplementary Material online).

In the transcript models deduced by DEXSeq, we observed that caste-specific alternative splicing events that affect ORFs occur predominantly in the 3'-ends of *dsx* via caste-specific




Fig. 4.—Estimated exon usage of *dsx* in *S. invicta* (A), *V. emeryi* (B), and *W. auropunctata* (C). Exon usage is color coded for the three castes. Annotated transcript models are shown below for each species, with boxes representing exons and lines for introns. Significant differences in exon usage are shown in purple (P -value < 0.05; [supplementary table S6, Supplementary Material](#) online). The positions of start codons are indicated by black arrowheads. The predicted alternatively spliced C-terminus starting from the sex-specific skipped exon is highlighted in yellow. In these regions, exon-usage changes (absolute value of log₂ fold-changes, >1) of male-to-queen and worker-to-queen are shown as blue and red numbers, respectively.

exon skipping, resulting in the production of alternative transcripts containing different distal exons (fig. 4). The patterns of alternative splicing were similar for the three species, with the skipped exons (exons 9–11 in *Sinv_dsx*; exons 13–15 in

Veme_dsx; exon 6 in *Waur_dsx*) showing less expression in males than in either the queens or workers, and the distal exons (*Sinv_dsx*, exons 12–13; *Veme_dsx*, exons 16–18; *Waur_dsx*, exons 7–9) showing the highest expression in

males among the three castes (statistical analyses are given in [supplementary table S6, Supplementary Material](#) online). These results are consistent with the conserved splicing pattern for insect *dsx*: Female-specific exons are skipped in males, and the male-specific transcripts are more extended at the 3'-terminus. For *Sinv_dsx*, there were no significant expression differences for any exons between queens and workers. In contrast, in *Veme_dsx* and *Waur_dsx*, exon usage between queens and workers for some exons was different (*Veme_dsx*, exons 17 and 18; *Waur_dsx*, exon 7; [fig. 4 and supplementary table S6, Supplementary Material](#) online).

Taken together, our results suggest that although *dsx* lacks the OD2 domain, it is still alternatively spliced in ants, producing different C-terminal sequences with no similarity to known proteins. Furthermore, it is notable that both *Veme_dsx* and *Waur_dsx* exhibit caste-specific splicing between the two female castes.

Discussion

Evolution of *Dmrt* Genes in Ants

The *Dmrt* gene family is conserved among metazoans (Bellefroid et al. 2013). Our survey of four *Dmrt* genes in Hymenopterans by annotation and phylogenetic analyses revealed variation in the degree of conservation for *Dmrt* genes. All four *Dmrt* genes were primarily comprised single-copy genes, but there may be more gene gains and losses in ants than other Hymenopteran species.

Phylogenetic analyses of *dsx* in ants and vespids revealed long branch lengths compared with the other three *Dmrt* genes, indicating that *dsx* experienced the highest evolutionary rate among the four *Dmrt* genes. Accordingly, a selective pressure test revealed positively selected sites in *dsx* but not in the three other *Dmrt* genes. Moreover, the evolutionary rate of *dsx* genes varies in different ant subfamilies. PAML results suggest no positive selection on *dsx* in the phylogenetically basal subfamily Ponerinae but positive selection in the “higher ant” subfamilies Myrmicinae and Formicinae. Given that most Myrmicinae and Formicinae display a clear reproductive division of labor and a sophisticated communication systems (Hölldobler and Wilson 1990, 2009), in contrast to the Ponerinae, this raised the possibility of a role of *dsx* in the evolution of complex social organization.

Ant *dsx* Genes Are Alternatively Spliced despite Losing OD2

In our research, sequences homologous to the OD2 domain were only found in the Poneroid clade, a basal lineage of ants (Moreau and Bell 2013). Given that the OD2 domain was also present in the outgroup Vespidae, we propose that the OD2 domain was lost prior to the early radiations of the two advanced ant subfamilies (Myrmicinae and Formicinae) (Moreau and Bell 2013). Lineage-specific losses of OD2 have also been

reported in Hemiptera, Embioptera, Isoptera, and Blattaria, which are hemimetabolous insects (Geuverink and Beukeboom 2014; Price et al. 2015). However, there was no previous evidence of the OD2 domain being lost in holometabola.

The OD1 and OD2 domains in DSX promote oligomerization independently. OD1 and DNA-binding domain, which together compose the DM domain, are often present in all splice variants of DSX and together form a dimeric DNA-binding unit (An et al. 1996; ErdmAn et al. 1996). OD1 dimerization is important for DSX to bind to DNA, and OD2 provides additional dimerization energy, promoting and stabilizing dimeric interactions between DSX monomers (Cho and Wensink 1998).

The lack of OD2 in these ants may indicate that DSX in these species interacts with target DNAs in a monomeric form. However, although it has been reported that DSX could bind to DNA *in vitro* as monomer (Zhu et al. 2000), DSX binds to target DNA as dimer predominantly under conditions containing enough free DNA (Cho and Wensink 1996; Zhu et al. 2000). An alternative hypothesis is that the ant DSXs may bind to DNA in a dimeric form using the weak oligomerization energy provided by OD1. But the OD1–OD1 interaction alone may be insufficient to maintain an optimal dimeric structure, reducing DNA-binding affinity and specificity. Further studies are needed to investigate the consequence of the absence of the OD2 domain in DSX on its functional structure.

OD2 has other important roles than dimerization. The sex-specific OD2 confers upon *Drosophila* DSX the ability to regulate sex-specific gene expression. OD2 dimerization mediates the recruitment of other transcriptional cofactors such as Intersex (Garrett-Engel et al. 2002), increases DNA binding of the DM domain (Cho and Wensink 1998), and enhances binding site recognition (Cho and Wensink 1997). Changes that abolish OD2 dimerization result in an intersexual phenotype (ErdmAn et al. 1996). As a result, it seems possible that the absence of OD2 in ants may alter the specific transcriptional regulatory activity of DSX on its target genes.

However, although the *dsx* of higher ants lacks OD2, it is still alternatively spliced. There are sex- and caste-specific exon usages of ant *dsx* genes, without the OD2 domain. These results suggest that novel sex- or caste-specific DSX C-termini are generated by alternative splicing in these ants. Rather than completely losing their function, ant DSXs could have retained the ability to recruit different regulatory complexes in different sexes or castes. More functional experiments are needed to explore how DSX protein–protein interactions may be different in ants.

Unusual Expression of *dsx* in Two Genetic Caste Determination Ants

Our comparative transcriptomic analysis across eight Hymenopteran species revealed a male-biased expression pattern of *dsx* in six species. Male-biased expression of *dsx* was

also found in the ant *C. obscurior* (Klein et al. 2016). Homologs of *dsx* in arthropods outside of Hexapoda, such as water fleas and mites, also display male-biased expression (Kato et al. 2011; Pomerantz and Hoy 2015). However, the ants *W. auropunctata* and *V. emeryi* did not show the typical male-biased expression. In contrast to *S. invicta* and other well-studied insects that only generate different *dsx* isoforms between sexes, these two species express *dsx* transcripts differently both between sexes (queen and male) and among castes (worker and queen).

The distinctive *dsx* expression patterns in *W. auropunctata* and *V. emeryi* may be related to their special caste determination system. While most ants have an environmental caste determination system where the queen and worker castes are determined by environmental stimuli, *W. auropunctata* and *V. emeryi* have a genetic caste determination system, where caste differentiation is predicted by the individual's genotype (Schwander et al. 2010). In these two species, queens are produced by ameiotic parthenogenesis while workers are produced through normal sexual reproduction. Moreover, in both species males are produced clonally from their father, possibly as a result of the elimination of the maternal genome once the eggs have been fertilized (Fournier et al. 2005; Ohkawara et al. 2006). It would be of interest to study whether the noncanonical expression patterns of *dsx* in these two species are associated with their unusual mode of caste determination.

dsx May Be Related to Caste Differentiation in Ants

Another line of evidence that *dsx* may be associated with caste differentiation in ants is the correlation between *dsx* evolutionary patterns and the level of social organization. In contrast to most other ants, the workers of the phylogenetically basal ant species *H. saltator* and *Di. quadricaps* are able to mate and lay fertilized eggs. Both of these species retained the OD2-like domains in DSX, while all the more advanced ant species analyzed here have lost the OD2 domain, raising the possibility that changes in the ant *dsx* domain organization were possibly associated with the evolution of some crucial functions associated with eusocial life in ants, such as caste determination.

It is well known that the network comprising IIS and other pathways controls the development of different castes. These pathways integrate the different external stimuli such as nutrition and temperature, and ultimately initiate the different caste fates (queen and worker). However, the genetic mechanisms of developmental responses to IIS and other pathways are still unclear (Corona et al. 2016). In stag beetles, *dsx* interacts with nutrition-related pathways, such as the JH signaling pathway, to regulate sex-specific mandible growth (Gotoh et al. 2014). Recently, *dsx* was found to have sex- and morph-specific expression at different developmental stages in the ant *C. obscurior* (Klein et al. 2016). Similarly

RNA-seq data revealed that *dsx* was highly differentially expressed across different tissues of honey bee nurses and foragers, suggesting that *dsx* is a possible candidate gene for the key regulator of caste-specific gene expression (Johnson and Jasper 2016). Taken together, these studies combined with our evolutionary analysis suggest that *dsx* may function as a “connector” in the caste determination cascade. *dsx* may be sensitive to hormones, and mediate the signal transduction from the initial input of nutrition or other environmental cues to effectors that control development of different traits. Finally, this connector model indicates that large evolutionary changes (e.g., caste differentiation) can be achieved by changes in gene regulation without the need to involve novel gene formation. However, it cannot be ruled out that other mechanisms may be responsible for the lack of OD2 domains in ants. Future functional experiments will be needed to address the impact of the lack of the OD2 domain in ants.

In conclusion, this study suggests a specific pattern in the evolution of *dsx* that correlates with the evolution of eusociality in ants. In the ant species displaying an early evolutionary state of eusociality, *dsx* had ancestral gene structure. In contrast, the loss of the OD2 domain and positive selection on *dsx* indicate an extraordinarily high evolutionary rate in the advanced ant species. Besides, there are some exclusive patterns of expression of *Dmrt* genes in two genetic caste determination ants. All these results suggest that *Dmrt* genes may be involved in the development of caste-specific morphological features. More functional studies are necessary to more fully understand the role of *Dmrt* genes in ants.

Supplementary Material

Supplementary data are available at *Genome Biology and Evolution* online.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (31601854 to L.-Y.J., 31672336 to J.-H.X., and 31572313 to D.-W.H.). The computing resource was supported by HPC Platform, Scientific Information Center, Institute of Zoology, CAS, China.

Literature Cited

- Altschul SF. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25(17):3389–3402.
- An WQ, Cho SY, Ishii H, Wensink PC. 1996. Sex-specific and non-sex-specific oligomerization domains in both of the *doublesex* transcription factors from *Drosophila melanogaster*. *Mol Cell Biol.* 16(6):3106–3111.
- Baker BS, Wolfner MF. 1988. A molecular analysis of *doublesex*, a bifunctional gene that controls both male and female sexual-differentiation in *Drosophila melanogaster*. *Genes Dev.* 2(4):477–489.

- Bellefroid EJ, et al. 2013. Expanding roles for the evolutionarily conserved Dmrt sex transcriptional regulators during embryogenesis. *Cell Mol Life Sci.* 70(20):3829–3845.
- Bourke AFG. 1988. Worker reproduction in the higher eusocial Hymenoptera. *Q Rev Biol.* 63:291–311.
- Buschinger A. 1968. Untersuchungen an *Harpagoxenus sublaevis* Nyl. (Hymenoptera, formicidae). *Insectes Soc.* 15(1):89–104.
- Camacho C, et al. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.
- Cheng DF, Zhang ZL, He XF, Liang GW. 2013. Validation of reference genes in *Solenopsis invicta* in different developmental stages, castes and tissues. *PLoS One* 8(2):e57718.
- Cho S, Huang ZY, Zhang JZ. 2007. Sex-specific splicing of the honeybee *doublesex* gene reveals 300 million years of evolution at the bottom of the insect sex-determination pathway. *Genetics* 177(3):1733–1741.
- Cho S, Wensink PC. 1997. DNA binding by the male and female doublesex proteins of *Drosophila melanogaster*. *J Biol Chem.* 272(6):3185–3189.
- Cho S, Wensink PC. 1998. Linkage between oligomerization and DNA binding in *Drosophila* doublesex proteins. *Biochemistry* 37(32):11301–11308.
- Cho SY, Wensink PC. 1996. Purification and physical properties of the male and female double sex proteins of *Drosophila*. *Proc Natl Acad Sci U S A.* 93(5):2043–2047.
- Corona M, Libbrecht R, Wheeler DE. 2016. Molecular mechanisms of phenotypic plasticity in social insects. *Curr Opin Insect Sci.* 13:55–60.
- Crisuolo A. 2011. morePhyML: improving the phylogenetic tree space exploration with PhyML 3. *Mol Phylogenet Evol.* 61(3):944–948.
- Darriba D, Taboada GL, Doallo R, Posada D. 2011. ProtTest 3: fast selection of best-fit models of protein evolution. *Bioinformatics* 27(8):1164–1165.
- Duan JP, et al. 2013. New insights into the genomic organization and splicing of the *doublesex* gene, a terminal regulator of sexual differentiation in the silkworm *Bombyx mori*. *PLoS One* 8(11):e79703.
- Eddy SR. 1998. Profile hidden Markov models. *Bioinformatics* 14(9):755–763.
- Erdman SE, Burtis KC. 1993. The *Drosophila* doublesex proteins share a novel zinc finger related DNA binding domain. *EMBO J.* 12(2):527–535.
- Erdman SE, Chen HJ, Burtis KC. 1996. Functional and genetic characterization of the oligomerization and DNA binding properties of the *Drosophila* doublesex proteins. *Genetics* 144(4):1639–1652.
- Felsenstein J. 1978. Cases in which parsimony or compatibility methods will be positively misleading. *Syst Zool.* 27(4):401–410.
- Fournier D, et al. 2005. Clonal reproduction by males and females in the little fire ant. *Nature* 435(7046):1230–1234.
- Gadau J, et al. 2012. The genomic impact of 100 million years of social evolution in seven ant species. *Trends Genet.* 28(1):14–21.
- Garrett-Engel CM, et al. 2002. *intersex*, a gene required for female sexual development in *Drosophila*, is expressed in both sexes and functions together with *doublesex* to regulate terminal differentiation. *Development* 129(20):4661–4675.
- Geuverink E, Beukeboom LW. 2014. Phylogenetic distribution and evolutionary dynamics of the sex determination genes *doublesex* and *transformer* in insects. *Sex Dev.* 8(1–3):38–49.
- Goldman N, Yang ZH. 1994. Codon-based model of nucleotide substitution for protein-coding DNA-sequences. *Mol Biol Evol.* 11:725–736.
- Gotoh H, et al. 2014. Developmental link between sex and nutrition; *doublesex* regulates sex-specific mandible growth via juvenile hormone signaling in stag beetles. *PLoS Genet.* 10(1):e1004098.
- Hölldobler B, Wilson EO. 1990. *The ants*. London: Harvard University Press.
- Hölldobler B, Wilson EO. 2009. *The superorganism: the beauty, elegance, and strangeness of insect societies*. New York: W.W. Norton & Company.
- Helmkamp M, Cash E, Gadau J. 2015. Evolution of the insect *desaturase* gene family with an emphasis on social Hymenoptera. *Mol Biol Evol.* 32(2):456–471.
- Johnson BR, Jasper WC. 2016. Complex patterns of differential expression in candidate master regulatory genes for social behavior in honey bees. *Behav Ecol Sociobiol.* 70(7):1033–1043.
- Kato Y, et al. 2008. Molecular cloning and sexually dimorphic expression of DM-domain genes in *Daphnia magna*. *Genomics* 91(1):94–101.
- Kato Y, Kobayashi K, Watanabe H, Iguchi T. 2011. Environmental sex determination in the branchiopod crustacean *Daphnia magna*: deep conservation of a *Doublesex* gene in the sex-determining pathway. *PLoS Genet.* 7(3):e1001345.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 30(4):772–780.
- Klein A, et al. 2016. Evolution of social insect polyphenism facilitated by the sex differentiation cascade. *PLoS Genet.* 12(3):e1005952.
- Krieger MJB, Ross KG. 2002. Identification of a major gene regulating complex social behavior. *Science* 295(5553):328–332.
- Li YF, Rao XY, Mattox WW, Amos CI, Liu B. 2015. RNA-seq analysis of differential splice junction usage and intron retentions by DEXSeq. *PLoS One* 10(9):e0136653.
- Livramento K, Freitas N, Máximo W, Zanetti R, Paiva L. 2018. Gene expression profile analysis is directly affected by the selected reference gene: the case of leaf-cutting *Atta sexdens*. *Insects* 9(1):18.
- Marchler-Bauer A, et al. 2015. CDD: NCBI's conserved domain database. *Nucleic Acids Res.* 43(Database issue):D222–D226.
- Matson CK, Zarkower D. 2012. Sex and the singular DM domain: insights into sexual regulation, evolution and plasticity. *Nat Rev Genet.* 13(3):163–174.
- Moreau CS, Bell CD. 2013. Testing the museum versus cradle tropical biological diversity hypothesis: phylogeny, diversification, and ancestral biogeographic range evolution of the ants. *Evolution* 67(8):2240–2257.
- Nagoshi RN, McKeown M, Burtis KC, Belote JM, Baker BS. 1988. The control of alternative splicing at genes regulating sexual differentiation in *D. melanogaster*. *Cell* 53(2):229–236.
- Ohkawara K, Nakayama M, Satoh A, Trindl A, Heinze J. 2006. Clonal reproduction and genetic caste differences in a queen-polymorphic ant, *Vollenhovia emeryi*. *Biol Lett.* 2(3):359–363.
- Ottolenghi C, Fellous M, Barbieri M, McElreavey K. 2002. Novel paralogy relations among human chromosomes support a link between the phylogeny of *doublesex*-related genes and the evolution of sex determination. *Genomics* 79(3):333–343.
- Peeters C, Liebig J, Hölldobler B. 2000. Sexual reproduction by both queens and workers in the ponerine ant *Harpegnathos saltator*. *Insectes Soc.* 47(4):325–332.
- Pomerantz AF, Hoy MA. 2015. Expression analysis of *Drosophila doublesex*, *transformer-2*, *intersex*, *fruitless-like*, and *vitellogenin* homologs in the parhaploid predator *Metaseiulus occidentalis* (Chelicerata: Acari: Phytoseiidae). *Exp Appl Acarol.* 65(1):1–16.
- Price DC, Egizi A, Fonseca DM. 2015. The ubiquity and ancestry of insect *doublesex*. *Sci Rep.* 5:13068.
- R Core Team. 2000. R language definition. Vienna (Austria): R Foundation for Statistical Computing.
- Rideout EJ, Dornan AJ, Neville MC, Eadie S, Goodwin SF. 2010. Control of sexual differentiation and behavior by the *doublesex* gene in *Drosophila melanogaster*. *Nat Neurosci.* 13(4):458–466.
- Sanchez L, Gorfinkiel N, Guerrero I. 2001. Sex determination genes control the development of the *Drosophila* genital disc, modulating the response to *Hedgehog*, *Wingless* and *Decapentaplegic* signals. *Development* 128(7):1033–1043.
- Scharlaken B, et al. 2008. Reference gene selection for insect expression studies using quantitative real-time PCR: the head of the honeybee, *Apis mellifera*, after a bacterial challenge. *J Insect Sci.* 8(33):1–10.

- Schwander T, Lo N, Beekman M, Oldroyd BP, Keller L. 2010. Nature versus nurture in social insect caste differentiation. *Trends Ecol Evol.* 25(5):275–282.
- Shirangi TR, Dufour HD, Williams TM, Carroll SB. 2009. Rapid evolution of sex pheromone-producing enzyme expression in *Drosophila*. *PLoS Biol.* 7(8):e1000168.
- Toyota K, et al. 2013. Molecular cloning of *doublesex* genes of four Cladocera (water flea) species. *BMC Genomics* 14:239.
- Trapnell C, et al. 2012. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc.* 7(3):562–578.
- Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* 25(9):1105–1111.
- Verhulst EC, van de Zande L. 2015. Double nexus—*Doublesex* is the connecting element in sex determination. *Brief Funct Genomics.* 14(6):396–406.
- Volff JN, Zarkower D, Bardwell VJ, Schartl M. 2003. Evolutionary dynamics of the DM domain gene family in metazoans. *J Mol Evol.* 57:S241–S249.
- Ward PS, Brady SG, Fisher BL, Schultz TR. 2015. The evolution of myrmicine ants: phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae). *Syst Entomol.* 40(1):61–81.
- Wexler JR, Plachetzki DC, Kopp A. 2014. Pan-metazoan phylogeny of the DMRT gene family: a framework for functional studies. *Dev Genes Evol.* 224(3):175–181.
- Yang ZH. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol.* 24(8):1586–1591.
- Yang ZH, Nielsen R, Goldman N, Pedersen AMK. 2000. Codon-substitution models for heterogeneous selection pressure at amino acid sites. *Genetics* 155:431–449.
- Zhang JZ, Nielsen R, Yang ZH. 2005. Evaluation of an improved branch-site likelihood method for detecting positive selection at the molecular level. *Mol Biol Evol.* 22(12):2472–2479.
- Zhang M, et al. 2014. Cloning, expression and cellular localization of the *Doublesex* gene in the water flea, *Daphnia carinata*, during different developmental stages. *Gene* 550(2):185–192.
- Zhu LY, et al. 2000. Sexual dimorphism in diverse metazoans is regulated by a novel class of intertwined zinc fingers. *Genes Dev.* 14(14):1750–1764.

Associate editor: Ya-Ping Zhang