

Serveur Académique Lausannois SERVAL [serval.unil.ch](http://serval.unil.ch)

## Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

**Title:** Rituximab as monotherapy for the treatment of chronic active antibody-mediated rejection after kidney transplantation.

**Authors:** Muller YD, Ghaleb N, Rotman S, Vionnet J, Halfon M, Catana E, Golshayan D, Venetz JP, Aubert V, Pascual M

**Journal:** Transplant international : official journal of the European Society for Organ Transplantation

**Year:** 2018 Apr

**Issue:** 31

**Volume:** 4

**Pages:** 451-455

**DOI:** [10.1111/tri.13111](https://doi.org/10.1111/tri.13111)

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

DR. YANNICK MULLER (Orcid ID : 0000-0003-0513-7156)

Article type : Letter to the Editor

## Rituximab as monotherapy for the treatment of chronic active antibody-mediated rejection after kidney transplantation

Yannick D. Muller (1), Ghaleb Nseir (1), Samuel Rotman (2), Julien Vionnet (1), Mathieu Halfon (1), Emmanuelle Catana (1), Déla Golshayan (1), Jean-Pierre Venetz (1), Vincent Aubert (3), Manuel Pascual (1)

(1) Transplantation Center, Lausanne University Hospital, Lausanne, Switzerland

(2) Service of Clinical Pathology, Lausanne University Hospital, Lausanne, Switzerland

(3) Service of Immunology and Allergy, Lausanne University Hospital, Lausanne, Switzerland

**Competing interests:** The authors have no financial disclosure nor any conflicts of interest.

**Funding information:** none

**Authorship:** Y.D.M, G.N, M.P designed research study. Y.D.M, G.N, S.R, E.C, J.V, M.H, D.G, J-P.V and V.A performed research/study and collected the data. Y.M, G.N, S.R, V.A and M.P analyzed the data, Y.D.M and M.P wrote the paper.

### **Corresponding author contact information:**

Manuel Pascual, Centre de Transplantation d'Organes, Centre Hospitalier Universitaire Vaudois (CHUV) Rue du Bugnon 11, 1011 Lausanne. Tel : +41 795563720. manuel.pascual@chuv.ch

Yannick Muller, Centre de Transplantation d'Organes, Centre Hospitalier Universitaire Vaudois (CHUV). yannick.muller@chuv.ch

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/tri.13111

This article is protected by copyright. All rights reserved.

**Running Title:** Rituximab and chronic active antibody-mediated rejection

**Keywords:** chronic active antibody-mediated rejection, humoral rejection, rituximab, kidney, transplantation.

**Abbreviations:** Chronic active antibody-mediated rejection (caAMR), donor-specific anti-HLA antibodies (DSA), mean fluorescence intensity (MFI), intravenous immunoglobulins (IVIg).

**Ethical protocol:** this study was reviewed and accepted by the “Commission cantonale d'éthique de la recherche sur l'être humain” (CER-VD, protocol 2017-01077).

**Dear Editors,**

Chronic active antibody-mediated rejection (caAMR) is a major cause of allograft loss after kidney transplantation (1). The BANFF 2013 classification redefined caAMR by the presence of donor-specific anti-HLA antibodies (DSA) together with immuno-histopathological evidence for active vascular lesions of the endothelium (C4d deposits, glomerulitis, peritubular capillaritis) as well as evidence of chronic tissue injury (transplant glomerulopathy, peritubular capillary basement membrane multilayering or arterial intimal fibrosis) (2,3). Humoral immunity, detected by the presence of DSA, and B cells are considered pivotal in the development of caAMR. Gosset et al. showed that circulating DSA are responsible for accelerated allograft fibrosis independently of acute AMR (1). Recent evidence suggested that B cells also mediate chronic allograft rejection independently of DSA (4) and that rituximab, a chimeric IgG1-antibody targeting human CD20, impair B-cell regulation of T-cells (5-7). However, the efficacy of rituximab in patients with caAMR remains controversial (8).

We retrospectively analyzed 12 kidney-transplant recipients followed at our institution between January 2007 and December 2012 who were diagnosed with biopsy-proven caAMR (Table 1). All patients were treated with a regimen of 2 doses of rituximab (375 mg/m<sup>2</sup>) at one week interval, as monotherapy, together with a transient increase of 20 mg/day oral prednisone, progressively tapered to 5 mg at three months. Peripheral B-cell depletion (<10 B-cells/ $\mu$ l) after rituximab administration was confirmed by flow cytometry. In seven out of 12 patients, there was a decrease in DSA levels below 2000 mean fluorescence intensity (MFI) (Luminex, Austin, TX) or a >75% DSA level decrease after rituximab (Figure 1A). Serum creatinine and proteinuria levels were not significantly different after rituximab. Regarding active antibody-mediated vascular lesions, 4/6 patients with positive C4d deposits, 6/8 with peritubular capillaritis and 3/4 with glomerulitis, had a decrease in the activity score at 1-year post rituximab (Figure 1B). However, chronic histological lesions did not improve one year after rituximab (Figure 1C). We further analyzed clinical responders defined as patients who improved or maintained stable kidney function at 12 months post-rituximab (<20% change of the baseline creatinine) with a decrease of DSA titers below 2000 MFI or >75% DSA titers decrease. The results showed significantly more C4d deposition in patients with persistent DSA before and after rituximab (Figure 2A-B), whereas no significant differences were found between clinical responders and non-responders regarding capillaritis and glomerulitis (Figure 2C-D).

We performed CD20 and CD45 staining on all kidney allograft biopsies before and after rituximab. The results showed a significant reduction of CD20<sup>+</sup> B cells in the kidney allograft at one-year (Figure 3A-B). Interestingly, the absolute number of CD45<sup>+</sup> was not changed (Figure 3C-D). Peripheral B-cell depletion (<10 B-cells/ $\mu$ l) was present only in 5/12 patients after one-year and absolute B cell numbers were not different between responders and non-responders (data not shown). These results demonstrate that CD20<sup>+</sup> B-cells were targeted in the kidney allograft, although complete long-term depletion was not achieved.

Evidence for successful management of caAMR is currently lacking. The first randomized trial of eculizumab therapy for caAMR failed to show clinically-relevant efficacy on kidney function, DSA titers and chronic-active histopathological lesions (9). More recently, Moreso et al published a second randomized trial comparing patients with transplant glomerulopathy and DSA receiving four cures of intravenous immunoglobulins (IVIG) followed by one infusion (375mg/m<sup>2</sup>) of rituximab compared to placebo, but no improvement in histological lesions, DSA titers and kidney function was observed (10). Although both studies were underpowered, they demonstrated that untreated patients did not improve chronic active histological lesions and maintained high-titers DSA.

Despite the negative results of the two randomized trials, there is some evidence suggesting that B cell depletion is important in the management of caAMR. Thus, patients highly sensitized or receiving ABO-incompatible transplants were more protected against chronic rejection if treated with rituximab (11,12). In the setting of acute AMR and intensive immunosuppression regimen, rituximab seems also to be

beneficial (5,13-14). In the present series, rituximab administration was associated with a reduction in DSA levels and active histopathological lesions in most patients, although this did not translate in an improvement of allograft function, which was also confirmed by absence of improvement of chronic lesions in 1-year control biopsies. In the EudraCT trial, rituximab was administered only once, up to 6 months after that the diagnosis was made resulting in no improvement of DSA or histological lesions (10). The results of the prospective multicentric randomized trial (NCT00307125), aiming to compare the use of rituximab versus placebo for the treatment of *de novo* early DSA in kidney allograft recipients over a 3-year period time, are therefore expected with interest. Finally, it must be also emphasized that future protocols should consider the possibility to reinfuse rituximab every 6 months as B-cell depletion was found to be significantly reduced but not complete one year after rituximab.

The importance of chronic active T cell-mediated rejection in caAMR should not be neglected as T helper cells are key in priming B-cells. Earlier studies showed that reversal of DSA can be achieved in the absence of a “desensitization regimen”, but in general this was associated with some rescue therapy based on tacrolimus and mycophenolate mofetil switch and corticosteroids (15,16). Furthermore, allograft survival can be significantly improved after *de novo* DSA detection if levels of tacrolimus are maintained above 5.3 ng/ml (17). In our series, patients received transiently more oral prednisone, which possibly acted synergistically with rituximab against active lesions, although it must be acknowledged that in the absence of an untreated control group, the interpretation of the data must be taken with caution.

Tocilizumab and C1-inhibitor, have also been studied for caAMR. Thirty-six patients who were refractory to rituximab and IVIG treatment, received tocilizumab, an anti-IL-6 receptor monoclonal antibody, as rescue therapy, which resulted in reduction in DSA titers and in stabilization of allograft function at 2-years (18). Because rituximab depletes CD20<sup>+</sup> precursor B cells and tocilizumab disrupts the IL-6 supplying niches important for Th1/Tfollicular differentiation and long-lived plasma cells, a synergic beneficial action of both antibodies could be hypothesized as in Castleman's disease (19). Recently, plasma-derived human C1-inhibitor (20UI/kg/twice weekly), an inhibitor which targets the classical complement pathway was successfully administered for caAMR prevention in highly sensitized patients (20,21). Thus, in the future, non-responders to rituximab and/or IVIG may be candidates for combination therapy of B cell depleting agents with complement inhibitors or tocilizumab.

At our center, we do not perform routine protocol biopsies, so that we did not have the possibility to select well-matched controls. Even if most patients responded to rituximab, active lesions, B cells and in some cases DSA remained present. This was also reported by Gupta *et al.* who targeted B cells using a combination of rituximab, IVIG and bortezomib (22), showing that 48% of the patients with late AMR were non-responders (22). In the future, more detailed analysis is warranted on the potential relationship between DSA, mean fluorescence intensity, HLA types (IgG-IgM-IgA), sub-classes (IgG 1/2/3/4), HLA class I or II specificity, epitope specificity, complement binding (C1q) DSA ability and clinical responses to rituximab. Moreover, distinctions between patients with preexisting DSA and *de novo* DSA would also be interesting (23). Overall, it would be important to better define which transplant recipients can benefit from rituximab, alone or in combination with other therapies.

## References

1. Gosset C, Viglietti D, Rabant M et al. Circulating donor-specific anti-HLA antibodies are a major factor in premature and accelerated allograft fibrosis. *Kidney Int.* 2017
2. Haas M, Sis B, Racusen LC et al. Banff 2013 meeting report: inclusion of c4d-negative antibody-mediated rejection and antibody-associated arterial lesions. *Am J Transplant.* 2014;14:272-283.
3. Haas M. The Revised (2013) Banff Classification for Antibody-Mediated Rejection of Renal Allografts: Update, Difficulties, and Future Considerations. *Am J Transplant.* 2016;16:1352-1357.
4. Zeng Q, Ng YH, Singh T et al. B cells mediate chronic allograft rejection independently of antibody production. *J Clin Invest.* 2014;124:1052-1056.
5. Barnett AN, Hadjianastassiou VG, Mamode N. Rituximab in renal transplantation. *Transpl Int.* 2013;26:563-575.
6. Bouaziz JD, Calbo S, Maho-Vaillant M et al. IL-10 produced by activated human B cells regulates CD4(+) T-cell activation in vitro. *Eur J Immunol.* 2010;40:2686-2691.
7. de Wit J, Jorritsma T, Makuch M et al. Human B cells promote T-cell plasticity to optimize antibody response by inducing coexpression of T(H)1/T(FH) signatures. *J Allergy Clin Immunol.* 2015;135:1053-1060.
8. Mulley WR, Huang LL, Ramessur Chandran S et al. Long-term graft survival in patients with chronic antibody-mediated rejection with persistent peritubular capillaritis treated with intravenous immunoglobulin and rituximab. *Clin Transplant.* 2017



9. Kulkarni S, Kirkiles-Smith NC, Deng YH et al. Eculizumab Therapy for Chronic Antibody-Mediated Injury in Kidney Transplant Recipients: A Pilot Randomized Controlled Trial. *Am J Transplant.* 2017;17:682-691.
10. Moreso F, Crespo M, Ruiz JC et al. Treatment of chronic antibody mediated rejection with intravenous immunoglobulins and rituximab: A multicenter, prospective, randomized, double-blind clinical trial. *Am J Transplant.* 2017
11. Loupy A, Suberbielle-Boissel C, Zuber J et al. Combined posttransplant prophylactic IVIg/anti-CD 20/plasmapheresis in kidney recipients with preformed donor-specific antibodies: a pilot study. *Transplantation.* 2010;89:1403-1410.
12. Kohei N, Hirai T, Omoto K, Ishida H, Tanabe K. Chronic antibody-mediated rejection is reduced by targeting B-cell immunity during an introductory period. *Am J Transplant.* 2012;12:469-476.
13. Sautenet B, Blancho G, Büchler M et al. One-year Results of the Effects of Rituximab on Acute Antibody-Mediated Rejection in Renal Transplantation: RITUX ERAH, a Multicenter Double-blind Randomized Placebo-controlled Trial. *Transplantation.* 2016;100:391-399.
14. Zarkhin V, Li L, Kambham N, Sigdel T, Salvatierra O, Sarwal MM. A randomized, prospective trial of rituximab for acute rejection in pediatric renal transplantation. *Am J Transplant.* 2008;8:2607-2617.
15. Stegall MD, Simon M, Wachs ME, Chan L, Nolan C, Kam I. Mycophenolate mofetil decreases rejection in simultaneous pancreas-kidney transplantation when combined with tacrolimus or cyclosporine. *Transplantation.* 1997;64:1695-1700.
16. Theruvath TP, Saidman SL, Mauiyyedi S et al. Control of antidonor antibody

- production with tacrolimus and mycophenolate mofetil in renal allograft recipients with chronic rejection. *Transplantation*. 2001;72:77-83.
17. Béland MA, Lapointe I, Noël R et al. Higher calcineurin inhibitor levels predict better kidney graft survival in patients with de novo donor-specific anti-HLA antibodies: a cohort study. *Transpl Int*. 2017;30:502-509.
  18. Choi J, Aubert O, Vo A et al. Assessment of Tocilizumab (Anti-Interleukin-6 Receptor Monoclonal) as a Potential Treatment for Chronic Antibody-Mediated Rejection and Transplant Glomerulopathy in HLA-Sensitized Renal Allograft Recipients. *Am J Transplant*. 2017
  19. Louis C, Vijgen S, Samii K et al. TAFRO Syndrome in Caucasians: A Case Report and Review of the Literature. *Front Med (Lausanne)*. 2017;4:149.
  20. Berger M, Baldwin WM, Jordan SC. Potential Roles for C1 Inhibitor in Transplantation. *Transplantation*. 2016;100:1415-1424.
  21. Vo AA, Zeevi A, Choi J et al. A phase I/II placebo-controlled trial of C1-inhibitor for prevention of antibody-mediated rejection in HLA sensitized patients. *Transplantation*. 2015;99:299-308.
  22. Gupta G, Abu Jawdeh BG, Racusen LC et al. Late antibody-mediated rejection in renal allografts: outcome after conventional and novel therapies. *Transplantation*. 2014;97:1240-1246.
  23. Aubert O, Loupy A, Hidalgo L et al. Antibody-Mediated Rejection Due to Preexisting versus De Novo Donor-Specific Antibodies in Kidney Allograft Recipients. *J Am Soc Nephrol*. 2017;28:1912-1923.

Table 1

Patient's clinical characteristics. Abbreviation: MMF, mycophenolate mofetil; Pred, prednisone; Tac, tacrolimus; Csa, ciclosporine; SRL sirolimus. TX- transplantation date. DSA, donor-specific antibodies, RTX, rituximab, ACR acute cellular rejection.

#### Figure 1

(A) Mean fluorescence intensity (MFI) of *de novo* class I and II donor-specific antibodies (DSA) before and 12 months after rituximab. (B) Active histological lesions including capillary C4d deposits and microvascular inflammation score calculated as the combination score of glomerulitis (g) and peritubular capillaritis (ptc). (C) Chronic histopathological lesions including chronic glomerulopathy score, peritubular capillary (PTC) lamellation score and chronic arteriopathy score. All renal biopsies were evaluated by the same experienced pathologist. Two-tailed student t tests were used. Mean and standard deviations are shown. \* P value < 0.05.

#### Figure 2

(A) Representative CD20 staining. (B) Allograft CD20 positive cells were stained, slides were scanned (Zeiss, Oberkochen, Germany) and cells were counted (XX software) before and one year after rituximab. (C) Representative CD45 staining. (D) Allograft CD45 positive cells were stained, slides were scanned and cells were counted before and one year after rituximab. Immunohistochemistry was performed on paraffin-fixed tissue for mouse anti-human CD20 (1/400, clone L26, Novocastra/Leica, Wetzlar, Germany), and mouse anti-human CD45 (1/1000, clone 2B11, DAKO, Ely, UK) with automat (Ventana) per our clinical standardized protocol. The slides were thereafter scanned (Mirax, Zeiss, Oberkochen, Germany) and cells

were quantified using the DIH Image analysis software and the measure-stained-cells Algorithm (Slidepath software, Leica, Wetzlar, Germany). \*\* Pvalue < 0.01.

### Figure 3

(A) C4d deposition score was compared in clinical responders and non-responders before (A) and 12 months after rituximab. (B) The combined peritubular capillaritis and glomerulitis score was assessed in both group before (C) and 12 months after (D) rituximab. \* P value < 0.05. \*\* P value < 0.01.

**Table 1:** Baseline clinical and biological data of all patients

Patient	Age	Sex	Hypertension	Diabetes mellitus	2d TX	Anterior ACR	IS regimen	Interval TX-DSA (months)	Interval DSA-RTX (months)
1	47	M	Yes	No	No	No	Tac/MMF/Pred	6	1
2	31	M	Yes	No	No	Yes	SRL/MMF/Pred	14	2
3	21	F	No	No	No	Yes	Tac/MMF/Pred	20	0
4	30	F	Yes	No	No	No	CsA/MMF/Pred	30	3
5	20	M	No	No	No	Yes	Tac/MMF	132	0
6	40	F	Yes	No	No	No	SRL/MMF/Pred	84	1
7	61	F	Yes	No	No	No	Tac/MMF	74	3
8	20	F	Yes	No	No	Yes	Tac/MMF/Pred	7	1
9	55	M	Yes	No	Yes	No	Tac/MMF/Pred	0	20
10	46	M	No	Yes	No	No	Tac/MMF	58	4
11	39	M	Yes	No	No	Yes	Tac/MMF	?	9
12	22	F	Yes	Yes	No	Yes	Tac/MMF/Pred	26	2

Table 1 (continued)

Patient	Serum Creatinine $\mu\text{mol/l}$ before RTX	Serum Creatinine $\mu\text{mol/l}$ after RTX	Proteinuria Prot/creat g/mol before RTX	Proteinuria Prot/creat g/mol after RTX	Likely non-compliant	Tac ( $\mu\text{g/L}$ ) CsA ( $\mu\text{g/L}$ ) level at diagnosis
1	149	123	2770	530	N	7.6
2	115	120	800	120	N	5.8
3	125	127	400	300	Y	<2
4	215	228	100	190	Y	239
5	90	107	10	0	N	5.5
6	160	149	380	147	N	-
7	112	113	150	0	N	6.2
8	178	290	34	90	Y	6.2
9	118	119	2360	3500	N	10
10	100	141	140	3000	N	7
11	180	197	700	770	N	5.4
12	240	291	130	430	Y	5.5





