

**Serveur Académique Lausannois SERVAL [serval.unil.ch](http://serval.unil.ch)**

## **Author Manuscript**

**Faculty of Biology and Medicine Publication**

**This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.**

Published in final edited form as:

**Title:** Role of homologous ASP334 and GLU319 in human non-gastric H,K- and Na,K-ATPases in cardiac glycoside binding.

**Authors:** Radkov R, Kharoubi-Hess S, Schaer D, Modyanov NN, Geering K, Horisberger JD

**Journal:** Biochemical and biophysical research communications

**Year:** 2007 Apr 27

**Volume:** 356

**Issue:** 1

**Pages:** 142-6

**DOI:** [10.1016/j.bbrc.2007.02.119](https://doi.org/10.1016/j.bbrc.2007.02.119)

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

Published in final edited form as:

*Biochem Biophys Res Commun.* 2007 April 27; 356(1): 142–146.

## Role of homologous ASP334 and GLU319 in human non-gastric H,K- and Na,K-ATPases in cardiac glycoside binding

Rossen Radkov<sup>1</sup>, Solange Kharoubi-Hess<sup>2</sup>, Danièle Schaer<sup>2</sup>, Nikolai N. Modyanov<sup>1</sup>, Käthi Geering<sup>2</sup>, and Jean-Daniel Horisberger<sup>2</sup>

<sup>1</sup>*Department of Physiology, Pharmacology, Metabolism, and Cardiovascular Sciences, University of Toledo College of Medicine, 3035 Arlington Ave, Toledo, Ohio, 43614 USA* <sup>2</sup>*Department of Pharmacology and Toxicology of the University of Lausanne, Lausanne, Switzerland*

### Summary

Cardiac steroids inhibit Na,K-ATPase and the related non-gastric H,K-ATPase, while they do not interact with gastric H,K-ATPase. Introducing an arginine, the residue present in the gastric H,K-ATPase, in the second extracellular loop at the corresponding position 334 in the human non-gastric H,K-ATPase (D334R mutation) rendered it completely resistant to 2 mM ouabain. The corresponding mutation (E319R) in  $\alpha 1$  Na,K-ATPase produced a ~2 fold increase of the ouabain IC<sub>50</sub> in the ouabain-resistant rat  $\alpha 1$  Na,K-ATPase and a large decrease of the ouabain affinity of human  $\alpha 1$  Na,K-ATPase, on the other hand this mutation had no effect on the affinity for the aglycone ouabagenin. These results provide a strong support for the orientation of ouabain in its binding site with its sugar moiety interacting directly with the second extracellular loop.

### Keywords

H; K-ATPase; Na; K-ATPase; cardiac steroid; cardiac glycoside; ouabain affinity; site directed mutagenesis

### Introduction

According to the classification proposed by Axselsen & Palmgren [1], the group IIc of P-type ATPases, contains three types of ion motive ATPases composed of an  $\alpha$  and a  $\beta$  subunit. Included in this group are the four isoforms ( $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$  and  $\alpha 4$ ) of the  $\alpha$  subunit of Na,K-ATPase, the  $\alpha$  subunit of gastric proton pump or H,K-ATPase ( $\alpha$ HKg) and the  $\alpha$  subunit of the third member of this group, the non-gastric H,K-ATPase ( $\alpha$ HKng, initially named ATP1AL1 in human), colonic H,K-ATPase in rat or H,K-ATPase  $\alpha 2$  subunit, the product of the ATP12A gene. This protein is named H,K-ATPase even if its precise ion specificity and transport stoichiometry are not yet exactly known. All these proteins share highly related sequences with more than 60% amino acid identity.

The  $\alpha$  subunit of the non gastric H,K-ATPase assembles physiologically with the  $\beta 1$  subunit of Na,K-ATPase [2] although functional expression can be obtained with other  $\beta$  isoforms

Corresponding author: J.-D. Horisberger, Institut de Pharmacologie et de Toxicologie, rue du Bugnon 27, CH-1005 LAUSANNE, Switzerland. Phone +41 21 692 5362, FAX +41 21 692 5355, E-Mail Jean-Daniel.Horisberger@unil.ch

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

[3]. In this work we have used the  $\beta$  subunit of the rabbit gastric H,K-ATPase with which we obtain good functional expression of the non gastric H,K-ATPase in *Xenopus* oocytes.

All Na,K-ATPase  $\alpha$  isoforms and  $\alpha$ HKng can be inhibited by ouabain (and a group of related compounds known as cardiac steroids), although with a wide range of sensitivity ranging from nanomolar for highly sensitive isoforms (Na,K-ATPase  $\alpha 2$ ,  $\alpha 3$ , and  $\alpha 1$  in most species), to micromolar for moderately sensitive (Na,K-ATPase  $\alpha 4$ ) and submillimolar for poorly sensitive isoforms (Na,K-ATPase  $\alpha 1$  of rodents, Na,K-ATPase  $\alpha 1$  of the toad *Bufo marinus*, and the non-gastric H,K-ATPases). In contrast, the gastric H,K-ATPase is insensitive to ouabain. Resistance to ouabain in rodents and *Bufo marinus*  $\alpha 1$  isoform is known to be related to the presence of charged residues in the first extracellular loop [4] while the low sensitivity of human and other nongastric H,K-ATPase or the total resistance of gastric H,K-ATPase must be due to other primary structure differences as these proteins do not have charged residues at homologous locations.

Cardiac steroids are used for more than 200 years to treat congestive heart failure and are known for 50 years to be specific inhibitors of Na,K-ATPase. Understanding their binding site and mechanisms is important first because of their wide clinical use but also because of the presence of endogenous related compound(s) that may act as hormonal agents to control the activity of Na,K-ATPase and be implicated in the pathogenesis of conditions such as arterial hypertension or heart failure [5–7]. Another point of interest is the potential role of cardiac steroids as signaling molecules and the signal transducer role of Na,K-ATPase [8], and the potential anti-tumor activity of analogues of cardiac steroids [9;10]. High affinity binding of ouabain to Na,K-ATPase has been studied in details and major progress have been made recently by determining several residues directly involved in the structure of the ouabain binding site by the study of mutants in which residues belonging to the gastric and non gastric H,K-ATPases were exchanged with those of a sensitive form of Na,K-ATPase [11;12]. Besides the well known role of charged residues in the first extracellular loop [13], a small number of residues in the second (transmembrane segments M3-M4) and third (M5-M6) hairpin were shown to be essential for high affinity binding. A structural model of ouabain in its binding site was proposed in which several of these residues make H-bonds with ouabain [11;12]. Several aspects of this model remain hypothetical and some residues that are functionally important are not predicted to make direct contact with the ouabain molecule. Thus more experimental data are needed to obtain a complete and safe model of the ouabain binding site.

To better understand the mechanism of ouabain binding, in particular to the human non gastric H,K-ATPase, we have selected a number of residues, located on the extracytoplasmic surface or in the transmembrane segments, which are divergent between Na,K-ATPase and non-gastric or gastric H,K-ATPase (see table 1), and studied ouabain inhibition of ion transport functions in wild type and mutant  $\alpha$ HKng and  $\alpha 1$  Na,K-ATPase proteins in which residues were exchanged.

## Materials and Methods

### Mutant constructions

Five mutants of human  $\alpha$ HKng were created in which a residue was replaced by the amino acid present at the homologous position in the sequence of rat  $\alpha 1$  (K329E, L942E) or  $\alpha 2$  or  $\alpha 3$  (K329G), in both Na,K- or gastric H,K-ATPase (A800P) or in the gastric H,K-ATPase (D334R). Similarly, 4 mutants of rat Na,K-ATPase  $\alpha 1$  were created in which a residue was replaced by the amino acid present at the homologous position in the sequence of human  $\alpha$ HKng (E314K, P785A, and V927E) or in gastric H,K-ATPase (E319R). Finally a mutant of the human  $\alpha 1$  Na,K-ATPase (E319R) was created in the same way (table1).

Mutagenesis was performed using Quickchange II XL site-Directed Mutagenesis Kit (Stratagene), following the manufacturer's instructions. All mutations were confirmed by sequencing. cRNA of  $\alpha$  and  $\beta$  subunits was prepared by *in vitro* transcription.

### Expression in *Xenopus* oocytes

*Xenopus laevis* were anaesthetized by immersion in MS 222 (2 g/l; Sandoz, Basel, Switzerland) and ovarian tissue was removed. Frogs were killed by decapitation under MS 222 anesthesia. Stage V/VI oocytes were collected from the ovarian fragments treated with collagenase as described earlier [14]. cRNA of mutant and wild type  $\alpha$  subunit (8–10 ng) and  $\beta$  subunit (1 ng) was injected into the vegetal pole of the oocytes. The rat  $\beta 1$  subunit was expressed with the rat  $\alpha 1$  subunit and rabbit  $\beta$ HK was expressed with the human  $\alpha$ HKng. Two to 3 days later the oocytes were loaded with  $\text{Na}^+$  by exposure to a  $\text{K}^+$ -free solution containing 0.2  $\mu\text{M}$  ouabain in order to inhibit the oocyte endogenous Na,K-ATPase. This concentration of ouabain inhibits more than 95% of the endogenous oocyte Na,K-ATPase while it has practically no effect on resistant isoforms such as rat  $\alpha 1$  Na,K-ATPase or non gastric H,K-ATPase [3;15;16].

### Electrophysiological measurements

The functional expression of wild type and mutant Na,K-ATPase was first studied by the 2-electrode voltage clamp technique starting in the following control  $\text{K}^+$  free solution : (in mM)  $\text{Na}^+$  87.4,  $\text{Ca}^{++}$  0.4,  $\text{Mg}^{++}$  0.8, N-methyl-D-glucamine (NMDG) 5,  $\text{Cl}^-$  86.0,  $\text{HCO}_3^-$  2.4, 4-(2-hydroxyethyl)-1-piperazineethanesulphonic acid (HEPES) 10.0. The current induced by addition of 10 mM  $\text{K}^+$  ( $I_{\text{K}^+}$ ) was recorded as well as the current inhibited by 2 mM ouabain added to the solution containing 10 mM  $\text{K}^+$  solution.

For the measurements of ouabain inhibition constant ( $K_{\text{I}}$ ) in oocytes expressing rat  $\alpha 1/\beta 1$  Na,K-ATPase or human  $\alpha$ HKng/rabbit  $\beta$ HK, the current was recorded in the 10 mM  $\text{K}^+$  solution and then during exposure to increasing concentrations of ouabain (50, 200, 600  $\mu\text{M}$  and 2.0 mM).

For measurements of ouabain  $K_{\text{I}}$  in human  $\alpha 1\beta 1$  Na,K-ATPase which has a much higher affinity for ouabain (close to that of the *Xenopus* oocyte Na,K-ATPase) it was not possible to inhibit the endogenous component by exposure to low doses of ouabain. and also, because of the slow association rate constant of ouabain, it was not possible to use acute exposure to increasing concentration of ouabain. We therefore started with  $\text{Na}^+$  loaded oocyte injected either with the human  $\beta 1$  alone or wild type or mutant  $\alpha 1 + \beta 1$  human Na,K-ATPase. We then exposed groups of oocytes to various concentrations of ouabain during 1 hour to reach equilibrium binding and then measured rapidly (within 1 min) the current induced by 10 mM  $\text{K}^+$  and then the current inhibited by a high concentration (2 mM) of ouabain. This concentration is expected to inhibit very close to 100 % of the sensitive human  $\alpha 1\beta 1$  Na,K-ATPase. Ouabain  $K_{\text{I}}$  was obtained by analyzing the relationship between the remaining ouabain-sensitive current and the concentration of ouabain during the 1 h exposure. To obtain the affinity of the exogenous component the mean current measured in the oocytes expressing the  $\beta$  subunit alone (endogenous component) was subtracted at each corresponding ouabain concentration.

### $^{86}\text{Rb}$ uptake measurements

$^{86}\text{Rb}$  uptake was performed as previously described [17]. In brief, oocytes were injected either with 8 ng ATP1AL1 cRNA or with 8 ng rat Na,K-ATPase  $\alpha 1$  subunit cRNA in combination with 1 ng of the  $\beta$  subunit of the rabbit gastric H,K-ATPase ( $\beta$ HK) or 1 ng rat  $\beta 1$  cRNA, respectively. After loading of the oocytes with  $\text{Na}^+$  (see above) and recovery in a solution containing 90 mM NaCl, 2 mM  $\text{CaCl}_2$ , 5 mM  $\text{BaCl}_2$ , 10 mM MOPS, pH 7.4, and 0.2  $\mu\text{M}$  ouabain to inhibit the endogenous oocyte Na,K-ATPase, oocytes were transferred to a solution containing 5 mM KCl, 90 mM NaCl, 1 mM  $\text{CaCl}_2$ , 5 mM  $\text{BaCl}_2$ , 1 mM  $\text{MgCl}_2$ , 10 mM Hepes,

pH 7.4, containing 10  $\mu\text{M}$  bumetanide. After addition of 5  $\mu\text{Ci/ml}$   $^{86}\text{Rb}$  (Amersham), oocytes were incubated for 12 min at room temperature with or without ouabain, before washing in a solution containing 90 mM NaCl, 1.0 mM  $\text{CaCl}_2$ , 1.0 mM  $\text{MgCl}_2$  and 10 mM Hepes, pH 7.4. Individual oocytes were dissolved in 0.5% SDS and counted.

## Results

### Inhibition by ouabain of human $\alpha\text{HKng}$ mutants

$^{86}\text{Rb}$  uptake was measured in wild type rat  $\alpha 1\beta 1$  Na,K-ATPase and in wild type and mutant human  $\alpha\text{HKng}$  coexpressed with the  $\beta$  subunit of the rabbit gastric H,K-ATPase ( $\beta\text{HK}$ ), in the absence and in the presence of ouabain. In all cases, the endogenous Na,K-ATPase was inhibited by previous exposure to 0.2  $\mu\text{M}$  ouabain. As shown in figure 1, wild type ATPases and all mutants were well expressed with a robust  $^{86}\text{Rb}$  uptake after subtraction of the uptake observed in the oocytes expressing only the  $\beta$  subunit. Two  $\alpha\text{HKng}$  mutants (K329E and D334R) showed a slightly reduced transport function to about 60% of the wild type  $\alpha\text{HKng}$ . Ouabain, at a 2 mM concentration, inhibited more than 80% of the  $^{86}\text{Rb}$  uptake in wild type rat  $\alpha 1\beta 1$  Na,K-ATPase and  $\alpha\text{HKng}/\beta\text{HK}$ , a value compatible with published values of ouabain inhibition constant ( $K_I$ ) for these enzymes. All the mutants were inhibited to a similar extent by 2 mM ouabain, except D334R for which there was no significant difference between the conditions with and without ouabain. This result indicates a very high level of resistance to ouabain inhibition, similar to that observed in the gastric H,K-ATPase. Because of the difficulty to solubilize ouabain at concentration much higher than 2 mM it would not be possible to obtain direct  $K_I$  measurement with this mutant.

### Inhibition by ouabain of Na,K-ATPase mutants

The current activated by addition of 10 mM  $\text{K}^+$  to a previously  $\text{K}^+$ -free solution ( $I_K$ ) and the current inhibited by 2 mM ouabain ( $I_{\text{ouab}}$ ) added in the presence of 10 mM  $\text{K}^+$  were measured in the wild type and mutant Na,K-ATPase. All the mutants had a good electrogenic functional expression but 2 of them (E319R and V927E) had a significantly reduced expression to about 50% and 70% of wild type, respectively. In all cases the current inhibited by 2 mM ouabain was similar to the current activated by  $\text{K}^+$ , indicating that no major change of ouabain affinity had occurred.

To assess more precisely the affinity for ouabain, the effect of progressively higher concentrations of ouabain (0, 0.5, 0.2, 0.6 and 2 mM) were applied in the presence of 10 mM  $\text{K}^+$  resulting in the inhibition curves shown in figure 2. Ouabain  $K_I$  values were similar in the wild type Na,K-ATPase and in the E314K and V927E mutants, 153, 173 and 157  $\mu\text{M}$ , respectively, while the apparent affinity for ouabain was decreased about two-fold ( $K_I$  328  $\mu\text{M}$ ) for the E319R mutant, and increased about two-fold ( $K_I$  75  $\mu\text{M}$ ) for the P785A mutant.

### Inhibition of the E319R mutant of the human Na,K-ATPase $\alpha$ subunit by ouabain and ouabagenin

The lower affinity of E319R for ouabain is consistent with the results recently reported by Qiu et al. (2005) who showed that this residue is directly involved in the structure of the ouabain binding site. These authors propose a structural model in which E319 directly interacts with the sugar moiety of ouabain through a H-bond. This hypothesis requires that ouabain is oriented in its binding site with its sugar part between the extracellular part of M4 and that of the M1–M2 hairpin. To test for a direct link between E319 and the rhamnose part of ouabain, we compared the inhibition of wild type and E319R mutant Na,K-ATPase by the glycoside ouabain and its aglycone ouabagenin. Because of resistance to ouabain and the potentially even lower affinity of the aglycone, we could not define precisely the affinity of ouabagenin for the rat  $\alpha 1$  Na,K-ATPase. We have therefore generated the E319R mutant of the human  $\alpha 1$  subunit

of Na,K-ATPase and performed these studies by measuring steady state inhibition after 1-h incubation at various concentrations of ouabain or ouabagenin, as described in the Methods section.

As shown in figure 3, there was a large difference in the apparent affinity between wild type human  $\alpha 1$  Na,K-ATPase and the E319R mutant ( $P < 0.001$ ), while there was no significant difference for ouabagenin. These results support the presence of a direct interaction between E319 and the sugar moiety of ouabain.

## Discussion

Major progress have been made recently in the characterization of the high affinity cardiac steroid binding site present at the extracellular surface of Na,K-ATPase, particularly by the use of amino acid exchange between Na,K-ATPase and the related gastric or non-gastric H,K-ATPases [11;12]. The determinants of the low affinity ouabain inhibition of the non-gastric H,K-ATPase is less well understood and was addressed in the present work.

Qiu et al. [12] have shown that the position of E319 (E312 in their numbering) position is one of the most important residues for high affinity ouabain binding of Na,K-ATPase. Here we show that mutating this glutamic acid residue in  $\alpha$ HKng to arginine (mutant D334R) renders the non gastric H,K-ATPase completely resistant to ouabain. Arginine-330 is a highly conserved residue present in all known gastric H,K-ATPase sequences at the position homologous to D334 in  $\alpha$ HKng. It seems thus likely that this sequence divergence between gastric and non-gastric H,K-ATPase is crucial to the difference in low affinity ouabain binding between these two related proteins.

In Na,K-ATPase, the E319R mutant had also a lower apparent affinity for ouabain, an observation consistent with those recently reported by Qiu et al. (2005) who showed that a similar mutant had a reduced ouabain binding and proposed that it is directly involved in the structure of the ouabain binding site and interacts with ouabain through a H-bound. The difference was however only a two-fold change in the case of Na,K-ATPase compared to a much larger change for the homologous mutation in  $\alpha$ HKng.

Two main orientations of ouabain in its binding site have been proposed. The possibility to label a cysteine in M1 with the steroid derivative HDMA (in which the reactive group is located at the position of the sugar of ouabain) suggested that the ouabain was oriented with its sugar part towards the first hairpin formed by the M1 and M2 transmembrane segments [18]. A more precise position of ouabain in its binding site has been proposed by Qiu et al. [11;12] who proposed that E319, a residue located at the extracellular end of M4, directly interacts with the rhamnose part of ouabain through a H-bound. Other authors [19] have proposed a rather opposite orientation of ouabain in its binding site with the sugar moiety pointing rather to the transmembrane segments 9 and 10. Our results show that a mutation of E319 has a large effect on the affinity for the glycoside ouabain but does not influence the binding of the corresponding aglycone supporting the hypothesis put forward by Qiu et al. [11;12].

Because P785 of  $\alpha 1$  Na,K-ATPase is located close to the cation binding site and rather distant from the putative ouabain binding site, the small effect of the P785A mutation in rat  $\alpha 1$  may be explained by an allosteric type of effect due to a change in cation affinity or conformational equilibrium.

In summary, our results confirm the importance of a charged residue in the extracellular part of the 4<sup>th</sup> transmembrane segment of  $\alpha 1$  Na,K-ATPase and show that this position is also crucial for the effect of ouabain on the human non-gastric H,K-ATPase. In addition, we provide

experimental data supporting a direct interaction of the sugar moiety of ouabain with the second extracellular loop allowing to define the orientation of the ouabain molecule in its binding site.

### Acknowledgements

This work was supported by the Swiss National Fund grant #3100A0-107513/1 to K.G. and grant #31-65441.02 to J.-D. H. and by National Institutes of Health Grant HL-36573 to N. N. M.


### Reference List

1. Axelsen KB, Palmgren MG. Evolution of substrate specificities in the P-type ATPase superfamily. *J Mol Evol* 1998;46:84–101. [PubMed: 9419228]
2. Pestov NB, Korneenko TV, Radkov R, Zhao H, Shakhparonov MI, Modyanov NN. Identification of the beta-subunit for nongastric H-K-ATPase in rat anterior prostate. *Am J Physiol Cell Physiol* 2004;286:C1229–C1237. [PubMed: 14749213]
3. Crambert G, Horisberger JD, Modyanov NN, Geering K. Human nongastric H<sup>+</sup>-K<sup>+</sup>-ATPase: transport properties of ATP1a1 assembled with different beta-subunits. *Am J Physiol Cell Physiol* 2002;283:C305–C314. [PubMed: 12055100]
4. Lingrel JB, Kuntzweiler TA. Na<sup>+</sup>,K<sup>+</sup>-ATPase. *J Biol Chem* 1994;269:19659–19662. [PubMed: 8051040]
5. Nicholls MG, Richards AM, Lewis LK, Yandle TG. Welcome to ouabain--A new steroid hormone. *Lancet* 1991;338:543–544. [PubMed: 1678804]
6. Nicholls MG, Richards AM, Lewis LK, Yandle TG. Ouabain - a new steroid hormone. *Lancet* 1995;346:1381–1382. [PubMed: 7475817]
7. Blaustein MP. Endogenous ouabain - role in the pathogenesis of hypertension. *Kidney Int* 1996;49:1748–1753. [PubMed: 8743490]
8. Xie ZJ, Askari A. Na<sup>+</sup>/K<sup>+</sup>-ATPase as a signal transducer. *European Journal of Biochemistry* 2002;269:2434–2439. [PubMed: 12027880]
9. Stenkvist B. Cardenolides and cancer. *Anti-Cancer Drugs* 2001;12:635–636. [PubMed: 11487722]
10. Lin H, Juang JL, Wang PS. Involvement of Cdk5/p25 in digoxin-triggered prostate cancer cell apoptosis. *J Biol Chem* 2004;279:29302–29307. [PubMed: 15123618]
11. Qiu LY, Krieger E, Schaftenaar G, Swarts HGP, Willems PHGM, De Pont JJHH, Koenderink JB. Reconstruction of the complete ouabain-binding pocket of Na,K-ATPase in gastric H,K-ATPase by substitution of only seven amino acids. *J Biol Chem* 2005;280:32349–32355. [PubMed: 16051601]
12. Qiu LY, Swarts HGP, Tonk ECM, Willems PHGM, Koenderink JB, De Pont JJHH. Conversion of the low affinity ouabain-binding site of non-gastric H,K-ATPase into a high affinity binding site by substitution of only five amino acids. *J Biol Chem* 2006;281:13533–13539. [PubMed: 16531406]
13. Price EM, Rice DA, Lingrel JB. Structure-function studies of Na,K-ATPase. Site-directed mutagenesis of the border residues from the H1–H2 extracellular domain of the  $\alpha$  subunit. *J Biol Chem* 1990;265:6638–6641. [PubMed: 2157705]
14. Jaunin P, Horisberger JD, Richter K, Good PJ, Rossier BC, Geering K. Processing, intracellular transport and functional expression of endogenous and exogenous  $\alpha$ - $\beta$ 3 Na,K-ATPase complexes in *Xenopus* oocytes. *J Biol Chem* 1992;267:577–585. [PubMed: 1309755]
15. Burnay M, Crambert G, Kharoubi-Hess S, Geering K, Horisberger JD. Electrogenicity of Na,K- and H,K-ATPase activity and presence of a positively charged amino acid in the fifth transmembrane segment. *J Biol Chem* 2003;278:19237–19244. [PubMed: 12637496]
16. Horisberger JD, Kharoubi-Hess S. Functional differences between  $\alpha$  subunit isoforms of the rat Na,K-ATPase expressed in *Xenopus* oocytes. *J Physiol(London)* 2002;539:669–680. [PubMed: 11897839]
17. Beggah AT, Jaunin P, Geering K. Role of glycosylation and disulfide bond formation in the  $\beta$  subunit in the folding and functional expression of Na,K-ATPase. *J Biol Chem* 1997;272:10318–10326. [PubMed: 9092584]
18. Antolovic R, Schoner W, Geering K, Canessa CM, Rossier BC, Horisberger J-D. Labeling of a cysteine in the cardiotonic glycoside binding site by the steroid derivative HDMA. *FEBS Lett* 1995;368:169–17. [PubMed: 7615075]

19. Keenan SM, Delisle RK, Welsh WJ, Paula S, Ball WJ. Elucidation of the Na<sup>+</sup>, K<sup>+</sup>-ATPase digitalis binding site. *Journal of Molecular Graphics & Modelling* 2005;23:465–475. [PubMed: 15886034]


## Abbreviations

<b><math>\alpha</math>HKng</b>	$\alpha$ subunit of the non gastric H,K-ATPase
<b><math>\alpha</math>HKg</b>	$\alpha$ subunit of the gastric H,K-ATPase
<b>M4, M5</b>	4 <sup>th</sup> , 5 <sup>th</sup> transmembrane segments




**Figure 1.**

Transport function of wild type and mutant  $\alpha$ HKng expressed in *Xenopus* oocytes.  $^{86}\text{Rb}$  uptake was measured as described in the text after inhibition of the endogenous oocyte Na,K-ATPase by overnight exposure to 0.2  $\mu\text{M}$  ouabain. White and black columns represent the uptake observed in the absence and in the presence of 2 mM ouabain, respectively. The number of measurements is between 18 and 22 in each group.



**Figure 2.** Steady state ouabain concentration/inhibition curves of the  $K^+$  activated outward current in oocytes expressing wild type and mutant rat  $\alpha 1/\beta 1$  Na,K-ATPase. Exposure to ouabain was performed in the presence of 10 mM  $K^+$  extracellular concentrations. Solid lines show the best fitting normalized current ( $I_{norm}$ ) curves to the equation

$$I_{norm} = 1 - 1 / \left( 1 + \frac{K_I}{[ouabain]} \right)$$



**Figure 3.** Ouabain and ouabagenin inhibition constants ( $K_I$ ) in wild type and E319R mutant  $\alpha 1/\beta 1$  human Na,K-ATPase. The results are the mean of 9 to 11 measurements in each group.

**Table 1**

Alignment of human  $\alpha 1$  Na,K-ATPase,  $\alpha$  non-gastric H,K-ATPase and  $\alpha$  gastric H,K-ATPase sequences in the regions concerned by the mutations. On the right are shown the mutants generated and studied in this work.

[[wild type sequences]]	[[[]]]	[[mutants]]
[[== M4 ==]]	[[[]]]	
$\alpha 1$ NaK SLILE <sub>314</sub> YTWLE <sub>319</sub> AVIF	[[ $\alpha 1$ NaK]]	<b>E<sub>314</sub>K E<sub>319</sub>R</b>
$\alpha$ HKng AVSLK <sub>329</sub> YQVLD <sub>334</sub> SIIF	[[ $\alpha$ HKng]]	<b>K<sub>329</sub>E K<sub>329</sub>G D<sub>334</sub>R</b>
$\alpha$ HKg AMCIG <sub>325</sub> YTFLR <sub>330</sub> AMVF	[[[]]]	
[[===== M5 =====]]	[[[]]]	
$\alpha 1$ NaK LTSNIP <sub>785</sub> EIT <sub>788</sub> PFLI	[[ $\alpha 1$ NaK]]	<b>P<sub>785</sub>A</b>
$\alpha$ HKng LTKNIA <sub>800</sub> ELC <sub>803</sub> PFLI	[[ $\alpha$ HKng]]	<b>A<sub>800</sub>P</b>
$\alpha$ HKg LTKNIP <sub>796</sub> ELT <sub>799</sub> PYLI	[[[]]]	
[[===== M8 =====]]	[[[]]]	
$\alpha 1$ NaK AFFVSIV <sub>927</sub> VVQWAD	[[ $\alpha 1$ NaK]]	<b>V<sub>927</sub>E</b>
$\alpha$ HKng AFFVGI <sub>942</sub> VQIAD	[[ $\alpha$ HKng]]	<b>L<sub>942</sub>E</b>
$\alpha$ HKg VFFISIE <sub>938</sub> VCQIAD	[[[]]]	