

## RESEARCH PAPER

# Managing argatroban in heparin-induced thrombocytopenia: A retrospective analysis of 729 treatment days in 32 patients with confirmed heparin-induced thrombocytopenia

Matteo Marchetti<sup>1,2</sup>  | Stefano Barelli<sup>1</sup>  | Tobias Gleich<sup>3</sup>  | Francisco J. Gomez<sup>1</sup>  | Matthew Goodyer<sup>4</sup>  | Francesco Grandoni<sup>1</sup>  | Lorenzo Alberio<sup>1,2</sup> 

<sup>1</sup>Division of Hematology and Central Laboratory of Hematology, CHUV, Lausanne University Hospital, Lausanne, Switzerland

<sup>2</sup>Faculty of Biology and Medicine, UNIL, University of Lausanne, Lausanne, Switzerland

<sup>3</sup>Division of Transfusion Medicine and Immunohematology, CHUV, Lausanne University Hospital, Lausanne, Switzerland

<sup>4</sup>Division of Hematology and Laboratory of Hematology, Institut Central des Hôpitaux Valaisans, Sion, Switzerland

## Correspondence

Lorenzo Alberio, Service et Laboratoire Central d'hématologie, CHUV Centre Hospitalier Universitaire Vaudois, Rue du Bugnon 46, CH – 1011 Lausanne, Switzerland. Email: [lorenzo.alberio@chuv.ch](mailto:lorenzo.alberio@chuv.ch)

## Summary

Argatroban is a first-line anticoagulant for patients with heparin-induced thrombocytopenia (HIT). Published data on practical aspects of its use in HIT are lacking. We aim to establish recommendations based on our experience. This cohort of 32 patients is the largest describing cases of HIT confirmed by a functional assay treated with argatroban. Among patients with normal liver function, median starting argatroban doses (SAD) of 0.54, 0.98, and 1.27 µg/kg/min reached steady-state plasmatic argatroban concentrations (PAC) of 0–0.39, 0.40–0.99, and 1.00–1.5 µg/ml, respectively. Median argatroban dose increases (ADI) induced similar median steady-state PAC increases ( $\Delta$  µg/kg/min  $\approx$   $\Delta$  µg/ml). PAC measurements performed more than 240 min after SAD or ADI were significantly higher compared to earlier controls. Quantitative PAC measurements and thrombin time (TT) appeared adequate for monitoring. Thirty-eight percent of the thrombotic events were preceded by PAC below 0.4 µg/ml. Four hours after argatroban discontinuation, median international normalised ratio (INR) decrease was –1.2. We suggest: (i) monitoring argatroban with PAC or TT at least 240 min after SAD and/or AID; (ii) using SAD of 1.0 µg/kg/min and ADI of at least 0.2 µg/kg/min when liver function is normal; (iii) targeting therapeutic PAC of 0.5–1.0 µg/ml; and (iv) targeting INR of 3.5–4.5 when bridging argatroban with vitamin K antagonists.

## KEYWORDS

anticoagulation, argatroban, argatroban monitoring, heparin-induced thrombocytopenia, plasmatic argatroban concentration

## INTRODUCTION

Heparin-induced thrombocytopenia (HIT) is an adverse effect of unfractionated (UFH) and low-molecular-weight heparin (LMWH), occurring with variable risk depending on the type and dose of heparin and the clinical context.<sup>1–4</sup> HIT is characterised by a severe limb- and life-threatening

pro-thrombotic state requiring immediate cessation of all heparin and a switch to an alternative non-heparin anticoagulant.<sup>5–9</sup>

Argatroban is an effective first-line therapy for HIT<sup>5,10</sup> and was described in a recent meta-analysis as being superior to other parenteral drugs used for the management of HIT.<sup>11</sup> Argatroban is a synthetic, reversible and competitive

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *British Journal of Haematology* published by British Society for Haematology and John Wiley & Sons Ltd.

direct thrombin inhibitor, which binds thrombin on its catalytic site<sup>12</sup> and is hepatically metabolised.<sup>13</sup>

Less is known, however, about several practical aspects of anticoagulation with argatroban. The magnitude of starting doses and dose adaptations, as well as the degree of dose reduction required among patients with different types of hepatic dysfunction remain unclear and are performed empirically. Furthermore, despite recent guidelines,<sup>10</sup> the best timing of, and laboratory assays for, argatroban monitoring still remain open to discussion. The aim of our study was to investigate how argatroban therapy is conducted at our institution and to develop evidence-based in-house recommendations for an optimal anticoagulation with argatroban in patients with HIT.

## METHODS

### Study design

This was a retrospective observational derivation study. Data were collected between October 2018 and February 2019 and analysed between February 2019 and August 2021.

### Patient cohort

Between August 2014 and February 2019, 32 patients with confirmed HIT<sup>14</sup> [positive anti-PF4/heparin antibodies by


HemosIL-Acustar-HIT-IgG (Instrumentation Laboratory GmbH, Munich, Germany) (a chemiluminescent immunoassay, CLIA) and ID-H/PF4-PaGIA (Bio-Rad, DiaMed SA, Basel, Switzerland) (a particle gel immunoassay, PaGIA) and positive heparin-induced platelet aggregation (HIPA, University Hospital, Greifswald, Germany)] received argatroban at our institution and were included in this study.

### Laboratory monitoring of argatroban


We performed the following assays on an automated coagulometer (Sysmex CS-5100; Siemens Healthineers, Erlangen, Germany): (i) activated partial thromboplastin time (aPTT; Pathromtin SL®; Siemens Healthineers, Erlangen, Germany); (ii) thrombin time (TT; Thromboclotin®, Siemens Healthineers, Erlangen, Germany) with a final thrombin concentration of 1.25 U/ml; and (iii) assessment of the plasma argatroban concentration by a commercial diluted TT assay (Hemoclot Thrombin Inhibitors; Hyphen Biomed, Neuville-sur-Oise, France).

### Definition of patients with impaired hepatic function

As argatroban is mainly metabolised in the liver, we divided the studied patients into four categories according to their



**FIGURE 1** Laboratory parameters. The figure depicts daily platelet counts (G/L, grey), D-dimers (ng/ml, blue) and haemoglobin levels (g/l, red) before and under argatroban therapy in 32 heparin-induced thrombocytopenia (HIT) patients. Points represent median daily values; error bars represent SEM daily values. Note that day 0 is the first day with argatroban treatment [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com)]




**FIGURE 2** Plasma argatroban concentrations measured after different argatroban starting doses (SAD). (A) Intended SAD of 0.5 µg/kg/min in patients with normal liver function. (B) Intended SAD of 1.0 µg/kg/min in patients with normal liver function

estimated liver function. (i) Normal liver function: serum alanine aminotransferase (ALT) levels less than 180 U/l [three times the upper limit of normal (ULN)] and serum total bilirubin levels less than 25.5 µmol/l.<sup>13</sup> When a patient previously had ALT higher than 180 U/l and/or serum total bilirubin higher than 25.5 µmol/l and recovered, he was included into the 'normal liver function group' five days after recovery of ALT less than 180 U/l and serum total bilirubin less than 25.5 µmol/l. (ii) Liver cytolysis: ALT higher than 180 U/l and serum total bilirubin less than 25.5 µmol/l. (iii) Impaired bilirubin excretion: ALT less than 180 U/l and serum total

bilirubin higher than 25.5 µmol/l. (iv) Liver cytolysis associated with impaired bilirubin excretion: ALT higher than 180 U/l and serum total bilirubin higher than 25.5 µmol/l.

### Statistics


Median and interquartile range (IQR) values were calculated with Excel (Microsoft Schweiz GmbH, Zürich-Flughafen, Switzerland). We employed MedCalc (version 15.11.0; MedCalc Software Ltd, Ostend, Belgium) to perform



**FIGURE 3** Effect of magnitude of dose increase and time of monitoring on plasma argatroban concentration in patients with normal liver function. (A) Plasmatic argatroban concentrations (PAC, µg/ml) measured 3–6 h after different magnitudes of argatroban dose increase (ADI, µg/kg/min). We noted significant plasmatic argatroban concentration (PAC) increases with a median of 0.06 µg/ml [interquartile range (IQR) 0–0.17] after argatroban dose increase (ADI) of 0.01–0.10 µg/kg/ml ( $p$ -value = 0.0080), 0.15 µg/ml (IQR 0.07–0.26) after ADI of 0.11–0.20 µg/kg/ml ( $p$ -value = 0.0016), and 0.21 µg/ml (IQR 0.15–0.28) after ADI of 0.21–0.30 µg/kg/ml ( $p$ -value < 0.0001). Significant differences in median PAC increases were observed between ADI of 0.01–0.10 versus 0.21–0.30 µg/kg/min ( $p$ -value = 0.0007) and ADI of 0.01–0.10 versus 0.31–0.40 µg/kg/min ( $p$ -value = 0.047). (B) Plasmatic argatroban concentrations (PAC, µg/ml) measured before or after 3 h after the same magnitude of ADI (µg/kg/min). Median PAC increase in the group with controls performed 3–6 h (median, 240 min) after ADI was 0.19 µg/ml [interquartile range (IQR) 0.14–0.27] and was significantly higher than the median PAC increase in the group with control within 3 h (median, 128 min) after ADI (median, 0.08 µg/ml, IQR 0.04–0.12,  $p$ -value = 0.009)

Passing–Bablock regression analysis. Sigma Plot (version 13.0; Systat Software GmbH, Erkrath, Germany) was used to create Figure 1 and GraphPad Prism (version 8; GraphPad Software, San Diego, USA) was used to create Figures 2–8 and

to perform Mann–Whitney tests ( $t$ -test, unpaired values, non-parametric distribution), Wilcoxon tests ( $t$ -test, paired values, non-parametric distribution) and Spearman correlation measures.



**FIGURE 4** Effect of liver function on argatroban steady-state doses and plasma concentration. Argatroban doses at steady state and different plasma concentrations in patients with different patterns of liver function. (A) Prophylactic plasmatic argatroban concentrations (PAC) (0–0.39 µg/ml). (B) Standard therapeutic PAC (0.40–0.99 µg/ml). (C) High therapeutic PAC (1.00–1.5 µg/ml)

## Ethics

The study complies with the guidelines of the Institutional Ethical Board (Commission Cantonale Vaudoise d’Ethique de la Recherche sur l’Être Humain, CER-VD, protocol number 497/95) and was accepted for quality control assessment of laboratory and clinical management practice.

## RESULTS


### Patient population

Thirty-two patients with confirmed HIT (15 of them presenting with HIT and thrombosis, HIT-T) were treated with

argatroban. Twenty-seven patients (84%) only received argatroban and five (16%) received multiple treatments (Table 1). Overall, we studied 729 argatroban treatment days. Median duration of argatroban therapy was 17.5 days (IQR 10–29.5).

### Complications

Compared to the cohort published by Tardy-Poncet *et al.*,<sup>15</sup> we registered a higher rate of death related to HIT ( $n = 1/32$  vs.  $0/16$ ), a similar rate of thromboembolic complications ( $n = 13/32$ , 41% vs.  $n = 7/16$ , 44%), and less major bleeding ( $n = 4/32$ , 13% vs.  $n = 3/16$ , 19%) and overall death rates ( $n = 4/32$ , 13% vs.  $n = 4/16$ , 25%). Individual clinical courses are detailed in Table 2.



**FIGURE 5** Serum total bilirubin levels, steady-state argatroban doses and plasma concentration. Serum total bilirubin levels and steady-state argatroban doses among patients that had a ‘prophylactic’, ‘standard therapeutic’, and ‘high therapeutic’ plasmatic argatroban concentrations (PAC). (A) Serum total bilirubin levels and argatroban dose at steady state among patients that had a ‘prophylactic’ PAC (0–0.39 µg/ml). (B) Serum total bilirubin levels and argatroban dose at steady state among patients that had a ‘standard therapeutic’ PAC (0.4–0.99 µg/ml). (C) Serum total bilirubin levels and argatroban dose at steady state among patients that had a ‘high therapeutic’ PAC (≥1.0 µg/ml)


We observed 13 thromboembolic events among nine different patients (Table 3). Seven were venous and six were arterial thromboses. Five out of 13 (38%) thromboembolic events were related to an anticoagulation that was probably inadequate. Indeed, PAC had been below 0.4 µg/ml in the 24 h preceding diagnosis in four cases and three days before diagnosis in one patient. Additionally, other risk factors were present: recent surgery and immobilisation ( $n = 11$ ), systemic infection ( $n = 8$ ), active cancer ( $n = 6$ ), body mass index (BMI) higher than 30 kg/m<sup>2</sup> ( $n = 2$ ), intravascular devices ( $n = 5$ ), and/or poor haemodynamic conditions ( $n = 4$ ).

Concerning the five bleeding events (Table 4), four were considered as major according to the International Society for Thrombosis and Haemostasis (ISTH) definitions.<sup>16,17</sup> PAC measured in the 24 h preceding these bleeding events were within target ranges in 4/5 patients. Among the major bleeding events, two intracranial haemorrhages occurred

after ischaemic strokes and were considered as secondary to anatomical brain lesions. The other two major bleeding events occurred under concomitant double antiplatelet therapy.

### Outcomes

Four (13%) patients underwent limb amputation and four patients died (Table 2). Among patients that underwent amputation, three had established advanced peripheral arterial disease. The fourth patient suffered acute hepatic necrosis with ischaemic limb necrosis<sup>18</sup> prior to HIT onset and underwent amputation during argatroban therapy (unique patient number, UPN 15.103). Among the four deceased patients, three were still under argatroban therapy at the time of death. In one patient (UPN 16.193), death was directly related to HIT. In two cases, advanced comorbidities may



**FIGURE 6** Argatroban monitoring with routine coagulation assays. The figure depicts thrombin time (TT, s) and activated partial thromboplastin time (aPTT, s) intervals with the corresponding median plasmatic argatroban concentrations (PAC). (A) TT. (B) aPTT. Error bars represent the 90th (upper) and 10th (lower) percentiles

have contributed to the fatal outcome (UPN 18.148 and UPN 15.001).


### Evolution of laboratory parameters under argatroban therapy

As depicted in Figure 1, we recorded a median platelet count nadir of 51 G/l on the second day of argatroban therapy, which was followed by full recovery (>150 G/l) on the seventh

to eighth day of treatment. In parallel, median D-dimers levels progressively decreased.

### Argatroban starting dose

Every starting argatroban dose (SAD, µg/kg/min) and the corresponding PAC measurements (µg/ml) at time  $t'$  (min) after SAD were analysed. Among patients with normal liver function, we divided all recorded SAD into two categories,



**FIGURE 7** Argatroban clearance from plasma. Argatroban plasma concentrations measured after its discontinuation are depicted as percentage of the last steady state concentration. Panel A: Patients with normal liver function

according to an intended SAD of 0.5 or 1.0  $\mu\text{g}/\text{kg}/\text{min}$ , respectively.

As illustrated in [Figure 2A](#), after an overall median SAD of 0.5  $\mu\text{g}/\text{kg}/\text{min}$  (IQR 0.48–0.58), median PAC among controls performed within 240 min reached 0.26  $\mu\text{g}/\text{ml}$  (IQR 0.20–0.35) vs. 0.44  $\mu\text{g}/\text{ml}$  (IQR 0.29–0.53) in controls at least 240 min after argatroban initiation ( $p$ -value = 0.0275). Of note, median SAD administered in patients with monitoring performed after at least 240 min (0.53  $\mu\text{g}/\text{kg}/\text{min}$ ) did not differ significantly from SAD applied in controls less than 240 min (0.50  $\mu\text{g}/\text{kg}/\text{min}$ ;  $p$ -value = 0.1594).

[Figure 2B](#) shows data for an intended SAD of 1  $\mu\text{g}/\text{kg}/\text{min}$  (median, 1.04, IQR 0.94–1.09). This led to median PAC of 0.30  $\mu\text{g}/\text{ml}$  (IQR 0.14–0.63) among controls less than 240 min versus 0.77  $\mu\text{g}/\text{ml}$  (IQR 0.74–0.97) in controls at least 240 min after argatroban initiation, respectively ( $p$ -value = 0.0749). Again, median SAD administered in patients with monitoring after at least 240 min (0.98  $\mu\text{g}/\text{kg}/\text{min}$ ) did not significantly differ from SAD applied in controls after less than 240 min (1.06  $\mu\text{g}/\text{kg}/\text{min}$ ,  $p$ -value >0.9999).

### Magnitude and effect of argatroban dose adaptations on plasma concentration

The most frequent dose adaptations in our cohort were ADI motivated by clinico-biological features, such as monitoring


parameters below target ranges, insufficient improvement of platelet count and/or D-dimers, and thromboembolic events. Among patients with normal liver function, recorded ADI were divided into four intervals (+0.01–0.10, 0.11–0.20, 0.21–0.30, and 0.31–0.40  $\mu\text{g}/\text{kg}/\text{min}$ ). [Figure 3A](#) depicts PAC measured 3–6 h after ADI (median: 240 min, IQR: 217–297), showing that median PAC increases ( $\mu\text{g}/\text{ml}$ ) corresponded roughly to the respective ADI ( $\mu\text{g}/\text{kg}/\text{min}$ ).

### Effect of timing of monitoring on plasma argatroban concentration

Among patients with normal liver function, ADI within a magnitude of 0.11–0.40  $\mu\text{g}/\text{kg}/\text{min}$  were divided into two groups, based on the timing of monitoring: within 3 h (median, 128 min) or between 3 and 6 h (median, 240 min) after ADI. [Figure 3B](#) shows that late monitoring was associated with significantly higher PAC compared to early monitoring, despite the fact that, paradoxically, the median ADI performed in the group with PAC control within 3 h was significantly higher compared to the group with PAC control performed after 3–6 h (median, 0.22 vs. 0.21  $\mu\text{g}/\text{kg}/\text{min}$ , IQR 0.21–0.26 vs. 0.14–0.22,  $p$ -value = 0.048).

### Argatroban dosing adapted to liver function

Steady-state PAC was defined as no argatroban dose modification in the 6 h preceding monitoring. Patients were



**FIGURE 8** Impact of argatroban on the international normalized ratio (INR). (A) INR under vitamin K antagonists (VKA) before and 3 to less than 6 h after argatroban discontinuation. (B) Median INR decrease after argatroban discontinuation. Long bar represents median, short bars represent 95% confidence intervals

divided into four categories according to their estimated liver function. For each group, we created PAC intervals according to the treatment aim: (i) 'prophylactic' PAC, 0–0.39  $\mu\text{g}/\text{ml}$  (Figure 4A); (ii) 'standard therapeutic' PAC, 0.40–0.99  $\mu\text{g}/\text{ml}$  (Figure 4B); and (iii) 'high therapeutic' PAC, 1.00–1.5  $\mu\text{g}/\text{ml}$  (Figure 4C). At steady state, patients

with normal liver function required median (IQR) argatroban doses of 0.54  $\mu\text{g}/\text{kg}/\text{min}$  (0.47–1.0), 0.98  $\mu\text{g}/\text{kg}/\text{min}$  (0.61–1.32), and 1.27  $\mu\text{g}/\text{kg}/\text{min}$  (0.90–1.45) to reach prophylactic, standard therapeutic, or high therapeutic PAC, respectively. Patients with liver cytolysis required argatroban doses of 0.25  $\mu\text{g}/\text{kg}/\text{min}$  (0.24–0.29), 0.78  $\mu\text{g}/\text{kg}/\text{min}$

**TABLE 1** Epidemiologic and demographic features of the 32 heparin-induced thrombocytopenia (HIT) patients

Feature	
Females, % (n/N)	47 (15/32)
Age, years; median (IQR; range)	67.4 (58.3–75.1; 33.5–88.9)
Hospital ward	
Cardiac surgery	11 <sup>a</sup>
Vascular surgery	6 <sup>a</sup>
Orthopaedics and traumatology surgery	0
Thoracic surgery	4
Neurosurgery	1
Intensive care unit	4
Internal medicine	2
Cardiology	5
Heparin treatment and regimen	
LMWH prophylactic	9
LMWH therapeutic	4
UFH prophylactic	25
UFH therapeutic	20
HIT diagnosis features	
4 T score; median (IQR; range)	5 (5–5; 3–7)
CLIA, U/ml; median (IQR; range)	8.64 (1.96–21.69; 0.15–129)
PaGIA, titre; median (IQR; range)	8 (8–32; 2–64)
HIPA, number of positive	32
Type of HIT	
HIT without thrombosis, n (%)	17 (53)
HIT-T, n (%)	15 (47)
Treatments	
Argatroban only	27
Danaparoid prior to argatroban	3 (UPN 17.173, UPN 17.079, UPN 15.103)
IVIG	3 (UPN 17.173, UPN 17.042 <sup>b</sup> , UPN 17.222)
Plasmapheresis	1 (UPN 17.042 <sup>b</sup> )
Argatroban treatment, days	729
Median duration (days)	17.5
IQR duration (days)	10; 29.5
Range duration (days, min; max)	3; 109
Complications under argatroban	
Arterial thromboembolic events, n (%)	6 (19)
Venous thromboembolic events, n (%)	7 (25)
Bleeding events, minor, n (%)	1 (3)
Bleeding events, major, n (%)	4 (13)

(Continues)

(0.53–1.10), and 1.08 µg/kg/min (0.69–1.08) respectively, and patients with hyperbilirubinaemia, argatroban doses of 0.12 µg/kg/min (0.11–0.16), 0.16 µg/kg/min (0.12–0.18), and 0.51 µg/kg/min (0.10–0.51).

**TABLE 1** (Continued)

Feature	
Outcomes	See also <a href="#">Tables 2–4</a>
Limb amputation, n (%)	4 (13)
In-hospital mortality, n (%)	4 (13)

Abbreviations: CLIA, chemiluminescent immunoassay; HIPA, heparin-induced platelet aggregation; HIT, heparin-induced thrombocytopenia; IQR, interquartile range; IVIG, intravenous immunoglobulin; LMWH, low-molecular-weight heparin; PaGIA, particle gel immunoassay; UFH, unfractionated heparin.

<sup>a</sup>One patient underwent two operations (cardiac and vascular surgery) on the same day.

<sup>b</sup>One patient received plasmapheresis and then IVIG after argatroban discontinuation.

Median argatroban doses were significantly lower in patients with hyperbilirubinaemia compared to patients with normal liver function or liver cytolysis and this was irrespective of the PAC interval. Moreover, among patients with liver cytolysis, median argatroban doses were significantly lower compared to those in patients with normal liver function in the ‘prophylactic’ and ‘high therapeutic’ PAC intervals.

### Total bilirubin levels and argatroban doses

Figure 5 depicts argatroban doses at steady state and concomitant bilirubin level, according to the targeted PAC. Among patients that had ‘prophylactic’ (0–0.39 µg/ml; Figure 5A) and ‘standard therapeutic’ (0.40–0.99 µg/ml; Figure 5B) PAC, serum total bilirubin levels and argatroban doses at steady state were strongly inversely correlated [Spearman *r* (95% confidence interval (CI)): –0.8203 (–0.9156; –0.6376), *p*-value: <0.0001, *n* = 28 pairs and –0.8728 (–0.9185; –0.8041), *p*-value: <0.0001, *n* = 77 pairs, respectively]. Inverse correlation was lower among patients that had a ‘high therapeutic’ (≥1.0 µg/ml; Figure 5C) PAC [Spearman *r* (95% CI): –0.5334 (–0.7644; –0.1809), approximate *p*-value: 0.0042, *n* = 27 pairs].

### Monitoring of argatroban plasma concentration with routine assays

Three different assays for argatroban anticoagulation monitoring were available: aPTT (s), TT (s), and PAC (µg/ml). PAC and corresponding simultaneous TT and aPTT were analysed. PAC and corresponding single TT values showed the highest Spearman correlation [*p* (95% CI): 0.726 (0.689–0.7599); *n* = 749 pairs], while aPTT and simultaneous PAC showed a lower correlation [*p* (95% CI): 0.590 (0.548–0.630); *n* = 1031 pairs]. As shown in Figure 6A, increasing TT intervals showed a linear trend, with 80% of TT values higher than 79 s reaching at least a therapeutic PAC (>0.4 µg/ml). Of note, while aPTT intervals within 30–69 s showed a linear trend with the corresponding PAC, aPTT intervals higher than 70 s plateaued (Figure 6B).

**TABLE 2** Patients' individual clinical course

	Sex	Age	Care unit; underlying main disease(s) and management at our institution	Type of heparin and dose	Platelet count nadir value (G/l)	4T score	HIT versus HIT-T	HIT-T: Type of thrombo-embolic event at diagnosis	Diagnostic work-up
1. UPN: 18.130	F	63	Thoracic surgery; idiopathic pulmonary fibrosis and bilateral lung transplant	Prophylactic UFH, then prophylactic LMWH	51	5	HIT-T	Pulmonary embolism and internal jugular vein thrombosis (secondary to catheter & HIT)	Chest CT scan with contrast product ultrasonography by an angiologist
2. UPN: 18.022	F	52	Cardiac surgery; pheochromocytoma and right adrenalectomy vena cava and right atrium thrombi thrombectomies	Therapeutic UFH, then therapeutic LMWH	111	5	HIT	None	N/A
3. UPN: 17.234	M	69	Cardiac and vascular surgery; ischaemic heart disease with triple coronary bypass; left carotid stenosis with carotid endarterectomy (two simultaneous operations)	Prophylactic UFH, then prophylactic LMWH	48	4	HIT	None	N/A
4. UPN: 17.222	M	72	Cardiac surgery; severe aortic valve stenosis with bioprothetic aortic replacement valve	Prophylactic UFH, then prophylactic LMWH	18	5	HIT	None	N/A
5. UPN: 17.206	M	89	Cardiac surgery; ischaemic heart disease with triple coronary bypass; severe aortic valve stenosis and bioprothetic aortic replacement valve	Prophylactic UFH, then prophylactic LMWH, then prophylactic UFH	54	5	HIT-T	Fibular artery thrombosis	Ultrasonography and plethysmography by an angiologist
6. UPN: 17.173	F	29	Internal medicine; systemic lupus erythematosus with antiphospholipid antibodies syndrome and heparin anticoagulation	Prophylactic UFH, then therapeutic UFH, then therapeutic LMWH	5	5	HIT	None	N/A
7. UPN: 17.081	F	75	Cardiology; acute heart failure and anticoagulation for mechanical mitral valve	Therapeutic UFH, then prophylactic UFH	79	4	HIT	None	N/A
8. UPN: 17.079	M	68	Intensive care unit; septic shock secondary to Balthazar E pancreatitis	Prophylactic UFH	16	4	HIT	None	N/A
9. UPN: 18.148	M	89	Vascular surgery; bilateral advanced arterial peripheral artery disease with two arterio-arterial bypasses, one Fogarty thrombo-embolectomy and one thrombolysis	Therapeutic UFH	49	6	HIT-T	External iliac artery and commune femoral artery thrombosis	Ultrasonography and plethysmography by an angiologist; arterial CT scan with contrast product
10. UPN: 17.042	M	75	Cardiology; ischaemic heart disease with cardiogenic shock and angioplasty with stenting	Therapeutic and then prophylactic UFH	30	4	HIT-T	Internal jugular vein thrombosis (secondary to catheter & HIT)	Ultrasonography by an angiologist

CLIA (U/ml)	PaGIA (titre)	Type of treatment and duration (days)	Thrombo-embolic events during treatment YES/NO *see Table 3	Bleeding events YES/NO *see Table 4	Limb outcome: Amputation and day of Argatroban treatment	Final outcome: Discharge xxx days after diagnosis of HIT versus hospital transfer
7.03	8	Argatroban; 11 days Central venous catheter (jugular) already removed at time of HIT diagnosis	No	No	No amputation	Hospital transfer 11 days after diagnosis of HIT
11.70	16	Argatroban; 10 days	YES: 1* (SVT LLL)	NO	No amputation	Discharged 14 days after diagnosis of HIT
28.00	>16	Argatroban; 18 days	NO	NO	No amputation	Discharged 24 days after diagnosis of HIT
12.56	>16	Argatroban; 18 days IVIG	NO	YES; 1* (minor bleeding; spontaneous haematoms on the arms and forearms)*	No amputation	Hospital transfer 18 days after diagnosis of HIT
70.88	16	Argatroban; 14 days	NO	NO	No amputation	Discharged 23 days after diagnosis of HIT
1.40	16	Danaparoid; 2 days then argatroban; 19 days IVIG	NO	NO	No amputation	Discharged 22 days after diagnosis of HIT
10.24	8	Argatroban; 24 days	NO	NO	No amputation	Discharged 32 days after HIT diagnosis
12.42	>32	Danaparoid; 8 days argatroban; 47 days	NO	NO	No amputation	Discharged 104 days after HIT diagnosis
>128	8	Argatroban; 44 days; prosthetic bypass confection; lower left limb	YES: 4* (1× arterial prosthetic bypass: 1× PE 1× DVT 1× arterial; lower left proximal limb)	NO	Gritti amputation of right lower limb 16 days after HIT diagnosis	Deceased 44 days after HIT diagnosis (pneumonia with cardiac failure)
2.14	004	Argatroban; 28 days, IVIG plasmapheresis because of discontinuation of argatroban (haemorrhagic shock)	NO	YES; 1* (major bleeding; haemorrhagic shock after self-snatch of urinary catheter in the context of confusional state)	No amputation	Discharged 69 days after HIT diagnosis

**TABLE 2** (Continued)

	Sex	Age	Care unit; underlying main disease(s) and management at our institution	Type of heparin and dose	Platelet count nadir value (G/l)	4T score	HIT versus HIT-T	HIT-T: Type of thrombo-embolic event at diagnosis	Diagnostic work-up
11. UPN: 16.269	F	73	Cardiac surgery; ischaemic heart disease, ascendant aorta aneurysm with moderate aortic valve regurgitation. Triple coronary bypass, bioprosthetic aortic replacement valve and ascendant aorta replacement	Therapeutic and then prophylactic UFH	62	5	HIT	None	N/A
12. UPN: 16.193	M	64	Thoracic surgery; mediastinal sarcoma with vena cava superior infiltration and tumour resection	Prophylactic UFH, then therapeutic UFH and then prophylactic LMWH	75	5	HIT-T	Bilateral pulmonary embolism	Chest CT scan with contrast product
13. UPN: 16.199	M	59	Thoracic surgery; empyema and thoracoscopy with pulmonary decortication	Prophylactic LMWH	134	5	HIT-T	Pulmonary embolism	Chest CT scan with contrast product
14. UPN: 16.151	M	81	Cardiac surgery; severe aortic valve stenosis and bioprosthetic aortic replacement valve	Therapeutic LMWH and then prophylactic UFH	71	5	HIT	None	N/A
15. UPN: 16.225	M	71	Vascular surgery; abdominal aorta aneurysm and vascular endoprosthesis	Prophylactic UFH	32	5	HIT	None	N/A
16. UPN: 16.351	F	48	Intensive care unit; second-degree burns on 14% of body surface area, including sub-glottic burn	Prophylactic LMWH and then therapeutic UFH (HIT-T)	79	6	HIT-T	Deep vein and superficial vein thrombosis	Ultrasonography by an angiologist
17. UPN: 17.201	F	67	Cardiac surgery; ascendant and descendant thoracic, supra-renal abdominal aortic ecstasia with prosthetic replacement of ascending aorta and thoracic aorta endoprosthesis	Prophylactic UFH	25	5	HIT	None	N/A
18. UPN: 16.305	M	50	Cardiology; anterior STEMI with angioplasty and stenting	Prophylactic, then therapeutic, then prophylactic UFH	56	7	HIT-T	Bilateral pulmonary embolism and bilateral deep vein thrombosis	Chest CT scan with contrast product ultrasonography by an angiologist
19. UPN: 15.103	F	53	Intensive care unit; perimyocarditis with cardiogenic shock, septic shock and disseminated intravascular coagulation with heparin anticoagulation	Prophylactic, then therapeutic UFH	19	5	HIT	None	N/A

CLIA (U/ml)	PaGIA (titre)	Type of treatment and duration (days)	Thrombo-embolic events during treatment YES/NO *see Table 3	Bleeding events YES/NO *see Table 4	Limb outcome: Amputation and day of Argatroban treatment	Final outcome: Discharge xxx days after diagnosis of HIT versus hospital transfer
2.98	N.A.	Argatroban; 10 days	NO	NO	No amputation	Discharged 13 days after HIT diagnosis
2.05	>32	Argatroban; 3 days	YES: 1* (internal carotid artery thrombosis and stroke)	YES: 1* (intracranial bleeding secondary to ischaemic stroke)	No amputation	Deceased 3 days after diagnosis of HIT (massive PE, pneumonia, intracranial bleeding with cerebral herniation)
20.71	>32	Argatroban; 8 days	NO	NO	No amputation	Discharged 8 days after HIT diagnosis
1.70	8	Argatroban; 10 days	NO	NO	No amputation	Discharged 15 days after HIT diagnosis
>128	>32	Argatroban; 29 days	NO	NO	No amputation	Discharged 31 days after HIT diagnosis
1.09	>16	Argatroban; 13 days	NO	NO	No amputation	Discharged 14 days after HIT diagnosis
1.00	2	Argatroban; 49 days	NO	NO	No amputation	Discharged 93 days after HIT diagnosis
33.15	>16	Argatroban; 31 days	NO	NO	No amputation	Discharged 79 days after HIT diagnosis
0.74	8	Danaparoid; 6 days and then argatroban; 109 days	NO	NO	1: Left lower limb Gritti amputation 13 days after HIT diagnosis 2: Right heel necrosectomy 43 days after HIT diagnosis 3: Left fingers 2,3,4,5 amputations 50 days after HIT diagnosis 4: Right fingers 2,3,4 amputation and right foot toes 1,3,4,5 amputation 104 days after HIT diagnosis NB: patient suffered acute hepatic necrosis and ischaemic limb necrosis before onset of HIT	Discharged 155 days after HIT diagnosis

(Continues)

**TABLE 2** (Continued)

	Sex	Age	Care unit; underlying main disease(s) and management at our institution	Type of heparin and dose	Platelet count nadir value (G/l)	4T score	HIT versus HIT-T	HIT-T: Type of thrombo-embolic event at diagnosis	Diagnostic work-up
20. UPN: 14.038	M	61	Thoracic surgery; metastatic neuro-endocrine tumour and exploratory thoracotomy with oncologic surgery	Prophylactic, then therapeutic UFH	25	4	HIT	None	thrombin time N/A
21. UPN: 14.049	F	78	Internal medicine; metastatic small-cell lung carcinoma and chemotherapy	Port-a-cath flushing with UFH, then therapeutic LMWH, then therapeutic UFH	23	4	HIT-T	Pulmonary embolism and internal jugular and subclavian veins thrombosis	Chest CT scan with contrast product ultrasonography by an angiologist
22. UPN: 14.072	F	83	Neurosurgery; glioblastoma and radiochemotherapy	Prophylactic UFH	63	5	HIT-T	Multiple pulmonary embolisms	Chest CT scan with contrast product
23. UPN: 15.001	M	72	Intensive care unit; metastatic prostate adenocarcinoma with pleural carcinosis and right thoracotomy with right pleural decortication	Prophylactic, then therapeutic UFH	56	5	HIT-T	Radial artery thrombosis (when radial catheter was removed)	Ultrasonography by an angiologist
24. UPN: 15.138	M	60	Vascular surgery; peripheral artery embolies with rapidly progressive lower limb ischaemia and lower limb bypass and thrombolysis	Therapeutic UFH	63	7	HIT-T	Aorto-iliac and infra-renal aorta aneurysm thrombosis	Ultrasonography by an angiologist; vascular CT scan with contrast product
25. UPN: 15.143	F	86	Cardiac surgery; ascendant aorta aneurysm and ascending aorta replacement	Prophylactic UFH, then prophylactic LMWH, then prophylactic UFH	35	5	HIT	None	N/A
26. UPN: 15.160	F	66	Vascular surgery; advanced bilateral peripheral artery disease and angioplasty with stenting, thrombolysis, Fogarty thrombectomy, bypass and endarterectomy	Prophylactic, then therapeutic UFH	82	6	HIT-T	Bilateral superficial femoral artery and venous popliteo-poplital bypass thrombosis	Ultrasonography by an angiologist
27. UPN: 15.166	F	54	Cardiac surgery; severe aortic valve stenosis associated with ascendant aorta dilatation. Mechanic aortic replacement valve and ascending aorta replacement	Prophylactic, then therapeutic UFH	14	6	HIT	None	N/A
28. UPN: 15.169	M	57	Cardiac surgery; coronary heart disease and coronary bypass	Prophylactic LMWH	23	6	HIT-T	Splanchnic vein thrombosis with mesenteric ischaemia	Chest and abdominal CT with contrast product

CLIA (U/ml)	PaGIA (titre)	Type of treatment and duration (days)	Thrombo-embolic events during treatment YES/NO *see Table 3	Bleeding events YES/NO *see Table 4	Limb outcome: Amputation and day of Argatroban treatment	Final outcome: Discharge xxx days after diagnosis of HIT versus hospital transfer
0.15	4	Argatroban; 17 days	YES: 1* (bilateral PE)	NO	No amputation	Discharged 28 days after HIT diagnosis
11.84	32	Argatroban; 8 days	Check; YES?	NO	No amputation	Discharged 8 days after HIT diagnosis
14.16	32	Cava filter on day -5 of treatment argatroban; 24 days	YES: 1* (vena cava filter and bilateral pelvic veins thrombosis)	NO	No amputation	Deceased 46 days after HIT diagnosis
24.62	32	Argatroban; 5 days	NO	NO	No amputation	Deceased 5 days after HIT diagnosis (metastatic prostatic carcinoma with pleural carcinosis and global respirator insufficiency)
2.77	32	Argatroban; 19 days	NO	NO	Gritti amputation of right lower limb 4 days after HIT diagnosis	Discharged 59 days after HIT diagnosis
28.7	32	Argatroban; 8 days	NO	NO	No amputation	Discharged 12 days after HIT diagnosis
1.37	8	Argatroban; 21 days Left lower limb prosthetic bypass between superficial femoral artery and popliteal artery	NO	YES* 1 major bleeding; hematochezia (argatroban; ASS cardio and clopidogrel) 1 minor bleeding; melena (argatroban and clopidogrel)	Burgess amputation of left lower limb 8 days after HIT diagnosis	Discharged 72 days after HIT diagnosis
17.97	8	Argatroban; 16 days	NO	NO	No amputation	Discharged 17 days after HIT diagnosis
26.16	8	Argatroban; 21 days median laparotomy; resection of 1.5 m of bowel	YES* thrombosis of principal left thumb artery	NO	No amputation resection of 1.5 m of ischaemic bowel 1 day after HIT diagnosis	Discharged 23 days after HIT diagnosis

(Continues)

**TABLE 2** (Continued)

	Sex	Age	Care unit; underlying main disease(s) and management at our institution	Type of heparin and dose	Platelet count nadir value (G/l)	4T score	HIT versus HIT-T	HIT-T: Type of thrombo-embolic event at diagnosis	Diagnostic work-up
29. UPN: 15.186	M	33	Cardiology; idiopathic dilatative cardiomyopathy with intra-ventricular thrombus and heparin anticoagulation	Therapeutic UFH	55	5	HIT	None	N/A
30. UPN: 18.225	F	65	Cardiac surgery; severe aortic and moderate mitral valves stenosis; bioprosthetic aortic and mitral replacement valves	Prophylactic UFH	16	5	HIT	None	N/A
31. UPN: 18.243	M	70	Cardiology; heart failure with reduced ejection fraction and dilatative cardiomyopathy of unknown origin and left ventricular assist device	UFH therapeutic	50	4	HIT	None	N/A
32. UPN: 19.001	F	79	Vascular surgery; advanced peripheral artery disease and thrombendarterectomy and Fogarty thromboembolectomy	Therapeutic, then prophylactic UFH	38	3	HIT-T	Arterial; common and external iliac arteries thrombosis	Ultrasonography by an angiologist

Note: As per inclusion criteria, all 32 subjects were 'HIPA-positive'. N/A, not applicable: no HIT-T.

Abbreviations: ASA, acetylsalicylic acid; CLIA, chemiluminescent immunoassay; CT, computed tomography; DVT, deep vein thrombosis; F, female; HIPA, heparin-induced platelet aggregation; HIT, heparin-induced thrombocytopenia; HIT-T, HIT and thrombosis; IVIG, intravenous immunoglobulin; LMWH, low-molecular-weight heparin; M, male; PaGIA, particle gel immunoassay; PE, pulmonary embolism; STEMI, ST-segment elevation myocardial infarction; UFH, unfractionated heparin; UPN, unique patient number.

## Plasma argatroban elimination

PAC was monitored after argatroban discontinuation in case of planned surgery, bleeding events, and switching to vitamin K antagonists (VKA). Residual PAC at time  $t'$  after argatroban discontinuation were calculated as percentage of the last steady-state concentration and represented with a one-phase decay curve. We estimate that argatroban had an approximate plasma half-life of 60 min among patients with normal hepatic function (Figure 7A). Because of data scarcity, we could not analyse PAC decay after argatroban discontinuation among patients with hyperbilirubinaemia.

## Impact of argatroban on values of the international normalised ratio

Figure 8 depicts median international normalized ratio (INR) among patients ( $n = 11$ ) treated with VKA and

argatroban. Median INR was 4.5 immediately before argatroban discontinuation and 2.9 measured 3–6 h (median, 263 min) following argatroban discontinuation (median INR decrease  $-1.2$ , IQR INR decrease  $-0.7$  to  $-1.7$ ).

## DISCUSSION

In order to develop evidence-based in-house recommendations for argatroban dosing and monitoring we analysed a cohort of 32 patients with confirmed HIT, about half of them ( $n = 15$ , 47%) presenting with HIT-T (Table 1). Median anticoagulation duration was 17.5 days (IQR 10–29.5, range 3–109) for a total of 729 treatment days. As depicted in Figure 1, argatroban was biologically effective, enabling progressive decrease of D-dimers and platelet count normalisation within 7–8 days of treatment, without causing a drop of haemoglobin levels. These results are in line with those presented in our previous publication.<sup>6</sup>

CLIA (U/ml)	PaGIA (titre)	Type of treatment and duration (days)	Thrombo-embolic events during treatment YES/NO *see Table 3	Bleeding events YES/NO *see Table 4	Limb outcome: Amputation and day of Argatroban treatment	Final outcome: Discharge xxx days after diagnosis of HIT versus hospital transfer
3.55	8	Argatroban; 18 days	YES* Intra-ventricular thrombus (already present before HIT) with multiples embolic strokes; thrombo-emboly into the anterior Interventricular coronary artery	YES; Major bleeding: intracranial bleeding secondary to ischaemic stroke	No amputation	Discharged 28 days after HIT diagnosis
3.19	16	Argatroban; 14 days	NO	NO	No amputation	Hospital transfer 19 days after HIT diagnosis (under AVK)
0.84	8	Argatroban; 53 days	YES* Thrombosis of aortic valve leaflet (patient with impaired opening of the aortic valve; i.e. opening 1/10 cardiac cycles)	NO	No amputation	Discharged 109 days after HIT diagnosis
5.59	4	Argatroban; 10 days; left iliac arteries and left superficial and deep femoral arteries Fogarty thromboembolectomy; angioplasty and stenting of superficial femoral artery; short femoro-femoral bypass, angioplasty and stenting of left distal external iliac artery	NO	NO	No amputation	Hospital transfer 10 days after HIT diagnosis

### The effect of timing of first monitoring on the plasma argatroban concentration

Our results do not support the recommendation of performing the first laboratory control 2 h after starting argatroban.<sup>10,15,19</sup> Among patients with normal liver function, a steady-state plasmatic concentration is reached at the earliest 4 h after starting argatroban, both in case of a SAD of 0.5 µg/kg/min (Figure 2A) and 1.0 µg/kg/min (Figure 2B). Based on our observations, we estimate a plasmatic argatroban half-life of approximately 1 h (Figure 7A), which agrees with published data.<sup>12</sup> Our results are in line with those of Tardy-Poncet *et al.* who noted that argatroban concentrations measured 2 h after SAD were lower than steady-state concentrations determined later (i.e. 0.39 ± 0.29 µg/ml vs. 0.61 ± 0.28 µg/ml respectively).<sup>15</sup> Based on these results and since four to five drug half-lives are necessary to reach plasma steady state after starting administration,<sup>20</sup> we

suggest performing the first PAC control at least 4 h after SAD.

### The effect of the dose adjustments and timing of monitoring on subsequent plasma argatroban concentrations

To our knowledge, there is only one publication that assessed the magnitude of argatroban dose adaptations and their impact on consecutive PAC.<sup>6</sup> Recent guidelines on management of patients with HIT did not give recommendations on this issue.<sup>10</sup> Our data show that: (i) ADI intervals of 0.01–0.10, 0.11–0.20, 0.21–0.30, and 0.31–0.4 µg/kg/min lead to significant PAC increases (Figure 3A); (ii) a given ADI induced a roughly similar median PAC increase (i.e., Δ ADI µg/kg/min ≈ Δ PAC µg/ml); and (iii) significant differences in PAC increases were observed between ADI of 0.01–0.10

TABLE 3 Thromboembolic events under argatroban anticoagulation

	Timing of the thromboembolic event;	Type and localisation	Argatroban dosage in the 24 h preceding the TE diagnosis; mean (detailed doses), µg/kg/min	PAC measured in the 24 h preceding the TE diagnosis; mean (µg/ml)	Minimal PAC measured in the 24 h preceding the TE diagnosis (µg/ml)	Additional thrombotic risk factors
HIT type	day of argatroban treatment					
1. UPN: 18.022	4th	Deep vein thrombosis	1.39 (1.33; 1.45)	0.59	0.55 <sup>b</sup>	High BMI; recent surgery; immobilisation; active cancer; systemic infection (urosepsis)
2. UPN: 18.148	2nd	Femoro-iliac arterial prosthetic bypass thrombosis	0.65 (1.0; 0.5; 0.45)	1.04	0.91	Recent surgery; immobilisation; systemic infection (pneumonia)
3. UPN: 18.148	28th	Bilateral lung emboli	0.5 (0.5)	0.45	0.43; NB: 0.26; there might be a diagnostic delay of the lung emboli <sup>bc</sup>	Recent surgery; immobilisation; systemic infection (pneumonia) <sup>a</sup>
4. UPN: 18.148	36th	Superficial femoral and popliteal arteries thrombosis	0.6 (0.5; 0.59; 0.7)	0.56	0.36 <sup>b</sup>	Recent surgery; immobilisation; systemic infection (pneumonia)
5. UPN: 16.193	2nd	Internal carotid artery thrombosis and subsequent ischaemic stroke	0.55 (0.5; 0.59)	0.53	0.21 <sup>b</sup>	High BMI; recent surgery; immobilisation; active cancer; systemic infection (pneumonia) <sup>a</sup>
6. UPN: 14.038	4th	Bilateral lung emboli	0.37 (0.12; 0.39; 0.6)	0.5	0.22 <sup>b</sup>	Recent surgery; immobilisation; active cancer
7. UPN: 14.049	4th	Subclavian vein and internal jugular vein thrombosis	0.53 (0.45; 0.48; 0.5; 0.52; 0.69)	0.20	0.11	Active cancer; subclavian catheter for chemotherapy
8. UPN: 14.072	8th	Vena cava filter thrombosis	1.32 (1.21; 1.42)	0.38	0.3	Recent surgery; immobilisation; active cancer
9. UPN: 14.072	14th	Vena cava filter thrombosis extension and bilateral pelvic veins thrombosis	1.86 (1.86)	0.66	0.63	Recent surgery; immobilisation; active cancer

**TABLE 3** (Continued)

	Timing of the thromboembolic event;	Argatroban dosage in the 24 h preceding the TE diagnosis;	PAC measured in the 24 h preceding the TE diagnosis;	Minimal PAC measured in the 24 h preceding the TE diagnosis	Additional thrombotic risk factors
HIT type	Type and localisation	mean (detailed doses), µg/kg/min	mean (µg/ml)	(µg/ml)	
10. UPN: 15.169	2nd day of argatroban treatment Main thumb artery thrombosis	<b>0.61</b> (0.92; 0; 0.92)	0.28	0	Recent surgery; systemic infection secondary to mesenteric ischaemia; radial artery catheter
11. UPN: 15.186	15th day of argatroban treatment Intra-cardiac thrombus with subsequent cardio-embolic occipital strokes	<b>2.29</b> (2.29)	0.58	0.49	Immobilisation; advanced heart failure (dilatative cardiomyopathy with intra-ventricular thrombus); coronary and ventricular multiple catheterisms
12. UPN: 18.243	3rd day of argatroban treatment Left coronary leaflet of the aortic valve thrombosis	<b>0.27</b> (0.23; 0.29; 0.25; 0.29)	0.50	0.37	Recent surgery; immobilisation; systemic infection (pneumonia with sepsis); heart failure with reduced EF (because of an aortic valve briefly and partially opening 1/10 cardiac cycles) and ventricular dilatation; left jugular central venous catheter
13. UPN: 18.243	27th day of argatroban treatment Internal jugular vein and subclavian vein thrombosis	<b>0.35</b> (0.35)	0.75	0.65	Recent surgery; immobilisation; systemic infection (pneumonia with sepsis); heart failure with reduced EF and ventricular dilatation; left jugular central venous catheter

Five thromboembolic events were related to a possibly insufficient anticoagulation (°). Among them, four events were preceded by a minimum PAC below 0.4 µg/ml (°) in the 24 h preceding the diagnosis. One event might have been diagnosed with delay, being preceded by a minimum PAC below 0.4 µg/ml three days before diagnosis confirmation (°). Three thromboembolic events were partially or fully related to poor haemodynamics; two events were associated with advanced cardiac failure with severely reduced ejection fraction (°) and one was associated with low blood flow through a prosthetic vascular bypass in a patient with advanced peripheral arterial disease (°). Five thromboembolic events were related to intravascular foreign material (°).

Abbreviations: BMI, body mass index; EF, ejection fraction; HIT, heparin-induced thrombocytopenia; HIT-T, HIT and thrombosis; PAC, plasmatc argatroban concentrations; TE, thrombotic event; UPN, unique patient number. Bold indicate mean values.

TABLE 4 Bleeding events under argatroban anticoagulation

	Timing of bleeding event;	HIT type	day of argatroban treatment	Type and localisation	Blood products administered	Argatroban dosage in the 24 h preceding the TE diagnosis;	Mean PAC ( $\mu\text{g/ml}$ ) measured in the preceding 24 h (day prior diagnosis and day of diagnosis of the bleeding event)	Maximal PAC ( $\mu\text{g/ml}$ ) measured in the preceding 24 h (day prior diagnosis and day of diagnosis of the bleeding event)	Additional bleeding risk factors (bleeding history—reduced haemoglobin—concomitant antiplatelet drug—renal impairment — anatomical lesions — recent surgery)	Suspected cause of bleeding events (excessive anticoagulation and antiaggregation versus anatomical lesion vs. undetermined)
1. UPN: 17.222	HIT	2nd	Minor bleeding; subcutaneous spontaneous haematoms on both upper limbs	0		<b>0.43</b> (0.25; 0.45; 0.58)	0.23	0.33	Recent aortic valve replacement; impaired renal function	Undetermined
2. UPN: 17.042	HIT-T	2nd	Major bleeding; macrohematuria and jugular bleeding with haemorrhagic shock <sup>b</sup>	3 units of packed red blood cells; 1 unit of fresh frozen plasma		<b>0.92</b> (0.92)	0.55	0.77	Vesical, prostatic and ureteral lesions secondary to self-removal of urinary catheter; internal jugular vein lesion secondary to self-snatch of central venous catheter clopidogrel and aspirin treatments (recent stenting of RIVA); renal impairment; reduced haemoglobin	Anatomical lesion; possibly 'excessive' antiaggregation
3. UPN: 16.193	HIT-T	3rd	Major bleeding; intracerebral bleeding with cerebral herniation <sup>a</sup>	0		<b>0.59</b> (0.59)	0.59	0.69	Recent ischaemic stroke secondary to thrombosis of the right internal carotid artery	Anatomical lesion
4. UPN: 15.160	HIT-T	9th	Major bleeding; hematochezia secondary to per-operative ischaemic colitis <sup>b</sup>	2 units of packed red blood cells		Mean: 0.39 (0; 0.41; 0.61; 0.52)	0.26	0.45	Clopidogrel and aspirin treatments; reduced haemoglobin due to digestive angiodysplasia	Anatomical lesion and possibly 'excessive' anti-aggregation
5. UPN: 15.186	HIT	20th	Major bleeding; asymptomatic bilateral frontal intracerebral bleedings and left intracerebellar bleeding <sup>a</sup>	0		<b>2.29</b> (2.29)	0.58	0.67	Multiple cardio-embolic cerebral and cerebellar strokes	Anatomical lesion

Note: We observed five bleeding events, four of them being major and one of them being minor, according to the ISTH definition.<sup>16,17</sup> Mean and maximal PAC ( $\mu\text{g/ml}$ ) measured in the 24 h preceding these five bleeding events seemed not to be excessive. Bold indicate mean values.

Abbreviations: HIT, heparin-induced thrombocytopenia; PAC, plasmatic argatroban concentrations; RIVA, ramus interventricularis anterior; TE, thrombotic event; UPN, unique patient number.

<sup>a</sup>Among the four major bleeding events, two were intracranial bleedings that occurred after ischaemic stroke.

<sup>b</sup>The two other major bleeding events occurred under concomitant double antiplatelet therapy (aspirin and clopidogrel) and were related to anatomical lesions.

**TABLE 5** Standardised in-house management of HIT patients with argatroban

	Normal liver function	Impaired liver function		
		Impaired bilirubin excretion	Isolated liver cytolysis	Multiorgan dysfunction <sup>a</sup>
Monitoring of platelets and D-dimers	Once daily	Once daily	Once daily	Once daily
Monitoring of hepatic function	Before SAD, then at discretion of physician	Before SAD, then at discretion of physician	Before SAD, then at discretion of physician	Before SAD, then at discretion of physician
SAD (µg/kg/min)	1.0	0.25 <sup>13</sup> versus contra-indicated <sup>b</sup>	0.75 <sup>c</sup>	0.50
Timing of monitoring after SAD (hours)	4 h; repeat twice at 4 h each if no ADA, afterwards once daily in ss	6 h; repeat twice at 6 h each if no ADA, afterwards once daily in ss	4 h; repeat twice at 4 h each if no ADA, afterwards once daily in ss	4 h; repeat twice at 4 h each if no ADA, afterwards once daily in ss
ss-PAC target (µg/ml)	0.5–1.0 <sup>d</sup>	0.5–1.0 <sup>d</sup>	0.5–1.0 <sup>d</sup>	0.5–1.0 <sup>d</sup>
ss-TT target (s)	80–120	80–120	80–120	80–120
ss-aPTT target (fold increase and s)	1.7–2.4 × baseline value (max. 70–75 s) <sup>e</sup>	1.7–2.4 × baseline value (max. 70–75 s) <sup>e</sup>	1.7–2.4 × baseline value (max. 70–75 s) <sup>e</sup>	1.7–2.4 × baseline value (max. 70–75 s) <sup>e</sup>
Magnitude of ADA (µg/kg/min)	Adapted to first PAC/TT and to ss-PAC-targeted; Δ ADA µg/kg/min ≈ Δ PAC µg/ml (at least 0.2 µg/kg/min)	Adapted to first PAC/TT and to ss-PAC-targeted; Δ ADA µg/kg/min ≈ 2Δ PAC µg/ml (0.05–0.1 µg/kg/min)	Adapted to first PAC/TT and to ss-PAC-targeted; Δ ADA µg/kg/min ≈ Δ PAC µg/ml (0.2 µg/kg/min)	Adapted to first PAC/TT and to ss-PAC-targeted; Δ ADA µg/kg/min ≈ Δ PAC µg/ml (0.2 µg/kg/min)
Timing of monitoring after ADA	4 h; repeat 4 h after each ADA	6 h; repeat 6 h after each ADA	4 h; repeat 4 h after each ADA	4 h; repeat 4 h after each ADA
Frequency of PAC/TT/aPTT monitoring	Once daily after 3 target ss-PAC/TT	Once daily after 3 target ss-PAC/TT	Once daily after 3 target ss-PAC/TT	Once daily after 3 target ss-PAC/TT
When bridging to AVK: target INR on argatroban	3.5–4.5	3.5–4.5	3.5–4.5	–
When bridging to AVK: timing of INR after argatroban stop	After 4 h	After more than 4 h	After 4 h	–

Abbreviations: ADA, argatroban dose adaptation; aPTT, activated partial thromboplastin time; AVK, vitamin K antagonist; INR, international normalised ratio; SAD, starting argatroban dose; ss, steady state; TT, thrombin time (see Methods).

<sup>a</sup>Multi-organ dysfunction without impaired bilirubin excretion.

<sup>b</sup>Use SAD of 0.25 µg/kg/min among patients with mild to moderate bilirubin excretion impairment and consider danaparoid for patients with severe bilirubin excretion impairment.

<sup>c</sup>See results in paragraph "Argatroban dosing adapted to liver function".

<sup>d</sup>Target higher PAC (1–1.5 µg/ml) in case of insufficient platelet count recovery and/or D-dimers level decrease.

<sup>e</sup>Reagent- and coagulometer-specific values, not generalisable.

µg/kg/min and 0.21–0.30 µg/kg/min and between ADI of 0.01–0.10 µg/kg/min and 0.31–0.40 µg/kg/min.

Correct timing of laboratory monitoring is crucial for obtaining a reliable result. However, to the best of our knowledge, there are no published data concerning this practical issue.<sup>10</sup> Similarly to what we have observed with initial monitoring (Figure 2), we show in Figure 3B that after ADI of 0.1–0.4 µg/kg/min, median PAC increase was significantly greater when laboratory control was performed 'late' (after 3 to up to 6 h) compared to 'early' controls (within 3 h). Hence, this underscores that timing of control after argatroban dose adaptation is critical<sup>15</sup> and we suggest waiting 4 h after dose adaptation among patients with normal liver function.

## The effect of liver function on argatroban dosing

The argatroban dose required to reach a given target PAC depends on liver function. As suggested by Levine *et al.*,<sup>13</sup> serum total bilirubin might be more useful than ALT to identify patients who require dose reduction. Based on the alteration of bilirubin and/or ALT, we defined four different hepatic impairment patterns (Figure 4). Patients with increased bilirubin levels required significantly lower argatroban steady-state doses than patients with normal liver function irrespective of the target PAC interval. Patients with liver cytolysis tended to require lower median argatroban doses at steady state than patients with normal liver

function, in order to reach prophylactic and high therapeutic PAC intervals. Our data confirm that impaired bilirubin excretion plays the most important role in decision-making for argatroban dosing and indicate that patients with liver cytolysis probably require smaller argatroban dose reductions compared to patients with impaired bilirubin excretion.

### The extent of bilirubinaemia and argatroban dosing

A total plasma bilirubin level higher than 25.5  $\mu\text{mol/l}$  has been proposed to identify patients with impaired bilirubin excretion, thus requiring argatroban dose reduction.<sup>13</sup> However, as presented in Figure 5, we could not identify a critical serum total bilirubin threshold, observing a progressive impact on the required dosage in order to achieve a prophylactic (Figure 5A), standard therapeutic (Figure 5B), or high therapeutic PAC (Figure 5C) already at bilirubin values within reference range. Based on our data, we suggest to consider reducing argatroban doses among patients with serum total bilirubin levels above 20  $\mu\text{mol/l}$ , which is lower than the threshold suggested by Levine *et al.*<sup>13</sup>

### Argatroban laboratory monitoring with aPTT and thrombin time

Activated partial thromboplastin time is the most widely available laboratory assay for monitoring argatroban anticoagulation and is currently recommended by the American Society of Hematology (ASH) guidelines.<sup>10</sup> However, multiple factors might affect its results, leading to inaccurate monitoring, which in turn may result in argatroban under-dosing and thrombosis progression.<sup>21</sup> Our group and others have shown that TT is more reliable and robust for the monitoring of argatroban.<sup>6,22</sup> Figure 6A confirms a linear relationship between PAC and corresponding TT intervals. In contrast, the relationship between increasing PAC and correspondent simultaneous aPTT intervals is not linear, reaching a plateau for aPTT values above 70 s (Figure 6B), underscoring that this coagulation assay is less accurate for monitoring argatroban anticoagulation.<sup>6,21,23</sup> We therefore suggest to monitor argatroban by TT when the assessment of PAC with a commercially diluted TT assay is not available. From an analytical point of view, it would be better to employ two TTs, e.g. with final thrombin concentrations of 1.5 and 5 U/ml as we did in our previous work.<sup>6,24</sup> However, our current experience indicates that also a single TT with a final thrombin concentration of 1.25 U/ml is adequate up to argatroban concentrations of about 1  $\mu\text{g/ml}$  (Figure 6A). Monitoring argatroban with thrombin-time-based assays is in line with current expert's opinions.<sup>21</sup>

### Argatroban half-life after discontinuation

We found an argatroban plasma half-life of about 1 h in patients with normal liver function (Figure 2A,B). Accordingly, Figure 7 confirms that the decay of PAC after argatroban discontinuation follows the same kinetics. We therefore suggest discontinuing argatroban at least 2 h before any invasive intervention among patients with normal liver function.

### The effect of argatroban on the INR

As well as direct oral anticoagulants,<sup>9</sup> VKA are still employed for HIT patients after the acute phase,<sup>5</sup> targeting INR values between 2 and 3 for optimal anticoagulation.<sup>5</sup> Of note, argatroban and other direct thrombin inhibitors increase INR values; this implies that during the overlap phase of anticoagulation with both argatroban and VKA, INR values greater than 2–3 have to be achieved before stopping argatroban.<sup>25</sup> Harder *et al.*<sup>26</sup> recommend to target INR values higher than 4 and Bartholomew *et al.*<sup>27</sup> recommend performing INR controls 6 h after argatroban discontinuation. In Figure 8, we show that INR values ranging from 3.5 to 4.5 during the overlapping phase are sufficient to obtain a target INR of 2–3 after argatroban discontinuation. Indeed, we observed a median INR decrease of 1.2 (range: –0.7; –1.7) when control was performed 3–6 h after argatroban discontinuation. These data are perfectly in line with those obtained by two other groups who observed median INR decreases of 1.2.<sup>6,28</sup>

### Limitations of our study

Our study has limitations. Firstly, this was a single-centre study. Secondly, our data were collected retrospectively. Thirdly, argatroban monitoring with a commercial diluted TT is not widely available. Nevertheless, all HIT cases were confirmed by a functional gold standard assay, we studied patients treated in a real-world clinical setting and analysed an argatroban treatment length of 729 days. Moreover, to the best of our knowledge, our cohort is the biggest ever described including patients with HIT confirmed by a functional gold standard assay who were treated with argatroban.<sup>6,11,29–32</sup>

### Practical implications of our study

Our results provide guidance on some practical aspects of anticoagulation of HIT patients with argatroban, such as: (i) the effect of various argatroban starting doses and dose adjustments on the following plasma concentrations; (ii) the effect of liver function on argatroban dosing; (iii) the effect of timing and laboratory assay on monitoring; and (iv) the effect of therapeutic argatroban on INR values. Based on these data we

have developed a standardised in-house management protocol (Table 5), which we are prospectively validating. Briefly, we do not differentiate between HIT and HIT-T and employ a SAD of 1.0 µg/kg/min in all HIT patients with normal liver function. In patients with impaired liver function, we decrease SAD to 0.75 µg/kg/min in presence of isolated hepatic cytolysis and to 0.25 µg/kg/min in case of hyperbilirubinaemia. We monitor treatment with a double pharmacokinetic and pharmacodynamic approach. On one side, we assess PAC at steady state employing a thrombin-time-based assay, which is more robust than aPTT. On the other side, we follow the course of platelet counts and D-dimer levels in order to fine-tune the anticoagulation intensity required in each individual patient.

## ACKNOWLEDGEMENTS

We thank the physicians caring for the patients, the technicians of the haemostasis laboratory, Prof. MD Michel Duchosal for support, and Prof. MD Pierre-Alexandre Bart for acting as expert of the MD thesis of Matteo Marchetti. This study has been partially presented at the Swiss Oncology and Haematology Congress (SOHC) in Zürich, Switzerland, June 2019. Open Access Funding provided by Université de Lausanne. [Correction added on 26 May 2022, after first online publication: CSAL funding statement has been added.]

## CONFLICT OF INTEREST

The authors do not have any conflict of interest to declare.

## AUTHOR CONTRIBUTIONS

Matteo Marchetti designed the research, collected and analysed data, and wrote the manuscript. Stefano Barelli designed the research and cowrote the manuscript. Tobias Gleich and Francisco J. Gomez collected data. Matthew Goodyer cowrote the manuscript. Francesco Grandoni was in charge of the patients, analysed data and cowrote the manuscript. Lorenzo Alberio is the corresponding author of the manuscript. He was in charge of the patients, designed the research, analysed data, and wrote the manuscript. All authors read and approved the final version of the manuscript.

## ORCID

Matteo Marchetti  <https://orcid.org/0000-0002-7741-0740>

Stefano Barelli  <https://orcid.org/0000-0001-6710-9738>

Tobias Gleich  <https://orcid.org/0000-0002-2145-0612>

Francisco J. Gomez  <https://orcid.org/0000-0002-8816-037X>

Matthew Goodyer  <https://orcid.org/0000-0001-5804-4397>

Francesco Grandoni  <https://orcid.org/0000-0003-2208-2070>

Lorenzo Alberio  <https://orcid.org/0000-0001-9686-9920>

## REFERENCES

- East JM, Cserti-Gazdewich CM, Granton JT. Heparin-induced thrombocytopenia in the critically ill patient. *Chest*. 2018;154(3):678–90.
- Warkentin TE, Levine MN, Hirsh J, Horsewood P, Roberts RS, Gent M, et al. Heparin-induced thrombocytopenia in patients treated with low-molecular-weight heparin or unfractionated heparin. *N Engl J Med*. 1995;332(20):1330–5.
- Warkentin TE, Sheppard JA, Horsewood P, Simpson PJ, Moore JC, Kelton JG. Impact of the patient population on the risk for heparin-induced thrombocytopenia. *Blood*. 2000;96(5):1703–8.
- Greinacher A. Clinical practice. Heparin-induced thrombocytopenia. *N Engl J Med*. 2015;373(3):252–61.
- Alatri A, Armstrong AE, Greinacher A, Koster A, Kozek-Langenecker SA, Lance MD, et al. Results of a consensus meeting on the use of argatroban in patients with heparin-induced thrombocytopenia requiring antithrombotic therapy - a European perspective. *Thromb Res*. 2012;129(4):426–33.
- Colucci G, Nagler M, Klaus N, Conte T, Giabani E, Alberio L. Practical guidelines for argatroban and bivalirudine in patients with heparin-induced thrombocytopenia. *J Transl Sci*. 2015;1(2):37–42.
- Chong BH, Gallus AS, Cade JF, Magnani H, Manoharan A, Oldmeadow M, et al. Prospective randomised open-label comparison of danaparoid with dextran 70 in the treatment of heparin-induced thrombocytopenia with thrombosis: a clinical outcome study. *Thromb Haemost*. 2001;86(5):1170–5.
- Schindewolf M, Steindl J, Beyer-Westendorf J, Schellong S, Dohmen PM, Brachmann J, et al. Frequent off-label use of fondaparinux in patients with suspected acute heparin-induced thrombocytopenia (HIT)--findings from the GerHIT multi-Centre registry study. *Thromb Res*. 2014;134(1):29–35.
- Warkentin TE, Pai M, Linkins LA. Direct oral anticoagulants for treatment of HIT: update of Hamilton experience and literature review. *Blood*. 2017;130(9):1104–13.
- Cuker A, Arepally GM, Chong BH, Cines DB, Greinacher A, Gruel Y, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: heparin-induced thrombocytopenia. *Blood Adv*. 2018;2(22):3360–92.
- Colarossi G, Maffulli N, Trivellas A, Schnoring H, Hatam N, Tingart M, et al. Superior outcomes with Argatroban for heparin-induced thrombocytopenia: a Bayesian network meta-analysis. *Int J Clin Pharmacol*. 2021;43(4):825–38.
- Swan SK, Hursting MJ. The pharmacokinetics and pharmacodynamics of argatroban: effects of age, gender, and hepatic or renal dysfunction. *Pharmacotherapy*. 2000;20(3):318–29.
- Levine RL, Hursting MJ, McCollum D. Argatroban therapy in heparin-induced thrombocytopenia with hepatic dysfunction. *Chest*. 2006;129(5):1167–75.
- Marchetti M, Barelli S, Zermatten MG, Monnin-Respen F, Matthey-Guirao E, Nicolas N, et al. Rapid and accurate Bayesian diagnosis of heparin-induced thrombocytopenia. *Blood*. 2020;135(14):1171–84.
- Tardy-Poncet B, Nguyen P, Thiranos JC, Morange PE, Biron-Andreani C, Gruel Y, et al. Argatroban in the management of heparin-induced thrombocytopenia: a multicenter clinical trial. *Crit Care*. 2015;19:396.
- Schulman S, Kearon C. Subcommittee on control of anticoagulation of the S, standardization Committee of the International Society on Thromb Haemost. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost*. 2005;3(4):692–4.
- Schulman S, Angeras U, Bergqvist D, Eriksson B, Lassen MR, Fisher W, et al. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in surgical patients. *J Thromb Haemost*. 2010;8(1):202–4.
- Siegal DM, Cook RJ, Warkentin TE. Acute hepatic necrosis and ischemic limb necrosis. *N Engl J Med*. 2012;367(9):879–81.
- Alberio L, Angelillo-Scherrer A, Asmis L, Casini A, Fontana P, Graf L, et al. Recommendations on the use of anticoagulants for the treatment of patients with heparin-induced thrombocytopenia in Switzerland. *Swiss Med Wkly*. 2020;150:w20210.
- Ito S. Pharmacokinetics 101. *Paediatr Child Health*. 2011;16(9):535–6.
- Warkentin TE. Anticoagulant failure in coagulopathic patients: PTT confounding and other pitfalls. *Expert Opin Drug Saf*. 2014;13(1):25–43.

22. Shepherd MF, Jacobsen JM, Rosborough TK. Argatroban therapy using enzymatic anti-factor IIa monitoring. *Ann Pharmacother.* 2011;45(3):422–3.
23. Keyl C, Zimmer E, Bek MJ, Wiessner M, Trenk D. Argatroban pharmacokinetics and pharmacodynamics in critically ill cardiac surgical patients with suspected heparin-induced thrombocytopenia. *Thromb Haemost.* 2016;115(6):1081–9.
24. Tschudi M, Lammler B, Alberio L. Dosing lepirudin in patients with heparin-induced thrombocytopenia and normal or impaired renal function: a single-center experience with 68 patients. *Blood.* 2009;113(11):2402–9.
25. Koster A, Fischer KG, Harder S, Mertzlufft F. The direct thrombin inhibitor argatroban: a review of its use in patients with and without HIT. *Biol Theory.* 2007;1(2):105–12.
26. Harder S, Graff J, Klinkhardt U, von Hentig N, Walenga JM, Watanabe H, et al. Transition from argatroban to oral anticoagulation with phenprocoumon or acenocoumarol: effects on prothrombin time, activated partial thromboplastin time, and Ecarin clotting time. *Thromb Haemost.* 2004;91(6):1137–45.
27. Bartholomew JR, Hursting MJ. Transitioning from argatroban to warfarin in heparin-induced thrombocytopenia: an analysis of outcomes in patients with elevated international normalized ratio (INR). *J Thromb Thrombolysis.* 2005;19(3):183–8.
28. Bain J, Meyer A. Comparison of bivalirudin to lepirudin and argatroban in patients with heparin-induced thrombocytopenia. *Am J Health-Syst Pharm.* 2015;72(17 Suppl 2):S104–9.
29. Lewis BE, Wallis DE, Berkowitz SD, Matthai WH, Fareed J, Walenga JM, et al. Argatroban anticoagulant therapy in patients with heparin-induced thrombocytopenia. *Circulation.* 2001;103(14):1838–43.
30. Begelman SM, Baghdasarian SB, Singh IM, Militello MA, Hursting MJ, Bartholomew JR. Argatroban anticoagulation in intensive care patients: effects of heart failure and multiple organ system failure. *J Intensive Care Med.* 2008;23(5):313–20.
31. Lewis BE, Wallis DE, Leya F, Hursting MJ, Kelton JG, Argatroban I. Argatroban anticoagulation in patients with heparin-induced thrombocytopenia. *Arch Intern Med.* 2003;163(15):1849–56.
32. Kuter DJ, Konkle BA, Hamza TH, Uhl L, Assmann SF, Kiss JE, et al. Clinical outcomes in a cohort of patients with heparin-induced thrombocytopenia. *Am J Hematol.* 2017;92(8):730–8.

## SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

**How to cite this article:** Marchetti M, Barelli S, Gleich T, Gomez FJ, Goodyer M, Grandoni F, Alberio L. Managing argatroban in heparin-induced thrombocytopenia: A retrospective analysis of 729 treatment days in 32 patients with confirmed heparin-induced thrombocytopenia. *Br J Haematol.* 2022;197:766–790. <https://doi.org/10.1111/bjh.18120>