

## **RUIN EXCURSIONS, THE $G/G/\infty$ QUEUE AND TAX PAYMENTS IN RENEWAL RISK MODELS**

H. ALBRECHER,\* *University of Lausanne*

S.C. BORST,\*\* *Eindhoven University of Technology*

O.J. BOXMA,\*\*\* *Eindhoven University of Technology*

J. RESING,\*\*\*\* *Eindhoven University of Technology*

### **Abstract**

In this paper we investigate the number and maximum severity of the ruin excursion of the insurance portfolio reserve process in the Cramér-Lundberg model with and without tax payments. We also provide a relation of the Cramér-Lundberg risk model with the  $G/G/\infty$  queue and use it to derive some explicit ruin probability formulas. Finally, the renewal risk model with tax is considered, and an asymptotic identity is derived that in some sense extends the tax identity of the Cramér-Lundberg risk model.

*Keywords:* classical risk model; ruin probability;  $G/G/\infty$  queue; tax; renewal model

2000 Mathematics Subject Classification: Primary 91B30

Secondary 60K30

### **1. Introduction**

Consider the classical Cramér-Lundberg model in risk theory to describe the surplus process  $\{R_t\}$  at time  $t$  of an insurance portfolio. Starting with an initial capital  $x$ , premium is collected according to a constant premium intensity (normalized to) 1.

---

\* Postal address: Department of Actuarial Science, Faculty of Business and Economics, University of Lausanne, Quartier UNIL-Dorigny, 1015 Lausanne, Switzerland

\*\* Postal address: Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands and Alcatel-Lucent, Bell Labs, 600 Mountain Avenue, P.O. Box 636, Murray Hill, NJ 07974-0636, USA

\*\*\* Postal address: Eindhoven University of Technology and EURANDOM, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

\*\*\*\* Postal address: Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

Claims occur according to a homogeneous Poisson process with intensity  $\lambda$  and are paid at the times of their occurrence. The claim sizes are independent and identically distributed random variables with distribution function  $H(\cdot)$ . Define  $\phi_0(x) = \mathbb{P}(R_t \geq 0 \text{ for all } t | R_0 = x)$  as the probability of survival and correspondingly the ruin probability as  $\psi_0(x) = 1 - \phi_0(x)$ . Let further  $V_{\max}$  be the maximum workload in an  $M/G/1$  queue with arrival rate  $\lambda$  and service time distribution  $H(\cdot)$ . Then the following relation between the Cramér-Lundberg risk model and the  $M/G/1$  queueing model is classical:

$$\phi_0(x) = e^{-\lambda \int_x^\infty \mathbb{P}(V_{\max} > y) dy}. \quad (1)$$

Let  $G(\cdot)$  denote the distribution function of  $V_{\max}$ . One way to show (1) is to use the well-known relation

$$G(u) = \mathbb{P}(V_{\max} \leq u) = 1 - \frac{1}{\lambda} \frac{d}{du} \ln \mathbb{P}(V \leq u), \quad (2)$$

where  $V$  is the stationary workload in the same  $M/G/1$  queue as described above, and use the sample path duality result  $\phi_0(x) = \mathbb{P}(V \leq x)$  (see e.g. Asmussen & Albrecher [5] for a recent survey). In [2] another more direct proof of (1) was given and subsequently used to establish a simple proof of the tax identity

$$\phi_\gamma(x) = \left( \phi_0(x) \right)^{\frac{1}{1-\gamma}} = e^{-\frac{\lambda}{1-\gamma} \int_x^\infty \mathbb{P}(V_{\max} > y) dy}, \quad (3)$$

where  $\phi_\gamma(x) = 1 - \psi_\gamma(x)$  is the survival probability in a Cramér-Lundberg model with tax rate  $0 \leq \gamma < 1$ , i.e. whenever the risk process is in its running maximum (and hence in a profitable position), a constant proportion  $\gamma$  of the incoming premium is paid as tax ( $\gamma = 0$  corresponds to the Cramér-Lundberg model without tax). For extensions of this identity in various directions see [1, 3, 4, 7, 10].

In this paper we will provide a relation of the Cramér-Lundberg risk model with the  $G/G/\infty$  queue, which will give rise to another view towards identity (1) and some explicit ruin probability formulas. Subsequently, we will consider the renewal risk model with tax, and establish an asymptotic identity that may be interpreted as an extension of the tax identity (3). We start with some refined results on the number and maximum severity of the ruin excursion in the Cramér-Lundberg model with and without tax.

## 2. Maximum severity of the ruin excursion

Consider the Cramér-Lundberg model with tax rate  $\gamma$ . Ruin can only occur during an ‘interruption’, i.e., a period in between running maxima. Denote the  $k$ th interruption period by  $P_k$ . Interruptions occur according to a Poisson process with intensity  $\lambda$ . The probability that no ruin occurs during an interruption that starts at surplus level  $z$  is given by  $G(z) = 1 - \bar{G}(z)$  (cf. (2)). Let  $R_{\min}$  be the lowest surplus value during the ruin excursion. Let further  $A_k(x, d)$  be the probability that ruin occurs during the  $k$ th interruption  $P_k$  and  $R_{\min} \leq -d$ , where  $d \geq 0$ . Then, for  $k \in \mathbb{N}$

$$A_k(x, d) = \int_{t=0}^{\infty} \lambda^k \frac{t^{k-1}}{(k-1)!} e^{-\lambda t} \left[ \int_{v=0}^t G(x + (1-\gamma)v) \frac{dv}{t} \right]^{k-1} \bar{G}(x + (1-\gamma)t + d) dt. \quad (4)$$

Here we have used that the sum of  $k$  independent exponential arrival intervals is Erlang( $k, \lambda$ ) distributed, and given that their sum is  $t$ , the interruption epochs are uniformly distributed on  $[0, t]$ .

**Proposition 2.1.** *Let  $A(x, d)$  be the probability that ruin occurs and the lowest surplus value of the ruin excursion is smaller than  $-d \leq 0$ . Then*

$$A(x, d) = \int_x^{\infty} \frac{\phi'_\gamma(w+d)}{\phi_\gamma(w+d)} \frac{\phi_\gamma(x)}{\phi_\gamma(w)} dw. \quad (5)$$

*Proof.* We have

$$\begin{aligned} A(x, d) &= \sum_{k=1}^{\infty} A_k(x, d) = \int_{t=0}^{\infty} \lambda e^{-\lambda t} \bar{G}(x + (1-\gamma)t + d) e^{\lambda \int_{v=0}^t G(x+(1-\gamma)v) dv} dt \\ &= \int_{t=0}^{\infty} \lambda \bar{G}(x + (1-\gamma)t + d) e^{-\lambda \int_{v=0}^t \bar{G}(x+(1-\gamma)v) dv} dt. \end{aligned} \quad (6)$$

Now the result follows from (2) and (3).  $\square$

**Remark 2.1.** Clearly  $d = 0$  gives  $A(x, 0) = 1 - \phi_\gamma(x) = \psi_\gamma(x)$ , so that in this case we indeed recover the usual ruin probability.

**Remark 2.2.** An alternative way to establish (6) is to use the joint distribution of the maximum surplus before ruin  $R_{\max} = \sup_{t \geq 0} R_t I_{\{R_u \geq 0 \text{ for all } u \in [0, t]\}}$  and the

maximum deficit of the ruin excursion  $R_{\min}$ . Concretely,

$$\begin{aligned} & \mathbb{P}(R_{\max} \in [y, y + dy]; R_{\min} \leq -d) \\ &= \frac{d}{dy} \left[ 1 - e^{-\frac{\lambda}{1-\gamma} \int_{v=x}^y \mathbb{P}(V_{\max} > v) dv} \right] \cdot \mathbb{P}(V_{\max} > y + d | V_{\max} > y) \\ &= \frac{\lambda}{1-\gamma} \mathbb{P}(V_{\max} > y + d) \cdot e^{-\frac{\lambda}{1-\gamma} \int_{v=x}^y \mathbb{P}(V_{\max} > v) dv}, \end{aligned}$$

which also yields (6) upon integration over  $y \geq x$ . Note in addition that the time spent in the running maximum until ruin is given by  $(R_{\max} - x)/(1 - \gamma)$ .

**Proposition 2.2.** *The generating function  $\Phi(z, x, d) := \sum_{k=1}^{\infty} z^k A_k(x, d)$  is given by*

$$\Phi(z, x, d) = z \int_x^{\infty} \frac{\phi'_\gamma(w + d)}{\phi_\gamma(w + d)} \left( \frac{\phi_\gamma(x)}{\phi_\gamma(w)} \right)^z e^{-\lambda(1-z)(w-x)/(1-\gamma)} dw. \quad (7)$$

*Proof.* From (4) it follows that

$$\begin{aligned} \Phi(z, x, d) &= z \int_{t=0}^{\infty} \lambda e^{-\lambda t} e^{z\lambda \int_{v=0}^t G(x+(1-\gamma)v) dv} \overline{G}(x + (1-\gamma)t + d) dt \\ &= z \int_{t=0}^{\infty} \lambda \overline{G}(x + (1-\gamma)t + d) e^{-\lambda(1-z)t} e^{-\lambda z \int_{v=0}^t \overline{G}(x+(1-\gamma)v) dv} dt, \end{aligned}$$

so that the assertion again follows from (2) and (3).  $\square$

Denote by  $K$  the number of the interruption that leads to ruin ( $K$  is a defective random variable on the positive integers). Then starting at (7) with  $d = 0$ , some elementary calculations lead to the following result:

**Corollary 2.1.**

$$\begin{aligned} \mathbb{E}[K | \text{Ruin occurs with } R_0 = x] &= \frac{\frac{\partial}{\partial z} \Phi(z, x, 0) \Big|_{z=1}}{\psi_\gamma(x)} \\ &= \ln \phi_\gamma(x) \left( 1 - \frac{1}{\psi_\gamma(x)} \right) - \frac{\lambda}{1-\gamma} \left( x - \frac{\phi_\gamma(x)}{\psi_\gamma(x)} \int_x^{\infty} \frac{w \phi'_\gamma(w)}{\phi_\gamma^2(w)} dw \right). \end{aligned}$$

On the other hand, one may rewrite (4) as follows:

$$\begin{aligned} A_k(x, d) &= \int_{t=0}^{\infty} \frac{\lambda}{(k-1)!} e^{-\lambda t} \left[ \lambda \int_{v=0}^t G(x + (1-\gamma)v) dv \right]^{k-1} \overline{G}(x + (1-\gamma)t + d) dt \\ &= \int_{t=0}^{\infty} \frac{e^{-\lambda t}}{(k-1)!} \left[ \lambda t - \int_{v=0}^t \frac{\phi'_0(x + (1-\gamma)v)}{\phi_0(x + (1-\gamma)v)} \right]^{k-1} \frac{\phi'_0(x + (1-\gamma)t + d)}{\phi_0(x + (1-\gamma)t + d)} dt \\ &= \int_{t=0}^{\infty} \frac{e^{-\lambda t}}{(k-1)!} \left[ \lambda t - \ln \frac{\phi_\gamma(x + (1-\gamma)t)}{\phi_\gamma(x)} \right]^{k-1} \frac{\phi'_0(x + (1-\gamma)t + d)}{\phi_0(x + (1-\gamma)t + d)} dt. \quad (8) \end{aligned}$$

Integrating over  $d$  and some elementary algebra then gives the following expressions:

**Corollary 2.2.** *The expected maximum severity of the ruin excursion, with ruin occurring at the  $k$ th interruption, is given by*

$$\begin{aligned} & \mathbb{E}[|R_{\min}| \cdot I_{\{\text{ruin at } P_k\}} | R_0 = x] \\ &= -\frac{1}{(1-\gamma)^k} \int_x^\infty \frac{e^{-\lambda(w-x)/(1-\gamma)}}{(k-1)!} \left[ \lambda(w-x) - \ln \frac{\phi_0(w)}{\phi_0(x)} \right]^{k-1} \ln \phi_0(w) \, dw. \end{aligned}$$

Furthermore, the expected maximum severity of the ruin excursion given that ruin occurs, is given by

$$\mathbb{E}[|R_{\min}| \mid \text{ruin occurs with } R_0 = x] = -\frac{\phi_\gamma(x)}{\psi_\gamma(x)} \int_x^\infty \frac{\ln \phi_\gamma(w)}{\phi_\gamma(w)} \, dw.$$

**Remark 2.3.** From the above formulas, it is straightforward to write down the probability that the ruin excursion stays above surplus level  $-d < 0$ , given that ruin occurs, as

$$\frac{A(x, 0) - A(x, d)}{\psi_\gamma(x)} = \frac{1}{\psi_\gamma(x)} \int_x^\infty \left[ \frac{\phi'_\gamma(w)}{\phi_\gamma(w)} - \frac{\phi'_\gamma(w+d)}{\phi_\gamma(w+d)} \right] \frac{\phi_\gamma(x)}{\phi_\gamma(w)} \, dw.$$

For the case without tax ( $\gamma = 0$ ), this formula can be compared with the following related classical formula for the distribution of the maximum severity  $M$  of ruin, where  $M$  is defined as the smallest value of the risk process between the time of ruin and the time of recovery to surplus level 0 (instead of the time that the running maximum is reached again):

$$P(M \leq d | R_0 = x \text{ and ruin occurs}) = \frac{\phi_0(x+d) - \phi_0(x)}{\phi_0(d)(1 - \phi_0(x))}$$

(see Picard [8]).

### 3. Relation with the $G/G/\infty$ queue

Consider the following situation. We have a sequence of pairs of random variables  $(X_1, Y_1), (X_2, Y_2), (X_3, Y_3), \dots$ , for which we want to calculate

$$\phi(x) = P \left( Y_i \leq x + \sum_{j=1}^i X_j \text{ for all } i = 1, 2, \dots \right). \quad (9)$$

As a first interpretation, the function  $\phi(x)$  is the survival probability in the risk model, if the  $X_i$ 's represent the increase of the surplus during periods in which the surplus process is in its running maximum (in the absence of tax payments, the  $X_i$ 's

equivalently represent the lengths of the periods during which the surplus process is in its running maximum) and the  $Y_i$ 's represent the maximal decreases of the surplus process in periods during which the surplus process is not in a profitable situation (i.e., the  $Y_i$ 's correspond to identically distributed copies of the random variable  $V_{\max}$ ).

A second interpretation of the function  $\phi(x)$  is as the steady-state probability that at an arrival instant in a  $G/G/\infty$  queue the residual service times of all the customers present in the system are less than  $x$ . Here, the  $X_i$ 's represent the interarrival times of the customers and the  $Y_i$ 's represent the service times of the customers.

Let us first assume that the random vectors  $(X_i, Y_i)$   $i = 1, 2, \dots$  are independent and identically distributed. Furthermore, assume that for each  $i = 1, 2, \dots$ , the random variables  $X_i$  and  $Y_i$  are independent.

**Remark 3.1.** These assumptions are satisfied in the Cramér-Lundberg risk model, where the claim arrival process is a Poisson process. However, when the claim arrival process is a general renewal process the random variables  $Y_i$  and  $X_{i+1}$  are dependent. In the related  $G/G/\infty$  queueing model this will mean that the service time of a customer depends on the previous interarrival time.

Let us denote by  $F(\cdot)$  the common distribution function of the random variables  $X_i$  (with corresponding probability density function  $f(\cdot)$ ). Furthermore, we denote by  $G(\cdot)$  the common distribution function of the random variables  $Y_i$ .

Conditioning on the value of  $X_1$  we obtain

$$\phi(x) = \int_{x_1=0}^{\infty} \phi(x + x_1)G(x + x_1)f(x_1)dx_1. \quad (10)$$

Iteration of this equation yields

$$\begin{aligned} \phi(x) &= \int_{x_1=0}^{\infty} \int_{x_2=0}^{\infty} \phi(x + x_1 + x_2)G(x + x_1 + x_2)G(x + x_1)f(x_2)f(x_1)dx_2dx_1 \\ &\quad \vdots \\ &= \lim_{M \rightarrow \infty} \int_{x_1=0}^{\infty} \dots \int_{x_M=0}^{\infty} \phi\left(x + \sum_{j=1}^M x_j\right) \prod_{i=1}^M \left\{ G\left(x + \sum_{j=1}^i x_j\right) f(x_i) \right\} dx_M \dots dx_1. \end{aligned}$$

**Example 3.1.** ( $X_i$ 's are deterministic.) If the  $X_i$ 's are deterministic, say  $X_i = w$ , we

have

$$\phi(x) = \phi(x+w)G(x+w) = \prod_{i=1}^{\infty} G(x+w \cdot i).$$

**Example 3.2.** ( $Y_i$ 's are deterministic.) If the  $Y_i$ 's are deterministic, say  $Y_i = v$ , we have

$$\phi(x) = \begin{cases} 1 & \text{for } x \geq v, \\ 1 - F(v-x) & \text{for } x < v. \end{cases}$$

**Example 3.3.** ( $X_i$ 's are exponential with parameter  $\lambda$ .) This is the case of the Cramér-Lundberg risk model. For an  $M/G/\infty$  queue it is well-known (see e.g. [9]) that the steady-state distribution of the number of customers is Poisson distributed and that the residual service times of the customers are all i.i.d. according to the excess lifetime distribution

$$G_e(x) := \frac{1}{\mathbb{E}[Y]} \int_0^x \bar{G}(y) dy.$$

Hence we find

$$\phi(x) = \sum_{n=0}^{\infty} \frac{(\lambda \mathbb{E}[Y])^n}{n!} e^{-\lambda \mathbb{E}[Y]} [G_e(x)]^n = e^{-\lambda \mathbb{E}[Y](1-G_e(x))} = e^{-\lambda \int_x^{\infty} \bar{G}(y) dy}, \quad (11)$$

which can be interpreted as yet another approach to establish formula (1). Of course, formula (11) can also be obtained from equation (10) which in this case takes the form

$$\phi(x) = \lambda \int_0^{\infty} \phi(x+x_1)G(x+x_1)e^{-\lambda x_1} dx_1.$$

Introducing  $T(x) := e^{-\lambda x} \phi(x)$  yields

$$T(x) = \lambda \int_x^{\infty} T(u)G(u) du,$$

which gives  $T'(x) = -\lambda G(x)T(x)$ . It follows that  $T(x) = Ce^{-\lambda \int_0^x G(y) dy}$ , so that  $\phi(x) = Ce^{\lambda \int_0^x \bar{G}(y) dy}$  with  $C$  some constant yet to be determined. Letting  $x \rightarrow \infty$ , we find  $C = e^{-\lambda \int_0^{\infty} \bar{G}(y) dy}$ , and hence  $\phi(x) = e^{-\lambda \int_x^{\infty} \bar{G}(y) dy}$ .

**Example 3.4.** ( $Y_i$ 's are exponential with parameter  $\nu$ .) For a  $G/M/\infty$  queue it is well-known (see e.g. [9]) that the steady-state probability for an arriving customer to find  $n$  customers in the system is given by

$$p_n = \sum_{r=n}^{\infty} (-1)^{r-n} \binom{r}{n} B_r,$$

where  $B_r$  is given by

$$B_r = \prod_{i=1}^r \left( \frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} \right)$$

and  $\tilde{F}(s)$  is the Laplace-Stieltjes transform of the interarrival time distribution. Exploiting the lack-of-memory property of the exponential distribution, we hence have

$$\begin{aligned} \phi(x) &= \sum_{n=0}^{\infty} p_n (1 - e^{-\nu x})^n \\ &= \sum_{n=0}^{\infty} \sum_{r=n}^{\infty} (-1)^{r-n} \binom{r}{n} \prod_{i=1}^r \left( \frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} \right) (1 - e^{-\nu x})^n \\ &= \sum_{r=0}^{\infty} \prod_{i=1}^r \left( \frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} \right) \sum_{n=0}^r \binom{r}{n} (-1)^{r-n} (1 - e^{-\nu x})^n \\ &= \sum_{r=0}^{\infty} \left( \prod_{i=1}^r \left( \frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} \right) \right) (-e^{-\nu x})^r. \end{aligned}$$

In the special case that the interarrival times are exponential as well (with parameter  $\lambda$ ), we have

$$\frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} = \frac{\lambda}{i\nu}$$

and correspondingly

$$\phi(x) = \sum_{r=0}^{\infty} \prod_{i=1}^r \left( \frac{\lambda}{i\nu} \right) (-e^{-\nu x})^r = \sum_{r=0}^{\infty} \left( -\frac{\lambda}{\nu} e^{-\nu x} \right)^r / r! = e^{-\frac{\lambda}{\nu} e^{-\nu x}} = e^{-\lambda \int_x^{\infty} e^{-\nu y} dy} \quad (12)$$

as before.

If on the other hand the interarrival times are Erlang(2,  $\lambda$ ) distributed, we have

$$\frac{\tilde{F}(i\nu)}{1 - \tilde{F}(i\nu)} = \frac{\lambda^2}{(i\nu)^2 + 2\lambda i\nu}$$

and consequently

$$\phi(x) = \sum_{r=0}^{\infty} \prod_{i=1}^r \left( \frac{\lambda^2}{(i\nu)^2 + 2\lambda i\nu} \right) (-e^{-\nu x})^r = \sum_{r=0}^{\infty} \left( \frac{\lambda}{\nu} \right)^{2r} \frac{1}{r!} \prod_{i=1}^r \left( \frac{1}{i + 2\frac{\lambda}{\nu}} \right) (-e^{-\nu x})^r.$$

Introducing  $\alpha = 2\lambda/\nu$  and using

$$\prod_{i=1}^r \left( \frac{1}{i + \alpha} \right) = \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha + r + 1)}$$

gives

$$\phi(x) = \Gamma(\alpha + 1) \sum_{r=0}^{\infty} \frac{\left[-\left(\frac{\lambda}{\nu}\right)^2 e^{-\nu x}\right]^r}{r! \Gamma(\alpha + r + 1)} = \frac{\Gamma(\alpha + 1)}{\left(\frac{\lambda}{\nu} e^{-\nu x/2}\right)^\alpha} \cdot J_\alpha\left(\alpha e^{-\nu x/2}\right) \quad (13)$$

where  $J_\alpha(\cdot)$  is the Bessel function of the first kind, defined by

$$J_\alpha(x) = \sum_{r=0}^{\infty} \frac{(-1)^r}{r! \Gamma(r + \alpha + 1)} \left(\frac{x}{2}\right)^{2r + \alpha}.$$

Formula (13) can also be obtained via equation (10): Plugging  $f(x) = \lambda^2 x e^{-\lambda x}$  and  $G(x) = 1 - e^{-\nu x}$  into (10), differentiating twice yields

$$(e^{-\lambda x} \phi(x))'' = \lambda^2 \phi(x) (1 - e^{-\nu x})$$

or equivalently

$$\phi''(x) - 2\lambda\phi'(x) + \lambda^2 e^{-\nu x} \phi(x) = 0.$$

This ordinary differential equation has the solution

$$\phi(x) = \left(\nu e^{\nu x/2} / \lambda\right)^\alpha \left[ C_1 \Gamma(1 + \alpha) J_\alpha(\alpha e^{-x\nu/2}) + C_2 \Gamma(1 - \alpha) J_{-\alpha}(\alpha e^{-x\nu/2}) \right],$$

where  $C_1, C_2$  are constants and again  $\alpha = 2\frac{\lambda}{\nu}$ . The boundary condition  $\lim_{x \rightarrow \infty} \phi(x) = 1$  then gives  $C_2 = 0$  and  $C_1 = 1$ , hence (13).

It is interesting to examine the asymptotic behavior of  $\phi(x_\lambda)$ , with  $x_\lambda := \kappa + \frac{1}{\nu} \log \lambda$ , as  $\lambda \rightarrow \infty$ . It is easily verified that

$$\begin{aligned} \lim_{\lambda \rightarrow \infty} \phi(x_\lambda) &= \lim_{\lambda \rightarrow \infty} \sum_{r=0}^{\infty} \left(\frac{\lambda}{\nu}\right)^{2r} \frac{1}{r!} \prod_{i=1}^r \left(\frac{1}{i + 2\frac{\lambda}{\nu}}\right) \left(\frac{-e^{-\kappa\nu}}{\lambda}\right)^r \\ &= \sum_{r=0}^{\infty} \frac{1}{r!} \left(\frac{-e^{-\kappa\nu}}{2\nu}\right)^r = e^{-\frac{1}{2}e^{-\kappa\nu}/\nu}. \end{aligned}$$

Note that this limit is the same as the value of  $\phi(x_\lambda)$  in the case of exponential interarrival times with parameter  $\lambda/2$  (cf. (12)).

#### 4. An asymptotic result for renewal risk models with tax

Assume that potential ‘catastrophes’ occur according to a delayed renewal process with initial delay  $T_0$  and interrenewal periods  $T_1, T_2, \dots$ . At time  $S_n := T_0 + \dots + T_n$ , an actual catastrophe occurs if  $V_n$  exceeds  $f(S_n)$ , with  $f(\cdot)$  some increasing function, and

$V_0, V_1, V_2, \dots$  a sequence of independent and identically distributed random variables. The random variables  $T_{n+1}$  and  $V_n$  may be dependent. Let the 0–1 variable  $I_n := \mathbb{I}_{\{V_n > f(S_n)\}}$  indicate whether or not an actual catastrophe occurs at time  $S_n$ , and denote

$$p(t) := \mathbb{P}\{V_n > f(t)\}.$$

We are interested in the probability of the event  $E_\tau$  that no actual catastrophe occurs during the time interval  $[0, \tau]$ , i.e.,

$$E_\tau = \cup_{n=-1}^{\infty} \{S_n \leq \tau < S_{n+1}; I_0 = \dots = I_n = 0\},$$

with the notational convention that  $S_{-1} := 0$ .

Now consider the surplus process in the Sparre Andersen risk model where claims of generic size  $Y$  occur according to a renewal process with generic interrenewal time  $X$ , and a marginal tax rate  $\gamma$  applies whenever the free surplus is at a running maximum. Let  $Q$  be a single-server queue with generic interarrival time  $X$  and generic service time  $Y$ . Let  $V_{\max}$  and  $T$  be a pair of random variables whose joint distribution is that of the maximum workload during a busy period of  $Q$  and the subsequent idle period. Further suppose that we take the joint distribution of  $T_{n+1}$  and  $V_n$  to be that of  $T$  and  $V_{\max}$ , and  $f(t) = x + (1 - \gamma)t$ . Then the probability of the event  $E_\tau$  with  $\tau = (v - x)/(1 - \gamma)$  equals the probability  $\phi_\gamma(x, v)$  that the surplus process reaches level  $v$ , starting from level  $x$ , before ruin occurs. In particular, the survival probability in the renewal model with tax is  $\phi_\gamma(x) = \mathbb{P}\{E_\infty\}$ , with  $E_\infty = \{V_n \leq x + (1 - \gamma)S_n \text{ for all } n = 0, 1, 2, \dots\}$ .

**Remark 4.1.** Following Section 3, the probability of the event  $E_\infty$  may also be interpreted as the probability that no customer with a remaining service time exceeding  $x$  is present in a  $G/G/\infty$  system where the joint distribution of the interarrival time and subsequent service time is that of  $(1 - \gamma)T_{n+1}$  and  $V_n$ , given that the past interarrival time is  $T_0$ .

In order to characterize the probability of interest, i.e.,  $\mathbb{P}\{E_\tau\}$ , we will consider a scenario where the interrenewal periods are relatively short (compared to the time interval  $[0, \tau]$ ), i.e., the number of potential catastrophes is relatively large, whereas the

probability that an actual catastrophe occurs is relatively small, such that the value of the ratio  $p(t)/\mathbb{E}\{T\}$  is moderate. More specifically, we assume an asymptotic regime where time is accelerated by a factor  $s$ , i.e., with interrenewal periods  $T^{(s)} := T/s$ , while the function  $f^{(s)}(\cdot)$  is simultaneously boosted in such a manner that the ratio  $p^{(s)}(t)/\mathbb{E}\{T^{(s)}\} = p(t)/\mathbb{E}\{T\}$ , i.e.,  $p^{(s)}(t) = p(t)/s$ . For each fixed value of  $s$ , denote the resulting event  $E_\tau$  by  $E_\tau^{(s)}$ .

The next theorem states the main result of this section.

**Theorem 4.1.** *Under the above-mentioned assumptions,*

$$\mathbb{P}\left\{E_\tau^{(s)}\right\} \rightarrow \exp\left(-\lambda \int_{t=0}^{\tau} p(t)dt\right) \quad (14)$$

as  $s \rightarrow \infty$ , with  $\lambda := 1/\mathbb{E}\{T\}$ .

**Remark 4.2.** Theorem 4.1 suggests that the expression on the right-hand side should provide a reasonable approximation for  $\mathbb{P}\left\{E_\tau^{(s)}\right\}$  in the above-described asymptotic regime where the interrenewal periods are relatively short compared to the time interval  $[0, \tau]$ . Note that (14) has a similar form as the earlier result (1) for the Cramér-Lundberg risk process.

In order to prove Theorem 4.1, we will establish lower and upper bounds for the unscaled process. Lemmas 4.2 and 4.4 below will show that these two bounds, while crude, coincide in the asymptotic regime under consideration.

For ease of notation, we henceforth drop the subscript  $\tau$  from the notation  $E_\tau^{(s)}$ , and simply write  $E^{(s)}$  or just  $E$ . Note that

$$\lim_{K \rightarrow \infty} \frac{\tau}{K} \sum_{k=1}^K p\left(k \frac{\tau}{K}\right) = \lim_{K \rightarrow \infty} \frac{\tau}{K} \sum_{k=1}^K p\left((k-1) \frac{\tau}{K}\right) = \int_{t=0}^{\tau} p(t)dt. \quad (15)$$

Let us now focus on the lower bound. Let  $K \geq 1$  and  $N \geq 1$  be integers and  $t_0 = 0 \leq t_1 \leq \dots \leq t_K = \tau$ . For any  $k = 1, \dots, K$ , define the events

$$D_k := \{S_{kN} > t_k\},$$

$$F_k := \{V_{(k-1)N} \leq f(t_{k-1}), \dots, V_{kN-1} \leq f(t_{k-1})\},$$

and

$$E^{\text{lower}} := \bigcap_{k=1}^K D_k \cap \bigcap_{k=1}^K F_k.$$

**Lemma 4.1.** *The event  $E^{\text{lower}}$  implies the event  $E$ .*

*Proof.* Suppose that the event  $E^{\text{lower}}$  occurs, i.e., all the events  $D_k$  and  $F_k$  occur. Let  $i$  be such that  $(k-1)N \leq i \leq kN-1$  for some  $k=1, \dots, K$ . The event  $D_k$  gives  $S_i \geq S_{(k-1)N} > t_{k-1}$ , while the event  $F_k$  implies  $V_i \leq f(t_{k-1})$ . Since the function  $f(\cdot)$  is increasing, it follows that  $V_i \leq f(S_i)$ . Hence  $I_i = 0$  for all  $i=0, \dots, KN-1$ . The event  $D_K$  implies that there exists an  $n \leq KN-1$  with  $S_n \leq \tau < S_{n+1}$ . Thus the event  $E$  occurs.  $\square$

**Lemma 4.2.**

$$\lim_{s \rightarrow \infty} \mathbb{P} \left\{ E^{(s)} \right\} \geq e^{-\lambda \int_{t=0}^{\tau} p(t) dt}. \quad (16)$$

*Proof.* Lemma 4.1 yields that

$$\begin{aligned} \mathbb{P} \{E\} &\geq \mathbb{P} \{E^{\text{lower}}\} = \mathbb{P} \left\{ \bigcap_{k=1}^K D_k \cap \bigcap_{k=1}^K F_k \right\} \geq \mathbb{P} \left\{ \bigcap_{k=1}^K F_k \right\} - \mathbb{P} \left\{ \overline{\bigcap_{k=1}^K D_k} \right\} \\ &\geq \prod_{k=1}^K \mathbb{P} \{F_k\} - \sum_{k=1}^K \mathbb{P} \{\overline{D_k}\} \\ &= \prod_{k=1}^K \mathbb{P} \{V_{(k-1)N} \leq f(t_{k-1}), \dots, V_{kN-1} \leq f(t_{k-1})\} - \sum_{k=1}^K \mathbb{P} \{S_{kN} \leq t_k\} \\ &= \prod_{k=1}^K (\mathbb{P} \{V \leq f(t_{k-1})\})^N - \sum_{k=1}^K \mathbb{P} \{S_{kN} \leq t_k\}. \end{aligned}$$

Choose now  $N = \lceil N(s) \rceil$ , with  $N(s) = (1 + \epsilon) \frac{\tau s}{K \mathbb{E}\{T\}}$ , and  $t_k = \frac{k\tau}{K}$ ,  $k=1, \dots, K$ .

Then

$$\begin{aligned} \mathbb{P} \{S_{kN} \leq t_k\} &= \mathbb{P} \left\{ T_0/s + T_1/s + \dots + T_{\lceil N(s) \rceil} / s \leq \frac{k\tau}{K} \right\} \\ &= \mathbb{P} \left\{ T_0 + T_1 + \dots + T_{\lceil N(s) \rceil} \leq \frac{kN(s) \mathbb{E}\{T\}}{1 + \epsilon} \right\}, \end{aligned}$$

which by the law of large numbers tends to zero as  $s \rightarrow \infty$ . Also,

$$\begin{aligned} \lim_{s \rightarrow \infty} \prod_{k=1}^K (\mathbb{P} \{V < f(t_{k-1})\})^{N(s)} &= \lim_{s \rightarrow \infty} \prod_{k=1}^K e^{-N(s)p^{(s)}(t_{k-1})} = e^{-\sum_{k=1}^K \lim_{s \rightarrow \infty} N(s)p^{(s)}(t_{k-1})} \\ &= e^{-\sum_{k=1}^K \frac{\tau p(t_{k-1})}{K \mathbb{E}\{T\}}} = e^{-\frac{\tau}{K \mathbb{E}\{T\}} \sum_{k=1}^K p(t_{k-1})}. \end{aligned}$$

We deduce that

$$\lim_{s \rightarrow \infty} \mathbb{P} \left\{ E^{(s)} \right\} \geq e^{-\frac{\tau}{K \mathbb{E}\{T\}} \sum_{k=1}^K p(t_{k-1})}$$

for any  $K \geq 1$ . Letting  $K \rightarrow \infty$  and applying (15), we obtain the lower bound (16).  $\square$

Next, we establish an upper bound that asymptotically matches the lower bound. Let  $K \geq 1$  and  $N \geq 1$  be integers and  $t_0 = 0 \leq t_1 \leq \dots \leq t_K = \tau$ . For any  $k = 1, \dots, K$ , define the events

$$G_k := \{V_{(k-1)N} \leq f(t_k), \dots, V_{kN-1} \leq f(t_k)\},$$

and

$$E^{\text{upper}} := \bigcup_{k=1}^K D_k \cup \bigcap_{k=1}^K G_k.$$

**Lemma 4.3.** *The event  $E$  implies the event  $E^{\text{upper}}$ .*

*Proof.* Suppose that the event  $E$  occurs, i.e., there exist an  $n(\tau)$  with  $S_{n(\tau)} \leq \tau < S_{n(\tau)+1}$  and  $I_0 = \dots = I_{n(\tau)} = 0$ . Also assume that all the events  $\bar{D}_k$  occur, i.e.,  $S_{kN} \leq t_k$  for all  $k = 1, \dots, K$ , because otherwise there is nothing to prove. This in particular implies that  $n(\tau) \geq KN - 1$ , and hence  $I_0 = \dots = I_{KN-1} = 0$ , i.e.,  $V_i \leq f(S_i)$  for all  $i = 0, \dots, KN - 1$ . Let  $i$  be such that  $(k-1)N \leq i \leq kN - 1$  for some  $k = 1, \dots, K$ , so that  $S_i \leq S_{kN}$ . Since the function  $f(\cdot)$  is increasing, it follows that  $V_i \leq f(t_k)$ , and thus all the events  $G_k$  occur, and hence the event  $E^{\text{upper}}$  occurs.  $\square$

**Lemma 4.4.**

$$\lim_{s \rightarrow \infty} \mathbb{P} \left\{ E^{(s)} \right\} \leq e^{-\lambda \int_{t=0}^{\tau} p(t) dt}. \quad (17)$$

*Proof.* Lemma 4.3 yields that

$$\begin{aligned} \mathbb{P} \{E\} &\leq \mathbb{P} \{E^{\text{upper}}\} \\ &= \mathbb{P} \left\{ \bigcup_{k=1}^K D_k \cup \bigcap_{k=1}^K G_k \right\} \\ &\leq \mathbb{P} \left\{ \bigcap_{k=1}^K G_k \right\} + \mathbb{P} \left\{ \bigcup_{k=1}^K D_k \right\} \\ &\leq \prod_{k=1}^K \mathbb{P} \{G_k\} + \sum_{k=1}^K \mathbb{P} \{D_k\} \\ &= \prod_{k=1}^K (\mathbb{P} \{V \leq f(t_k)\})^N + \sum_{k=1}^K \mathbb{P} \{S_{kN} > t_k\}. \end{aligned} \quad (18)$$

We now take  $N = \lceil N(s) \rceil$ , with  $N(s) = (1 - \epsilon) \frac{\tau s}{K \mathbb{E}\{T\}}$ , and  $t_k = \frac{k\tau}{K}$ ,  $k = 1, \dots, K$ , and proceed to evaluate the upper bound (18) in the asymptotic regime of interest. Note that

$$\begin{aligned} \mathbb{P}\{S_{kN} > t_k\} &= \mathbb{P}\left\{T_0/s + T_1/s + \dots + T_{\lceil N(s) \rceil}/s > \frac{k\tau}{K}\right\} \\ &= \mathbb{P}\left\{T_0 + T_1 + \dots + T_{\lceil N(s) \rceil} > \frac{kN(s)\mathbb{E}\{T\}}{1 - \epsilon}\right\}, \end{aligned}$$

which tends to zero as  $s \rightarrow \infty$  because of the law of large numbers. Also,

$$\begin{aligned} \lim_{s \rightarrow \infty} \prod_{k=1}^K (\mathbb{P}\{V \leq f(t_k)\})^{N(s)} &= \lim_{s \rightarrow \infty} \prod_{k=1}^K e^{-N(s)p^{(s)}(t_k)} = e^{-\sum_{k=1}^K \lim_{s \rightarrow \infty} N(s)p^{(s)}(t_k)} \\ &= e^{-\sum_{k=1}^K \frac{\tau p(t_k)}{K \mathbb{E}\{T\}}} = e^{-\frac{\tau}{K \mathbb{E}\{T\}} \sum_{k=1}^K p(t_k)}. \end{aligned}$$

We conclude that

$$\lim_{s \rightarrow \infty} \mathbb{P}\{E^{(s)}\} \leq e^{-\frac{\tau}{K \mathbb{E}\{T\}} \sum_{k=1}^K p(t_k)}$$

for any  $K \geq 1$ . Letting  $K \rightarrow \infty$  and invoking (15), we obtain the upper bound (17).

□

### Acknowledgement

H. Albrecher would like to acknowledge support from the Swiss National Science Foundation Project 200021-124635/1.

### References

- [1] H. Albrecher, A. Badescu and D. Landriault (2008) On the dual risk model with taxation. *Insurance: Mathematics and Economics* **42**, 1086–1094.
- [2] H. Albrecher, S.C. Borst, O.J. Boxma and J.A.C. Resing (2009) The tax identity in risk theory – a simple proof and an extension. *Insurance: Mathematics and Economics* **44**, 304–306.
- [3] H. Albrecher and C. Hipp (2007) Lundberg’s risk process with tax. *Blätter der DGVMF* **28**, 13–28.
- [4] H. Albrecher, J. Renaud and X. Zhou (2008) A Lévy insurance risk process with tax. *Journal of Applied Probability* **45**, 363–375.
- [5] S. Asmussen and H. Albrecher. *Ruin Probabilities*. 2nd edn. World Scientific, New Jersey, 2010.

- [6] J.M. Harrison and S.I. Resnick (1976) The stationary distribution and first exit probabilities of a storage process with general release rule. *Mathematics of Operations Research* **1**, 347–358.
- [7] A. Kyprianou and X. Zhou (2009) General tax structures and the Lévy insurance risk model. *Journal of Applied Probability* **46**, 1146–1156.
- [8] P. Picard (1994) On some measures of the severity of ruin in the classical Poisson model. *Insurance: Mathematics and Economics* **14**, 107–115.
- [9] L. Takács (1962) *Introduction to the Theory of Queues*. Oxford University Press, New York.
- [10] J. Wei, H. Yang and R. Wang (2010) On the Markov-modulated insurance risk model with tax. *Blätter der DGVMF* **31**, 65–78.