



UNIL | Université de Lausanne

Unicentre

CH-1015 Lausanne

<http://serval.unil.ch>

Year : 2016

Attitude des ambulanciers face à la vaccination antigrippale

MOSER Alexandre

MOSER Alexandre, 2016, Attitude des ambulanciers face à la vaccination antigrippale

Originally published at : Thesis, University of Lausanne

Posted at the University of Lausanne Open Archive <http://serval.unil.ch>

Document URN : urn:nbn:ch:serval-BIB_CA1D1F960B221

Droits d'auteur

L'Université de Lausanne attire expressément l'attention des utilisateurs sur le fait que tous les documents publiés dans l'Archive SERVAL sont protégés par le droit d'auteur, conformément à la loi fédérale sur le droit d'auteur et les droits voisins (LDA). A ce titre, il est indispensable d'obtenir le consentement préalable de l'auteur et/ou de l'éditeur avant toute utilisation d'une oeuvre ou d'une partie d'une oeuvre ne relevant pas d'une utilisation à des fins personnelles au sens de la LDA (art. 19, al. 1 lettre a). A défaut, tout contrevenant s'expose aux sanctions prévues par cette loi. Nous déclinons toute responsabilité en la matière.

Copyright

The University of Lausanne expressly draws the attention of users to the fact that all documents published in the SERVAL Archive are protected by copyright in accordance with federal law on copyright and similar rights (LDA). Accordingly it is indispensable to obtain prior consent from the author and/or publisher before any use of a work or part of a work for purposes other than personal use within the meaning of LDA (art. 19, para. 1 letter a). Failure to do so will expose offenders to the sanctions laid down by this law. We accept no liability in this respect.

UNIVERSITE DE LAUSANNE - FACULTE DE BIOLOGIE ET DE MEDECINE

Département des Centres Interdisciplinaires et de Logistique Médicale - DCILM
Service des Urgences

Attitude des ambulanciers face à la vaccination antigrippale

THESE

préparée sous la direction du Docteur Pierre-Nicolas CARRON, PD & MER
avec la co-direction du Professeur Bertrand YERSIN

et présentée à la Faculté de biologie et de médecine de
l'Université de Lausanne pour l'obtention du grade de

DOCTEUR EN MEDECINE

par

Alexandre MOSER

Médecin diplômé de la Confédération Suisse
Originaire de Landiswil (BE)

Lausanne

2016

Imprimatur

Vu le rapport présenté par le jury d'examen, composé de

Directeur de thèse *Monsieur le Docteur Pierre Carron*
Co-Directeur de thèse *Monsieur le Professeur Bertrand Yersin*
Expert *Monsieur le Professeur John-Paul Vader*
Directeur de l'Ecole *Monsieur le Professeur Niko Geldner*
doctorale

la Commission MD de l'Ecole doctorale autorise l'impression de la thèse de

Monsieur Alexandre Moser

intitulée

Attitude des ambulanciers face à la vaccination anti-grippale

Lausanne, le 19 avril 2016

*pour Le Doyen
de la Faculté de Biologie et de Médecine*


*Monsieur le Professeur John Prior
Vice-Directeur de l'Ecole doctorale*

Vaccination Against Seasonal or Pandemic Influenza in Emergency Medical Services

Alexandre Moser, MD;¹ Cédric Mabire, PhD;² Olivier Hugli, MD, MPH;³ Victor Dorribo, MD;⁴ Giorgio Zanetti, MD;^{5,6} Catherine Lazor-Blanchet, MD;⁶ Pierre-Nicolas Carron, MD³

1. Department of Internal Medicine, Hospital of Thun and Simmental (STS AG), Zweisimmen, Switzerland
2. University of Health Sciences (HESAV), Lausanne, Switzerland
3. Emergency Department, Lausanne University Hospital, Lausanne, Switzerland
4. Service of Occupational Medicine, Institute for Work and Health, Lausanne, Switzerland
5. Service of Infectious Diseases, Lausanne University Hospital, Lausanne, Switzerland
6. Service of Hospital Preventive Medicine, Lausanne University Hospital, Lausanne, Switzerland

Correspondence:

Alexandre Moser
Department of Internal Medicine
Hospital of Thun and Simmental (STS AG)
Karl Haueter-Strasse 21
CH – 3770 Zweisimmen
Switzerland
E-mail: alexandre.moser@h-ne.ch

Conflicts of interest: none

Keywords: Emergency Medical Services; EMS-workers; influenza; vaccination

Abbreviation:

EMS: Emergency Medical Service
EMS-W: Emergency Medical Service workers
HCW: health care workers
PI: pandemic influenza
PPE: personal protective equipment
SI: seasonal influenza

Received: August 11, 2015

Accepted: October 14, 2015

doi:10.1017/S1049023X16000121

Abstract

Introduction: Influenza is a major concern for Emergency Medical Services (EMS); EMS workers' (EMS-Ws) vaccination rates remain low despite promotion. Determinants of vaccination for seasonal influenza (SI) or pandemic influenza (PI) are unknown in this setting.

Hypothesis: The influence of the H1N1 pandemic on EMS-W vaccination rates, differences between SI and PI vaccination rates, and the vaccination determinants were investigated.

Methods: A survey was conducted in 2011 involving 65 Swiss EMS-Ws. Socio-professional data, self-declared SI/PI vaccination status, and motives for vaccine refusal or acceptance were collected.

Results: Response rate was 95%. The EMS-Ws were predominantly male ($n = 45$; 73%), in good health (87%), with a mean age of 36 (SD = 7.7) years. Seventy-four percent had more than six years of work experience. Self-declared vaccination rates were 40% for both SI and PI (PI + /SI +), 19% for PI only (PI + /SI -), 1.6% for SI only (PI - /SI +), and 39% were not vaccinated against either (PI - /SI -). Women's vaccination rates specifically were lower in all categories but the difference was not statistically significant. During the previous three years, 92% of PI + /SI + EMS-Ws received at least one SI vaccination; it was 8.3% in the case of PI - /SI - ($P = .001$) and 25% for PI + /SI - ($P = .001$). During the pandemic, SI vaccination rate increased from 26% during the preceding year to 42% ($P = .001$). Thirty percent of the PI + /SI + EMS-Ws declared that they would not get vaccination next year, while this proportion was null for the PI - /SI - and PI + /SI - groups. Altruism and discomfort induced by the surgical mask required were the main motivations to get vaccinated against PI. Factors limiting PI or SI vaccination included the option to wear a mask, avoidance of medication, fear of adverse effects, and concerns about safety and effectiveness.

Conclusion: Average vaccination rate in this study's EMS-Ws was below recommended values, particularly for women. Previous vaccination status was a significant determinant of PI and future vaccinations. The new mask policy seemed to play a dual role, and its net impact is probably limited. This population could be divided in three groups: favorable to all vaccinations; against all, even in a pandemic context; and ambivalent with a "pandemic effect." These results suggest a consistent vaccination pattern, only altered by exceptional circumstances.

Moser A, Mabire C, Hugli O, Dorribo V, Zanetti G, Lazor-Blanchet C, Carron PN. Vaccination against seasonal or pandemic influenza in Emergency Medical Services. *Prehosp Disaster Med.* 2016;31(2):1-8.

Introduction

In the United States, seasonal influenza (SI) is responsible for over 36,000 deaths and 134,000 hospitalizations.¹ Influenza is a major concern for Emergency Medical Services (EMS).² Since infected persons can spread the virus before signs of illness develop, exposure of EMS workers (EMS-Ws) can contribute to the spread of influenza into the community, resulting in increased morbidity and mortality, particularly among vulnerable patients.^{2,3} Nevertheless, most EMS-Ws underestimate this risk or are not aware of the recommendations for prevention.^{2,4} Furthermore, compliance of EMS-Ws with infection control recommendations is frequently suboptimal, particularly when it comes to the use of personal protective equipment (PPE), like surgical masks.⁵⁻⁸

Influenza vaccination is both a personal and an altruist intervention, limiting the spread of the disease and reducing morbidity and mortality in health care workers (HCWs) and in the community.⁹⁻¹² Influenza vaccination prevents between 70% and 90% of influenza cases¹³ and is considered cost-effective.^{14,15} Therefore, vaccination has long been recommended for all HCWs – including EMS-Ws involved in prehospital direct patient care¹⁶ – and has become mandatory in some settings.^{17,18} Nevertheless, vaccination rates among HCWs in acute care services remain low: between 30% to 65% among emergency department HCWs^{2,19} and usually between 20% and 50% for prehospital EMS-Ws.^{2,4} Vaccination is also recommended for Swiss EMS-Ws.²⁰ Since 2009, the vaccination strategy was associated with a dedicated new mandatory mask wearing policy requiring that unvaccinated EMS-Ws wear a surgical mask during patient care duties.²¹

In April 2009, swine influenza H1N1 was identified in North America. On April 29, the World Health Organization (WHO; Geneva, Switzerland) raised its pandemic alert level to Phase 5 (imminent pandemic)²² and the pandemic state was finally declared on June 11, 2009. In three months, more than 94,000 cases were confirmed worldwide.²³ The H1N1 pandemic influenza (PI) was responsible for serious complications and increased mortality, particularly among children and healthy young adults.²⁴

Only a few studies about vaccination rates have focused on EMS-Ws and most of these studies were prior to the H1N1 pandemic.^{2,4} Differences in vaccination rates for SI vs PI have been described in some studies,^{4,19,25} but variations and determinants of vaccination for either of these types of influenza aren't well-known in the prehospital EMS setting (Table 1). The main objective of this study was to observe if EMS-W vaccination rates had been influenced by the H1N1 pandemic and if vaccination rates were different for PI and SI. A second aim also was to identify potential specific factors promoting or limiting PI and SI vaccination rates.

Methods

Setting

This questionnaire-based survey was conducted from December 1, 2010 through January 21, 2011 in the two prehospital EMS sites serving the city of Lausanne, located in the French-speaking part of Switzerland, with a population of 130,000. The clinical guidelines, rules of operations, and safety regulations are enacted by the State authorities and are the same for the two EMS sites. For the purpose of analysis, the data of the two EMS were pooled.

According to the international guidelines, during the period under analysis, the local recommendations concerning influenza for EMS-Ws was to get vaccinated²⁰ and to wear a surgical mask as PPE during the two weeks after vaccination or, if vaccination was declined by the HCW, to wear a mask at all times when tending to patients.²¹

Study Design

A standardized, multiple-choice, anonymous questionnaire (Appendix 1; available online only) was developed to collect demographic data, self-declared vaccination status against the 2009-2010 SI and/or the 2009 PI, and to evaluate the factors associated with vaccination against the 2009-2010 SI and/or the 2009 PI. The questions were based on a previous standardized questionnaire addressing the intra-hospital determinants of vaccination,²⁵ and on a review of the literature for specific questions related to EMS and EMS-Ws. Motives for accepting or refusing vaccination were assessed by a Likert

Predictors of Vaccination Acceptance among EMS-Ws
Working in one urban EMS
"No waiting" vaccination at work and free vaccination
Painless vaccination (eg, nasal spray)
If bird flu was found in the country
Previous influenza infection
Perception of increased risk for contracting the flu
Believe in vaccine effectiveness
Employer vaccine recommendation
Age (older than 36 years old)
Obligation to use of PPE (mask) to limit exposure
Continuous educational campaign
Having family members who relied on them for support
Predictors of Vaccine Refusal among EMS-Ws
Concern about vaccine effectiveness
Concern about side effects
Allergy
Belief in personal health as a protector
Lack of employer mandate or recommendation
Trivialization of flu
No previous influenza vaccination
Concern about having the flu by getting vaccinated
Working in a rural EMS or for a fire-based service

Moser © 2016 Prehospital and Disaster Medicine

Table 1. Identified Vaccination Determinants of EMS-Ws^{2,4,29,30}

Abbreviations: EMS, Emergency Medical Services; EMS-W, EMS workers.

scale, with results expressed as a mean on a 10-point scale. Questions assessing motivations were analyzed according to self-reported vaccinations status. All paramedics of the two EMS were asked to complete the survey, regardless of their level of certification or level of employment (part-time vs full-time). Study participation was voluntary, and all participants signed an informed consent. The Lausanne University Ethics Committee (Lausanne, Switzerland) and the officials from both EMS sites approved the protocol.

Statistics

The motives for acceptance or refusal of vaccination were dichotomized for the purpose of analysis (claimed motives on one side, unclaimed motives and neutral opinion pooled on the other side). Continuous data are expressed as mean and standard deviation (SD), whereas categorical data are presented as numbers and percentages. Differences between vaccination groups in normally distributed data and independent continuous variable

	PI + / SI + (n = 25)	PI + / SI - (n = 12)	PI - / SI + (n = 1)	PI - / SI - (n = 24)	P Value ^a
Age, mean (SD)	36.9 (8.1)	36.0 (5.4)	22.0	36.8 (8.6)	.94 ^b
Women	4 (6.5%)	2 (3.2%)	0 (0%)	11 (18%)	.06 ^c
Work Experience					
0 – 3 years	3 (4.8%)	1 (1.6%)	1 (1.6%)	4 (6.5%)	.99 ^c
4 – 6 years	3 (4.8%)	1 (1.6%)	0 (0%)	3 (4.8%)	
6 – 9 years	3 (4.8%)	2 (3.25%)	0 (0%)	4 (6.5%)	
>9 years	16 (26%)	8 (13%)	0 (0%)	13 (21%)	
Employed during Pandemics	25 (40%)	11 (18%)	0 (0%)	22 (35%)	.19 ^c
Chronically Ill	5 (8.1%)	0 (0%)	1 (1.6%)	2 (3.2%)	.22 ^c
Living with a Person at Risk of Complicated SI	1 (1.6%)	0 (0%)	0 (0%)	1 (1.6%)	.99 ^c
Living with a Child Under 2 or a Pregnant Women	5 (8.1%)	4 (6.5%)	0 (0%)	3 (4.8%)	.34 ^c
Pregnancy (n, % of women EMS-Ws)	0 (0%)	0 (0%)	0 (0%)	2 (12%)	.66 ^c

Moser © 2016 Prehospital and Disaster Medicine

Table 2. Characteristics of the Responders

Abbreviations: EMS-W, Emergency Medical Services worker; PI, pandemic influenza; SI, seasonal influenza.

^aGroup PI-/SI+ was excluded from analysis.^bANOVA oneway with Bonferroni test.^cFisher's exact test.

were assessed using Student's *t*-test or using one-way analysis of variance (ANOVA). In skewed data, the Mann-Whitney test was used, and for dependent continuous variables, the Wilcoxon rank sum test was used. Differences in proportion for categorical variables were analyzed using Fischer's exact test. Differences for all analyses were considered statistically significant at two-tailed $P < .05$. Analysis was undertaken using Stata 13.1 (StataCorp; College Station, Texas USA).

Results

Population Description and Vaccination Rates

Sixty-five EMS-Ws were included in the study, with a 95% response rate (62/65). The population (Table 2) was composed of a majority of men (72%), with more than six years of work experience in 74%. Only 6.5% were not employed during the 2009 PI season. A minority of the EMS-Ws (13%) declared suffering from a chronic illness. While only two EMS-Ws (3.2%) declared living with a person at risk for an influenza complication, 19% ($n = 12$) lived with a child under two years of age or with a pregnant woman at the time of the PI season. Two female EMS-Ws were pregnant during the PI season.

Forty percent of the EMS-Ws declared to be vaccinated simultaneously for SI and PI (PI + /SI +), 19% for PI only (PI + /SI -), and 1.6% for SI only (PI - /SI +). Thirty-nine percent ($n = 24$) of the EMS-Ws were not vaccinated at all (PI - /SI -). Compared with men, women's vaccination rates were lower for both PI and SI, but these results were not statistically significant (PI + /SI +: 24% vs 47%; PI + /SI -: 12% vs 22%; PI - /SI -: 65% vs 29% in female and male EMS-Ws, respectively). The two pregnant women were not immunized for PI or SI. Amongst chronically ill EMS-Ws ($n = 8$),

five were PI + /SI + (63%) and two were PI - /SI - (25%). Only one of the two EMS-Ws who declared living with a person at risk of influenza complications was PI + /SI +.

Past and Future Vaccination Status

In terms of vaccination habits (Table 3), 92% of the PI + /SI + EMS-Ws had received at least one SI vaccination during the previous three years, whereas only 8.3% of the PI - /SI - group got one ($P = .001$). Seventy-five percent of PI + /SI - EMS-Ws didn't get any SI immunization during the previous three years.

Compared with the preceding year, the SI vaccination rate increased globally from 26% to 42% during the H1N1 pandemic year (+62%). Specifically, in the PI + /SI + group, the vaccination rate against SI increased by 40% in comparison with the preceding year.

Questioned about future vaccination, 30% of the PI + /SI + EMS-Ws declared that they would not get vaccination during the following year. In the PI - /SI - and PI + /SI - EMS-W groups, no one was willing to be vaccinated in the future.

Positive Determinants of Vaccination Status

For PI + /SI + EMS-Ws, "protection of close family or friends" and "protection of patients" were the two main reasons for getting vaccinated against SI (Table 4), with mean values of 8.9 on the Likert scale (protection of close family or friends) and of 8.6 (protection of patients), respectively. For them, "protection of close family or friends" and "protection of patients" were also the two main motivations for vaccination against PI, with mean values

	PI + / SI + (n = 25)	PI + / SI - (n = 12)	PI - / SI + (n = 1)	PI - / SI - (n = 24)	P Value ^a
SI Vaccination Previous Year					
Yes	15 (24%)	1 (1.6%)	0	0	.001 ^b
No	8 (13%)	11 (18%)	1 (1.6%)	24 (39%)	
Unknown	2 (3.2%)	0	0	0	
Number of Vaccination during the Past 3 Years					
One	4 (6.5%)	3 (4.8%)	1 (1.6%)	2 (3.2%)	.001 ^b
Two	11 (18%)	0	0	0	
Three	8 (13%)	0	0	0	
None	2 (3.2%)	9 (15%)	0	22 (36%)	
Will Get Vaccination the Following Year					
Yes	18 (29%)	0	1 (1.6%)	0	.001 ^b
No	7 (11%)	12 (19%)	0	23 (37%)	
Don't Know Yet	0	0	0	1 (1.6%)	

Moser © 2016 Prehospital and Disaster Medicine

Table 3. Vaccinations Habits

Abbreviations: PI, pandemic influenza; SI, seasonal influenza.

^aGroup PI-/SI+ was excluded from analysis.^bFisher's exact test.

of 9.1 on the Likert scale (protection of close family or friends) and of 8.4 (protection of patients), respectively.

A significant difference on the Likert scale ($P < .01$) was found for "self-protection against PI" as a motivation factor for the PI + / SI + group (7.7; SD = 0.4) vs PI + / SI - group (4.6; SD = 1.2).

For the EMS-Ws only vaccinated against PI (PI + / SI -), the discomfort from the mandatory surgical mask wearing policy was the most important motivation (mean Likert scale: 8.1; SD = 0.8). For the EMS-Ws with complete immunization (PI + / SI +), this discomfort appeared less important but was present (7.4; SD = 0.5).

Peer model ("encouraged by a colleague") had limited impact and was rated higher for PI vaccination (3.8; SD = 0.7) than for SI vaccination (2.0; SD = 0.6; $P < .01$).

The other potential positive determinants of vaccination (easy access, free vaccination, and information campaigns) were not considered relevant by EMS-Ws and were not statistically different between PI + and SI + groups.

Negative Determinants of Vaccination Status

In the PI - / SI - group, the "preference for barrier precaution" was the main justification to reject SI vaccination (8.0; SD = 0.5 on the Likert scale) or PI vaccination (7.8; SD = 0.6; Table 5). This negative determinant was less important for PI + / SI - EMS-Ws ($P < .01$). For the PI - / SI - EMS-Ws, the two other main negative determinants against SI were "avoidance of medication" (7.2; SD = 0.5) and "fear of adverse effects" (6.7; SD = 0.8). The main negative determinants for the pandemic influenza vaccination were "avoidance of drugs" (6.8; SD = 0.6), "fear of adverse effects" (6.0; SD = 0.8), and "vaccination deemed unsafe" (6.0; SD = 0.7). The "fear of adverse effects" was significantly less prominent for PI + / SI - EMS-Ws (3.8; SD = 1.0) than for PI - / SI - EMS-Ws (6.7; SD = 0.8; $P < .05$).

Finally, concerns about effectiveness ("vaccination deemed ineffective") and safety ("vaccination deemed unsafe") of vaccines were two other negative determinants, especially for PI - / SI - EMS-Ws (Table 5). The other negative determinants of vaccination were not considered relevant by EMS-Ws and were not statistically different between PI and SI vaccinations.

Additional Comments

Analysis of comments showed similar positive determinants (self protection, $n = 2$; and protection of close family, $n = 2$) and some additional positive arguments (peer pressure, $n = 4$; and free-of-charge vaccination, $n = 1$). Comments related to negative determinants included mistrust against the pharmaceutical industry ($n = 4$), fear of adverse effects ($n = 3$), and vaccination deemed unsafe or unnecessary ($n = 3$).

Attitude Towards the Scenario of a Hospitalized Relative

The PI - / SI - EMS-Ws largely agreed with the assertion that they would accept that a family member could be managed by an HCW who didn't get influenza vaccination (8.9; SD = 1.9; Table 6). The acceptance rate towards this assertion was lower for PI + / SI - EMS-Ws (6.9; SD = 2.6) and for the PI + / SI + EMS-Ws (5.4; SD = 2.6; $P = .01$).

For PI - / SI - EMS-Ws, the result remained unchanged if the HCW wore a surgical mask (9.0; SD = 1.8). In opposite, the acceptance rate improved for the PI + / SI + EMS-Ws if a facemask was worn (7.6; SD = 2.3; $P = .01$).

Fairness of a Mandatory Surgical Mask Wearing Policy

When asked about the fairness of the new surgical mask wearing policy for unvaccinated EMS-Ws, the PI + / SI - EMS-Ws scored only 3.5 (SD = 3.4) points on the Likert scale, significantly less than the PI - / SI - EMS-Ws (6.4; SD = 3.2; $P = .02$) or the

	PI + / SI + (n = 25)	PI + / SI - (n = 12)	PI - / SI + (n = 1)	PI - / SI - (n = 24)
If you were vaccinated against seasonal influenza this season: What were your motivations?				
Protection of Close Family/Friends	8.9 (0.3)		5	
Protection of Patients	8.6 (0.3)		5	
Self Protection	6.6 (3.1)		7.5	
Free and Available Vaccine	4.7 (0.6)		7.5	
Convinced by Campaign	3.0 (0.6)		0	
Encouraged by a Colleague ^a	2.0 (0.6)		5	
If you were vaccinated against pandemic influenza this season: What were your motivations?				
Self Protection ^b	7.7 (0.4)	4.6 (1.2)		
Protection of Patients	8.4 (0.4)	7.1 (0.9)		
Protection of Close Family/Friends	9.1 (0.3)	7.7 (0.4)		
Free and Available Vaccine	4.3 (0.6)	4.4 (1.3)		
Encouraged by a Colleague	3.8 (0.7)	3.1 (1.0)		
Convinced by Campaign	3.2 (0.7)	2.5 (1.1)		
Discomfort from the Mask	7.4 (0.5)	8.1 (0.8)		

Moser © 2016 Prehospital and Disaster Medicine

Table 4. Positive Determinants for SI and PI Vaccination

Data are expressed as mean of all opinion on the Likert scale and standard deviation (SD). $P < .01$ for peer example (« *Encouraged by a Colleague* ») as motivation to get vaccination PI vaccination in the PI + /SI + group. $P < .01$ for self protection motivation concerning the PI vaccination between both groups.

Abbreviations: PI, pandemic influenza; SI, seasonal influenza.

^at test q11 vs q17, for PI + /SI +.

^bt test q17 PI + /SI + vs q17 PI + /SI -.

PI + /SI + EMS-Ws (7.0; SD = 3.4; $P = .01$). All EMS-W groups agreed on the fact that the surgical mask was hard to wear on duty, without any statistical difference between vaccination statuses.

Discussion

Rates of Vaccination and Differences between PI and SI Vaccination

This study shows that a minority of EMS-Ws were vaccinated against SI (42%). The vaccination rate was higher for PI (60%). The rates of vaccination for PI and SI were lower in women EMS-Ws, but these differences were not statistically significant. A simultaneous vaccination against both types of influenza was observed in only 40% of the cases. Interestingly, 39% of the EMS-Ws declared not to be vaccinated against either.

The global vaccination rate against SI is low in this study, but remains within the range of the results observed in previous publications about EMS-Ws.^{2,4} Compared with the vaccination rates observed in the HCWs of the Lausanne University Hospital (Lausanne, Switzerland), or in other emergency services,^{2,25,26} the vaccination rates of these EMS-Ws were lower for both PI and SI.

In this study, EMS-Ws were mainly young, healthy men and only a minority of them reported living with a person at risk of complicated influenza. Nevertheless, one in five declared living with a child less than two years of age or with a pregnant woman. These results therefore suggest an incomplete knowledge and a potential misunderstanding, combined with a trivialization about

the risk, of influenza. These results are a source of concern as they pertain to HCWs involved in acute care of vulnerable, elderly, or pediatric patients.

Interestingly, compared with the previous year, the occurrence of the PI was associated with a 1.6-fold increase in the SI vaccination rate. This effect was observed predominantly in the PI + /SI + group, showing a 40% increase of the vaccination rate against SI in comparison with the previous year. This “pandemic effect” has been described previously in the literature^{27,28} and may be related to the fear of adverse events – directly in line with the mediatization of the PI – or with the facilitated access to free SI vaccination and to both SI and PI vaccinations. At the same time, a “pandemic peer pressure” was also identified as a significant positive determinant for PI vaccination.

Positive and Negative Determinants of Vaccination

The analysis of the positive determinants of vaccination revealed that altruism remained the main motivation to get SI or PI vaccination, a result that was consistent with results of the literature.^{2,4,29,30} “Self-protection against PI” was rated lower in the PI + /SI - group than in the PI + /SI + group. Again, these results raise the question of the trivialization of the influenza infection, particularly for PI, in the PI + /SI - EMS-Ws. This difference may also be explained by the surgical mask wearing policy. Getting the vaccination to avoid the surgical mask may

	PI + / SI + (n = 25)	PI + / SI - (n = 12)	PI - / SI + (n = 1)	PI - / SI - (n = 24)
If you were NOT vaccinated against seasonal influenza this season: Why were you demoted?				
Allergy		0 (0)		0.3 (0.2)
Contraindication(s)		0 (0)		0.2 (0.1)
Vaccination of Other HCW Deemed Sufficient		0.4 (0.3)		0.5 (0.3)
Not Prone to Flu		3.5 (0.9)		2 (0.5)
Fear of Adverse Effects ^a		3.8 (1.0)		6.7 (0.8)
Avoidance of Drugs		5.8 (0.8)		7.2 (0.5)
Vaccination Deemed Inefficient		5.8 (0.9)		5.9 (0.6)
Vaccination Deemed Unsafe		4.6 (0.9)		5.7 (0.6)
History of Adverse Effects		0 (0)		0.5 (0.4)
Barrier Precaution Preferred ^b		4.4 (1.0)		8 (0.5)
Fear of Injections		0.8 (0.5)		0.3 (0.2)
Lack of Time		0 (0)		0.2 (0.2)
Vaccination Forgotten		0 (0)		0.5 (0.4)
Use of Alternative Medicine		2.9 (1.1)		4.1 (0.8)

Moser © 2016 Prehospital and Disaster Medicine

Table 5. Negative Determinants of SI and PI Vaccination

Data are expressed as mean of all opinion on the Likert scale and standard deviation (SD). $P < .05$ for fear of adverse effects between both groups and $P < .01$ concerning barrier precaution (use of PPE) to avoid vaccination between both groups.

Abbreviations: HCW, health care worker; PI, pandemic influenza; PPE, personal protection equipment; SI, seasonal influenza.

^a $P < .05$.

^b $P < .01$.

	PI + / SI + (n = 25)	PI + / SI - (n = 12)	PI - / SI + (n = 1)	PI - / SI - (n = 24)	P Value
Would accept that one hospitalized relative (parents, child) during one influenza outbreak could be managed by someone who didn't get influenza vaccination.	5.4 (2.6)	6.9 (2.6)	7.5	8.9 (1.9)	.001 ^a
Would accept that one hospitalized relative (parents, child) during one influenza outbreak could be managed by someone who didn't get influenza vaccination, but who is wearing a mask.	7.6 (2.3)	8.5 (1.7)	7.5	9 (1.8)	.06 ^a
P Value	.01 ^b	.09 ^b	-	.85 ^b	

Moser © 2016 Prehospital and Disaster Medicine

Table 6. Attitude toward Immunization und Mask in the Scenario of a Hospitalized Relative

Abbreviations: PI, pandemic influenza; SI, seasonal influenza.

^aANOVA oneway test.

^b χ^2 test.

therefore appear as an indirect positive determinant of vaccination in the PI + /SI- subgroup. Others positive determinants, including an information campaign or a history of a previous influenza infection, weren't relevant for these EMS-Ws.

Even if the surgical mask policy played a positive role towards vaccination, it was also the main negative determinant of vaccination in this study, particularly for PI-/SI-EMS-Ws.

In this context, the surgical mask policy may impact the vaccination strategy adversely, providing an alternative to the vaccination. The other main negative determinants of vaccination included avoidance of medication, fear of adverse effects, and concerns about safety and effectiveness of the vaccination. These elements corroborate previous studies about HCWs,^{2,30-32} revealing serious doubts regarding adverse effects and distrust towards the

pharmaceutical industry. These reservations may have been enhanced by the presence of adjuvants in one of the PI vaccines. They illustrate a profound fear and anxiety to serve as “guinea pig” for a “new” vaccine, which only benefited from a short period of development.^{33,34}

EMS-W Vaccination Profiles

More than 90% of the PI+/SI+ EMS-Ws received at least one SI vaccination during the previous three years, whereas only 8.3% of the PI-/SI- group did. For the following influenza epidemics, 30% of the PI+/SI+ EMS-Ws declared that they would not get vaccinated, whereas in the EMS-Ws in the PI-/SI- group or in the PI+/SI-, no one was willing to be immunized. These results therefore confirm that the previous vaccination status is a significant determinant of future vaccination acceptance or refusal.

Analysis of behaviors toward vaccination reveals three distinct groups previously described in the literature.^{2,4,5} The first is globally favorable to the vaccination strategy (were vaccinated, are vaccinated during the period of observation, and according to official recommendations, will be vaccinated). In this study, they represent a disappointingly small group (40%) of the EMS-Ws. The second one is exactly at the opposite. This group more frequently involved women and shared a profound suspicion of the pharmaceutical industry. These EMS-Ws particularly are reluctant toward vaccinations, more prone to trivialization, and less easily reached by the recommendations and education policies about influenza vaccination. Coercion remains an option, but as demonstrated by the mask policy, may produce unpredictable results. The last one is composed of EMS-Ws who traditionally reject SI vaccination, but accept to be vaccinated against PI, and for convenience, against both PI and SI. This group is more heterogeneous and may be called the “pandemic effect group” (the majority of them didn’t get a vaccination during the previous years but at this point did due to the pandemic context). This group of people may also explain the significant increase in the global rate of SI vaccination, in comparison with the previous year. Unfortunately, most of the EMS-Ws of this group are not disposed to get vaccinated during the next flu season. This fact should be put into perspective with the limited number of PI cases and disproportionate media coverage. This group is more volatile and might be influenced more easily. The public health care challenge within this group is therefore to educate and to motivate them to get vaccinated in the next years, even without the pandemic context.

New Surgical Mask Wearing Policy

The new surgical mask wearing policy was a double-edged sword. On one side, it was a significantly positive determinant of PI

vaccination, but on the other side, offered an alternative to the vaccination for EMS-Ws tending to reject the influenza vaccination. Even if this policy had a positive impact on the vaccination rate, it isn’t considered fair by a majority of these EMS-Ws. Furthermore, the question of compliance with such a policy constitutes another challenge, as it’s well known that EMS-Ws are not compliant with infection control recommendations in general.^{5,7,8} This risk is strengthening by the fact that a large majority of these EMS-Ws expressed doubts about the acceptability of a surgical mask in the field.

Limitations

This study has some limitations. The small sample size and monocentric design reduce the external validity of the results; however, these findings are supported by studies in other settings.^{2,4,5,27-34} In the self-administered questionnaire, vaccination rates were self-reported and not verified by vaccination logs. Face-to-face interviews, on the other hand, could have led to a social desirability bias.

Conclusion

The EMS-W immunization rates observed in this study are not sufficient to prevent the spread of influenza to uninfected transported patients and can contribute to nosocomial influenza transmission. This issue needs to be addressed, as it implies a high-risk situation for the operation of the prehospital emergency medical system in this region, especially in case of a new pandemic. Therefore, one of the public health challenges for the coming years is to increase the proportion of EMS-Ws getting influenza vaccinations every year. One part of the solution is to develop specific educational campaign targeting EMS-Ws. A surgical mask wearing policy should be presented as a solution of last resort for EMS-Ws who have contra-indications to vaccination, versus an acceptable alternative.

Acknowledgement

The authors wish to acknowledge Dr. Geneviève Bois, MD, University of Montréal (Montréal, Québec, Canada), for her corrections of the script.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1049023X16000121>

References

1. Thompson WW, Shay DK, Weintraub E, et al. Influenza-associated hospitalizations in the United States. *JAMA*. 2004;292(11):1333-1340.
2. Rueckmann E, Shah MN, Humiston SG. Influenza vaccination among Emergency Medical Services and emergency department personnel. *Prehosp Emerg Care*. 2009;13(1):1-5.
3. Poland GA, Tosh P, Jacobson RM. Requiring influenza vaccination for health care workers: seven truths we must accept. *Vaccine*. 2005;23(17-18):2251-2255.
4. Hubble MW, Zontek TL, Richards ME. Predictors of influenza vaccination among Emergency Medical Services personnel. *Prehosp Emerg Care*. 2011;15(2):175-183.
5. Bledsoe BE, Sweeney RJ, Berkeley RP, Cole KT, Forred WJ, Johnson LD. EMS provider compliance with infection control recommendations is suboptimal. *Prehosp Emerg Care*. 2014;18(2):290-294.
6. Emanuelsson L, Karlsson L, Castrén M, Lindström V. Ambulance personnel adherence to hygiene routines: still protecting ourselves but not the patient. *Eur J Emerg Med*. 2013;20(4):281-285.
7. Goodman CS, Cone DC. Emergency Medical Services equipment hygiene practices. *Prehosp Emerg Care*. 2001;5(2):169-173.
8. Eustis TC, Wright SW, Wrenn KD, Fowlic EJ, Slovis CM. Compliance with recommendations for universal precautions among prehospital providers. *Ann Emerg Med*. 1995;25(4):512-515.
9. Carman WF, Elder AG, Wallace LA, et al. Effects of influenza vaccination of health-care workers on mortality of elderly people in long-term care: a randomized controlled trial. *Lancet*. 2000;355(9198):93-97.
10. Wilde JA, McMillan JA, Serwint J, Butta J, O’Riordan MA, Steinhoff MC. Effectiveness of influenza vaccine in health care professionals: a randomized trial. *JAMA*. 1999;281(10):908-913.
11. Nichol KL, Margolis KL, Wuorenma J, Von Sternberg T. The efficacy and cost effectiveness of vaccination against influenza among elderly persons living in the community. *N Engl J Med*. 1994;331(12):778-784.

12. Nichol KL, Lind A, Margolis KL, et al. The effectiveness of vaccination against influenza in healthy, working adults. *N Engl J Med.* 1995;333(14):889-893.
13. Bridges CB, Thompson WW, Meltzer MI, et al. Effectiveness and cost-benefit of influenza vaccination of healthy working adults: a randomized controlled trial. *JAMA.* 2000;284(13):1655-1663.
14. Saxén H, Virtanen M. Randomized, placebo-controlled, double blind study on the efficacy of influenza immunization on absenteeism of health care workers. *Pediatr Infect Dis J.* 1999;18(9):779-783.
15. Chan SS-W. Does vaccinating ED health care workers against influenza reduce sickness absenteeism? *Am J Emerg Med.* 2007;25(7):808-811.
16. Centers for Disease Control and Prevention (CDC). Prevention and control of influenza: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR.* 2007;56(RR06):1-54.
17. Caplan A. Time to mandate influenza vaccination in health-care workers. *Lancet.* 2011;378(9788):310-311.
18. Ottenberg AL, Wu JT, Poland GA, Jacobson RM, Koenig BA, Tilburt JC. Vaccinating health care workers against influenza: the ethical and legal rationale for a mandate. *Am J Public Health.* 2011;101(2):212-216.
19. King WD, Woolhandler SJ, Brown AF, et al. Brief report: influenza vaccination and health care workers in the United States. *J Gen Intern Med.* 2006;21(2):181-184.
20. Masserey E. Communication du 9 Novembre 2009 à l'intention des services d'ambulances et de leur médecin conseil. Service de la Santé Publique Vaudoise; 2009.
21. Boubaker K, Masserey E. Vaccination contre la grippe et port du masque. Service de la Santé Publique Vaudoise; 2010.
22. WHO. Influenza A(H1N1) [Internet]. WHO Web site. http://www.who.int/mediacentre/news/statements/2009/h1n1_20090429/en/index.html. Accessed January 18, 2014.
23. WHO. Pandemic (H1N1) 2009 - update 58 [Internet]. WHO Web site. http://www.who.int/csr/don/2009_07_06/en/index.html. Accessed April 8, 2013.
24. WHO. Clinical features of severe cases of pandemic influenza [Internet]. WHO Web site. http://www.who.int/csr/disease/swineflu/notes/h1n1_clinical_features_20091016/en/index.html. Accessed January 18, 2014.
25. Hugli O, Dorribo V, Lazor-Blanchet C, Zanetti G. Differential proportion of health care workers immunization against the seasonal and pandemic influenza in 2009 in an emergency department. *Acad Emerg Med.* 2010;18(suppl 1):S196.
26. Dorribo V, Hugli Oliviers, Lazor-Blanchet C, Zanetti G. Health care workers' influenza A(H1N1)2009 vaccination: motivations and mandatory mask policy. *Occupational Medicine.* 2015; (In Press).
27. Prematunge C, Corace K, McCarthy A, Nair RC, Pugsley R, Garber G. Factors influencing pandemic influenza vaccination of health care workers—a systematic review. *Vaccine.* 2012;30(32):4733-4743.
28. Bish A, Yardley L, Nicoll A, Michie S. Factors associated with uptake of vaccination against pandemic influenza: a systematic review. *Vaccine.* 2011;29(38):6472-6484.
29. Barnett DJ, Levine R, Thompson CB, et al. Gauging US Emergency Medical Services workers' willingness to respond to pandemic influenza using a threat- and efficacy-based assessment framework. *PLoS ONE.* 2010;5(3):e9856.
30. Goldstein AO, Kincade JE, Gamble G, Bearman RS. Policies and practices for improving influenza immunization rates among health care workers. *Infect Control Hosp Epidemiol.* 2004;25(11):908-911.
31. Salgado CD, Giannetta ET, Hayden FG, Farr BM. Preventing nosocomial influenza by improving the vaccine acceptance rate of clinicians. *Infect Control Hosp Epidemiol.* 2004;25(11):923-928.
32. Heimerberger T, Chang HG, Shaikh M, Crotty L, Morse D, Birkhead G. Knowledge and attitudes of health care workers about influenza: why are they not getting vaccinated? *Infect Control Hosp Epidemiol.* 1995;16(7):412-415.
33. Smith EC, Burkle FM Jr, Holman PF, Dunlop JM, Archer FL. Lessons from the front lines: the prehospital experience of the 2009 novel H1N1 outbreak in Victoria, Australia. *Disaster Med Public Health Prep.* 2009;3(Suppl 2): S154-S159.
34. Smith E. Paramedic Perception of Risk and Willingness to Work during Disasters: A Quantitative and Qualitative Analysis. Unpublished PhD thesis. 2009.