

Supplementary Material

The significance of *Anomalocaris* and other Radiodonta for understanding paleoecology and evolution during the Cambrian Explosion

Gaëtan J-M Potin^{1*}, Allison C Daley¹

* Correspondence: Corresponding Author: gaetan.jm.potin@outlook.fr

1 Full list of taxa included or linked to Radiodonta

Radiodonta: Liu et al. (2018) (Collins, 1996)

Ampectobeluidae: Pates et al. (2021a) (Vinther et al., 2014)

Ampectobelua: Cong et al. (2017) (Hou et al., 1995)

Ampectobelua stephenensis: Daley and Budd (2010)

Ampectobelua symbachiata: Cong et al. (2017) (Hou et al., 1995)

Guanshancaris Zhang et al. (2023)

Guanshancaris kunmingensis: Zhang et al. (2023) (Wang et al., 2013)

Lyrarapax: Liu et al. (2018) (Cong et al., 2014)

Lyrarapax trilobus: Cong et al. (2016)

Lyrarapax unguispinus: Cong et al. (2014)

Ramskoeldia: Cong et al. (2018)

Ramskoeldia platyacantha: Cong et al. (2018)

**Ramskoeldia consimilis*: Cong et al. (2018)

Anomalocarididae: (Raymond, 1935)

Anomalocaris: Collins (1996) (Whiteaves, 1892)

Anomalocaris canadensis: Daley and Edgecombe (2014) (Whiteaves, 1892)

Lenisicaris: Wu et al. (2021b)

Lenisicaris pennsylvanica: Pates and Daley (2019) (Resser, 1929), genus retribution in Wu et al. (2021b)

Lenisicaris lupata: Wu et al. (2021b)

Hurdiidae: Wu et al. (2022) and Moysiuk and Caron (2022) (Vinther et al., 2014)

Aegirocassis: Van Roy et al. (2015)

Aegirocassis benmoulai: Van Roy et al. (2015)

Buccaspinea: Pates et al. (2021b)

Buccaspinea cooperi: Pates et al. (2021b)

Cambroraster: Moysiuk and Caron (2019)

Cambroraster falcatus: Moysiuk and Caron (2019)

Cordaticaris: Sun et al. (2020)

Cordaticaris striatus: Sun et al. (2020)

Hurdia: Daley et al. (2013a) (Walcott, 1912)

Hurdia victoria: Daley et al. (2013a) (Walcott, 1912)

Hurdia triangulata: Daley et al. (2013a) (Walcott, 1912)

Pahvantia: Lerosey-Aubril and Pates (2018) (Robison and Richards, 1981)

Pahvantia hastata: Lerosey-Aubril and Pates (2018) (Robison and Richards, 1981)

Peytoia: Daley and Legg (2015) (Walcott, 1911)

Peytoia infercambriensis: Daley and Legg (2015) (Lendzion, 1975)

Peytoia nathorsti: Collins (1996) (Walcott, 1911), synonymised from *Laggania cambria* in Daley and Bergström (2012).

Stanleycaris: Moysiuk and Caron (2022) (Caron et al., 2010)

Stanleycaris hirpex: Moysiuk and Caron (2022) (Caron et al., 2010)

Titanokorys: Caron and Moysiuk (2021)

Titanokorys gainesi: Caron and Moysiuk (2021)

Ursulinacaris: Pates et al. (2019)

Ursulinacaris grallae: Pates et al. (2019)

**Pseudoangustidontus*: Van Roy and Tetlie (2006)

Pseudoangustidontus duplospineus: Van Roy and Tetlie (2006)

**Schinderhannes*: Kühl et al. (2009)

Schinderhannes bartelsi: Kühl et al. (2009)

Tamisiocarididae: Wu et al. (2021a) (Vinther et al., 2014), originally called Cetiocaridae

“*Anomalocaris*” *briggsii*: Daley et al. (2013b) (Nedin, 1995)

Houcaris: Wu et al. (2021a)

Houcaris saron: Wu et al. (2021a) (Hou et al., 1995)

Houcaris magnabasis: Pates et al. (2021a), genus reatribution in Wu et al. (2021a)

Tamisiocaris: Vinther et al. (2014) (Daley and Peel, 2010)

Tamisiocaris borealis: Pates and Daley (2019) (Daley and Peel, 2010)

Incertae sedis:

***Caryosyntrips*: Pates and Daley (2017) (Daley and Budd, 2010)

Caryosyntrips camurus: Pates and Daley (2017)

Caryosyntrips durus: Pates and Daley (2017)

Caryosyntrips serratus: Pates and Daley (2017) (Daley and Budd, 2010)

***Cucumericrus*: Hou et al. (1995)

Cucumericrus decoratus: Hou et al. (1995)

***Innovatiocaris*: Zeng et al. (2022)

Innovatiocaris maotianshanensis: Zeng et al. (2022)

“*Innovatiocaris*” *multispiniformis*: Zeng et al. (2022)

***Laminacaris*: Guo et al. (2019)

Laminacaris chimera: Guo et al. (2019)

***Paranomalocaris*: Jiao et al. (2021) (Wang et al., 2013)

Paranomalocaris multisegmentalis: Jiao et al. (2021) (Wang et al., 2013)

Paranomalocaris simplex: Jiao et al. (2021)

****Parapeytoia*: Hou et al. (1995)

Parapeytoia yunnanensis: Hou et al. (1995)

*: attribution to this family is uncertain

**: likely to be a radiodont, but unclear to which family it belongs

***: probably not a radiodont, but has been described as one previously and remains closely associated to radiodonts (not included in our counts of radiodont genera and species)

2 Radiodonta terminology

Supplementary Table 1. Non-exhaustive lists of anatomical terminology applied to Radiodonta in this review as compared to previous publications.

3 Radiodonta global and temporal distribution

Supplementary Table 2. Complete list of lower Paleozoic localities from which Radiodonta are described, and the taxa found therein.

4 References

Caron, J.-B., Gaines, R.R., Mángano, M.G., Streng, M., and Daley, A.C. (2010). A new Burgess Shale-type assemblage from the “thin” Stephen Formation of the southern Canadian Rockies. *Geology* 38(9), 811-814. doi: 10.1130/G31080.1.

Caron, J.-B., and Moysiuk, J. (2021). A giant nektobenthic radiodont from the Burgess Shale and the significance of hurdiid carapace diversity. *Royal Society open science* 8(9), 210664. doi: 10.1098/rsos.210664.

Collins, D. (1996). The “evolution” of *Anomalocaris* and its classification in the arthropod class Dinocarida (nov.) and order Radiodonta (nov.). *Journal of Paleontology* 70(2), 280-293. doi: 10.1017/S0022336000023362.

- Cong, P., Daley, A.C., Edgecombe, G.D., and Hou, X. (2017). The functional head of the Cambrian radiodontan (stem-group Euarthropoda) *Amplectobelua symbrachiata*. *BMC Evolutionary Biology* 17(1), 1-23.
- Cong, P., Daley, A.C., Edgecombe, G.D., Hou, X., and Chen, A. (2016). Morphology of the radiodontan *Lyrarapax* from the early Cambrian Chengjiang biota. *Journal of Paleontology* 90(4), 663-671. doi: 10.1017/jpa.2016.67.
- Cong, P., Ma, X., Hou, X., Edgecombe, G.D., and Strausfeld, N.J. (2014). Brain structure resolves the segmental affinity of anomalocaridid appendages. *Nature* 513(7519), 538-542. doi: 10.1038/nature13486.
- Cong, P.Y., Edgecombe, G.D., Daley, A.C., Guo, J., Pates, S., and Hou, X.G. (2018). New radiodonts with gnathobase-like structures from the Cambrian Chengjiang biota and implications for the systematics of Radiodonta. *Papers in Palaeontology* 4(4), 605-621. doi: 10.1002/spp2.1219.
- Daley, A.C., and Bergström, J. (2012). The oral cone of Anomalocaris is not a classic “peytoia”. *Naturwissenschaften* 99(6), 501-504.
- Daley, A.C., and Budd, G.E. (2010). New anomalocaridid appendages from the Burgess Shale, Canada. *Palaeontology* 53(4), 721-738. doi: 10.1111/j.1475-4983.2010.00955.x.
- Daley, A.C., Budd, G.E., and Caron, J.-B. (2013a). Morphology and systematics of the anomalocaridid arthropod *Hurdia* from the Middle Cambrian of British Columbia and Utah. *Journal of Systematic Palaeontology* 11(7), 743-787. doi: 10.1080/14772019.2012.732723.
- Daley, A.C., and Edgecombe, G.D. (2014). Morphology of Anomalocaris canadensis from the Burgess Shale. *Journal of Paleontology* 88(1), 68-91.
- Daley, A.C., and Legg, D.A. (2015). A morphological and taxonomic appraisal of the oldest anomalocaridid from the lower Cambrian of Poland. *Geological Magazine* 152(5), 949-955. doi: 10.1017/S0016756815000412.
- Daley, A.C., Paterson, J.R., Edgecombe, G.D., García-Bellido, D.C., and Jago, J.B. (2013b). New anatomical information on *Anomalocaris* from the Cambrian Emu Bay Shale of South Australia and a reassessment of its inferred predatory habits. *Palaeontology* 56(5), 971-990. doi: 10.1111/pala.12029.
- Daley, A.C., and Peel, J.S. (2010). A possible anomalocaridid from the Cambrian Sirius Passet Lagerstätte, north Greenland. *Journal of Paleontology* 84(2), 352-355. doi: 10.1666/09-136R1.1.
- Guo, J., Pates, S., Cong, P., Daley, A.C., Edgecombe, G.D., Chen, T., et al. (2019). A new radiodont (stem Euarthropoda) frontal appendage with a mosaic of characters from the Cambrian (Series 2 Stage 3) Chengjiang biota. *Papers in Palaeontology* 5(1), 99-110. doi: 10.1002/spp2.1231.
- Hou, X.G., Bergström, J., and Ahlberg, P. (1995). *Anomalocaris* and other large animals in the Lower Cambrian Chengjiang fauna of southwest China. *Gff* 117(3), 163-183. doi: 10.1080/11035899509546213.
- Jiao, D.-G., Pates, S., Leroey-Aubril, R., Ortega-Hernández, J., Yang, J., Lan, T., et al. (2021). The endemic radiodonts of the Cambrian Stage 4 Guanshan Biota of South China. *Acta Palaeontologica Polonica* 66(2). doi: 10.4202/app.00870.2020.

- Kühl, G., Briggs, D.E.G., and Rust, J. (2009). A great-appendage arthropod with a radial mouth from the Lower Devonian Hunsrück Slate, Germany. *Science* 323(5915), 771-773. doi: 10.1126/science.1166586.
- Lendzion, K. (1975). Fauna of the Mobergella zone in the Polish Lower Cambrian. *Geological Quarterly* 19(2), 237-242.
- Leroosey-Aubril, R., and Pates, S. (2018). New suspension-feeding radiodont suggests evolution of microplanktivory in Cambrian macronekton. *Nature Communications* 9(1), 3774. doi: 10.1038/s41467-018-06229-7.
- Liu, J., Leroosey-Aubril, R., Steiner, M., Dunlop, J.A., Shu, D., and Paterson, J.R. (2018). Origin of raptorial feeding in juvenile euarthropods revealed by a Cambrian radiodontan. *National Science Review* 5(6), 863-869. doi: 10.1093/nsr/nwy057.
- Moysiuk, J., and Caron, J.-B. (2019). A new hurdiid radiodont from the Burgess Shale evinces the exploitation of Cambrian infaunal food sources. *Proceedings of the Royal Society B* 286(1908), 20191079. doi: 10.1098/rspb.2019.1079.
- Moysiuk, J., and Caron, J.-B. (2022). A three-eyed radiodont with fossilized neuroanatomy informs the origin of the arthropod head and segmentation. *Current Biology* 32(15), 3302-3316. e3302. doi: 10.1016/j.cub.2022.06.027.
- Nedin, C. (1995). The Emu Bay Shale, a Lower Cambrian fossil Lagerstätte, Kangaroo Island, South Australia. *Memoirs of the Association of Australasian Palaeontologists* 18, 31-40.
- Pates, S., and Daley, A.C. (2017). *Caryosyntrips*: a radiodontan from the Cambrian of Spain, USA and Canada. *Papers in Palaeontology* 3(3), 461-470. doi: 10.1002/spp2.1084.
- Pates, S., and Daley, A.C. (2019). The Kinzers Formation (Pennsylvania, USA): the most diverse assemblage of Cambrian Stage 4 radiodonts. *Geological Magazine* 156(7), 1233-1246. doi: 10.1017/S0016756818000547.
- Pates, S., Daley, A.C., and Butterfield, N.J. (2019). First report of paired ventral endites in a hurdiid radiodont. *Zoological Letters* 5(1), 18. doi: 10.1186/s40851-019-0132-4.
- Pates, S., Daley, A.C., Edgecombe, G.D., Cong, P., and Lieberman, B.S. (2021a). Systematics, preservation and biogeography of radiodonts from the southern Great Basin, USA, during the upper Dyeran (Cambrian Series 2, Stage 4). *Papers in Palaeontology* 7(1), 235-262. doi: 10.1002/spp2.1277.
- Pates, S., Leroosey-Aubril, R., Daley, A.C., Kier, C., Bonino, E., and Ortega-Hernández, J. (2021b). The diverse radiodont fauna from the Marjum Formation of Utah, USA (Cambrian: Drumian). *PeerJ* 9, e10509. doi: 10.7717/peerj.10509.
- Raymond, P.E. (1935). *Leanchoilia and other mid-Cambrian Arthropoda*. Bulletin Museum of Comparative Zoology Harvard University.
- Resser, C.E. (1929). New lower and middle Cambrian Crustacea. *Proceedings of the United States National Museum* 76, 1-18.
- Robison, R.A., and Richards, B.C. (1981). Larger bivalve arthropods from the Middle Cambrian of Utah. *Kansas Paleontological Contributions* 106, 1-28.
- Sun, Z., Zeng, H., and Zhao, F. (2020). A new middle Cambrian radiodont from North China: implications for morphological disparity and spatial distribution of hurdiids.

Palaeogeography, Palaeoclimatology, Palaeoecology 558, 109947. doi: 10.1016/j.palaeo.2020.109947.

- Van Roy, P., Daley, A.C., and Briggs, D.E.G. (2015). Anomalocaridid trunk limb homology revealed by a giant filter-feeder with paired flaps. *Nature* 522(7554), 77-80. doi: 10.1038/nature14256.
- Van Roy, P., and Tetlie, O.E. (2006). A spinose appendage fragment of a problematic arthropod from the Early Ordovician of Morocco. *Acta Palaeontologica Polonica* 51(2), 239-246.
- Vinther, J., Stein, M., Longrich, N.R., and Harper, D.A.T. (2014). A suspension-feeding anomalocarid from the Early Cambrian. *Nature* 507(7493), 496-499. doi: 10.1038/nature13010.
- Walcott, C.D. (1911). Cambrian Geology and Paleontology II: Middle Cambrian Holothurians and Medusae. *Smithsonian Miscellaneous Collections* 57(3), 41-69.
- Walcott, C.D. (1912). *Middle Cambrian Branchiopoda, Malacostraca, Trilobita, and Merostomata*. Smithsonian Institution.
- Wang, Y., Huang, D., and Hu, S. (2013). New anomalocarid frontal appendages from the Guanshan biota, eastern Yunnan. *Chinese Science Bulletin* 58(32), 3937-3942. doi: 10.1007/s11434-013-5908-x.
- Whiteaves, J.F. (1892). Description of a new genus and species of phyllocarid crustacea from the middle Cambrian of Mount Stephen, B.C. *The Canadian Record of Science* V(4), 205-208.
- Wu, Y., Fu, D., Ma, J., Lin, W., Sun, A., and Zhang, X. (2021a). *Houcaris* gen. nov. from the early Cambrian (Stage 3) Chengjiang Lagerstätte expanded the palaeogeographical distribution of tamisiocaridids (Panarthropoda: Radiodonta). *PalZ* 95(2), 209-221. doi: 10.1007/s12542-020-00545-4.
- Wu, Y., Ma, J., Lin, W., Sun, A., Zhang, X., and Fu, D. (2021b). New anomalocaridids (Panarthropoda: Radiodonta) from the lower Cambrian Chengjiang Lagerstätte: Biostratigraphic and paleobiogeographic implications. *Palaeogeography, Palaeoclimatology, Palaeoecology* 569, 110333. doi: 10.1016/j.palaeo.2021.110333.
- Wu, Y., Pates, S., Ma, J., Lin, W., Wu, Y., Zhang, X., et al. (2022). Addressing the Chengjiang conundrum: A palaeoecological view on the rarity of hurdiid radiodonts in this most diverse early Cambrian Lagerstätte. *Geoscience Frontiers* 13(6), 101430. doi: 10.1016/j.gsf.2022.101430.
- Zhang, M., Wu, Y., Lin, W., Ma, J., Wu, Y., and Fu, D. (2023). Amplectobeluid Radiodont *Guanshancaris* gen. nov. from the Lower Cambrian (Stage 4) Guanshan Lagerstätte of South China: Biostratigraphic and Paleobiogeographic Implications. *Biology* 12(583). doi: 10.3390/biology12040583.
- Zeng, H., Zhao, F., and Zhu, M. (2022). *Innovatiocaris*, a complete radiodont from the early Cambrian Chengjiang Lagerstätte and its implications for the phylogeny of Radiodonta. *Journal of the Geological Society* 180(1), jgs2021-2164. doi: 10.11114/jgs2021-164.