

**Serveur Académique Lausannois SERVAL [serval.unil.ch](http://serval.unil.ch)**

## **Author Manuscript**

**Faculty of Biology and Medicine Publication**

**This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.**

Published in final edited form as:

**Title:** Light-regulated interactions with SPA proteins underlie cryptochrome-mediated gene expression.

**Authors:** Fankhauser C, Ulm R

**Journal:** Genes & development

**Year:** 2011 May 15

**Volume:** 25

**Issue:** 10

**Pages:** 1004-9

**DOI:** 10.1101/gad.2053911

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

## **Light-regulated interactions with SPA proteins underlie cryptochrome-mediated gene expression**

Christian Fankhauser<sup>1,3</sup> and Roman Ulm<sup>2</sup>

<sup>1</sup>Center for Integrative Genomics, Faculty of Biology and Medicine, University of Lausanne, CH-1015 Lausanne, Switzerland; <sup>2</sup>Department of Botany and Plant Biology, University of Geneva, Sciences III, CH-1211 Geneva 4, Switzerland

<sup>3</sup>Corresponding author.

EMAIL: christian.fankhauser@unil.ch; FAX: ++41 21 692 39 25

[*Keywords:* cryptochrome; COP1; SPA1; photomorphogenesis; photoreceptor; E3 ubiquitin ligase]

**Cryptochromes are a class of photosensory receptors that control important processes in animals and plants. They act primarily by regulating gene expression in a blue light-responsive manner. How photon absorption by cryptochromes leads to changes in gene expression has remained largely elusive. Now three publications demonstrate the crucial interaction of light-activated Arabidopsis cryptochromes with a class of regulatory components of E3 ubiquitin ligase complexes involved in plant photomorphogenesis. Negative regulation of specific protein degradation allows environmental control of the abundance of transcriptional regulators.**

Living organisms have evolved mechanisms to sense, respond and adapt to changes in their environment. Alterations in the light environment provide important cues regarding the time of the day, the season and the surrounding habitat (e.g. water depth, shade from the vegetation) (Kami et al. 2010; Zoltowski and Gardner 2010). Organisms in all kingdoms of life possess a variety of photoreceptors that trigger these light responses. Among these is the cryptochrome family of blue light sensors that are present in both animals and plants (Cashmore 2003; Liu et al. 2010). The founding member of this family was uncovered in Arabidopsis almost 20 years ago (Ahmad and Cashmore 1993).

In plants, cryptochromes control important developmental transitions. These include de-etiolation, a transition during which seedlings switch from using energy within seed reserves to photoautotrophic growth, and photoperiodic induction of floral initiation (Fig. 1A). In addition, cryptochromes are a component of the light input pathway that resets the circadian oscillator. In mammals and insects, cryptochromes are well known for their role in the circadian clock where they act as either light-regulated or light-independent components of the oscillator (Cashmore 2003). Interestingly, cryptochromes have also been implicated as magnetoreceptors and have been shown to be essential for the time-compensated sun compass of the monarch butterfly (Reppert et al. 2010).

### **Mechanism of cryptochrome light activation**

Cryptochromes are related to a class of DNA repair enzymes known as DNA photolyases. A key difference between DNA photolyases and cryptochromes is that in the former light energy is used to initiate DNA repair, whilst in the latter light is used as an informational cue (Liu et al. 2010). Proteins of the photolyase/cryptochrome superfamily share the N-terminal *Photolyase-Homologous Region* (PHR) that harbors the light-sensing chromophores. Photolyases possess an antenna pigment (e.g. methenyltetrahydrofolate) and a flavin cofactor (FADH<sup>-</sup>) that, upon light perception, transfers an electron to reverse crosslinking between bases for DNA repair. Cryptochromes also possess a primary FAD chromophore, but the presence and identity of a second antenna pigment is not always clearly established (Liu et al. 2010). Moreover, the photocycle of cryptochromes is not yet fully understood and this topic remains a matter of debate (Liu et al. 2010). Importantly, the redox state of the flavin chromophore is at the heart of light-induced events triggered upon cryptochrome excitation. Such a redox-based signaling system could also be influenced by other factors such as oxygen concentration, the cellular midpoint potential and temperature. Therefore, it is possible that cryptochrome signaling is also affected by additional factors other than light (Yang et al. 2008). Another distinguishing feature of cryptochromes is the presence of a C-terminal CCE (*Cryptochrome C-terminal Extension*) domain of varying length (Cashmore 2003; Liu et al. 2010). In *Arabidopsis* cryptochrome 1 (CRY1) and CRY2 the CCE appears to be largely unstructured yet is known to act as an effector domain (Yang et al. 2000; Partch et al. 2005; Kottke et al. 2006; Yu et al. 2007b; Liu et al. 2010; Zoltowski and Gardner 2010).

### **Light-controlled cryptochrome signaling in plants**

Cryptochromes are widespread in plants where they control important developmental transitions throughout the plant life cycle (Cashmore 2003; Liu et al. 2010). Higher plants typically possess two cryptochromes with clear functional diversification. This has been extensively studied in the model plant *Arabidopsis thaliana* where light-stable CRY1 mediates mostly high light responses while light-labile CRY2 specializes in responses to low light (Lin and Shalitin 2003). CRY1 and CRY2 are both nuclear proteins, but CRY1 is also present in the cytosol. Consistent with their role in blue-light regulated gene expression, cryptochromes act primarily in the nucleus (Wu and Spalding 2007; Yu et al. 2007a). Both *Arabidopsis* cryptochromes are phosphorylated upon blue light perception (Shalitin et al. 2002; Shalitin et al. 2003). For CRY2 this post-translational modification triggers proteasome-mediated degradation (Shalitin et al. 2002; Yu et al. 2007a). Phosphorylation has also been related to the activation of cryptochromes, but the link between phosphorylation and cryptochrome activity remains poorly understood (Shalitin et al. 2003; Yu et al. 2007a; Yu et al. 2007b). Despite the absence of a canonical kinase domain CRY1 binds to ATP and autophosphorylates *in vitro* (Liu et al. 2010). However, light-induced cryptochrome phosphorylation does not necessarily depend on the kinase activity of cryptochrome itself and the functional significance of this activity remains to be determined (Liu et al. 2010).

The CCE domain is central to the activity of plant cryptochromes. This was first demonstrated using plants expressing the CCE domain fused to a dimerization domain but lacking the PHR domain (Yang et al. 2000). These plants showed a phenotype referred to as *constitutively photomorphogenic*, where seedlings develop as they would normally under light conditions despite being grown in darkness. This result suggested that the CCE domain of plant cryptochromes is active in the absence of light when separated from the regulatory PHR domain. This hypothesis was further supported by reports demonstrating that the CCE domain undergoes light-induced conformational changes, suggesting that CCE domain activity is controlled by light (Partch et al. 2005; Kottke et al. 2006; Yu et al. 2007b; Liu et al. 2010; Zoltowski and Gardner 2010). The ensuing hunt for proteins interacting with the CCE domain rapidly identified the E3 ubiquitin ligase component CONSTITUTIVELY PHOTOMORPHOGENIC 1 (COP1) (Wang et al. 2001; Yang et al. 2001). COP1 exists both in animals and plants where it controls accumulation of important transcription factors (Yi and Deng 2005). In contrast to plant cryptochromes, there is currently no experimental evidence suggesting an interaction between animal cryptochromes and COP1. However, plant cryptochromes surprisingly interact with COP1 independently of light (Wang et al. 2001; Yang et al. 2001). Thus, although it was proposed that cryptochromes inhibit COP1 activity specifically upon light perception, the primary light-mediated mechanism through which COP1 activity is controlled has remained elusive. This review highlights three important papers that further our knowledge in this regard (Lian et al. 2011; Liu et al. 2011; Zuo et al. 2011).

### **Cryptochromes interact with SPA family proteins in a blue light-dependent manner to regulate COP1-mediated protein degradation**

COP1 is part of a multimeric E3 ubiquitin ligase complex that includes CULLIN 4 (CUL4), DAMAGED DNA BINDING PROTEIN 1 (DDB1), RING-BOX 1 (RBX1) and SUPPRESSOR OF PHYA (SPA) proteins (SPA1-SPA4 in Arabidopsis) (Chen et al. 2010). Consistent with the notion that SPAs and COP1 are part of one multimeric E3 ubiquitin ligase, the quadruple *spa* mutant displays a phenotype similar to *cop1* alleles. Members of this complex repress light responses in both seedlings and mature plants (Laubinger et al. 2004; Laubinger et al. 2006). COP1 is comprised of an N-terminal RING finger domain followed by a coiled-coil domain and C-terminal WD40 repeats. SPA proteins also have C-terminal WD40 repeats and a central coiled-coil domain but possess an N-terminus that resembles a protein kinase domain (Fig. 1B,C). COP1 interacts with substrates via its WD40 repeats, whereas the interaction between SPA and COP1 involves the coiled-coil domains of the two proteins (Hoecker and Quail 2001). Importantly, structure-function analyses have identified the coiled-coil domain and WD40 repeats of SPA1 as essential for its activity (Fittinghoff et al. 2006; Yang and Wang 2006).

COP1 acts downstream of multiple plant photoreceptors, such as cryptochromes, phytochromes and the UV-B receptor, and controls the light-regulated abundance of numerous transcription factors (Yi and Deng 2005; Favory et al. 2009). In most cases, target proteins are degraded in a COP1-dependent manner in the dark. Upon light exposure, these proteins then rapidly accumulate due to photoreceptor-mediated inhibition of COP1 (Yi and Deng 2005). The bZIP

transcription factor ELONGATED HYPOCOTYL5 (HY5), which is required for normal light-regulated seedling development under all light conditions, was the first known target of COP1-mediated degradation (Yi and Deng 2005). The mechanism by which phytochromes rapidly inhibit COP1 in response to red or far-red light is still largely unknown. However, in response to UV-B, COP1 interacts in a light-dependent manner with UVR8, a protein that selectively controls plant responses to UV-B exposure. The interaction with UVR8 presumably inhibits COP1-mediated degradation of target proteins and leads to the accumulation of HY5 in the presence of UV-B (Favory et al. 2009). The recently described process by which COP1 activity is controlled by cryptochromes bears some resemblance to the control mechanism identified for COP1 in response to UV-B radiation. However, in blue light signaling it is not COP1 but SPA1, another component of the multimeric E3 ligase complex, that exhibits light-mediated interaction with cryptochromes (Lian et al. 2011; Liu et al. 2011; Zuo et al. 2011).

#### *The blue light-dependent CRY1-SPA1 interaction inhibits COP1-SPA1 binding*

Two recent publications have shown that the light-regulated interaction between Arabidopsis CRY1 and SPA1 is mediated by the CCE domain of CRY1 that interacts with the WD40 domain of SPA1 (Lian et al. 2011; Liu et al. 2011) (Fig. 1B). This regulated protein-protein interaction presumably depends on light-induced conformational changes that take place in the CRY1 CCE domain (Partch et al. 2005; Kottke et al. 2006; Zoltowski and Gardner 2010). Consistent with this notion, the light-regulated CRY1-SPA1 interaction was reconstituted in a heterologous system (Lian et al. 2011; Liu et al. 2011). This raised the question of how the CRY1-SPA1 interaction is able to modulate ubiquitin ligase activity of the multimeric COP1 E3 ligase. It was previously shown that SPA1 modulated ubiquitin ligase activity of COP1 in vitro (Saijo et al. 2003; Seo et al. 2003) and there is genetic evidence that indicates SPA proteins are required for COP1 function in vivo (Laubinger et al. 2004). Moreover, there exists a strong interaction between SPA1 and COP1 in the dark whereas this interaction is inhibited by light (Saijo et al. 2003). Recent studies in the Lin and Yang laboratories, as published in this issue of *Genes and Development* (Lian et al. 2011; Liu et al. 2011), show that the light-induced interaction between CRY1 and SPA1 inhibits the interaction between SPA1 and COP1. This network of light-regulated interaction can be reconstituted in yeast using a three-hybrid assay, which indicates there are no other plant-specific factors needed for these interactions to take place (Lian et al. 2011; Liu et al. 2011). Furthermore, in *cry1* mutants the accumulation of downstream transcription factors is no longer regulated by blue light (Liu et al. 2011; Zuo et al. 2011). Taken together, these two studies suggest that CRY1-mediated disruption of the SPA1-COP1 interaction inhibits the activity of this multimeric E3 ligase (Fig. 1B).

#### *Blue light-dependent protein-protein interactions underlie CRY2-mediated flowering.*

In another study conducted in parallel in the Lin laboratory, a variation on the same theme is described for CRY2-regulated gene expression (Zuo et al. 2011). CRY2, COP1 and SPA1 play a very important role in a later stage of the plant life cycle, namely the initiation of flowering. The photoperiodic control of plant

reproduction requires an integration of information from both the circadian clock and day length (Turck et al. 2008). Central to this process is the B box-type zinc finger transcription factor CONSTANS (CO). Only when day length is appropriate does CO protein accumulate sufficiently to trigger flowering. Both COP1 and SPA1 are required for degradation of CO under short days, which do not promote flowering in Arabidopsis (Laubinger et al. 2006; Jang et al. 2008; Liu et al. 2008b). Under long days, a developmental switch takes place whereby proteolytic activity of the COP1/SPA1 E3 ligase is inhibited by CRY2 (Turck et al. 2008). Until recently, the mechanism through which CRY2 controls E3 ligase activity of the COP1/SPA1 complex had remained unknown. However, two publications now show that, similar to CRY1, CRY2 also interacts with SPA1 in a blue-light-dependent manner (Lian et al. 2011; Zuo et al. 2011). Interestingly, and in contrast to the CRY1/SPA1 mode of action, the light-regulated CRY2-SPA1 interaction does not affect the interaction of COP1 with SPA1 but rather strengthens the CRY2/COP1 complex (Fig. 1C). This enhanced interaction of COP1 with the photoreceptor is thought to suppress proteolytic activity of the COP1/SPA1 complex, thereby allowing accumulation of CO and initiation of flowering under long days (Zuo et al. 2011). In contrast to the light-regulated CRY1-SPA1 interaction described above, formation of the CRY2/SPA1 complex requires both the PHR domain of the photoreceptor and the N-terminus of SPA1 (Zuo et al. 2011) (Fig. 1B,C). Regardless of the notable differences between CRY1 and CRY2 modes of action, the light-regulated activity of both proteins result in accumulation of a key transcriptional regulator.

It should also be mentioned that CRY2 controls initiation of flowering by an additional mechanism. Light-activated CRY2 specifically interacts with the bHLH class transcription factor CIB1 (CRYPTOCHROME-INTERACTING bHLH) (Liu et al. 2008a). This mechanism seems analogous to the light-dependent interaction between phytochromes and PIFs (PHYTOCHROME INTERACTING PROTEINS), which are bHLH transcription factors related to CIB1 (Liu et al. 2008a). Interestingly, light-regulated interaction of CRY2 with either SPA1 or CIB1 requires the PHR domain (Kennedy et al. 2010; Zuo et al. 2011) whereas light-regulated interaction of CRY1 with SPA1 occurs via the CCE domain (Liu et al. 2011; Zuo et al. 2011) (Fig. 1B,C). CIB1 and related bHLH transcription factors then induce flowering in a CRY2-dependent but CO-independent mechanism (Liu et al. 2008a). However, similar to CO, CIB1 also interacts with the promoter of *FLOWERING LOCUS T (FT)*, inducing gene expression and thus flowering (Liu et al. 2008a; Turck et al. 2008). How interaction between the CRY2 PHR domain and CIB1 controls CIB1 activity remains unknown at present.

### **Additional roles of the SPA1-COP1 E3 ligase in light signaling**

To better understand how blue light controls plant growth and development, it is necessary to characterize the transcription factors controlled by cryptochromes via suppression of COP1 activity. In addition to HY5, as described above, several other transcription factors, including the bHLH factor HFR1, are required for de-etiolation to take place. Some of these transcription factors are also degraded in a COP1-dependent mechanism in the dark (Henriques et al. 2009). Interestingly, SPA1 also interacts with HFR1 and it has been shown that HFR1 is important for

cryptochrome-mediated de-etiolation, suggesting a similar mode of regulation for HFR1 as exists for HY5 (Henriques et al. 2009). The regulated abundance of multiple transcription factors during de-etiolation may thus contribute to the severe phenotype displayed by *cop1* mutants during young seedling development. In addition, given that COP1 controls accumulation of HFR1, which inhibits the activity of the skotomorphogenesis-promoting bHLH proteins from the PIF family, COP1-mediated degradation indirectly controls the activity of other transcription factors (Leivar et al. 2008; Hornitschek et al. 2009).

Interestingly, COP1-associated degradation machinery has also been implicated in feedback mechanisms controlling the level of activated light sensors. During phytochrome signaling, COP1 contributes to the degradation of the light-activated photoreceptor, indicating the presence of a feedback loop (Seo et al. 2004; Jang et al. 2010). In UV-B signaling, such a feedback loop has also been described but operating via a different mechanism. To begin with, two WD40 proteins related to COP1 are transcriptionally induced by UV-B. Once accumulated, these WD40 proteins then interact with UVR8 to negatively regulate its function (Gruber et al. 2010). It is currently unknown whether a related feedback loop mechanism exists in cryptochrome-regulated gene expression. Also, the exact role of COP1 in cryptochrome stability has not been elucidated.

Light controls activity of the SPA/COP1 E3 ligase via regulated protein-protein interactions (Fig. 1B,C). Additional layers of regulation result from the light-controlled abundance of SPA family members and the distinct functions among members of this family, which can all interact with COP1 (Chen et al. 2010). For example, *SPA1* transcript increases rapidly and transiently during de-etiolation (Fittinghoff et al. 2006). In contrast, *SPA2*, which is required in dark-grown seedlings but is dispensable in the light, is constitutively expressed (Fittinghoff et al. 2006). Moreover, both SPA1 and SPA2 proteins are degraded in a proteasome-dependent manner in response to light, although SPA2 is degraded more rapidly than SPA1 (Balcerowicz et al. 2010). Finally, SPA1 and SPA2 proteins have distinct roles with SPA2 being inherently incapable of repressing photomorphogenesis in the light (Balcerowicz et al. 2010). This difference is apparently not due to an inability of SPA2 to interact with CRY1, given that all members of the SPA family can interact with the CCE domain of CRY1 (Lian et al. 2011). However, the regulated abundance of SPA family members is predicted to lead to different COP1/SPA complexes that may have distinct functions (Chen et al. 2010).

### **Animal cryptochromes control degradation of transcriptional regulators**

The ability of light acting via the cryptochromes to reset the circadian clock is best understood in *Drosophila*. As in numerous species the circadian clock in *Drosophila* relies on a transcriptional negative feedback loop. CLOCK and CYCLE act as heterodimeric transcription factors to activate expression of the negative elements PERIOD and TIMELESS (Young and Kay 2001). The *Drosophila* type I cryptochrome dCRY synchronizes the circadian clock through light-regulated protein degradation events that have been elucidated only recently. This regulation mechanism has astonishing similarities with how plant

cryptochromes control transcription factor abundance. dCRY controls light-dependent degradation of TIMELESS in a manner requiring the E3 ubiquitin ligase JETLAG, which will also ultimately lead to dCRY degradation (Koh et al. 2006; Peschel et al. 2009; Ozturk et al. 2011). Both TIMELESS and JETLAG interact with dCRY in a light-dependent manner. The interaction between JETLAG and dCRY is enabled by a light-induced conformational change of the short dCRY CCE domain (Ozturk et al. 2011). Interestingly, a dCRY mutant lacking the CCE domain is constitutively active and interacts with JETLAG independently of light, highlighting the importance of this distinguishing feature of cryptochromes (Busza et al. 2004; Ozturk et al. 2011). Thus, in *Drosophila*, light stimulus causes a conformational change in the CCE domain of dCRY. This then leads to an interaction between dCRY and an E3 ubiquitin ligase, which controls abundance of the transcriptional regulator TIMELESS. The analogy between recently described plant mechanisms and the mode of action for dCRY is noteworthy. In both cases, light induces a conformational change in the CCE domain, which is central for cryptochrome activity, and also in both cases, this conformational change directly affects downstream degradation of transcription factors. However, in plants, constitutive signaling is conferred by the isolated CCE domain whereas, in *Drosophila*, the isolated PHR domain is constitutively active. Furthermore, in *Drosophila*, dCRY activation leads to degradation of TIMELESS whilst, in *Arabidopsis*, CRY activation leads to stabilization of HY5 and CO.

### **Concluding remarks**

Recent publications showing how plant cryptochromes inactivate the COP1 complex in response to blue light give rise to a number of further questions that need to be addressed: Can a similar molecular mechanism be projected to phytochrome and UV-B receptor pathways? How exactly do SPA proteins regulate E3 ubiquitin ligase activity of the COP1 complex? How can a light-induced interaction between SPA1 and the CRY2 PHR be reconciled with the constitutively active form of the CRY2 CCE domain? Finally, although the genetic and molecular experiments presented in the aforementioned publications show that the cryptochromes suppress COP1/SPA mediated degradation of HY5 and CO, it is surprising to observe that both proteins still accumulate in a blue-light dependent manner in *spa cry* mutants (Liu et al. 2011; Zuo et al. 2011). It will be interesting to uncover the mechanism behind this additional layer of light regulation.








## **Acknowledgements**

We apologize to colleagues whose work could not be cited in this brief perspective. We are grateful to Séverine Lorrain, Emilie Demarsy and Kimberley Tilbrook for their comments on this manuscript. Work in the Fankhauser laboratory is supported by the University of Lausanne, the Swiss National Science Foundation (grant n° 31003A-124747), the NCCR “Plant Survival in Natural and Agricultural Ecosystems” and the SystemsX.ch RTD “Plant growth in a changing environment”. Work in the Ulm laboratory is supported by the University of Geneva, the Emmy Noether Programme and the Swiss National Science Foundation (grant n° 31003A\_132902).

## Figure legend

Figure 1. A light-induced conformational change in the CCE domain of plant cryptochromes inhibits COP1-mediated degradation of transcription factors. (A) In plants the cryptochromes (CRY1 and CRY2 in Arabidopsis) control two major developmental transitions, which are controlled by the light-regulated abundance of transcription factors. In the dark, seedlings grow from their seed reserves. Upon transfer into the light, several photoreceptors, including the cryptochromes, promote photoautotrophic growth that is accompanied by major developmental adaptations such as development of chloroplasts, expansion of the leaves and inhibition of stem growth. Flowering in many plant species is controlled by day length. The cryptochromes, particularly CRY2 in Arabidopsis, are major contributors of photoperiodic induction of reproductive development. (B) A model depicting how plant CRY1 regulates activity of a multimeric E3 ubiquitin ligase composed by COP1, SPA proteins, RBX1, DDB1 and cullin 4 (the latter three are not shown on this figure). Schematic depiction of the domain structure of SPA1 and CRY1. KRD = Kinase Related Domain; CC = Coiled Coil domain; WD = WD40 repeat domain; PHR = Photolyase Homology Region; CCE = Cryptochrome C-terminal Extension. Parts of the proteins involved in the interaction are indicated. Cryptochromes act as dimers and are composed of two domains, the PHR and the CCE, the latter of which changes conformation upon light activation. In the dark, the CCE domain comes into direct contact with COP1 but not SPA proteins. However, COP1 and SPA interact with each other promoting E3 ubiquitin ligase activity of the complex leading to degradation of important transcription factors such as HY5. In the light, SPA interacts directly with the CRY1 CCE domain, preventing further interaction with COP1. This inhibits E3 ligase activity leading to the accumulation of HY5 during de-etiolation. (C) A model depicting the mechanism of CRY2 controlled CO accumulation. Schematic depiction of the domain structure of SPA1 and CRY2. KRD = Kinase Related Domain; CC = Coiled Coil domain; WD = WD40 repeat domain; PHR = Photolyase Homology Region; CCE = Cryptochrome C-terminal Extension. Parts of the proteins involved in the interaction are indicated. In short-day conditions the SPA/COP1 multimeric E3 ligase degrades the transcription factor CO. Increased day length leads to longer CRY2 activation and thus interaction between SPA and the PHR domain of CRY2. This leads to a tighter interaction between COP1 and CRY2 and subsequently, inhibition of ubiquitin E3 ligase activity resulting in stabilization of CO.

**A****B****C**

## References

- Ahmad M, Cashmore AR. 1993. HY4 gene of *A. thaliana* encodes a protein with characteristics of a blue-light photoreceptor. *Nature* **366**: 162-166.
- Balcerowicz M, Fittinghoff K, Wirthmueller L, Maier A, Fackendahl P, Fiene G, Koncz C, Hoecker U. 2010. Light exposure of *Arabidopsis* seedlings causes rapid de-stabilization as well as selective post-translational inactivation of the repressor of photomorphogenesis SPA2. *Plant J.*
- Busza A, Emery-Le M, Rosbash M, Emery P. 2004. Roles of the two *Drosophila* CRYPTOCHROME structural domains in circadian photoreception. *Science* **304**: 1503-1506.
- Cashmore AR. 2003. Cryptochromes: enabling plants and animals to determine circadian time. *Cell* **114**: 537-543.
- Chen H, Huang X, Gusmaroli G, Terzaghi W, Lau OS, Yanagawa Y, Zhang Y, Li J, Lee JH, Zhu D et al. 2010. *Arabidopsis* CULLIN4-damaged DNA binding protein 1 interacts with CONSTITUTIVELY PHOTOMORPHOGENIC1-SUPPRESSOR OF PHYA complexes to regulate photomorphogenesis and flowering time. *Plant Cell* **22**: 108-123.
- Favory JJ, Stec A, Gruber H, Rizzini L, Oravec A, Funk M, Albert A, Cloix C, Jenkins GI, Oakeley EJ et al. 2009. Interaction of COP1 and UVR8 regulates UV-B-induced photomorphogenesis and stress acclimation in *Arabidopsis*. *EMBO J* **28**: 591-601.
- Fittinghoff K, Laubinger S, Nixdorf M, Fackendahl P, Baumgardt RL, Batschauer A, Hoecker U. 2006. Functional and expression analysis of *Arabidopsis* SPA genes during seedling photomorphogenesis and adult growth. *Plant J* **47**: 577-590.
- Gruber H, Heijde M, Heller W, Albert A, Seidlitz HK, Ulm R. 2010. Negative feedback regulation of UV-B-induced photomorphogenesis and stress acclimation in *Arabidopsis*. *Proc Natl Acad Sci U S A* **107**: 20132-20137.
- Henriques R, Jang IC, Chua NH. 2009. Regulated proteolysis in light-related signaling pathways. *Curr Opin Plant Biol* **12**: 49-56.
- Hoecker U, Quail PH. 2001. The phytochrome A-specific signaling intermediate SPA1 interacts directly with COP1, a constitutive repressor of light signaling in *Arabidopsis*. *J Biol Chem* **276**: 38173-38178.
- Hornitschek P, Lorrain S, Zoete V, Michielin O, Fankhauser C. 2009. Inhibition of the shade avoidance response by formation of non-DNA binding bHLH heterodimers. *EMBO J* **28**: 3893-3902.
- Jang IC, Henriques R, Seo HS, Nagatani A, Chua NH. 2010. *Arabidopsis* PHYTOCHROME INTERACTING FACTOR proteins promote phytochrome B polyubiquitination by COP1 E3 ligase in the nucleus. *Plant Cell* **22**: 2370-2383.
- Jang S, Marchal V, Panigrahi KC, Wenkel S, Soppe W, Deng XW, Valverde F, Coupland G. 2008. *Arabidopsis* COP1 shapes the temporal pattern of CO accumulation conferring a photoperiodic flowering response. *EMBO J* **27**: 1277-1288.
- Kami C, Lorrain S, Hornitschek P, Fankhauser C. 2010. Light-regulated plant growth and development. *Curr Top Dev Biol* **91**: 29-66.

- Kennedy MJ, Hughes RM, Peteya LA, Schwartz JW, Ehlers MD, Tucker CL. 2010. Rapid blue-light-mediated induction of protein interactions in living cells. *Nat Methods* **7**: 973-975.
- Koh K, Zheng X, Sehgal A. 2006. JETLAG resets the *Drosophila* circadian clock by promoting light-induced degradation of TIMELESS. *Science* **312**: 1809-1812.
- Kottke T, Batschauer A, Ahmad M, Heberle J. 2006. Blue-light-induced changes in *Arabidopsis* cryptochrome 1 probed by FTIR difference spectroscopy. *Biochemistry* **45**: 2472-2479.
- Laubinger S, Fittinghoff K, Hoecker U. 2004. The SPA quartet: a family of WD-repeat proteins with a central role in suppression of photomorphogenesis in *Arabidopsis*. *Plant Cell* **16**: 2293-2306.
- Laubinger S, Marchal V, Le Gourrierec J, Wenkel S, Adrian J, Jang S, Kulajta C, Braun H, Coupland G, Hoecker U. 2006. *Arabidopsis* SPA proteins regulate photoperiodic flowering and interact with the floral inducer CONSTANS to regulate its stability. *Development* **133**: 3213-3222.
- Leivar P, Monte E, Oka Y, Liu T, Carle C, Castillon A, Huq E, Quail PH. 2008. Multiple phytochrome-interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness. *Curr Biol* **18**: 1815-1823.
- Lian H-L, He S-B, Zhang Y-C, Zhu D-M, Zhang J-Y, Sun S-X, Li L, Yang H-Q. 2011. Blue light-dependent interaction of cryptochrome 1 with SPA1 defines a dynamic signaling mechanism. *Genes and Development* **in press**.
- Lin C, Shalitin D. 2003. Cryptochrome structure and signal transduction. *Annu Rev Plant Biol* **54**: 469-496.
- Liu B, Liu H, Zhong D, Lin C. 2010. Searching for a photocycle of the cryptochrome photoreceptors. *Curr Opin Plant Biol* **13**: 578-586.
- Liu B, Zuo Z, Liu H, Liu X, Lin C. 2011. *Arabidopsis* cryptochrome 1 interacts with SPA1 to suppress COP1 activity in response to blue light. *Genes and Development*.
- Liu H, Yu X, Li K, Klejnot J, Yang H, Lisiero D, Lin C. 2008a. Photoexcited CRY2 interacts with CIB1 to regulate transcription and floral initiation in *Arabidopsis*. *Science* **322**: 1535-1539.
- Liu LJ, Zhang YC, Li QH, Sang Y, Mao J, Lian HL, Wang L, Yang HQ. 2008b. COP1-mediated ubiquitination of CONSTANS is implicated in cryptochrome regulation of flowering in *Arabidopsis*. *Plant Cell* **20**: 292-306.
- Ozturk N, Selby CP, Annayev Y, Zhong D, Sancar A. 2011. Reaction mechanism of *Drosophila* cryptochrome. *Proc Natl Acad Sci U S A* **108**: 516-521.
- Partch CL, Clarkson MW, Ozgur S, Lee AL, Sancar A. 2005. Role of structural plasticity in signal transduction by the cryptochrome blue-light photoreceptor. *Biochemistry* **44**: 3795-3805.
- Peschel N, Chen KF, Szabo G, Stanewsky R. 2009. Light-dependent interactions between the *Drosophila* circadian clock factors cryptochrome, jetlag, and timeless. *Curr Biol* **19**: 241-247.
- Reppert SM, Gegear RJ, Merlin C. 2010. Navigational mechanisms of migrating monarch butterflies. *Trends Neurosci* **33**: 399-406.
- Saijo Y, Sullivan JA, Wang H, Yang J, Shen Y, Rubio V, Ma L, Hoecker U, Deng XW. 2003. The COP1-SPA1 interaction defines a critical step in phytochrome A-mediated regulation of HY5 activity. *Genes Dev* **17**: 2642-2647.

- Seo HS, Watanabe E, Tokutomi S, Nagatani A, Chua NH. 2004. Photoreceptor ubiquitination by COP1 E3 ligase desensitizes phytochrome A signaling. *Genes Dev* **18**: 617-622.
- Seo HS, Yang JY, Ishikawa M, Bolle C, Ballesteros ML, Chua NH. 2003. LAF1 ubiquitination by COP1 controls photomorphogenesis and is stimulated by SPA1. *Nature* **423**: 995-999.
- Shalitin D, Yang H, Mockler TC, Maymon M, Guo H, Whitelam GC, Lin C. 2002. Regulation of Arabidopsis cryptochrome 2 by blue-light-dependent phosphorylation. *Nature* **417**: 763-767.
- Shalitin D, Yu X, Maymon M, Mockler T, Lin C. 2003. Blue light-dependent in vivo and in vitro phosphorylation of Arabidopsis cryptochrome 1. *Plant Cell* **15**: 2421-2429.
- Turck F, Fornara F, Coupland G. 2008. Regulation and identity of florigen: FLOWERING LOCUS T moves center stage. *Annu Rev Plant Biol* **59**: 573-594.
- Wang H, Ma LG, Li JM, Zhao HY, Deng XW. 2001. Direct interaction of Arabidopsis cryptochromes with COP1 in light control development. *Science* **294**: 154-158.
- Wu G, Spalding EP. 2007. Separate functions for nuclear and cytoplasmic cryptochrome 1 during photomorphogenesis of Arabidopsis seedlings. *Proc Natl Acad Sci U S A* **104**: 18813-18818.
- Yang HQ, Tang RH, Cashmore AR. 2001. The signaling mechanism of Arabidopsis CRY1 involves direct interaction with COP1. *Plant Cell* **13**: 2573-2587.
- Yang HQ, Wu YJ, Tang RH, Liu D, Liu Y, Cashmore AR. 2000. The C termini of Arabidopsis cryptochromes mediate a constitutive light response. *Cell* **103**: 815-827.
- Yang J, Wang H. 2006. The central coiled-coil domain and carboxyl-terminal WD-repeat domain of Arabidopsis SPA1 are responsible for mediating repression of light signaling. *Plant J* **47**: 564-576.
- Yang YJ, Zuo ZC, Zhao XY, Li X, Klejnot J, Li Y, Chen P, Liang SP, Yu XH, Liu XM et al. 2008. Blue-light-independent activity of Arabidopsis cryptochromes in the regulation of steady-state levels of protein and mRNA expression. *Mol Plant* **1**: 167-177.
- Yi C, Deng XW. 2005. COP1 - from plant photomorphogenesis to mammalian tumorigenesis. *Trends Cell Biol* **15**: 618-625.
- Young MW, Kay SA. 2001. Time zones: a comparative genetics of circadian clocks. *Nat Rev Genet* **2**: 702-715.
- Yu X, Klejnot J, Zhao X, Shalitin D, Maymon M, Yang H, Lee J, Liu X, Lopez J, Lin C. 2007a. Arabidopsis cryptochrome 2 completes its posttranslational life cycle in the nucleus. *Plant Cell* **19**: 3146-3156.
- Yu X, Shalitin D, Liu X, Maymon M, Klejnot J, Yang H, Lopez J, Zhao X, Bendehakalu KT, Lin C. 2007b. Derepression of the NC80 motif is critical for the photoactivation of Arabidopsis CRY2. *Proc Natl Acad Sci U S A* **104**: 7289-7294.
- Zoltowski BD, Gardner KH. 2010. Tripping the Light Fantastic: Blue-Light Photoreceptors as Examples of Environmentally Modulated Protein-Protein Interactions. *Biochemistry*.

Zuo Z, Liu H, Liu B, Liu X, Lin C. 2011. Arabidopsis cryptochrome 2 undergoes blue light-dependent interaction with the SPA1-COP1 complex to regulate floral initiation in plants. *Current Biology* **in press**.