

Serveur Académique Lausannois SERVAL [serval.unil.ch](http://serval.unil.ch)

## Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

**Title:** Pasteurella multocida zoonotic ascending infection: an unusual cause of tubo-ovarian abscess.

**Authors:** Baud D, Bizzini A, Jaton K, Achtari C, Prod'homme G, Greub G

**Journal:** Vector borne and zoonotic diseases

**Year:** 2012 Jan

**Volume:** 12

**Issue:** 1

**Pages:** 84-5

**DOI:** 10.1089/vbz.2011.0610

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

Vector-Borne and  
Zoonotic Diseases

Vector-Borne and Zoonotic Diseases: <http://mc.manuscriptcentral.com/vbz>

**PASTEURELLA MULTOCIDA ZOOBOTIC ASCENDING  
INFECTION:  
AN UNUSUAL CAUSE OF TUBOOVARIAN ABSCESS**

Journal:	<i>Vector-Borne and Zoonotic Diseases</i>
Manuscript ID:	VBZ-2011-0610.R1
Manuscript Type:	Short Communication
Date Submitted by the Author:	n/a
Complete List of Authors:	Baud, David; University Hospital Centre and University of Lausanne, Maternity, Obstetrics and Gynecology Bizzini, Alain; University Hospital Centre and University of Lausanne, Institute of Microbiology Jaton, Katia; University Hospital Centre and University of Lausanne, Institute of Microbiology Achtari, Chahin; University Hospital Centre and University of Lausanne, Maternity, Obstetrics and Gynecology Prod'hom, Guy; University Hospital Centre and University of Lausanne, Institute of Microbiology Greub, Gilbert; University Hospital Centre and University of Lausanne, Institute of Microbiology; University Hospital Centre and University of Lausanne, Infectious diseases Service
Keyword:	Zoonotic
Abstract:	We report a tubo-ovarian abscess due to <i>Pasteurella multocida</i> . This zoonotic infection was likely of ascending origin, since <i>Pasteurella</i> was also isolated from vaginal swab.

SCHOLARONE™  
Manuscripts

1  
2  
3 ***PASTEURELLA MULTOCIDA* ZOOONOTIC ASCENDING INFECTION:**  
4  
5 **AN UNUSUAL CAUSE OF TUBOOVARIAN ABSCESS**  
6  
7  
8  
9

10 **David Baud<sup>1\*</sup>, Alain Bizzini<sup>2\*</sup>, Katia Jatton<sup>2</sup>, Chahin Ahtari<sup>1</sup>, Guy Prod'hom<sup>2</sup> and**  
11 **Gilbert Greub<sup>2,3</sup>**  
12  
13

14  
15 \* Equally contributing authors  
16  
17  
18  
19

20 <sup>1</sup> Department of Obstetrics and Gynaecology, University Hospital of Lausanne, Maternity-  
21 CHUV, Lausanne, Switzerland  
22  
23

24 <sup>2</sup> Institute of Microbiology, University Hospital Centre and University of Lausanne,  
25 Lausanne, Switzerland  
26  
27

28 <sup>3</sup> Infectious diseases Service, University Hospital Centre and University of Lausanne,  
29 Lausanne, Switzerland  
30  
31  
32  
33  
34  
35  
36

37 **Corresponding author:** Gilbert Greub  
38  
39 Institute of Microbiology  
40  
41 University Hospital Center and University of Lausanne  
42  
43 Bugnon 48  
44  
45 1011 Lausanne  
46  
47 SWITZERLAND  
48  
49  
50  
51 Phone: (00) 41 21 314 49 79  
52  
53 Fax: (00) 41 21 314 40 60  
54  
55 Email: [gilbert.greub@chuv.ch](mailto:gilbert.greub@chuv.ch)  
56  
57  
58  
59  
60

**Word count:** 487 (main text), 25 (abstract)

**ABSTRACT:**

We report a tubo-ovarian abscess due to *Pasteurella multocida*. This zoonotic infection was likely of ascending origin, since *Pasteurella* was also isolated from vaginal swab.

**Keywords:**

*Pasteurella multocida*, tubo-ovarian abscess, zoonotic infection, urogenital infection

**TEXT**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

A 34-year old woman, gravida 1, para 1, with an uneventful past medical and surgical history, presented after two months of worsening low abdominal pain and flu-like syndrome. She exhibited no urinary, digestive or pulmonary symptoms. She had normal regular menstrual cycles and did not present any past gynaecological problems. A copper intra-uterine device (IUD) was inserted three years ago after the birth of her first child. She was living alone and denied any sexual partners for 2 years.

Her admission vital signs were normal, with temperature of 37°C. Physical examination showed diffuse tenderness of the low abdomen. Cervical examination revealed bilateral pelvic tenderness. Vaginal discharge appeared physiologic. The patient denied any contact with pets or other animals, except her parents' two dogs, who she regularly takes care of. However, she did not present any skin lesions or scratches.

Laboratory analysis showed a white cell count of 20'000 cell/ $\mu$ l and a CRP of 276 mg/l. The patient immediately received amoxicillin-clavulanate, metronidazole and doxycycline intravenously, but developed spiking fever after treatment initiation. Thoraco-abdominal CT-scan revealed a 7 cm right-sided abdominal mass and multiple retro-peritoneal adenopathies. Vaginal ultrasound confirmed the presence of an organised tubo-ovarian abscess (Figure 1A), without free abdominal liquid. The patient underwent laparoscopy, which showed a pelvic mass adherent to bowel, the omentum and to the posterior wall of the uterus. Despite inflammation of the peritoneum, there was neither intraperitoneal pus nor adhesions. There was no evidence of diverticulitis or appendicitis. During laparoscopic adhesiolysis, this adherent mass ruptured, showing a caseous thick content (Figure 1B).

Postoperatively, the patient defervesced rapidly under the same intravenous antibiotics. Serologies (HIV, HBV and HCV) were all negative. Histopathology showed a tubo-ovarian empyema. Blood, urine and IUD cultures remained sterile. All samples were PCR negative

1  
2  
3 for *Chlamydia trachomatis*, *Neisseria gonorrhoeae* and *Mycobacterium tuberculosis*. A  
4  
5 broad-spectrum 16SrRNA eubacterial PCR performed on the abdominal washing and on the  
6  
7 caseous tubo-ovarian abscess revealed *Pasteurella multocida* DNA. Later, conventional  
8  
9 cultures of these samples demonstrated the presence of *P. multocida* after enrichment step and  
10  
11 two colonies were recovered from vaginal swab primocultures. The patient was discharged on  
12  
13 day 5 post-surgery and underwent a 14-day course of oral amoxicillin-clavulanate, with a  
14  
15 favourable outcome.  
16  
17

18  
19 *Pasteurella multocida* is a gram-negative coccobacillus present in oral flora of domestic and  
20  
21 wild animals [1]. Human infection is rare and mostly seen in children or immuno-  
22  
23 compromised patients [2]. Zoonotic transmission results from pet bites or scratches [3] and  
24  
25 can cause severe infections, including pneumonia, necrotizing fasciitis, septic shock and  
26  
27 meningitis. *Pasteurella multocida* infection of the uro-genital tract has been reported [4-10]  
28  
29 related to miscarriage in pregnant women [11].  
30  
31

32  
33 Previously described *Pasteurella multocida* tubo-ovarian abscesses have been associated with  
34  
35 bacteremia secondary to pet scratches or bites [4, 6, 7]. A *Pasteurella multocida* bacteremia  
36  
37 may lead to a secondary tuboovarian localization and subsequently to vaginal discharge  
38  
39 containing the bacterium. In our case, *Pasteurella multocida* was recovered from both the  
40  
41 vaginal swab and tubo-ovarian abscess. Since the patient had neither skin bite, scratches nor a  
42  
43 positive history of such events, an ascending vaginal *Pasteurella* infection may have  
44  
45 occurred. Tubo-ovarian abscess secondary to such an uncommon organism, likely of zoonotic  
46  
47 origin, should be considered in absence of other more common etiology in the differential  
48  
49 diagnosis of pelvic inflammatory disease.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## ACKNOWLEDGMENTS

We warmly thanks Olivier Cottier for laparoscopic images, Malgorzata Kempf Haber for ultrasound images and the laboratory technicians of the diagnostic microbiology laboratory of the University Hospital of Lausanne, Switzerland. We thank Paulina Cybulska for critical review of the manuscript. David Baud is supported by the “Société Académique Vaudoise” through the “Paul Blanc” grant, the SICPA Foundation, “Air Canada Travel Grant” and the Foundation for Research and Development in Gynecology and Obstetrics of Lausanne, Switzerland. Gilbert Greub is supported by the Leenards Foundation through a career award entitled Bourse Leenards pour la relève académique en médecine clinique à Lausanne.

## DISCLOSURE STATEMENT

No competing financial interests exist.


## Reference List

- 1  
2  
3  
4  
5  
6  
7  
8 (1) Oehler RL, Velez AP, Mizrachi M, Lamarche J, Gompf S. Bite-related and septic  
9 syndromes caused by cats and dogs. *Lancet Infect Dis* 2009 Jul; 9(7):439-47.
- 10  
11  
12 (2) Kristinsson G. *Pasteurella multocida* infections. *Pediatr Rev* 2007 Dec; 28(12):472-3.
- 13  
14  
15 (3) Iaria C, Cascio A. Please, do not forget *Pasteurella multocida*. *Clin Infect Dis* 2007  
16 Oct 1; 45(7):940.
- 17  
18  
19 (4) Teng FY, Cardone JT, Au AH. *Pasteurella multocida* tubo-ovarian abscess in a virgin.  
20 *Obstet Gynecol* 1996 May; 87(5 Pt 2):883.
- 21  
22  
23 (5) Lukban JC, Baker MS. *Pasteurella multocida* isolation from a tuboovarian abscess. A  
24 case report. *J Reprod Med* 1995 Aug; 40(8):603-5.
- 25  
26  
27 (6) Rowe J, Mikuta J. Cat-scratch salpingitis. *N Engl J Med* 1992 Nov 5; 327(19):1395-6.
- 28  
29  
30 (7) Myckan KA, Booth CM, Mocarski E. *Pasteurella multocida* bacteremia and  
31 tuboovarian abscess. *Obstet Gynecol* 2005 Nov; 106(5 Pt 2):1220-2.
- 32  
33  
34 (8) Liu W, Chemaly RF, Tuohy MJ, LaSalvia MM, Procop GW. *Pasteurella multocida*  
35 urinary tract infection with molecular evidence of zoonotic transmission. *Clin Infect*  
36 *Dis* 2003 Feb 15; 36(4):E58-E60.
- 37  
38  
39 (9) Loiez C, Wallet F, Husson MO, Courcol RJ. *Pasteurella multocida* and intrauterine  
40 device: a woman and her pets. *Scand J Infect Dis* 2002; 34(6):473.
- 41  
42  
43 (10) Riley UB, De P. *Pasteurella multocida*--an uncommon cause of obstetric and  
44 gynaecological sepsis. *J Infect* 1995 Jul; 31(1):51-3.
- 45  
46  
47 (11) Waldor M, Roberts D, Kazanjian P. In utero infection due to *Pasteurella multocida* in  
48 the first trimester of pregnancy: case report and review. *Clin Infect Dis* 1992 Feb;  
49 14(2):497-500.
- 50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




**FIGURE 1**

Ultrasound (A) and laparoscopic (B) pictures showing the tubo-ovarian abscess and its caseous content.




**A**



**B**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Ultrasound (A) and laparoscopic (B) pictures showing the tubo-ovarian abscess and its caseous content.  
183x84mm (300 x 300 DPI)