
**UNIVERSITÉ DE LAUSANNE – FACULTÉ DE BIOLOGIE ET DE MÉDECINE
DEPARTEMENT DES CENTRES INTERDISCIPLINAIRES
ET LOGISTIQUE MEDICALE**

Service de médecine intensive adulte

Chef de service : Professeur René Chioléro

**Intravenous fish oil blunts the physiological response
to endotoxin in healthy subjects**

THESE

**présentée à la Faculté de biologie et de médecine de
l'Université de Lausanne pour l'obtention du grade de**

DOCTEUR EN MEDECINE

par

Thomas-Thi Pluess

BMTE 3449

Médecin diplômé de la Confédération Suisse

Originaire de Murgenthal (AG)

Lausanne

2007

*VVB
410
plv*

Intravenous fish oil blunts the physiological response to endotoxin in healthy subjects

Résumé :

L'administration par voie orale d'acides gras polyinsaturés de type ω -3 contenus dans l'huile de poisson exerce des effets bénéfiques sur la réponse métabolique et inflammatoire chez des sujets sains soumis à une injection d'endotoxine. Ce modèle expérimental a été validé pour l'investigation clinique. Il simule un sepsis et induit une réponse comparable à un état grippal, accompagné de modifications métaboliques et inflammatoires.

L'objectif de cette étude est de déterminer les effets de l'huile de poisson administré par voie intraveineuse sur la réponse à l'endotoxine chez le sujet sain. L'hypothèse est qu'il sera possible de réduire le temps de latence en comparaison avec la voie orale. Pour ce faire, nous avons inclus dans une étude prospective randomisée 16 volontaires sains âgés de 16 à 35 ans et les avons répartis en 2 groupes : l'un recevant une émulsion lipidique contenant les acides gras polyinsaturés EPA et DHA et l'autre, sans traitement, constituant le groupe contrôle. Huit sujets reçoivent une perfusion continue de 0.5g/kg d'huile de poisson durant 6h, 48h et 24h avant la journée test. Lors de cette journée test, tous les volontaires ont reçu une dose d'endotoxine (2ng/kg) au temps t0. Les paramètres vitaux sont monitorés et enregistrés : fréquence cardiaque, respiratoire, pression artérielle, saturation artérielle en oxygène, ainsi que température. Des prises de sang sont effectuées à intervalles réguliers pour déterminer :

- 1) l'incorporation membranaire des thrombocytes en EPA et DHA ;
- 2) le taux plasmatique d'hormones (insuline, glucagon, cortisol, ACTH et catécholamines), de marqueurs inflammatoires (TNF- α , IL-6, hsCRP), ainsi que de substrats énergétiques (glucose, lactate, acides gras libres et triglycérides). La dépense énergétique est déterminée par calorimétrie indirecte.

L'analyse statistique est effectuée par analyse de variance (ANOVA).

Les résultats montrent une incorporation significative de EPA et DHA au niveau membranaire des thrombocytes. L'huile de poisson induit une atténuation significative de la réponse neuro-endocrinienne et inflammatoire en réponse à l'injection d'endotoxine avec diminution de la fièvre (-0.7°C), ainsi que du taux plasmatique d'ACTH (-68%), TNF- α (-63%) et de noradrénaline (-%) dans le groupe huile de poisson.

En conclusion, cette étude montre que la supplémentation de 2 doses d'huile de poisson par voie intraveineuse modifie la composition phospholipidique des membranes des thrombocytes et diminue la réaction inflammatoire et neuroendocrinienne en réponse à l'endotoxine. Ces résultats positifs ouvrent la perspective d'une supplémentation parentérale préopératoire en acides gras polyinsaturés ω -3 pour diminuer le stress lié à la chirurgie majeure.

Thomas-Thi Pluess
Daniel Hayoz
Mette M. Berger
Luc Tappy
Jean-Pierre Revelly
Burkhard Michaeli
Yvon A. Carpentier
René L. Chioloro

Intravenous fish oil blunts the physiological response to endotoxin in healthy subjects

Received: 13 June 2006
Accepted: 21 February 2007
Published online: 22 March 2007
© Springer-Verlag 2007

T.-T. Pluess · M. M. Berger · J.-P. Revelly ·
B. Michaeli · R. L. Chioloro (✉)
CHUV, Department of Intensive Care
Medicine,
1011 Lausanne, Switzerland
e-mail: rene.chioloro@chuv.ch
Tel.: +41-21-3142002
Fax: +41-21-3143045

D. Hayoz
CHUV, Division of Hypertension and
Vascular Medicine,
1011 Lausanne, Switzerland

L. Tappy
University of Lausanne, Institute of Human
Physiology,
1011 Lausanne, Switzerland

Y. A. Carpentier
Université Libre de Bruxelles, Laboratory
of Experimental Surgery,
Brussels, Belgium

Abstract *Objective:* To assess the effects of intravenous fish oil fat emulsion on the metabolic alterations induced by lipopolysaccharide (LPS) challenge in healthy volunteers. *Design:* Two groups of eight healthy subjects were randomized to receive either two pharmacological doses of intravenous FO fat emulsion or no treatment. The FO group received twice 0.5 g/kg 10% emulsion (Omegaven) 48 and 24 h before investigation. LPS (2 ng/kg) was injected as a bolus on the investigation day. Systemic parameters, indirect calorimetry, heart rate variability, and platelet membrane phospholipid composition were measured. *Results:* Basal EPA and DHA content in platelet phospholipids was low (0.28% and 2.54%, respectively) and increased significantly after FO to 1.68% and 3.32%. LPS induced reproducible

effects in all subjects. Fever was higher in the FO group than in controls; the difference was significant from t_{120} until t_{360} . FO blunted the neuroendocrine response: the rise in plasma norepinephrine was sevenfold lower at t_{120} while the ACTH peak was fourfold lower. Tumor necrosis factor α was significantly lower between t_{360} and t_{180} in the FO group. *Conclusions:* Two doses of intravenous FO fat emulsion modified the phospholipid composition of platelets in healthy subjects. FO blunted fever and increased the neuroendocrine and the inflammatory responses to LPS.

Keywords Endotoxin challenge · Lipopolysaccharide · Fish oil supplement · Cytokines · Inflammatory response · Neuroendocrine response

Introduction

There are two main families of polyunsaturated fatty acids (PUFAs) n-6 and n-3. Among the n-3 PUFA family EPA and DHA are the most biologically active. Feeding with fish oil (FO) increases the proportion of EPA and DHA in the cell membrane phospholipids, including inflammatory cells. This decreases the synthesis of inflammatory eicosanoids by numerous mechanisms [1]: (a) substrate competition, (b) inhibition of membrane arachidonic acid release, (c) synthesis of less potent eicosanoids, and (d) downregulation of proinflammatory cytokines, adhesion molecules, and of genes involved in inflammation.

It is generally considered that the oral administration of n-3 PUFA is required for several weeks to produce biological effects, precluding its use in acute conditions. However, it has recently been observed that intravenous administration of fat emulsions enriched with n-3 PUFA leads to rapid incorporation of n-3 fatty acids into phospholipids of platelets or monocytes [2]. Modulation of the inflammatory response in septic patients is desirable as excessive release of proinflammatory cytokines is associated with multiple organ failure; FO may thus prove to be a new therapeutic tool [3, 4].

Endotoxin (LPS) administration is an experimental, reproducible model of sepsis, mimicking some important as-

pects of sepsis, and has the advantage of being short lived. This study examined whether two doses of intravenous FO downregulates the neuroendocrine and systemic responses to endotoxin in healthy volunteers in a prospective, randomized controlled study.

Materials and methods

Subjects

After approval by the institutional ethics committee and written informed consent we enrolled 16 men aged 18–35 years. Inclusion criteria were: good health, no medication, low fish consumption (less than twice a week), no nutritional supplementation, and particularly no FO supplements. A complete history, physical examination, and 12-lead electrocardiography were performed. Subjects had a mean age of 26.2 ± 2.7 years, body weight of 77.4 ± 7.2 kg, and body mass index of 23.4 ± 1.9 .

Study design and intervention

Subjects were randomized to receive two pharmacological doses of intravenous FO 48 and 24 h before investigation, a dose of 5 ml/kg during 6 h each time (100 ml Omegaven 10%: 10 g FO, providing 1.25–2.82 g EPA (n-3); 1.44–3.09 g DHA (n-3), 0.1–0.7 g linoleic acid (n-6), max. 0.2 g α -linolenic acid (n-3) and 15.0–29.6 mg α -tocopherol; Fresenius Kabi, Bad Homburg, Germany). On the investigation day subjects came after an overnight fast, and were requested to void (urine discarded). Thereafter they remained recumbent for the next 8 h. Two venous cannulas were inserted into the forearms: one for labeled glucose infusion and one for blood sampling. Blood samples were collected every 30–60 min. Endotoxin (2 ng/kg, USP Rockville, Md., USA; lot no. 2) was administered as an intravenous bolus at time t_0 .

Blood collection and analysis

In subjects who received FO fat emulsion two series of blood samples were collected (EDTA tubes): one before the first infusion and the second 48 h later, before LPS. Plasma was separated by low-speed centrifugation, and supernatant was centrifuged at 3,000/min for 10 min at 4 °C. After separation of platelet lipid components by thin-layer chromatography the fatty acid pattern was analyzed only in the phospholipid fraction since mainly phospholipid fatty acids affect cell function. Triacylglycerol, phospholipids, and cholesterylesters were first separated by a two-dimension thin-layer chromatogra-

phy. Fatty acids were analyzed by gas chromatography (Agilent, GC system, 6890A).

On the day of experiment blood was sampled during the protocol to determine hormone levels (insulin, glucagon, cortisol, ACTH, catecholamines), inflammatory markers [tumor necrosis factor (TNF) α , interleukin (IL) 6, hsCRP], glucose, lactate, free fatty acids (FFA), and triglycerides. Plasma aliquots were stored at -20 °C or -80 °C (for FFA determination). Plasma insulin (Adaltis kit, Casalecchio di Reno, Italy), glucagon (Linco Research, St. Charles, Mo., USA) and cortisol (Diagnostic Products, Los Angeles, Calif., USA) were determined by radioimmunoassay; ACTH by chemiluminescence (Nichols Institute Diagnostics, San Juan Capistrano, Calif., USA); TNF- α by photometric enzyme-linked immunosorbent assay in streptavidin-coated microtiter plates (Roche, Mannheim, Germany); IL-6 by enzyme immunoassay using an enzyme-linked polyclonal antibody specific for IL-6 (Quantikine, R&D Systems, Germany); hsCRP by turbidimetry on a Hitachi 917 (Roche). Plasma glucose and lactate concentrations were measured enzymatically (YSI 2300 STAT PLU, Yellow Spring, Ohio, USA); plasma FFA concentrations with a colorimetric method (Wako, Neuss, Germany). Urine was sampled from spontaneous voiding. Indirect calorimetry was performed using a canopy (Deltatrac2 model Datex, Kuopio, Finland). Resting energy expenditure (REE) was calculated from oxygen consumption (VO_2) and CO_2 production (VCO_2) using the Livesey and Elia [5] equation.

Vital functions

Heart rate, respiratory rate, rectal temperature (Hellige, Servomed, Germany), noninvasive arterial blood pressure (Critikon Dinamap, Tampa, Fla., USA), cardiac output (thoracic bioimpedance, NCCOM3 cardiodynamic monitor, BoMed, Irvine Calif., USA) and transcutaneous oxygen saturation (SpO_2 , Pulse oximeter Biox 3740, Ohmeda, Engelwood, Calif., USA) were monitored. Subjective symptoms such as muscle and head ache, and nausea/vomiting were recorded.

Baroreflex sensitivity and heart rate variability

Baroreflex sensitivity was derived from the spontaneous fluctuations in systolic blood pressure (SBP) and pulse rate. We used the sequence technique, which focuses on the baroreflex response to pressure transients [6, 7]. Blood pressure (BP) was measured continuously by a Finapres BP Monitor (Ohmeda). SBP and pulse series were scanned automatically in search for spontaneous sequences of three or more consecutive heart beats in which a progressive SBP increase was followed, with

a one-beat lag, by a progressive pulse rate lengthening or vice versa. The slope of the regression line between SBP and pulse interval values forming each sequence was taken as an estimation of the sensitivity of baroreflex cardiac modulation.

ECG, BP, and respiration were recorded at a sampling frequency of 200 Hz. Beat-to-beat series of R-R interval were derived for heart rate variability (HRV) analysis as were series of SBP and pulse rate for baroreflex sensitivity estimation. Signals were edited manually from artifacts such as movements and noise. Beat-to-beat series were interpolated and resampled at 5 Hz to obtain an evenly sampled series and then split into data segments of 120 s. Segments were overlapped by 50% and a fast Fourier transform spectrum was computed in each window. All fast Fourier transform spectra were averaged to obtain a single final spectrum, which was integrated over the low (LF; 0.05–0.15 Hz) and high (HF; 0.15–0.50 Hz) frequency bands to characterize the LF and HF powers and LF/HF power ratio. Measures were recorded three times (30 min): at baseline and at 120 and 360 min.

Statistical analysis

Results are given as means \pm SD. One-way and two-way analyses of variance for repeated measures (time, group, and interaction) were used to compare groups where appropriate. Differences with *p* value less than 0.05 were considered statistically significant. Analyses used the statistical package JMP 5.1 (SAS Institute, Cary, N.C., USA).

Results

Physiological response to endotoxin

LPS induced reproducible systemic, endocrine, and metabolic effects in both groups, starting about 75 min after LPS injection (Table 1). All subjects complained of headache, nausea, and muscle pain.

Systemic effects

Significant increases in body temperature ($+1.9^\circ\text{C}$, $p < 0.0001$), heart rate ($+52\%$, $p < 0.0001$), and cardiac output ($+40\%$, $p < 0.0001$), peaking at t_{240} after LPS were observed in control subjects. Systolic, diastolic, mean blood pressure, and SpO₂ did not change significantly.

Hormones, cytokines, and metabolism

LPS induced a marked endocrine response in control subjects (Table 2). At t_{240} ACTH levels were increased 15 times ($p < 0.0001$), cortisol 3 times ($p < 0.0001$), and glucagon 60% ($p = 0.012$). Both plasma norepinephrine (peak at t_{120} , $+252\%$, $p < 0.001$) and epinephrine (peak at t_{240} , $+117\%$, $p < 0.002$) increased strongly. Insulin levels showed a temporary decrease at t_{120} and then reached basal levels. Following LPS TNF- α levels increased 110 times at t_{120} ($p < 0.0001$) while IL-6 levels peaked at t_{180} ($p < 0.0001$). Plasma glucose

Table 1 Effect of endotoxin on the systemic variables, baroreflex sensitivity, and heart rate variability

	t_0	t_{30}	t_{60}	t_{120}	t_{180}	t_{240}	t_{360}
Control							
Temperature ($^\circ\text{C}$)	36.8 ± 0.3	–	36.9 ± 0.2	$38.0 \pm 0.5^{**}$	$38.6 \pm 0.5^{**}$	$38.7 \pm 0.3^{**}$	$38.0 \pm 0.3^{**}$
Heart rate (bpm)	59 ± 7	–	64 ± 6	$75 \pm 8^{**}$	$87 \pm 9^{**}$	$88 \pm 9^{**}$	$82 \pm 9^{**}$
Cardiac output (%)	100	–	96 ± 12	118 ± 16	$131 \pm 33^*$	$140 \pm 45^{**}$	$125 \pm 30^*$
Respiratory rate (bpm)	15 ± 1	–	15 ± 2	$18 \pm 2^{**}$	$17 \pm 3^*$	15 ± 2	13 ± 1
Baroreflex sensitivity ($\text{m s}^{-1} \text{mmHg}^{-1}$)	–	12.25 ± 3.67	–	11.23 ± 4.61	–	–	5.78 ± 2.04
LF (m s^{-2})	–	1499 ± 1486	–	2134 ± 1226	–	–	$627 \pm 538^*$
HF (m s^{-2})	–	797 ± 845	–	753 ± 517	–	–	$144 \pm 117^*$
LF/HF ratio	–	2.0 ± 0.8	–	$3.7 \pm 1.5^*$	–	–	$4.7 \pm 1.6^{**}$
Fish oil							
Temperature ($^\circ\text{C}$)	36.7 ± 0.1	–	36.8 ± 0.2	$37.3 \pm 0.3^{**}$	$37.9 \pm 0.3^{**}$	$38.1 \pm 0.2^{**}$	$37.7 \pm 0.3^{**}$
Heart rate (bpm)	63 ± 10	–	64 ± 7	$78 \pm 7^{**}$	$89 \pm 4^{**}$	$88 \pm 5^{**}$	$78 \pm 10^{**}$
Cardiac output (%)	100	–	101 ± 7	113 ± 18	$137 \pm 27^{**}$	$134 \pm 17^*$	$121 \pm 17^*$
Respiratory rate (bpm)	14 ± 2	–	14 ± 1	16 ± 2	16 ± 2	14 ± 1	14 ± 1
Baroreflex sensitivity ($\text{m s}^{-1} \text{mmHg}^{-1}$)	–	12.37 ± 3.04	–	10.07 ± 5.68	–	–	8.03 ± 2.42
LF (m s^{-2})	–	1480 ± 1169	–	1884 ± 1966	–	–	$754 \pm 592^*$
HF (m s^{-2})	–	897 ± 780	–	751 ± 1068	–	–	$301 \pm 394^*$
LF/HF ratio	–	1.7 ± 0.7	–	$3.2 \pm 1.1^*$	–	–	$3.8 \pm 1.7^*$

* $p < 0.05$, ** $p < 0.01$ vs. baseline

Table 2 Effect of endotoxin on hormones, cytokines and energy metabolism

	t ₀	t ₆₀	t ₁₂₀	t ₁₈₀	t ₂₄₀	t ₃₆₀
Control						
ACTH (ng/l)	23 ± 7	–	152 ± 66*	–	347 ± 204*	36 ± 28
Cortisol (nmol/l)	273 ± 92	–	551 ± 123*	–	678 ± 208*	458 ± 222*
Epinephrine (pg/ml)	37 ± 21	–	68 ± 30*	–	79 ± 51*	44 ± 15
Norepinephrine (pg/ml)	186 ± 51	–	653 ± 276*	–	169 ± 35	176 ± 51
Insulin (mU/ml)	7.7 ± 2.1	8.1 ± 2.4	7.2 ± 1.1	–	7.7 ± 1.6	7.6 ± 1.5
Glucagon (ng/l)	66 ± 22	65 ± 20	94 ± 45*	–	103 ± 28*	86 ± 28*
TNF- α (pg/ml)	5.8 ± 8.2	413.1 ± 267.1*	686.9 ± 554.7*	296.2 ± 206.4*	96.2 ± 41.6	–
IL-6 (pg/ml)	–	19 ± 30	540 ± 634*	617 ± 642*	237 ± 151	–
hs-CRP (mg/l)	0.4 ± 0.2	–	–	–	–	3.0 ± 0.9
Energy expenditure (kcal/min)	1.1 ± 0.2	1.1 ± 0.2	1.4 ± 0.2*	1.5 ± 0.3*	1.5 ± 0.2*	1.3 ± 0.2*
Glycemia (mg/dl)	89.3 ± 4.7	88.0 ± 6.0	80.4 ± 8.5*	88.5 ± 6.1	92.3 ± 6.0	93.0 ± 6.3
Lactate (mmol/l)	0.9 ± 0.2	0.9 ± 0.3	1.6 ± 0.4*	1.3 ± 0.4	1.0 ± 0.3	0.9 ± 0.2
Free fatty acids (mmol/l)	0.40 ± 0.12	0.46 ± 0.15	0.75 ± 0.28*	0.84 ± 0.30*	0.97 ± 0.25*	1.03 ± 0.22*
Triglycerides (mmol/l)	0.61 ± 0.21	0.55 ± 0.17	0.58 ± 0.21	0.64 ± 0.25	0.70 ± 0.31	0.60 ± 0.25
Fish oil						
ACTH (ng/l)	29 ± 17	–	109 ± 80*	–	110 ± 98*	20 ± 12
Cortisol (nmol/l)	205 ± 38	–	441 ± 114*	–	529 ± 219*	281 ± 180
Epinephrine (pg/ml)	18 ± 11	–	37 ± 20	–	38 ± 21	33 ± 13
Norepinephrine (pg/ml)	195 ± 77	–	266 ± 78	–	179 ± 56	194 ± 112
Insulin (mU/ml)	10.7 ± 4.0	10.8 ± 3.5	9.9 ± 3.0	–	9.2 ± 2.4	9.4 ± 2.5
Glucagon (ng/l)	65 ± 12	63 ± 11	80 ± 20	–	81 ± 21	72 ± 18
TNF- α (pg/ml)	7.1 ± 8.4	130.5 ± 98.3	255.4 ± 221.5*	118.6 ± 67.2	47.8 ± 29.1	–
IL-6 (pg/ml)	–	4.8 ± 10	522 ± 680*	595 ± 900*	227 ± 356	–
hs-CRP (mg/l)	0.5 ± 0.3	–	–	–	–	2.6 ± 1.4
Energy expenditure (kcal/min)	1.1 ± 0.1	1.1 ± 0.1	1.2 ± 0.2	1.4 ± 0.3*	1.3 ± 0.2*	1.2 ± 0.2
Glycemia (mg/dl)	92.2 ± 5.6	90.6 ± 4.5	85.9 ± 3.2*	91.4 ± 6.5	94.7 ± 8.2	93.8 ± 8.6
Lactate (mmol/l)	1.1 ± 0.3	0.9 ± 0.2	1.2 ± 0.4	1.2 ± 0.6	1.1 ± 0.4	1.0 ± 0.4
Free fatty acids (mmol/l)	0.24 ± 0.11	0.28 ± 0.11	0.33 ± 0.18	0.46 ± 0.18*	0.62 ± 0.26*	0.64 ± 0.22*
Triglycerides (mmol/l)	0.99 ± 0.73	0.87 ± 0.54	0.84 ± 0.42	0.81 ± 0.39*	0.76 ± 0.36*	0.60 ± 0.31*

* $p < 0.05$, ** $p < 0.01$ vs. baseline

concentration changes after LPS were not statistically significant. Plasma lactate increased from t₆₀ and peaked at t₁₂₀ (+76%, $p = 0.09$). A sustained increase in REE (+38%, $p < 0.0001$) starting at t₉₀ was observed. FFA increased significantly from baseline until the end of the test (+162%, $p < 0.0001$). Net fat oxidation increased significantly (+210%, $p = 0.001$).

Baroreflex sensitivity and heart rate variability

Baroreflex sensitivity decreased significantly between baseline and the end of the test ($p = 0.002$). LF power tended to increase from t₃₀ to t₁₂₀ and then markedly decreased at t₃₆₀ ($p = 0.04$ from t₁₂₀ to t₃₆₀). HF power remained stable between t₃₀ and t₁₂₀ and then significantly decreased at t₃₆₀ ($p = 0.012$). The LF/HF ratio, a surrogate for sympathovagal balance, increased significantly over time ($p < 0.0001$).

Effect of FO on the physiological response to LPS

Baseline plasma lipids did not differ: cholesterol 4.4 ± 0.3 mmol/l (control) vs. 4.2 ± 0.3 (FO) and triglycerides 0.99 ± 0.72 vs. 0.61 ± 0.21 mmol/l. The triglyceride profile differed during the study between groups ($p < 0.001$), with a progressive decrease in the control groups and stable levels in the FO group in relation to lipid administration (Table 2). The composition of platelet membrane was affected by FO infusions, with significant incorporation of EPA ($p = 0.0002$) and DHA ($p = 0.02$, Fig. 1). Significant differences were observed in body temperature and the neuroendocrine response between the FO and control groups. The increase in body temperature was attenuated in the FO group after LPS: peak temperature reached 38.1 °C (+1.4 °C) in the FO group vs. 38.7 °C (+1.9 °C) in controls. The difference was significant from t₁₂₀ until t₃₆₀ ($p = 0.003$ to $p = 0.022$; Fig. 2). The adrenocortical and sympathoadrenal responses

Fig. 1 Composition of platelet membrane before and after fish oil infusion expressed as percentage of membrane fatty acids: n-3 PUFA are incorporated into cell membrane phospholipids after two doses of fish oil. (Data available for only five volunteers due to technical problems)






Fig. 2 Course of plasma TNF- α and temperature in the fish oil and control groups after LPS injection



to LPS were blunted by FO: the rise in plasma norepinephrine was sevenfold lower ($p = 0.02$; Fig. 3). Plasma epinephrine was unchanged. ACTH peak at t_{240} was four times lower in the FO group than in controls ($p = 0.02$). Plasma cortisol was lower in the FO group but the difference was not statistically significant ($p = 0.07$). TNF- α was also significantly lower in the FO group

between t_{60} and t_{180} ($p = 0.022$ to $p = 0.038$), but IL-6 values did not differ between groups. REE variability was substantial in both groups, particularly in the subjects with marked shivering (trend to lower REE in the FO group). Heart rate, cardiac output, respiratory rate, arterial pressure, baroreflex sensitivity, and HRV response to LPS did not differ between groups.

Fig. 3 Plasma concentrations of ACTH and norepinephrine in the fish oil and control groups after LPS injection



Discussion

In this human model of intravenous LPS FO lipid emulsion blunted important aspects of the response. Fever was reduced while the hypothalamo-pituitary adrenal and sympathetic responses were blunted. Some aspects of the inflammatory response as well as the sympathovagal balance were altered.

Effects of LPS injection in healthy subjects

Intravenous LPS induces an acute inflammatory state in healthy humans and animals, which mimics some pathophysiological aspects of sepsis [8]: a flulike state rapidly develops, with fever, muscle ache, tachycardia, sweating, and nausea. Mononuclear cells are activated to release proinflammatory cytokines, while the complement and coagulation cascades are activated [9, 10]. A typical neuroendocrine response transiently occurs. Resting metabolic rate, glucose turnover, and protein catabolism are increased [11]. LPS also exerts a demonstrable effect on HRV, a surrogate parameter of sympathetic and parasympathetic control on heart rate. Healthy subjects receiving endotoxin present reduced HRV, as do critical care patients suffering sepsis [12]. Analysis within the frequency domain indicates that the LF/HF power ratio was increased. This observation is usually interpreted as a marker of sympathetic nerve activity. HRV was affected by LPS: HF decreased slightly compared to baseline, suggesting a small change in vagal activity. By contrast, LF power markedly increased, indicating a consistent increase in sympathetic tone. At the end of the test HF power

decreased, demonstrating a large reduction in parasympathetic activity. Changes in baroreflex sensitivity similarly reflect the modifications in sympathovagal balance. It is interesting to observe that baroreflex sensitivity changed only slightly at t_{120} , whereas it was markedly reduced at t_{360} . This suggests that under these experimental conditions baroreflex sensitivity is more dependent on changes in vagal tone than on changes in sympathetic activity.

Effects of n-3 PUFAs on acute inflammation

Various strategies have been used to affect the course of sepsis in addition to eradication of the septic or inflammatory sites. Recent research has focused on the development of specific nutrients influencing the organ and system functions as well as the responses to critical illness [13], such as n-3 PUFAs [1]. FO infusion in guinea pigs enhances survival following lethal LPS dose [14]. Intravenous FO has been shown to prevent the LPS-induced reduction in blood flow to the gut and to reduce bacterial translocation after exposure to live bacteria in the rat [15]. FO in guinea pigs and rats is associated with better perfusion in vital organs, improved metabolic adaptation, and reduced fever and weight loss [15–19]. In a previous study in healthy subjects our group showed that prolonged supplementation with oral FO blunts the neuroendocrine and inflammatory responses to LPS challenge [20]. Perioperative administration of n-3 fatty acids may favorably affect the postoperative outcome of surgical patients [21]. Recent data in critically ill patients suggest that FO supplemented parenteral nutrition improves survival, infection rate, and length of stay [4].

Central effects: stress and fever

The response to stress has traditionally been viewed as a consequence of the neuroendocrine activation, initiated by tissue injury and by cortical and noncortical signals from the central nervous system [22]. This response is characterized by the activation of the sympathoadrenal system and the hypothalamopituitary axis [23]. In healthy subjects 3–4 weeks of oral FO inhibited the adrenal activation elicited by mental stress [24], confirming previous data in subjects subjected to typhoid vaccine after 6 weeks of enteral FO [25]. However, the difference in body temperature between FO and saline was not significant, which limits the evidence. In our experiments FO given before LPS significantly attenuated fever and stress hormones production. This is an argument in favor of a central effect of n-3 PUFAs, downregulating temperature and sympathetic and adrenocortical axes.

The finding of decreased sympathetic nervous system activity in our study associated with a lower parasympathetic inhibition after FO may translate into a reduced arrhythmogenic risk. Failure to reach statistical significance with HRV and baroreflex variability may be related to the fact these variables are indirect evaluations of sympathetic activity, with relatively low sensitivity compared to more direct approaches such as microneurography or plasma norepinephrine turnover. This result is in agreement with the significant suppression by FO of plasma norepinephrine, a component of sympathetic activation. The relative inhibition of the central and sympathetic activity and the adrenal medulla could not be assessed with the present study design.

Fever is mediated by complex peripheral and central mechanisms, involving prostaglandin E₂, a mediator derived from arachidonic acid. The latter is produced in response to cytokines and other mediators. Both in experimental and clinical settings n-3 PUFA have been shown to downregulate the temperature response to inflammation [18, 20, 25]. In our study prostaglandin and their derivatives were not determined, and therefore we can only speculate that this might have been one of the underlying mechanisms of fever attenuation.

Inflammatory response

The n-3 PUFAs may modulate the inflammatory response by complex mechanisms involving the activity of transcriptional factors, such as nuclear factor κ B, peroxisome proliferator activated receptors, and sterol regulatory element binding protein as well as Toll-like receptors [26]. They also downregulate proinflammatory cytokine production by human monocytes *in vitro* and *in vivo* [27]. In patients undergoing abdominal surgery

provision of parenteral nutrition supplemented with n-3 PUFAs induced decreased levels of plasma TNF- α and IL-6 in comparison with n-6 fatty acids [28]. In the present study TNF- α concentrations increase was blunted in the FO group, confirming an anti-inflammatory effect of FO. By contrast, IL-6 plasma levels did not differ between groups, probably due to the short lasting effect of endotoxin.

Influence of the route of administration

The composition and characteristics of cell membranes is dependent on the diet composition. In Western populations the n-3 PUFA content of mononuclear cell membranes is low, as observed in our healthy volunteers. This may affect many important cell functions such as membrane fluidity, opening of ion channels, recognition and function of membrane receptors, eicosanoid synthesis, second messenger synthesis, and many signaling pathways [29]. In our study FO was infused to achieve rapid membrane incorporation, as shown by the rapid increase in the EPA and DHA cell membrane content. This corresponds to a pharmacological effect where nutrients are administered to modulate organ and system functions.

Such rapid delivery of specific compound to the target cells and organs is difficult to achieve with enteral administration of triacylglycerols containing long-chain PUFAs such as DHA and EPA since the hydrolysis of these compounds by the pancreatic lipase is rather slow. Furthermore, hydrolysis of lipoprotein triacylglycerols by lipoprotein lipase is also a slow process with a low efficiency. As a consequence the use of the intravenous route which bypasses the intestinal process of absorption and therefore decreases the delay of action seems attractive. In our study only two doses of a lipid emulsion containing 10% of FO significantly modified the cell membrane composition and the responses to endotoxin. Such a rapid effect may lead to potential indications for early intervention with FO.

Limitations of the study

The study was designed as a pharmacological investigation comparing FO lipid emulsion as a drug with no treatment. Such solution contains glycerol, phosphatidyl-choline, and some linoleic and linolenic acid and tocopherol in addition to EPA and DHA. Although these PUFAs were the principal components, and their incorporation into platelets could be confirmed, our results may also be interpreted as an effect of either of the components of the lipid emulsion. Finally lipid emulsions have been shown to blunt the response to LPS *in vitro* and *in vivo* [30]. This protection is achieved through the binding and neutralization of LPS

by circulating lipoproteins [31]. In our trial the lipoprotein profile was not determined, and we therefore cannot conclude on this aspect. Another theoretical limitation is related to the absence of saline perfusion in the control subjects, to avoid unnecessary cannulations before the study day. It seems very unlikely, however, that this design affected the results considering that basal values were measured on the day of the test only.

In conclusion, our results indicate that acute intravenous FO lipid emulsion administration was rapidly incorporated into cell membrane and downregulated the neuroendocrine and metabolic responses to LPS in

healthy humans. This suggests that intravenous FO exerts beneficial effects in acute medicine by modulating the stress and inflammatory responses.

Acknowledgements. The authors thank Mrs. Marie-Christine Cayeux (RN) and Mrs. Ludivine Soguel Alexander (RD) for their help, Mrs. A. Dufour and A. Chwalik (Université Libre de Bruxelles, Brussels, Belgium) for fatty acid analyses in platelet membranes, Ing. Paolo Castiglioni (PhD.) for helpful discussions, and Mr. P. Frascarolo (PhD.) for help with statistical outwork. The present study was performed with the support of the Swiss Fond National de Recherche, grant no. 3200B0-102064. Experimental emulsions were kindly provided by Fresenius Kabi AG (Bad Homburg, Germany).

References

- Calder PC (2004) n-3 fatty acids, inflammation, and immunity-relevance to postsurgical and critically ill patients. *Lipids* 39:1147–1161
- Mayer K, Gokorsch S, Fegbeutel C, Hattar K, Rosseau S, Walrath D, Seeger W, Grimminger F (2003) Parenteral nutrition with fish oil modulates cytokine response in patients with sepsis. *Am J Respir Crit Care Med* 167:1321–1328
- Calder PC (2006) Use of fish oil in parenteral nutrition: rationale and reality. *Proc Nutr Soc* 65:264–277
- Heller AR, Rossler S, Litz RJ, Stehr SN, Heller SC, Koch R, Koch T (2006) Omega-3 fatty acids improve the diagnosis-related clinical outcome. *Crit Care Med* 34:972–979
- Livesey G, Elia M (1988) Estimation of energy expenditure, net carbohydrate utilization, and net fat oxidation and synthesis by indirect calorimetry: evaluation of errors with special reference to the detailed composition of fuels. *Am J Clin Nutr* 47:608–628
- Parati G, Di Rienzo M, Bertinieri G, Pomidossi G, Casadei R, Groppelli A, Pedotti A, Zanchetti A, Mancia G (1988) Evaluation of the baroreceptor-heart rate reflex by 24-hour intra-arterial blood pressure monitoring in humans. *Hypertension* 12:214–222
- Bertinieri G, Di Rienzo M, Cavallazzi A, Ferrari A, U, Pedotti A, Mancia G (1988) Evaluation of baroreceptor reflex by blood pressure monitoring in unanesthetized cats. *Am J Physiol* 254:H377–H383
- Lin E, Lowry SF (1998) The human response to endotoxin. *Sepsis* 2:255–262
- Van der Poll T, Romijn JA, Endert E, Born JJJ, Büller HR, Sauerwein HP (1991) Tumor necrosis factor mimics the metabolic response to acute infection in healthy humans. *Am J Physiol* 261:H457–H465
- Spinas GA, Keller U, Brockhaus M (1992) Release of soluble receptors for tumor necrosis factor (TNF) in relation to circulating TNF during experimental endotoxemia. *J Clin Invest* 90:533–536
- Bloesch D, Keller U, Spinas GA, Kury D, Girard J, Stauffacher W (1993) Effects of endotoxin on leucine and glucose kinetics in man: contribution of prostaglandin E2 assessed by a cyclooxygenase inhibitor. *J Clin Endocrinol Metab* 77:1156–1163
- Schmidt H, Müller-Werdan U, Hoffmann T, Francis DP, Piepoli MF, Rauchhaus M, Prondzinsky R, Loppnow H, Buerke M, Hoyer D, Werdan K (2005) Autonomic dysfunction predicts mortality in patients with multiple organ dysfunction syndrome of different age groups. *Crit Care Med* 33:1994–2002
- Chioloro R, Kinney JM (2001) Metabolic and nutritional support in critically ill patients: feeding the whole body or individual organs? *Curr Opin Clin Nutr Metab Care* 4:127–130
- Mascioli E, Leader L, Flores E, Trimbo S, Bistran B, Blackburn G (1988) Enhanced survival to endotoxin in guinea pigs fed IV fish oil emulsion. *Lipids* 23:623–625
- Pscheidl E, Schywalsky M, Tschalkowsky K, Böke-Pröls T (2000) Fish oil-supplemented parenteral diets normalize splanchnic blood flow and improve killing of translocated bacteria in a low-dose endotoxin rat model. *Crit Care Med* 28:1489–1496
- Ellis EF, Police RJ, Dodson LY, McKinney JS, Holt SA (1992) Effect of dietary n-3 fatty acids on cerebral microcirculation. *Am J Physiol* 262:H1379–H1386
- Pomposelli JJ, Flores E, Hirschberg Y, Teo TC, Blackburn GL, Zeisel SH, Bistran BR (1990) Short-term TPN containing n-3 fatty acids ameliorate lactic acidosis induced by endotoxin in guinea pigs. *Am J Clin Nutr* 52:548–552
- Pomposelli JJ, Mascioli EA, Bistran BR, Lopes SM, Blackburn GL (1989) Attenuation of the febrile response in guinea pigs by fish oil enriched diets. *JPN J Parenter Enteral Nutr* 13:136–140
- Teo TC, Selleck KM, Wan JM, Pomposelli JJ, Babayan VK, Blackburn GL, Bistran BR (1991) Long-term feeding with structured lipid composed of medium-chain and N-3 fatty acids ameliorates endotoxic shock in guinea pigs. *Metabolism* 40:1152–1159
- Michaeli B, Berger MM, Revelly JP, Tappy L, Chioloro R (2007) Effects of fish oil on the neuro-endocrine responses to an endotoxin challenge in healthy volunteers. *Clin Nutr* 26:70–77
- Weiss G, Meyer F, Matthies B, Pross M, Koenig W, Lippert H (2002) Immunomodulation by perioperative administration of n-3 fatty acids. *Br J Nutr* 87 Suppl 1:S89–S94
- Assenmacher I, Barbanel G, Gaillet S, Givalois L, Ixart G, Malaval F, Mekaouche M, Siaud P, Szafarczyk A (1995) Central regulation of ACTH release in stress. *Ann N Y Acad Sci* 771:41–54
- Chrousos GP (1995) The hypothalamic-pituitary-adrenal axis and immune-mediated inflammation. *N Engl J Med* 332:1351–1362
- Delarue J, Matzinger O, Binnert C, Schneiter P, Chioloro R, Tappy L (2003) Fish-oil prevents the adrenal activation elicited by mental stress in healthy men. *Diabet Med* 29:289–295

-
25. Copper AL, Gibbons L, Horan MA, Little RA, Rothwell NJ (1993) Effect of dietary fish oil supplementation on fever and cytokine production in human volunteers. *Clin Nutr* 12:3212–3328
 26. Lee JY, Plakidas A, Lee WH, Heikkinen A, Chanmugam P, Bray G, Hwang DH (2003) Differential modulation of Toll-like receptors by fatty acids: preferential inhibition by n-3 polyunsaturated fatty acids. *J Lipid Res* 44:479–486
 27. Endres S, Ghorbani R, Kelley VE, Georgilis K, Lonnemann G, van der Meer JW, Cannon JG, Rogers TS, Klempner MS, Weber PC et al (1989) The effect of dietary supplementation with n-3 polyunsaturated fatty acids on the synthesis of interleukin-1 and tumor necrosis factor by mononuclear cells. *N Engl J Med* 320:265–271
 28. Wachtler P, Konig W, Senkal M, Kemen M, Koller M (1997) Influence of a total parenteral nutrition enriched with omega-3 fatty acids on leukotriene synthesis of peripheral leukocytes and systemic cytokine levels in patients with major surgery. *J Trauma* 42:191–198
 29. Carpentier YA, Dupont IE (2002) Optimizing intravenous supply of functional lipid components. *Nestle Nutr Workshop Ser Clin Perform Programme* 7:87–98
 30. Harris HW, Johnson JA, Wigmore SJ (2002) Endogenous lipoproteins impact the response to endotoxin in humans. *Crit Care Med* 30:23–31
 31. Feingold KR, Funk JL, Moser AH, Shigenaga JK, Rapp JH, Grunfeld C (1995) Role for circulating lipoproteins in protection from endotoxin toxicity. *Infect Immun* 63:2041–2046