

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Sudden cardiac death is associated both with epilepsy and with use of antiepileptic drugs.

Authors: Bauer PR, Novy J, Keezer MR, Bell GS

Journal: Heart (British Cardiac Society)

Year: 2015 Jan

Issue: 101

Volume: 1

Pages: 83

DOI: [10.1136/heartjnl-2014-306760](https://doi.org/10.1136/heartjnl-2014-306760)

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

CORRESPONDENCE

Sudden cardiac death is associated both with epilepsy and with use of antiepileptic drugs

To the Editor, We read with interest the recent paper by Bardai et al,¹ which reports that epilepsy and antiepileptic drugs (AEDs) were independently associated with sudden cardiac death (SCD). We are unconvinced that such a clear distinction between disease and drug effects can be made in this study as all people with epilepsy were by definition taking AEDs. We believe this is why SCD risk in those with epilepsy (table 2) and in AED users with epilepsy (table 3) was the same: OR 2.8 (95% CI 1.4 to 5.3). It is unclear whether these numbers reflect disease effects, drug effects or both. It is reported that people with epilepsy are at increased risk of SCD, yet in this population there is a major diagnostic alternative: sudden unexpected death in epilepsy, an autopsy-negative mostly seizure-related type of sudden death with cardiac and noncardiac causes.² We do not believe that sudden death in people with epilepsy can be assumed to be of cardiac origin (vs, for example neurogenic respiratory depression) without documentation of a cardiac mechanism of death (eg, by ECG). Carbamazepine and gabapentin were the only individual AEDs associated with increased SCD risk. The authors attribute this to the supposed sodium channel blocking properties of these drugs, but we believe that indication bias cannot be excluded. Carbamazepine is the drug of choice in people with focal seizures and stroke is a leading cause of this type of epilepsy, particularly in the elderly.³ An important indication for gabapentin is diabetic chronic neuropathic pain.⁴ It is possible that a worse cardiovascular status rather than sodium channel blocking properties explains the higher SCD risk in users of these AEDs. Despite the use of a large database, many numbers are small, so the demonstrated significant effect for sodium channel blocking AEDs but not for nonsodium channel blocking AEDs may reflect limited sample size rather than differences in effect. There was a consistent trend; all reported AED ORs were greater than the null value. This paper is important in that it confirms that epilepsy in the community is associated with an increased risk of sudden natural death. Future studies with additional documentation of death mechanisms, correction for indication bias and larger sample size are needed to explain the cause of this excess risk and clarify the role of AEDs.

Prisca R Bauer, Jan Novy, Mark R Keezer,

Gail S Bell

NIHR University College London Hospitals Biomedical
Research Centre, Clinical & Experimental Epilepsy, UCL
Institute of Neurology, London, UK

Correspondence to Prisca R Bauer, NIHR University
College London Hospitals Biomedical Research Centre,
UCL Institute of Neurology, Queen Square, London
WC1N 3BG, UK; prisca.bauer.12@ucl.ac.uk

Contributors All authors were involved in writing the
final manuscript.

Competing interests None.

To cite Bauer PR, Novy J, Keezer MR, et al. Heart
Published Online First: [please include Day Month Year]
doi:10.1136/heartjnl-2014-306760
► <http://dx.doi.org/10.1136/heartjnl-2014-305664>
Heart 2014;0:1. doi:10.1136/heartjnl-2014-306760

REFERENCES

- 1 Bardai A, Blom MT, van Noord C, et al. Sudden cardiac death is associated both with epilepsy and with use of antiepileptic drugs. Heart Published Online First: 16 Jul 2014 doi:10.1136/heartjnl-2014-305664
- 2 Shorvon S, Tomson T. Sudden unexpected death in epilepsy. Lancet 2011;378:2028–38.
- 3 Ryvlin P, Montavont A, Nighoghossian N. Optimizing therapy of seizures in stroke patients. Neurology 2006;67(12 Suppl 4):S3–9.
- 4 Moore RA, Wiffen PJ, Derry S, et al. Gabapentin for chronic neuropathic pain and fibromyalgia in adults.