

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Microsurgical disconnection of a ruptured intracranial pial arteriovenous fistula guided by indocyanine green video-angiography.

Authors: Signorelli F, Messina R, Maduri R, Barbagallo G, Tacconi L

Journal: World neurosurgery

Year: 2019 Mar 13


DOI: [10.1016/j.wneu.2019.03.045](https://doi.org/10.1016/j.wneu.2019.03.045)

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

Accepted Manuscript

Microsurgical disconnection of a ruptured intracranial pial arteriovenous fistula guided by indocyanine green video-angiography

Francesco Signorelli, MD, MSc, Raffaella Messina, MD, Rodolfo Maduri, MD, Giuseppe Barbagallo, MD, Leonello Tacconi, MD



PII: S1878-8750(19)30671-0

DOI: <https://doi.org/10.1016/j.wneu.2019.03.045>

Reference: WNEU 11723

To appear in: *World Neurosurgery*

Received Date: 29 January 2019

Revised Date: 4 March 2019

Accepted Date: 5 March 2019

Please cite this article as: Signorelli F, Messina R, Maduri R, Barbagallo G, Tacconi L, Microsurgical disconnection of a ruptured intracranial pial arteriovenous fistula guided by indocyanine green video-angiography, *World Neurosurgery* (2019), doi: <https://doi.org/10.1016/j.wneu.2019.03.045>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Microsurgical disconnection of a ruptured intracranial pial arteriovenous fistula guided by indocyanine green video-angiography

Francesco Signorelli, MD, MSc^{*}, Raffaella Messina, MD^{*}, Rodolfo Maduri, MD[§], Giuseppe Barbagallo, MD[‡], Leonello Tacconi, MD[#]

^{*}Division of Neurosurgery, Department of Basic Medical Sciences, Neurosciences and Sense Organs, University “Aldo Moro” of Bari, Italy

[§]Division of Neurosurgery, University Hospital of Vaud, Lausanne, Switzerland

[‡]Division of Neurosurgery, Policlinico "G. Rodolico" University Hospital, Catania, Italy

[#]Division of Neurosurgery, University Hospital of Trieste, Trieste, Italy

The authors have no conflicts of interest or personal financial or institutional interest in any of the drugs, materials, or devices described in this article. No funding has been received for the conduct of this study and/or preparation of this manuscript.

Corresponding Author: Francesco Signorelli, MD, MSc. Division of Neurosurgery, Department of Basic Medical Sciences, Neurosciences and Sense Organs, University “Aldo Moro” of Bari, Piazza Giulio Cesare, 11 70124 Bari, Italy.

Email: francesco.signorelli@uniba.it

Key words: brain vascular malformation, pial arteriovenous fistula, indocyanine green, video-angiography.

Intracranial arteriovenous fistulas, rare causes of spontaneous intracerebral bleeding, are direct communications between an arterial feeder and an arterialized vein that drains also normal brain. Arteriovenous disconnection is the only effective treatment for this type of vascular malformation, which is often reached microsurgically due to the difficult endovascular access. Intraoperative indocyanine green video-angiography (IGC-VA) is a valuable help in identifying the arterialized draining vein and its direct communication with the arterial feeder and in confirming real time the interruption of the fistula.

We describe the case of a 46 years old man presenting with a sudden onset of headache, left arm motor and sensory deficit associated with a fronto-parietal hematoma evacuated one week earlier in another institution.

Digital subtraction angiography showed a direct communication between an anterior parietal branch of the right middle cerebral artery and a parietal vein. Given the difficulty to reach endovascularly the point of fistula because of the small caliber and tortuosity of the arterial feeder and the short and relatively rapid flow through the arteriovenous communication, we decided to proceed with the microsurgical treatment. Under intraoperative neurophysiological monitoring the fistula was located with the aid of ICG-VA and interrupted. Both control ICG-VA and postoperative angiogram confirmed the resolution of the fistula. At a 3-month follow up the patient had a complete neurological recovery.

indocyanine green video-angiography (IGC-VA)

Article Title Microsurgical disconnection of a ruptured intracranial pial arteriovenous fistula guided by indocyanine green video-angiography

Francesco Signorelli, MD, MSc^{*}, Raffaella Messina, MD^{*}, Rodolfo Maduri, MD, Giuseppe Barbagallo, MD, Leonello Tacconi, MD

I, Francesco Signorelli, corresponding author, certify that there is no actual or potential conflict of interest in relation to this article.

Francesco Signorelli, MD, MSc, on behalf of all authors Date: March 4, 2019