

REVIEW

Probing the functional significance of wild animal microbiomes using omics data

Sarah F. Worsley¹  | Elin Videvall²  | Xavier A. Harrison³  | Johannes R. Björk^{4,5}  | Florent Mazel⁶  | Klara M. Wanelik⁷ 

¹School of Biological Sciences, University of East Anglia, Norwich Research Park, Norfolk, UK; ²Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden; ³Centre for Ecology and Conservation, University of Exeter, Penryn, UK; ⁴Department of Gastroenterology and Hepatology, University Medical Center Groningen, Groningen, The Netherlands; ⁵Department of Genetics, University of Groningen, Groningen, The Netherlands; ⁶Department of Fundamental Microbiology, University of Lausanne, Lausanne, Switzerland and ⁷School of Biosciences, University of Surrey, Guildford, UK

Correspondence

Sarah F. Worsley

Email: s.worsley@uea.ac.uk

Funding information

University of Surrey; Leverhulme Trust, Grant/Award Number: ECF-2023-433 and RPG-2020-320; Natural Environment Research Council, Grant/Award Number: NE/S010939/1; Vetenskapsrådet, Grant/Award Number: 2020-00259; Faculté de Biologie et de Médecine, Université de Lausanne

Handling Editor: Johannes Rousk

Abstract

1. Host-associated microbiomes are thought to play a key role in host physiology and fitness, but this conclusion mainly derives from systems biased towards animal models and humans.
2. While many studies on non-model and wild animals have characterised the taxonomic diversity of their microbiomes, few have investigated the functional potential of these microbial communities.
3. Functional 'omics' approaches, such as metagenomics, metatranscriptomics and metabolomics, represent promising techniques to probe the significance of host-associated microbiomes in the wild.
4. In this review, we propose to (1) briefly define the main available functional omics tools along with their strengths and limitations, (2) summarise the key advances enabled by omics tools to understand microbiome function in human and animal models, (3) showcase examples of how these methods have already brought invaluable insights into wild host microbiomes and (4) provide guidelines on how to implement these tools to address outstanding questions in the field of wild animal microbiomes.
5. To conclude, we suggest that, building on knowledge derived from cheaper, more traditional approaches (e.g. 16S metabarcoding and qPCR), functional omics tools represent a promising approach to test hypotheses regarding the ecological and evolutionary significance of the resident microbiota in wild animals.

KEYWORDS

culturomics, metabolomics, metagenomics, metaproteomics, metatranscriptomics, microbiome function, multi-omics, wild host

Florent Mazel and Klara M. Wanelik co-last authors.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Functional Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

1 | INTRODUCTION

Host-associated microbiomes represent complex ecosystems that often make fundamental contributions to host health (Sommer & Bäckhed, 2013; but see Hammer et al., 2019). For example, members of the vertebrate gut microbiome play an important role in host nutrient acquisition (Nicholson et al., 2012), disease resistance (Round & Mazmanian, 2009) and cognition (Davidson et al., 2020). Some host functions are thus fundamentally dependent on the microbes that colonise them and microbiomes are starting to be considered as an important contributor to the functional trait make up macro-organisms.

The vast majority of research evidencing these contributions has been conducted using humans and laboratory animals. Fewer studies have focussed on animals living in natural populations where hosts are exposed to complex and highly dynamic environmental pressures. However, with the advent of affordable high-throughput sequencing methodologies and non-invasive sampling techniques, there has been a recent proliferation of studies investigating the microbiome of wild animals. Through the amplification of microbial marker genes, usually the bacterial 16S rRNA gene, such studies have uncovered immense microbial taxonomic diversity which varies dynamically across host species (Knowles et al., 2019; Song et al., 2020; Thomas et al., 2016), populations (Grieneisen et al., 2019; Uren Webster et al., 2018) and individuals (Björk et al., 2022; Risely et al., 2022; Worsley et al., 2021).

Numerous factors have been shown to contribute to this variation in the wild. For example, across taxa, particularly in mammals, microbiome compositional dissimilarity is often correlated with host phylogenetic distance (a pattern termed 'phylosymbiosis') (Brooks et al., 2016; Groussin et al., 2017; Nishida & Ochman, 2018; Song et al., 2020; Youngblut et al., 2019). Diet has also emerged as a major driver of gut microbiome differences among host species (Ingala et al., 2021; Song et al., 2020; Youngblut et al., 2019). Within populations, a suite of ecological factors have been linked to microbiome dynamics, including habitat differences (Fackelmann et al., 2021) and seasonal fluctuations in dietary intake (Baniel et al., 2021; Björk et al., 2022; Michel et al., 2018). Furthermore, inter-individual microbiome variation is also correlated with a range of host factors including differences in host genotype (Davies et al., 2022; Thomas et al., 2016), relatedness (Baniel et al., 2022; Ren et al., 2017; Wanelik et al., 2023), social behaviour (Raulo et al., 2021; Tung et al., 2015) and age (Reese et al., 2021; Sadoughi et al., 2022).

However, aside from identifying which microbes are found in particular hosts and how their abundances vary, a key aim of microbiome research is to uncover the functional importance of microbial communities and their impact on host health and fitness (Suzuki, 2017). Despite this aim, few wildlife microbiome studies have integrated information on microbial function into their analyses (but see Levin et al., 2021; Youngblut et al., 2020). A variety of analytical ('omics') tools now exist that enable data on microbial functions to be collected (see Table 1), giving an additional layer of

resolution and understanding to microbiome studies. That is, instead of asking 'what is there?' we can now begin to ask questions like 'what does it do?'

While the gold standard for evaluating host-associated microbial function remains experimental manipulation in model systems, significant advances have been made by coupling non-invasive sampling with multi-omics approaches in non-experimental systems. For example, in the field of human microbiomes, it has been possible to identify specific microbial genes, transcripts and metabolites that are linked to, and may causally influence, specific disease states such as inflammatory bowel disease (IBD; Lloyd-Price et al., 2019), diabetes (Heintz-Buschart et al., 2016) and cancer (Wang et al., 2020). Metagenomic studies have also shed light on the functional and fine-grained phylogenetic divergence of human and non-human primate gut microbiomes providing insight into the processes shaping their evolution (Rühlemann et al., 2024; Sanders et al., 2023). Recent attempts to construct functionally annotated gut microbiome databases across wild hosts using shotgun metagenomics have revealed enormous functional diversity that varies across animal taxa (Levin et al., 2021; Youngblut et al., 2020) and according to key host traits such as diet, social structure and lifespan (Levin et al., 2021). Some of these microbial functions (e.g. toxin degradation by microbes associated with carrion-eaters) have been directly linked to host ecology through follow-up experiments (Levin et al., 2021). Thus, we believe that functional techniques could be applied to wild animal microbiomes to give a better understanding of the mechanisms by which microbes influence host ecology and evolution, and the relevance of the huge microbial taxonomic diversity observed within and across wild systems.

In this review, we have four main aims:

1. Highlight different types of functional data that can be generated (Table 1)
2. Illustrate how these have been applied to provide insight into the microbiomes of humans and model organisms.
3. Propose how these techniques might be transferred into the field of wild animal microbiomes to better understand how the microbiome, as a functional trait, can influence host health, ecology and fitness.
4. Discuss the challenges and potential solutions associated with collecting and processing functional data from understudied, non-model animal systems.

2 | GENERATING FUNCTIONAL DATA SETS

A variety of omics technologies now exist that enable the characterisation of host-associated microbiomes (Table 1). Each of these methods can provide different perspectives on microbiome functionality. In Table 1, we provide an overview of the main technologies available, including their advantages and limitations. We also outline these briefly below.

TABLE 1 Functional 'omics' approaches for studying the microbiome.

Omic technology	Description	Pros	Cons	Examples of analytical tools	Key reviews
DNA metabarcoding Aim: Infer functional potential based on homology to annotated taxa	Marker gene amplicon sequencing (e.g. the 16S rRNA gene for bacteria) can be combined with taxonomic assignment using microbial databases (e.g. SILVA, Quast et al., 2012). This approach provides information on the identity and relative abundance of microbes in a sample. Some tools (see examples) also use information on the homology of marker gene sequences to functionally annotated genomes in reference databases to provide rough estimates of the metabolic capabilities of microbial communities	Cheap per-sample cost Samples can be multiplexed Less likely to be affected by host contamination or non-target reads compared to other omics methods as a specific microbial genomic region is amplified Easy to implement (standardised wet lab and bioinformatic procedures exist)	Low taxonomic resolution (microbes often only identified to family or genus level) Limited functional information, particularly if there are no close relatives in reference databases Difficult to compare studies that have sequenced different regions of the same marker gene, or used different bioinformatic workflows	Picrust2 (Douglas et al., 2020) FUNGuild (Nguyen et al., 2016) FungalTraits (Pöhlme et al., 2020) Tax4Fun2 (Wemheuer et al., 2020) MifFunPred (Mongad et al., 2021)	Djemieli et al. (2022)
Metagenomics Aim: Infer functional potential based on gene content	Shotgun sequencing of total genomic DNA in a sample provides information on the genes, and potentially genomes, of the microbes present. Mapping sequences to reference databases ('reference-based profiling') provides information on the taxonomic identity of microbes (potentially to strain level), and the functional potential of the microbiome, such as the types of enzymes and metabolic pathways present High-coverage metagenomic data may also allow metagenome-assembled genomes (MAGs) to be constructed; this can enable the identification of genes carried by individual microbial strains, including those that are currently uncharacterised	Provides direct estimates of microbial gene content & 'functional potential' (e.g. KEGG & MetaCyc pathways, virulence factors and antibiotic resistance genes) Possible to identify microbial taxa to species or even strain level with deep sequencing Can reconstruct microbial genomes if coverage is sufficient; this can identify taxa that have not yet been isolated or functionally characterised Costs can be reduced by using shallow sequencing Potential to assay all microbial groups in a sample e.g. bacteria, fungi, viruses	Expensive, particularly if deep sequencing Analytically more complex than metabarcoding Host contamination can be an issue for some sample types Pipelines for reconstruction of MAGs not yet standardised Presence of gene content does not imply genes are transcribed and translated into functional products	<i>Reference-based profiling:</i> Kraken 2 taxonomic profiler (Wood et al., 2019) MetaPhlan 4 taxonomic profiler (Blanco-Miguez et al., 2023) HUMAN 3 functional profiler (Beghini et al., 2021) <i>Genome assembly approaches (MAGs):</i> MEGAHIT, for assembly (Li et al., 2015) MetaBAT 2, for binning (Kang et al., 2019) Prokka, for gene prediction/annotation (Seemann, 2014)	Meyer et al. (2022), Quince et al. (2017), Szczyrba et al. (2017), Yang et al. (2021)

(Continues)

TABLE 1 (Continued)

Omic technology	Description	Pros	Cons	Examples of analytical tools	Key reviews
<p><i>Metatranscriptomics</i></p> <p>Aim: Infer functional potential based on gene expression</p>	<p>RNA is converted to complementary DNA (cDNA) before subjecting it to shotgun sequencing. The resulting data provide information on the functional activity of the microbiome, including which genes are expressed by microbial members (at the time of sampling) and at what levels. It is also possible to use these data to identify genetic variation among microbial members</p>	<p>Moves beyond descriptions of gene content by providing information on which genes are actively transcribed by microbes in a sample</p> <p>Can provide information on taxonomic composition</p> <p>Potential to assay all microbial groups in a sample e.g. bacteria, fungi, viruses</p> <p>Possible to examine host and microbial responses in tandem (e.g. to stressors/experimental manipulation) using dual RNA sequencing</p>	<p>Expensive, notably due to the higher abundance of ribosomal RNA than messenger RNA molecules, meaning high sequencing depth (or rRNA depletion) is needed</p> <p>Requires different sample storage and handling to DNA-based methods, and samples can be unstable due to the presence of RNases</p> <p>No consensus on standardised bioinformatic pipeline</p> <p>Host transcripts may overwhelm microbial signals when host and microbial RNA is co-extracted from samples</p> <p>When microbial taxonomic composition is unknown, pairing metatranscriptomics with metagenomics is needed to determine whether differences in transcript abundance are driven by expression level changes or gene copy number differences (e.g. due to compositional change)</p>	<p>MetaTrans (Martinez et al., 2016)</p> <p>HUMAN3 (Beghini et al., 2021)</p> <p>FMAP (Kim et al., 2016)</p>	<p>Niu et al. (2017), Shakya et al. (2019), Zhang et al. (2021)</p>
<p><i>Metaproteomics</i></p> <p>Aim: Infer functional potential based on protein content</p>	<p>High resolution mass spectrometry techniques can be used to identify and quantify the microbial proteins present in a microbiome sample. In addition, recently developed metaproteomic tools can allow per-species biomass to be quantified which provides information on community structure and carbon sources available for microbial members</p>	<p>Information on which genes are translated by microbes in a sample, resulting in stronger evidence of function than transcription data</p> <p>Ability to measure both taxonomic composition and function</p> <p>Can examine both host and microbial responses (e.g. to stressors/experimental manipulation)</p> <p>Potential to assay all microbial groups in a sample e.g. bacteria, fungi, viruses</p>	<p>Expensive</p> <p>No consensus on standardised bioinformatic pipeline</p> <p>Extensive sample preparation and extraction steps</p> <p>Lower throughput than other omics methods due to long run times per sample</p> <p>Requires a priori knowledge of the functional properties of target proteins</p>	<p>Unipept (Verschaffelt et al., 2021)</p> <p>MetaProteomAnalyzer (Muth et al., 2015)</p> <p>ProPHAnE (Schneider et al., 2011)</p>	<p>Kleiner (2019), Peters et al. (2019), Sajulga et al. (2020), Schiebenhoefer et al. (2020), Wilmes and Bond (2006)</p>

TABLE 1 (Continued)

Omic technology	Description	Pros	Cons	Examples of analytical tools	Key reviews
<p>Metabolomics</p> <p>Aim: Infer functional potential based on metabolite content</p>	<p>Identifies and quantifies the small molecules (metabolites) present in a microbiome sample. Thus, provides information on which metabolites are being produced, or consumed, by microbes. Metabolomics is usually performed by coupling either gas chromatography or liquid chromatography to mass spectrometry. Signal peaks of different intensities are produced depending on metabolite mass to charge ratio (m/z); these peaks can sometimes be matched with those of known metabolites. Lipidomics, a subfield of metabolomics, involves the identification and quantification of lipid molecules (fatty acids). The functional activity of microbial lipids may reflect the community's ability to metabolise different nutrients, adapt to environmental stressors or interact with host cells</p>	<p>Provides information on microbial metabolites present in a sample which may have direct functional effects in hosts</p> <p>No a priori knowledge of metabolomic targets required</p> <p>Imaging mass spectrometry (IMS) allows measurement of the spatial distribution of metabolites/biomolecules</p>	<p>No consensus on standardised bioinformatic pipeline</p> <p>Expensive and can require sample-specific optimisation prior to a full run</p> <p>High numbers of unknown metabolites</p> <p>Analytically challenging to identify which microbe(s) produce sets of metabolites</p> <p>Distinguishing between host and microbial metabolites can be challenging</p> <p>No robust workflow for determining the taxonomic composition of the sample</p>	<p>MetaboAnalyst (Chong et al., 2018)</p> <p>XCMS (Huan et al., 2017)</p> <p>MZmine 2 (Pluskal et al., 2010)</p> <p>iMAP (Zhou et al., 2021)</p>	<p>Bhosle et al. (2022), Peters et al. (2019), Yasuda et al. (2020)</p>
<p>Culturomics</p> <p>Aim: Characterising microbes through culturing, experimentation and functional assays</p>	<p>For this technique, microbes are isolated from target environments (e.g. host, soil, water) and identified using sequencing—most often Sanger sequencing of the full-length 16S rRNA gene for bacteria. Isolates can then be tested using in vitro assays that directly assess their functional capacity e.g. inhibition of pathogen growth, ability to digest substrates or antimicrobial resistance. Isolates can also be whole-genome sequenced to provide high quality genome sequences to serve as references</p>	<p>Provides direct functional information for traits of interest</p> <p>Can differentiate isolates based on function even if they have the same taxonomic assignment</p> <p>Can scale up from single isolates to multi-isolate consortia to probe diversity–function relationships or test for interaction among microbes</p> <p>Cheap to identify microbes using Sanger sequencing e.g. via full-length sequencing of 16S rRNA gene for bacteria</p> <p>Full genome sequencing of isolates can identify genes underpinning variation in functional assays</p> <p>Isolates with desirable functional properties in vitro can be validated in vivo with probiotic experiments</p> <p>Essential for developing high quality reference databases</p>	<p>Requires that microbes in a sample can be cultured</p> <p>Culturing and sequencing microbial isolates is time-intensive</p> <p>Assays of function in vitro may not represent the functional phenotype of that microbe within a community</p> <p>Difficult/time-intensive to track microbial dynamics in multi-isolate consortia</p> <p>Requires genome sequence to determine why taxonomically identical microbes have different functional profiles</p>	<p>NA</p>	<p>Antwis and Harrison (2018), Lagier et al. (2018), Renwick et al. (2021)</p>

2.1 | DNA metabarcoding

Currently, the most widely used approach for characterising host-associated microbiomes, particularly in wild systems, is DNA metabarcoding (Table 1). Here, full length or partial segments of marker genes (such as ribosomal RNA genes) are amplified and sequenced, then taxonomically assigned using microbial databases. This methodology provides information on the identity of microbes, and their relative abundances, within a sample. However, tools have also been developed that enable microbial functions to be inferred from these data sets (reviewed in Djemiel et al., 2022). Here, functions are estimated based on sequence homology to annotated microbial genomes in reference databases. For example, PICRUSt2 (Douglas et al., 2020) enables preliminary functional prediction from 16S rRNA gene metabarcoding data sets by providing rough estimates of KEGG pathways and the metabolic diversity present in the overall microbial community. However, the accuracy of DNA metabarcoding and tools such as PICRUSt2 can be limited, because phylogenetically similar microbial taxa can provide different functions, even within species (e.g. *Escherichia coli*, Rasko et al., 2008).

2.2 | Shotgun metagenomics

Shotgun metagenomics (Table 1), whereby total genomic DNA is sequenced, can provide information on the taxonomic composition and the presence of microbial genes within a sample (reviewed in Quince et al., 2017). This, in turn, enables detailed inference about the types of enzymes and metabolic pathways that may be expressed by the microbiome. High-coverage metagenomic sequencing can also enable metagenome-assembled genomes (MAGs) to be constructed, potentially allowing the identification of taxa (and their genes) that are currently uncharacterised (Quince et al., 2017). A growing number of metagenomic data sets are becoming publicly available for wild animals (e.g. Levin et al., 2021; Rühlemann et al., 2024; Sanders et al., 2023; Youngblut et al., 2019).

2.3 | Metatranscriptomics, metaproteomics & metabolomics

The presence of a gene does not necessarily mean that it is expressed or translated in situ and other tools are needed to determine whether this is the case. Metatranscriptomics (Table 1) can be used to evaluate which genes are actually expressed by microbial members (at time of sampling), while metaproteomics (Table 1) and metabolomics (Table 1) enable the identification and quantification of microbial proteins and metabolites, respectively.

2.4 | Culturomics

A final, often overlooked, approach that can be used to define microbial function is culturomics (Table 1). This technique involves

isolating and culturing microbes associated with host animals and identifying them using sequencing approaches. Isolates can then be characterised using in vitro assays that directly test their functional capacity (e.g. their ability to inhibit the growth of pathogens) (Antwis & Harrison, 2018; Lagier et al., 2018). Assays can also be paired with whole-genome sequencing of isolates potentially allowing the identification of genes underlying functions of interest (e.g. Holmes et al., 2016; Kwong et al., 2014).

While all methodologies come with limitations (Table 1), combining multiple omics tools (i.e. a 'multi-omics' approach) can provide detailed information on microbiome functionality and genetic attributes. Combining such information with data on the genetics and ecology of hosts (a so-called 'hologenomics' approach) provides a powerful means to unpick complex host-microbe interactions (Alberdi et al., 2022). In the sections below, we highlight examples of where functional approaches have provided additional insight and how similar methodologies could be applied in non-model systems to probe the functional significance of the microbiome.

3 | FUNCTIONAL INSIGHTS FROM STUDIES OF MODEL ORGANISMS AND HUMANS

Over the past decade, functional omics tools (Table 1) have been widely employed in a variety of research fields, particularly those aiming to understand how human gut microbiomes assemble and impact health (e.g. reviewed in Heintz-Buschart & Wilmes, 2018; Rojo et al., 2017; Zhang et al., 2019). Our goal here is not to give an exhaustive overview of this work. Instead, we aim to highlight key examples of how such data sets can provide a deeper understanding of host-microbiome interactions, beyond what could normally be achieved using traditional metabarcoding techniques.

One key advantage of using functional omics tools is the potential to move beyond descriptions of taxonomic structure and glean detailed information about the microbial genes, pathways, proteins and metabolites that might directly impact host health and disease. Such information can provide mechanistic insight and has added to our understanding of the contribution of the gut microbiome to various diseases, including IBD (Jacobs et al., 2023; Lee et al., 2021; Lloyd-Price et al., 2019), diabetes (Heintz-Buschart et al., 2016) and cancer (Wang et al., 2020). Combining multiple omics tools has proven to be a particularly powerful approach. For example, a large-scale, longitudinal study of IBD patients used metagenomics, metatranscriptomics, metaproteomics and metabolomics to determine functional changes in the gut microbiome associated with periods of increased disease activity (Lloyd-Price et al., 2019). Alongside a characteristic increase in the abundance of facultative anaerobic bacteria, this study also identified disrupted microbial transcription (particularly amongst *Clostridia*), as well as corresponding changes to the abundance of key metabolites (such as key bile acids and butyrate) that were potentially linked to gut inflammation (Lloyd-Price et al., 2019).

Functional omics tools are also able to provide a more accurate approximation of the functional plasticity of a microbiome than estimates from metabarcoding. This is evident from human metatranscriptomic studies showing that different disease states may be associated with changes in microbial gene expression even when there are no, or very few, changes to the taxonomic composition of the microbiome (Franzosa et al., 2014; Gosalbes et al., 2011; Heintz-Buschart et al., 2016). Similarly, the reconstruction of microbial draft genomes (MAGs) via shotgun metagenomics has revealed that microbiome-associated disease outcomes are often a consequence of specific microbial strains that, because of their unique gene content, perform different functions within the gut ecosystem (Hajjo & Geva-Zatorsky, 2021; Hall et al., 2017; Ward et al., 2016). For example, certain bacterial strains are enriched in the gut microbiome of IBD patients and carry unique genes involved in bacterial stress responses and nutrient utilisation (Hall et al., 2017). Other studies have shown that strain-level genetic diversity can contribute to the digestion of different dietary components (De Filippis et al., 2019), enable microbes to resist antibiotics (Roodgar et al., 2021) and result in different immunomodulatory effects on the host (Hajjo & Geva-Zatorsky, 2021). Adding to this complexity, new strains can rapidly evolve in situ without significant changes to a species' relative abundance (Dapa et al., 2023; Roodgar et al., 2021). Since marker genes are often identical across strains that are highly divergent at other loci across the genome, such complexities would go undetected in metabarcoding studies.

Conversely, functional studies may also highlight instances where metabarcoding studies overestimate the potential for function divergence among groups due to functional redundancy (Doolittle & Booth, 2017; Louca et al., 2018). This is when different microbial taxa perform the same functional role within an ecosystem (Louca et al., 2018). Functional studies on various hosts, including humans (Turnbaugh et al., 2009; Vieira-Silva et al., 2016) and ruminants (Taxis et al., 2015), have provided evidence that taxonomically dissimilar gut microbes can share many metabolic pathways. Such redundancy has been hypothesised to increase community resilience in the face of perturbation (Lozupone et al., 2012; Vieira-Silva et al., 2016). These ideas have not been widely tested in the wild animal microbiome literature. However, it is possible that at least some of the enormous microbiome variation observed across species, populations and individuals could be explained by functional redundancy.

In summary, research from other fields has shown that the integration of multiple omics tools can be a powerful way to decipher the compositional and functional characteristics of host-associated microbiomes. Furthermore, these tools provide a means to better understand the mechanisms by which microbiome variation may impact host health. Crucially, the results of such studies also serve as an important hypothesis generating tool by identifying possible targets for future experimental studies and medical intervention (Zhang et al., 2019). Such tools could serve a similar purpose in studies of the wild animal microbiome. For example, the integration of multi-omics data sets could provide insight into the functional relevance

of the taxonomic differences frequently identified between the microbiome of different host species, populations and individuals. The results of these studies could also inform manipulative experiments to confirm the contribution of specific microbiome components to host health or fitness.

4 | HOW AND WHERE HAVE FUNCTIONAL DATA SETS BEEN USED TO ADDRESS QUESTIONS IN THE FIELD OF WILD ANIMAL MICROBIOMES?

Microbiomes are usually assumed to play a key role in host physiology and fitness. However, the functional importance of host-associated microbes has only been causatively demonstrated in a handful of model organisms such as corals (Bourne et al., 2016), bees (Motta & Moran, 2023), ruminants (O'Hara et al., 2020), mice (Clavel et al., 2016) and termites (Scharf & Peterson, 2021), although it can reasonably be assumed for many other animal taxa. In some cases, particularly in some invertebrates, the microbiome might be of less importance to the host (reviewed in Hammer et al., 2017). An important question thus arises: Is it common that animal hosts rely on a microbiome for certain functions? To highlight how omics tools could be applied to address this question in non-model animals, we summarise some key examples of microbial functions that have been dissected using omics tools in different wild hosts.

4.1 | The gut microbiome and host physiology: Diet as a key example

Many studies are starting to demonstrate the value of generating functional data sets to gain mechanistic insights into the role of the microbiome (Table 2). This has been particularly exemplified by research aimed at understanding the contribution of microbes to host dietary adaptation, and especially herbivory. Historically, this has been intensively studied in domesticated ruminants (Mizrahi et al., 2021) whereby metagenomics data have helped to better document the diverse community of fibre degrading bacteria, and their functions, in the rumen microbiome (Xie et al., 2021). Recently, these approaches have also been expanded to wild systems. For example, Cabral et al. (2022) used a combination of omics tools to elucidate the gut microbial community composition, enzymatic systems and metabolic pathways involved in depolymerisation and utilisation of plant fibres in wild capybaras (*Hydrochoerus hydrochaeris*). The study initially described microbial taxonomic community composition using metabarcoding data, then used metagenomics and metatranscriptomics to investigate the genes and gene pathways associated with plant fibre degradation, and to identify the microbial agents involved in depolymerisation. Finally, using metabolomic data, they were able to confirm that these microbial taxa play a role in the conversion of free sugars into energy that is potentially usable by the

TABLE 2 Examples of wild animal studies which have applied functional omics tools and how they have added to our understanding of the wild animal microbiome.

Host taxa	Function studied	Functional omics used	Added value of functional data	Reference
Capybara (<i>Hydrochoerus hydrochaeris</i>)	Host diet digestion	MGX, MTX, MBX	Identified microbial taxa involved in depolymerisation and utilisation of plant fibres, as well as the enzymatic systems and metabolic pathways involved	Cabral et al. (2022)
Midwife toad (<i>Alytes obstetricans</i>)	Host skin disease mitigation	MGX, MBX	Identified functionally relevant taxa driving disease outcome (see also Box 1)	Bates et al. (2022)
Giant panda (<i>Ailuropoda melanoleuca</i>)	Host diet digestion	MGX, MPX	Identified functionally relevant taxa driving differences in body condition	Huang et al. (2022)
Mammals	Host diet digestion	MGX, MTX	Showed that mammalian gut microbiota composition and functionality is partially shaped by host dietary differences	Milani et al. (2020)
Ruddy Turnstone (<i>Arenaria interpres</i>)	Host metabolism	MTX	Showed that taxonomic composition of the microbiome and functional aspects of the metatranscriptome shift during weight gain	Grond et al. (2023)
Common marmoset (<i>Callithrix jacchus</i>)	Various functions sensitive to environmental change	MGX, MTX	Found that differences in microbial gene expression were more dynamic and sensitive to environmental shifts than the abundances of microbial genes	Uehara et al. (2022)
Wild (and captive) non-human primates	Host diet digestion	MGX, MTX	Identified that certain microbial taxa and functions were enriched in primate species with different diets	Zhang et al. (2023)
Baleen whale (Mysticeti)	Host diet digestion	MGX	Showed that baleen whales, which prey on fish and crustaceans, harbour unique gut microbiomes with surprising parallels (in terms of functional capacity and taxonomy) to those of terrestrial herbivores	Sanders et al. (2015)
Black howler monkeys (<i>Alouatta pigra</i>)	Host diet shift	MBX	Showed that faecal metabolite profiles differed between seasons and were strongly associated with changes in plant metabolite consumption. However, faecal metabolite composition was not strongly associated with microbiome taxonomic composition	Mallott et al. (2022)
Turtle ants (<i>Cephalotes</i>)	Nitrogen recycling (from diet and host excretions)	MGX, MBX	Used MBX to show that a specialised core gut microbiome provisions nitrogen for the host by recycling N-rich compounds (e.g. urea from diet). MGX then showed that this phenomenon is likely conserved across many ant species	Hu et al. (2018)
Migratory bird species	Host metabolism and resistome	MGX	Showed that 10 migratory bird species (predominantly waterfowl) vary greatly in gut bacterial composition but are similar in their microbiome metabolism and function	Cao et al. (2020)
Wild vertebrates, including fish, birds and mammals	Various functions including host diet digestion, toxin production and bioluminescence	MGX	Identified host traits that correlate with the composition, diversity, and functional content of their microbiota. For example, they identified proteases, some previously undescribed, from the gut of griffon vultures that degrade toxins present in their carrion diet	Levin et al. (2021)
Vampire bats (<i>Desmodus rotundus</i>)	Host diet digestion	MGX	Compared the taxonomy and function of sanguivorous and non-sanguivorous bat gut microbiomes. The microbiome of vampire bats was functionally, but not taxonomically, distinct from non-sanguivorous bats suggesting that is highly specialised to its nutrient-poor diet	Zepeda Mendoza et al. (2018)

TABLE 2 (Continued)

Host taxa	Function studied	Functional omics used	Added value of functional data	Reference
Myrmecophagous mammals	Host diet digestion	MGX	Reconstructed bacterial genomes from the gut microbiome of nine myrmecophagous mammals. Many genomes carried chitinase genes and several were shared across host orders, suggesting convergent recruitment and highlighting their potential role in the digestion of insect prey	Teuliet et al. (2023)
Spotted hyenas (<i>Crocuta crocuta</i>)	Host diet digestion	MGX	Reconstructed bacterial genomes from the gut microbiome of 4 hyenas across multiple years. Functional profiles were individual specific and varied with prey abundance	Rojas et al. (2023)

Abbreviations: MBX, metabolomics; MGX, metagenomics; MPX, metaproteomics; MTX, metatranscriptomics.

host, focussing predominantly on short-chain fatty acids, a set of key microbial metabolites involved in host health.

Similar functional approaches have been used to better understand the contribution of the microbiota to sanguivory (Zepeda Mendoza et al., 2018) and myrmecophagy (Teuliet et al., 2023) in other mammalian hosts, as well as nitrogen cycling in ants (Hu et al., 2018). The broad finding of these studies is that host-associated microbial communities harbour some of the enzymatic toolkit needed to degrade host dietary molecules (e.g. chitin in insect-eaters) and/or synthesise key molecules unavailable from food (as in the nutrient-poor sanguivorous diet of bats, or in the nitrogen-poor diet of turtle ants). A key outstanding question in many animal systems is the extent to which microbial degradation of the host diet actually contributes to host diet energy budget: Is it relatively low as suggested for humans (1.8%–12.1%) and mice (21.4%) (Arnoldini et al., 2024) or extremely high as in termites (Scharf & Peterson, 2021) or ruminants (>70%, Bergman, 1990; Mizrahi et al., 2021)? This question is also of interest for diets not mentioned here like carnivory, corallivory (in fish) and detritivory. Furthermore, although DNA metabarcoding studies have demonstrated microbial taxonomic discrepancies between host taxa with different dietary adaptations (e.g. herbivores vs. carnivores and omnivores) (Ingala et al., 2021; Song et al., 2020; Youngblut et al., 2019), attempts to compare microbial function across host dietary guilds are currently lacking (but see Levin et al., 2021).

4.2 | How does microbiome variation relate to host fitness differences?

Aside from providing mechanistic insight into host adaptations, functional data sets could also improve our understanding of the significance of the inter-individual microbiome variation present within wild populations. Several studies have found correlations between microbiome taxonomic variation and host fitness components in wild animals, including disease susceptibility (Bates et al., 2022; Risely et al., 2023), condition (Huang et al., 2022), survival (Davidson et al., 2021; Worsley et al., 2021, 2022) and reproduction (Leclaire et al., 2022). In many cases, however, it is not known whether shifts in the microbiome play a causal role in fitness outcomes or are indirect consequences of changes to host physiology or health status. Functional omics tools applied on data collected in situ could provide (1) mechanistic insight into the relevance of microbial taxonomic differences in relation to host health and fitness outcomes, and (2) could also generate hypotheses about causal relationships that can be further tested in laboratory experiments.

For example, Huang et al. (2022) used such an approach to better understand how seasonal gut microbiome variation in wild giant pandas (*Ailuropoda melanoleuca*) is associated with changes in host body mass. Combining in situ metagenomic and metaproteomic techniques with experimental faecal transplants in mice, they attributed wild giant panda body mass gain to butyrate-producing microbes. Similarly, Bates et al. (2022) integrated metabarcoding, metagenomic and metabolomic analyses to better understand links

between the skin microbiome and *Batrachochytrium dendrobatidis* (*Bd*) infection in wild amphibians. Metabarcoding and metagenomic data established differences in microbial taxonomic community composition associated with *Bd* infection. Microbe–metabolite integration analysis was then used to identify functionally relevant taxa driving disease outcomes. These examples demonstrate how using a multi-omics approach can shed light on the functional relevance of shifts in the microbiome for the health outcomes of hosts. Such insights could inform wildlife disease monitoring and conservation strategies (see **Box 1** for more details about amphibians as a model for host–microbe–pathogen interactions).

Understanding how microbiome variation is associated with host fitness, and if and how microbiome changes might contribute to host adaptation in the wild (Alberdi et al., 2016; Rennison et al., 2019), is essential if we are to understand the evolutionary and ecological importance of the microbiome. Furthermore, since hosts are likely to be under strong natural selection to shape their microbiome to be functionally beneficial, and microbial functions evolve to enable microbes to compete and survive within the host, studying microbial functions provides an important evolutionary perspective that may give insight into microbiome assembly mechanisms (Foster et al., 2017; Rühlemann et al., 2024). Although functional approaches

BOX 1 Amphibians as a model for host–microbiome–pathogen interactions

Microbial communities associated with amphibians have been the focus of an enormous amount of research effort in the last decade, principally because of their role in defending their hosts from lethal pathogens such as *Bd* (reviewed in Rebollar et al., 2020) and ranavirus. For example, variation in the composition of skin-associated bacterial communities can predict the intensity of *Bd* outbreaks in natural systems (e.g. Bates et al., 2018). Furthermore, recent research has revealed the importance of the early-life gut bacterial microbiota for defence against infection by ranavirus in adulthood (Warne et al., 2019), and that skin bacterial diversity correlates with survival following ranavirus infection (Harrison et al., 2019). Although the majority of research to date has relied on metabarcoding approaches (see **Table 1**), combining multi-omic tools has yielded significant insight into the mechanism(s) by which amphibian hosts, resident microbes and wildlife pathogens interact.

A key component of amphibian immune defence is the production of antimicrobial peptides (AMPs) by the host, which can prevent infection by inactivating ranavirus virions (Chinchar et al., 2004) and *Bd* zoospores (reviewed in Varga et al., 2019) (**Figure 1a**). Alongside these host derived defences, metabolites produced by resident bacteria and fungi (Kearns et al., 2017) on the skin can also prevent infection (**Figure 1b**; see Woodhams et al., 2023). Indeed, experimental augmentation of the skin microbiome with a bacterium known to produce anti-*Bd* metabolites improved survival following *Bd* challenge by 40% (Kueneman et al., 2016). Furthermore, in vitro bacterial challenge assays have shown that more diverse bacterial communities can provide stronger or more consistent protection against a broad array of *Bd* genotypes (Harrison et al., 2020; Piovita-Scott et al., 2017). Other omics tools have provided further insight into these microbial defence mechanisms. Using a combination of metabarcoding & shotgun metagenomics Rebollar et al. (2018) quantified taxonomy–function associations in skin microbial communities of the amphibian host, *Craugastor fitzingerii*. They found that taxonomic similarity from metabarcoding data predicted similarity in functional profiles from metagenomics, and identified genes linked to the production of anti-*Bd* secondary metabolites in skin metagenomes (**Figure 1b**). Bates et al. (2022) integrated metabarcoding, metagenomics and metabolomics to investigate drivers of variation in *Bd* infection outbreak severity and dynamics in several populations of *Alytes obstetricans*; they found that both skin metagenome composition and metabolome varied depending on whether populations showed epizootic dynamics (associated with mass die offs) or more mild enzootic dynamics. Finally, metatranscriptomic data from a study investigating host transcriptional responses to viral infection in *Rana temporaria* have been used to discover a novel virus, illustrating the utility of omic data sets for wildlife epidemiology (Parry et al., 2023).

Collectively these studies highlight the power of multi-omic studies for identifying the mechanisms by which host-associated microbes respond to pathogens. However, several outstanding questions remain. Chief among these is determining how and when host-derived AMPs and microbial metabolites interact to determine infection probability and disease severity (**Figure 1c**). It has been hypothesised that amphibian hosts may ‘select’ for specific microbes through modulation of their AMP profiles, and that resident microbes may stimulate the production of certain AMPs through cross-talk with the host immune system (**Figure 1c**), though data for the latter are especially limited (Woodhams et al., 2023). Crucially, Rebollar et al. (2018) found several genes involved in AMP resistance in metagenomes derived from the *C. fitzingerii* microbiome, illustrating a genomic mechanism by which symbiotic microbes can persist alongside host-derived immune defences. These insights would not be possible without functional multi-omic data. Unravelling complex interactions between host and microbial genomes, and how gene expression may shift in response to pathogens, requires that we continue to integrate omic approaches (Rebollar et al., 2016). Amphibians and their pathogens continue to provide a model system for understanding the interplay between hosts and their microbial symbionts. Using these systems, we can generate testable hypotheses about the mechanisms by which microbes drive variation in host life history in other wildlife species.




FIGURE 1 Simplified schematic of interactions between amphibian hosts, resident skin microbial communities and pathogens. Both host-derived antimicrobial peptides (AMPs) (a) and metabolites from skin-associated microbes (b) can inactivate pathogens and either prevent infection or reduce burdens of infection. Although driven in part by host genomic variation, AMP composition in the 'mucosome' can select for particular suites of resident microbes, and microbe-host crosstalk may stimulate the production of AMPs (c). Red connecting lines represent inhibition, while blue lines represent promotion. Figure created in the Mind the Graph platform (mindthegraph.com).

have begun to yield valuable insights into the mechanism(s) by which microbes shape the life histories of their hosts, several outstanding questions remain to be addressed (see [Box 2](#)).

5 | GENERATING FUNCTIONAL DATA SETS FOR NON-MODEL ORGANISMS: CHALLENGES AND POTENTIAL SOLUTIONS

5.1 | Study design and sample collection

The enormous variability of wild systems means that additional factors may need to be considered when designing a microbiome study in natural populations. For example, host-associated microbiomes can vary according to host factors (e.g. age and sex, [Reese et al., 2021](#); [Stoffel et al., 2020](#)) and may also exhibit seasonal ([Baniel et al., 2021](#); [Björk et al., 2022](#)) and circadian dynamics ([Risely et al., 2021](#); [Schmid et al., 2023](#)). If such variables are recorded, it is often possible to control for their effects in downstream analyses. However, when sample size is limited by the economic cost of omics approaches, high levels of variability can mean statistical analyses become underpowered. Furthermore, in some cases, microbiome composition and host traits (e.g. reproductive fitness) may co-correlate with another variable (e.g. differences in abiotic conditions across years) making it difficult to

disentangle their relative effects. Thus, for certain research questions, it could be beneficial to try to minimise ecological variability through careful study design. For example, where practical, the impact of abiotic variation could be minimised by collecting samples for all individuals in the same time frame or during the same study period. Furthermore, when longitudinal microbiome dynamics are of interest, collecting a greater number of repeat samples from fewer individuals (rather than fewer samples from a wider range of individuals) could reduce additional noise and lead to a more powerful analysis.

Once sampling design has been determined, a critical step in any microbiome workflow is preserving collected samples prior to further processing and analysis. Techniques that have been developed for laboratory systems or human samples may be helpful in some cases, however, collecting microbiome samples from wild animals comes with its own set of challenges. Often, sampling is undertaken in remote field locations without access to specialist equipment, and could take place days, or even months, before the samples are processed. As such, maintaining sample integrity can be a significant issue ([Menke et al., 2015](#)), particularly for functional studies that rely on biomolecules that are very sensitive to degradation, such as RNA or metabolites.

Freezing samples immediately (at -20°C or lower) is often considered to be the gold standard for preserving microbiome samples of all types ([Cardona et al., 2012](#); [Song et al., 2016](#)). Indeed, if culturomics is the end goal, samples need to be kept cold for short periods in the field (e.g. using ice or a cryoshipper) and frozen as soon as possible for long-term storage, potentially using a cryoprotectant, such as glycerol, to retain cellular integrity ([Bircher et al., 2018](#)). However, access to freezers is not always possible at remote field sites. As such, other alternatives have been extensively evaluated, particularly for DNA-based studies of faecal samples. For example, there is some evidence that freeze-drying DNA samples have negligible impacts on microbiome community measures when compared to frozen samples ([Blekhman et al., 2016](#); [Risely et al., 2023](#)). Multiple studies also suggest that storage in 95%–100% ethanol or RNAlater can be an effective alternative if immediate freezing is not possible or specialist equipment is not available ([Blekhman et al., 2016](#); [Hale et al., 2015](#); [Song et al., 2016](#); [Vlčková et al., 2012](#)). RNAlater has the added advantage that it can be used to preserve RNA, DNA and proteins simultaneously ([Mordant & Kleiner, 2021](#)). Such buffers could be particularly useful if limited sample material is available and to improve cost-effectiveness. Similarly, pilot studies on human faecal samples have shown that buffers such as OMNImet-GUT and OMNIgene-GUT can simultaneously preserve both DNA and faecal metabolome profiles at room temperature for several days in a way that is comparable to flash-freezing ([Lim et al., 2020](#); [Ramamoorthy et al., 2021](#); [Wang et al., 2018](#)). However, further studies are needed to establish the efficacy of these techniques in field-based studies.

The most appropriate method of sample storage will vary depending on the context, the availability of resources and the research question. However, given that virtually all studies on sample storage have evidenced considerable variation in the ability of

BOX 2 Outstanding questions for ecologists and evolutionary biologists in functional microbiome research

Adopting functional omics approaches to study microbial communities, and how they interact with both their wild animal hosts and the environment, unlocks a range of questions inaccessible to researchers using metabarcoding approaches alone. Here, we list some key unanswered questions that should be addressed as an immediate priority by ecologists and evolutionary biologists studying host-microbe interactions, covering both fundamental and applied topics.

1. Is it common for wild animals to rely on a microbiome for a certain function?
2. How much of a given function is performed by each microbe under different abiotic or biotic conditions? For example, does a stressed or perturbed condition favour microbes which are metabolically independent (i.e. taxa that can independently complete all metabolic pathways defining a function)?
3. How do microbiome functions contribute to key host traits such as diet, thermoregulation, immunity and behaviour (e.g. via the gut-brain axis)?
4. Is the microbiome causally related to variation in host fitness components (condition, survival, reproductive success) among individuals and which microbial functions contribute to this relationship?
5. How much functional redundancy is there in host-associated microbiotas, whereby taxonomically distinct communities converge on similar functional profiles?
6. Does functional redundancy contribute to microbiome resilience in the face of perturbation in the wild?
7. Is high functional diversity beneficial? For example, does it confer resistance against infection by pathogens, or greater tolerance of infection?
8. What does microbiome functional diversity represent in terms of host ecosystem services? Are microbiomes that are taxonomically rich more likely to contain the one species performing the beneficial function, or is microbiome function an emergent property of multiple synergistic interactions among microbes, including bacteria, viruses and fungi?
9. How does microbiome functional diversity and composition change across animals? For example, can taxonomic patterns of phyllosymbiosis be extended to functions in different vertebrate groups (e.g. fish, mammals, birds), or does redundancy lead to functional homogenisation across taxa?
10. How important is strain-level diversity for microbiome function in wild animals? How do strains evolve in situ?
11. How plastic is microbiome function? For example, do changes in microbial transcription enable hosts to rapidly respond to abiotic/biotic challenges? Or is compositional turnover and/or strain evolution more important?
12. How can the functional properties of a microbiota be manipulated, for example, to improve host health outcomes and achieve conservation aims?

different methods to preserve sample integrity and information (Carruthers et al., 2019; Choo et al., 2015; Hale et al., 2015; Mordant & Kleiner, 2021; Vargas-Pellicer et al., 2019), using one method consistently within a study and/or system is an important consideration when planning a wild microbiome study to ensure results are comparable (Blekhman et al., 2016; Vargas-Pellicer et al., 2019).

5.2 | Processing samples

Aside from high quality material, shotgun metagenomic sequencing pipelines also require high yields of DNA to be extracted (often tens of nanograms). This can be difficult for samples with low biomass, such as oral, cloacal or skin swabs. While studies have explored whole genome amplification methods to improve yield prior to sequencing such samples (Lasken, 2009; Wang et al., 2022), the extent to which these techniques impact upon subsequent community analysis remains unclear. As such, amplification methods are not widely applied in metagenomic library preparation pipelines. This means that microbiome samples that yield high

quantities of molecular material may be better suited for metagenomic sequencing.

A further consideration for methods that use untargeted shotgun sequencing approaches (e.g. metagenomics and metatranscriptomics) is the possibility of host contamination. Untargeted sequencing methods produce reads in proportion to the relative concentrations of DNA or RNA present within a sample; this means that, if co-extracted, host molecules can significantly overwhelm signals from the microbiome and result in poor sensitivity to detect low abundance microbial taxa due to poor coverage (Pereira-Marques et al., 2019). The amount of host contamination often varies depending on the sampling site. For example, in human studies, skin, saliva and vaginal samples suffer from greater levels of host contamination than faecal samples (Marotz et al., 2018). A high sequencing depth may be necessary to overcome host contamination issues; however, various chemical approaches have also been developed that can deplete host DNA sequences (Ahannach et al., 2021; Heravi et al., 2020; Marotz et al., 2018). Such depletion methods have mainly been applied to human clinical samples and, while these may be transferable to other vertebrate systems, their general applicability to samples

from non-mammalian hosts remains to be tested. It should also be noted that while it may be desirable to remove host molecules, these sequences could add valuable information to a study and dedicated tools have been developed for this purpose (Dylus et al., 2024). For example, it may be possible to reconstruct host DNA for phylogenetic analysis. This approach was recently demonstrated in a study on termites (*Isoptera*) whereby shotgun metagenomics enabled both functional profiling of the gut microbiome and reconstruction of near-complete host mitochondrial genomes across termite species (Arora et al., 2022).

5.3 | Practicality of sequencing samples

Untargeted shotgun sequencing approaches can provide greater taxonomic resolution and functional insight compared to traditional metabarcoding techniques. However, the vast increase in sequencing depth required to achieve this resolution (tens to hundreds of millions of read pairs per sample, compared to just thousands of reads for metabarcoding studies) means that these approaches are also much more expensive (Ranjan et al., 2016; Stothart et al., 2022). In wild studies, where large sample sizes may be needed to detect small effects due to the noisy nature of the data, such costs can be prohibitive. As such, less expensive alternatives could be used to gain coarse-level insight into the microbiome before carrying out more detailed functional analyses. For example, methods such as qPCR and 16S rRNA gene metabarcoding could be used to first establish the abundance of microbial communities and broadly characterise taxonomic and functional microbiome differences among groups of wild individuals. The results could then inform hypotheses to be further investigated in metagenomic studies on a more limited sample set. Such an approach has been used in several wild metagenomics studies to assess specific functional differences once an initial group effect has been established using metabarcoding (e.g. Bates et al., 2022; Rojas et al., 2023; Sanders et al., 2015).

As discussed, a disadvantage of the above approach is that metabarcoding data sets do not always reveal the full complexity of microbiome differences among samples. This was demonstrated in a recent study comparing the gut microbiota of two honey bee species; while 16S rRNA gene metabarcoding suggested that both host species carried the same, highly conserved bacterial phylotypes, metagenomic sequencing revealed that bacterial species and strains differed between the two hosts and that this had implications for polysaccharide degradation (Ellegaard et al., 2020).

Low coverage ('shallow') metagenomic sequencing has been proposed as an economical alternative to deep metagenomic sequencing. While this approach provides less resolution than deep metagenomic sequencing and does not enable the construction of MAGs, it provides the capability to carry out species-level microbial characterisation and enables functional inference via information on microbial gene content (Hillmann et al., 2018; La Reau et al., 2023). A study using faecal samples from wild Sable Island horses (Stothart et al., 2022) showed that shallow metagenomic

sequencing yielded near identical patterns of microbial alpha and beta diversity across samples when compared to a successively rarefied deep metagenomic sequencing data set. This result was obtained despite read depths only reaching hundreds of thousands (compared to tens of millions) in the shallow data set. Shallow shotgun sequencing also recapitulated microbiome taxonomic patterns but yielded further functional insight than the metabarcoding data set (Stothart et al., 2022). Thus, the accuracy and the comparative cost-effectiveness of shallow shotgun sequencing suggest that it may be a promising alternative to metabarcoding in wild animal studies. However, further studies are needed to validate shallow sequencing across other host taxa and sample types.

5.4 | Analysis pipelines

In contrast to metabarcoding techniques where relatively well-standardised pipelines are in place (e.g. Bolyen et al., 2019), numerous equivalent and competing bioinformatic pipelines for the analysis of more complex omic data have been developed. Choosing the most appropriate analysis for a given question and data set is therefore not trivial. While we provide examples of such pipelines and tools for each omic data type in Table 1, we stress that the choice of pipeline requires a good understanding of the underlying methods, especially in non-model hosts where benchmarking of tools is usually lacking. Numerous standardised initiatives and studies have emerged to compare and benchmark tools and we refer readers to these to optimise their analytical tool choice (e.g. Jurado-Rueda et al., 2023; Meyer et al., 2022; Sczyrba et al., 2017; Ye et al., 2019). Several pipelines are also accompanied by detailed tutorials including the Web of Life Toolkit App (Woltka—<https://github.com/qiyunzhu/woltka>) (Zhu et al., 2022) and tools available from the Huttenhower lab (<https://huttenhower.sph.harvard.edu/tools/>) which have been collated into reproducible bioBakery workflows (Beghini et al., 2021; McIver et al., 2018). There are also a wealth of educational resources available (e.g. <https://datacarpentry.org/>, <https://methods-in-microbiomics.readthedocs.io/en/latest/index.html>) that can help the non-specialist to navigate through the jungle of non-standardised pipelines for omics data.

Taxonomic and functional profiling techniques rely on aligning reads, assembled genomes or metabolic profiles to reference databases. This means that all omics tools are subject to database biases and incompleteness (Bundy et al., 2009; Jiang et al., 2016; Sun et al., 2020). While this is a limitation facing all microbiome studies, it poses a particular issue for understudied wild taxa, given that databases are typically biased towards hosts of clinical or economic interest, such as humans, murine models and ruminants (Sun et al., 2020). This is evidenced by the vast amounts of taxonomic novelty identified in recent large-scale metagenomics studies of wild animal microbiomes (Levin et al., 2021; Youngblut et al., 2020). These results demonstrate that microbial databases lack information on the genetic diversity of microbes associated with most wild host species and suggest that some wild animals may be more amenable to functional studies than

others. Indeed, host species that are closely related to those that are well represented in databases (e.g. wild primates and mice) may be better candidates.

Certain bioinformatic approaches can be used to reduce the impact of database incompleteness on results. For example, in metagenomic studies, assembly-based (vs. raw-read classification) approaches allow recovery of microbial sequences that have not yet been isolated or characterised and, thus, are absent from reference databases (Tamames et al., 2019). Assembly may also yield more accurate results because longer sequences are advantageous when assessing distant homology (Tamames et al., 2019). However, assembly-based methods are still influenced by database completeness to some extent and greater computational resources and higher coverage are required to achieve good assemblies. This, in turn, increases the costs per sample. Analytical tools are continuously being developed to improve the accuracy and resolution of taxonomic and functional profiling. For example, MetaPhlan 4 now incorporates information on genome-sequenced isolates and metagenome assemblies from diverse environments (human and non-human hosts, as well as non-host associated) to enable more comprehensive microbiome taxonomic profiling (Blanco-Míguez et al., 2023).

6 | CONCLUSION

Huge progress has been made in characterising the taxonomic composition and diversity of wild animal microbiomes across many host species. However, despite this, we still have a relatively poor understanding of the functional characteristics, and importance, of the microbiome in non-model and wild hosts. This is largely because traditional marker gene profiling only provides limited taxonomic resolution and rough estimates of microbial functions.

Building on knowledge derived from more traditional (and less expensive) approaches (e.g. 16S, qPCR), microbial functions can be probed with multi-omics tools to gain a deeper mechanistic understanding of host-microbiome interactions. The continued development of sample storage techniques, cheaper sequencing methods, analytical tools and reference databases will further widen the applicability of these approaches to non-model organisms enabling key outstanding questions to be answered.

AUTHOR CONTRIBUTIONS

All authors were involved in developing ideas and concepts for the paper. SFW led the writing of the manuscript with significant input from FM, EV, XAH, JRB and KMW. All authors contributed critically to the drafts and gave final approval for publication.

ACKNOWLEDGEMENTS

This manuscript evolved from a symposium at the European Society for Evolutionary Biology (ESEB) Congress in Prague, 2022 and a special topic network (Wild Animal Microbiomes and Evolution) funded by ESEB. SFW was funded by the Natural Environmental Research Council (NE/S010939/1) and a Leverhulme Trust Early Career

Fellowship (ECF-2023-433). EV was funded by the Swedish Research Council (2020-00259). XAH was funded by the Leverhulme Trust (RPG-2020-320). FM was funded by the University of Lausanne. KMW was funded by a University of Surrey Future Fellowship. We would like to thank Amy Sweeney for feedback on the final manuscript draft.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

No new data are associated with this manuscript.

ORCID

Sarah F. Worsley  <https://orcid.org/0000-0003-4736-0938>

Elin Videvall  <https://orcid.org/0000-0002-9998-3689>

Xavier A. Harrison  <https://orcid.org/0000-0002-2004-3601>

Johannes R. Björk  <https://orcid.org/0000-0001-9768-1946>

Florent Mazel  <https://orcid.org/0000-0003-0572-9901>

Klara M. Wanelik  <https://orcid.org/0000-0003-1485-0340>

REFERENCES

- Ahannach, S., Delanghe, L., Spacova, I., Wittouck, S., Van Beeck, W., De Boeck, I., & Lebeer, S. (2021). Microbial enrichment and storage for metagenomics of vaginal, skin, and saliva samples. *iScience*, 24(11), 103306. <https://doi.org/10.1016/j.isci.2021.103306>
- Alberdi, A., Aizpurua, O., Bohmann, K., Zepeda-Mendoza, M. L., & Gilbert, M. T. P. (2016). Do vertebrate gut metagenomes confer rapid ecological adaptation? *Trends in Ecology & Evolution*, 31(9), 689–699. <https://doi.org/10.1016/j.tree.2016.06.008>
- Alberdi, A., Andersen, S. B., Limborg, M. T., Dunn, R. R., & Gilbert, M. T. P. (2022). Disentangling host-microbiota complexity through hologenomics. *Nature Reviews Genetics*, 23(5), 281–297. <https://doi.org/10.1038/s41576-021-00421-0>
- Antwis, R. E., & Harrison, X. A. (2018). Probiotic consortia are not uniformly effective against different amphibian chytrid pathogen isolates. *Molecular Ecology*, 27(2), 577–589. <https://doi.org/10.1111/mec.14456>
- Arnoldini, M., Sharma, R., Moresi, C., Chure, G., Chabbey, J., Slack, E., & Cremer, J. (2024). Quantifying the daily harvest of fermentation products from the human gut microbiota. *bioRxiv*. <https://doi.org/10.1101/2024.01.05.573977>
- Arora, J., Kinjo, Y., Šobotník, J., Buček, A., Clitheroe, C., Stiblík, P., Roisin, Y., Žifčáková, L., Park, Y. C., Kim, K. Y., Sillam-Dussès, D., Hervé, V., Lo, N., Tokuda, G., Brune, A., & Bourguignon, T. (2022). The functional evolution of termite gut microbiota. *Microbiome*, 10(1), 78. <https://doi.org/10.1186/s40168-022-01258-3>
- Baniel, A., Amato, K. R., Beehner, J. C., Bergman, T. J., Mercer, A., Perlman, R. F., Petruccio, L., Reitsema, L., Sams, S., Lu, A., & Snyder-Mackler, N. (2021). Seasonal shifts in the gut microbiome indicate plastic responses to diet in wild geladas. *Microbiome*, 9(1), 26. <https://doi.org/10.1186/s40168-020-00977-9>
- Baniel, A., Petruccio, L., Mercer, A., Reitsema, L., Sams, S., Beehner, J. C., Bergman, T. J., Snyder-Mackler, N., & Lu, A. (2022). Maternal effects on early-life gut microbiota maturation in a wild nonhuman primate. *Current Biology*, 32, 4508–4520.e6. <https://doi.org/10.1016/j.cub.2022.08.037>
- Bates, K. A., Clare, F. C., O'Hanlon, S., Bosch, J., Brookes, L., Hopkins, K., McLaughlin, E. J., Daniel, O., Garner, T. W. J., Fisher, M. C., & Harrison, X. A. (2018). Amphibian chytridiomycosis outbreak

- dynamics are linked with host skin bacterial community structure. *Nature Communications*, 9(1), 693. <https://doi.org/10.1038/s41467-018-02967-w>
- Bates, K. A., Sommer, U., Hopkins, K. P., Shelton, J. M. G., Wierzbicki, C., Sergeant, C., Tapley, B., Michaels, C. J., Schmeller, D. S., Loyau, A., Bosch, J., Viant, M. R., Harrison, X. A., Garner, T. W. J., & Fisher, M. C. (2022). Microbiome function predicts amphibian chytridiomycosis disease dynamics. *Microbiome*, 10(1), 44. <https://doi.org/10.1186/s40168-021-01215-6>
- Beghini, F., McIver, L. J., Blanco-Míguez, A., Dubois, L., Asnicar, F., Maharjan, S., Mailyan, A., Manghi, P., Scholz, M., Thomas, A. M., Valles-Colomer, M., Weingart, G., Zhang, Y., Zolfo, M., Huttenhower, C., Franzosa, E. A., & Segata, N. (2021). Integrating taxonomic, functional, and strain-level profiling of diverse microbial communities with bioBakery 3. *eLife*, 10, e65088. <https://doi.org/10.7554/eLife.65088>
- Bergman, E. N. (1990). Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiological Reviews*, 70(2), 567–590. <https://doi.org/10.1152/physrev.1990.70.2.567>
- Bhosle, A., Wang, Y., Franzosa, E. A., & Huttenhower, C. (2022). Progress and opportunities in microbial community metabolomics. *Current Opinion in Microbiology*, 70, 102195. <https://doi.org/10.1016/j.mib.2022.102195>
- Bircher, L., Geirnaert, A., Hammes, F., Lacroix, C., & Schwab, C. (2018). Effect of cryopreservation and lyophilization on viability and growth of strict anaerobic human gut microbes. *Microbial Biotechnology*, 11(4), 721–733. <https://doi.org/10.1111/1751-7915.13265>
- Björk, J. R., Dasari, M. R., Roche, K., Grieneisen, L., Gould, T. J., Grenier, J.-C., Yotova, V., Gittel, N., Jansen, D., Gesquiere, L. R., Gordon, J. B., Learn, N. H., Wango, T. L., Mututua, R. S., Kinyua Warutere, J., Siodi, L., Mukherjee, S., Barreiro, L. B., Alberts, S. C., ... Archie, E. A. (2022). Synchrony and idiosyncrasy in the gut microbiome of wild baboons. *Nature Ecology & Evolution*, 6, 955–964. <https://doi.org/10.1038/s41559-022-01773-4>
- Blanco-Míguez, A., Beghini, F., Cumbo, F., McIver, L. J., Thompson, K. N., Zolfo, M., Manghi, P., Dubois, L., Huang, K. D., Thomas, A. M., Nickols, W. A., Piccinno, G., Piperni, E., Punčochář, M., Valles-Colomer, M., Tett, A., Giordano, F., Davies, R., Wolf, J., ... Segata, N. (2023). Extending and improving metagenomic taxonomic profiling with uncharacterized species using MetaPhlan 4. *Nature Biotechnology*, 41(11), 1633–1644. <https://doi.org/10.1038/s41587-023-01688-w>
- Blekhman, R., Tang, K., Archie, E. A., Barreiro, L. B., Johnson, Z. P., Wilson, M. E., Kohn, J., Yuan, M. L., Gesquiere, L., Grieneisen, L. E., & Tung, J. (2016). Common methods for fecal sample storage in field studies yield consistent signatures of individual identity in microbiome sequencing data. *Scientific Reports*, 6, 31519. <https://doi.org/10.1038/srep31519>
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., ... Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology*, 37(8), 852–857. <https://doi.org/10.1038/s41587-019-0209-9>
- Bourne, D. G., Morrow, K. M., & Webster, N. S. (2016). Insights into the coral microbiome: Underpinning the health and resilience of reef ecosystems. *Annual Review of Microbiology*, 70(1), 317–340. <https://doi.org/10.1146/annurev-micro-102215-095440>
- Brooks, A. W., Kohl, K. D., Brucker, R. M., Van Opstal, E. J., & Bordenstein, S. R. (2016). Phyllosymbiosis: Relationships and functional effects of microbial communities across host evolutionary history. *PLoS Biology*, 14(11), e2000225. <https://doi.org/10.1371/journal.pbio.2000225>
- Bundy, J. G., Davey, M. P., & Viant, M. R. (2009). Environmental metabolomics: A critical review and future perspectives. *Metabolomics*, 5(1), 3–21. <https://doi.org/10.1007/s11306-008-0152-0>
- Cabral, L., Persinoti, G. F., Paixão, D. A. A., Martins, M. P., Morais, M. A. B., Chinaglia, M., Domingues, M. N., Sforca, M. L., Pirolla, R. A. S., Generoso, W. C., Santos, C. A., Maciel, L. F., Terrapon, N., Lombard, V., Henrissat, B., & Murakami, M. T. (2022). Gut microbiome of the largest living rodent harbors unprecedented enzymatic systems to degrade plant polysaccharides. *Nature Communications*, 13(1), 629. <https://doi.org/10.1038/s41467-022-28310-y>
- Cao, J., Hu, Y., Liu, F., Wang, Y., Bi, Y., Lv, N., Li, J., Zhu, B., & Gao, G. F. (2020). Metagenomic analysis reveals the microbiome and resistome in migratory birds. *Microbiome*, 8(1), 26. <https://doi.org/10.1186/s40168-019-0781-8>
- Cardona, S., Eck, A., Cassellas, M., Gallart, M., Alastrue, C., Dore, J., Azpiroz, F., Roca, J., Guarner, F., & Manichanh, C. (2012). Storage conditions of intestinal microbiota matter in metagenomic analysis. *BMC Microbiology*, 12(1), 158. <https://doi.org/10.1186/1471-2180-12-158>
- Carruthers, L. V., Moses, A., Adriko, M., Faust, C. L., Tukaheba, E. M., Hall, L. J., Ranford-Cartwright, L. C., & Lambertson, P. H. L. (2019). The impact of storage conditions on human stool 16S rRNA microbiome composition and diversity. *PeerJ*, 7, e8133. <https://doi.org/10.7717/peerj.8133>
- Chinchar, V. G., Bryan, L., Silphadaung, U., Noga, E., Wade, D., & Rollins-Smith, L. (2004). Inactivation of viruses infecting ectothermic animals by amphibian and piscine antimicrobial peptides. *Virology*, 323(2), 268–275. <https://doi.org/10.1016/j.virol.2004.02.029>
- Chong, J., Soufan, O., Li, C., Caraus, I., Li, S., Bourque, G., Wishart, D. S., & Xia, J. (2018). MetaboAnalyst 4.0: Towards more transparent and integrative metabolomics analysis. *Nucleic Acids Research*, 46(W1), W486–W494. <https://doi.org/10.1093/nar/gky310>
- Choo, J. M., Leong, L. E., & Rogers, G. B. (2015). Sample storage conditions significantly influence faecal microbiome profiles. *Scientific Reports*, 5(1), 16350. <https://doi.org/10.1038/srep16350>
- Clavel, T., Lagkouvardos, I., Blaut, M., & Stecher, B. (2016). The mouse gut microbiome revisited: From complex diversity to model ecosystems. *International Journal of Medical Microbiology*, 306(5), 316–327. <https://doi.org/10.1016/j.ijmm.2016.03.002>
- Dapa, T., Wong, D. P., Vasquez, K. S., Xavier, K. B., Huang, K. C., & Good, B. H. (2023). Within-host evolution of the gut microbiome. *Current Opinion in Microbiology*, 71, 102258. <https://doi.org/10.1016/j.mib.2022.102258>
- Davidson, G. L., Raulo, A., & Knowles, S. C. L. (2020). Identifying microbiome-mediated behaviour in wild vertebrates. *Trends in Ecology & Evolution*, 35(11), 972–980. <https://doi.org/10.1016/j.tree.2020.06.014>
- Davidson, G. L., Somers, S. E., Wiley, N., Johnson, C. N., Reichert, M. S., Ross, R. P., Stanton, C., & Quinn, J. L. (2021). A time-lagged association between the gut microbiome, nestling weight and nestling survival in wild great tits. *Journal of Animal Ecology*, 1365-2656, 13428. <https://doi.org/10.1111/1365-2656.13428>
- Davies, C. S., Worsley, S. F., Maher, K. H., Komdeur, J., Burke, T., Dugdale, H. L., & Richardson, D. S. (2022). Immunogenetic variation shapes the gut microbiome in a natural vertebrate population. *Microbiome*, 10(1), 41. <https://doi.org/10.1186/s40168-022-01233-y>
- De Filippis, F., Pasoli, E., Tett, A., Tarallo, S., Naccarati, A., De Angelis, M., Neviani, E., Cocolin, L., Gobbetti, M., Segata, N., & Ercolini, D. (2019). Distinct genetic and functional traits of human intestinal *Prevotella copri* strains are associated with different habitual diets. *Cell Host & Microbe*, 25(3), 444–453.e3. <https://doi.org/10.1016/j.chom.2019.01.004>
- Djemiel, C., Maron, P.-A., Terrat, S., Dequiedt, S., Cottin, A., & Ranjard, L. (2022). Inferring microbiota functions from taxonomic genes: A review. *GigaScience*, 11, giab090. <https://doi.org/10.1093/gigascience/giab090>

- Doolittle, W. F., & Booth, A. (2017). It's the song, not the singer: An exploration of holobiosis and evolutionary theory. *Biology & Philosophy*, 32(1), 5–24. <https://doi.org/10.1007/s10539-016-9542-2>
- Douglas, G. M., Maffei, V. J., Zaneveld, J. R., Yurgel, S. N., Brown, J. R., Taylor, C. M., Huttenhower, C., & Langille, M. G. I. (2020). PICRUSt2 for prediction of metagenome functions. *Nature Biotechnology*, 38(6), 685–688. <https://doi.org/10.1038/s41587-020-0548-6>
- Dylus, D., Altenhoff, A., Majidian, S., Sedlazeck, F. J., & Dessimoz, C. (2024). Inference of phylogenetic trees directly from raw sequencing reads using Read2Tree. *Nature Biotechnology*, 42(1), 139–147. <https://doi.org/10.1038/s41587-023-01753-4>
- Ellegaard, K. M., Suenami, S., Miyazaki, R., & Engel, P. (2020). Vast differences in strain-level diversity in the gut microbiota of two closely related honey bee species. *Current Biology*, 30(13), 2520–2531.e7. <https://doi.org/10.1016/j.cub.2020.04.070>
- Fackelmann, G., Gillingham, M. A. F., Schmid, J., Heni, A. C., Wilhelm, K., Schwensow, N., & Sommer, S. (2021). Human encroachment into wildlife gut microbiomes. *Communications Biology*, 4(1), 800. <https://doi.org/10.1038/s42003-021-02315-7>
- Foster, K. R., Schluter, J., Coyte, K. Z., & Rakoff-Nahoum, S. (2017). The evolution of the host microbiome as an ecosystem on a leash. *Nature*, 548(7665), 43–51. <https://doi.org/10.1038/nature23292>
- Franzosa, E. A., Morgan, X. C., Segata, N., Waldron, L., Reyes, J., Earl, A. M., Giannoukos, G., Boylan, M. R., Ciulla, D., Gevers, D., Izard, J., Garrett, W. S., Chan, A. T., & Huttenhower, C. (2014). Relating the metatranscriptome and metagenome of the human gut. *Proceedings of the National Academy of Sciences of the United States of America*, 111(22), E2329–E2338. <https://doi.org/10.1073/pnas.1319284111>
- Gosalbes, M. J., Durbán, A., Pignatelli, M., Abellan, J. J., Jiménez-Hernández, N., Pérez-Cobas, A. E., Latorre, A., & Moya, A. (2011). Metatranscriptomic approach to analyze the functional human gut microbiota. *PLoS One*, 6(3), e17447. <https://doi.org/10.1371/journal.pone.0017447>
- Grieneisen, L. E., Charpentier, M. J. E., Alberts, S. C., Blekhan, R., Bradburd, G., Tung, J., & Archie, E. A. (2019). Genes, geology and germs: Gut microbiota across a primate hybrid zone are explained by site soil properties, not host species. *Proceedings of the Royal Society B: Biological Sciences*, 286(1901), 20190431. <https://doi.org/10.1098/rspb.2019.0431>
- Grond, K., Louyakis, A. S., & Hird, S. M. (2023). Functional and compositional changes in the fecal microbiome of a shorebird during migratory stopover. *mSystems*, 8, e01128-22. <https://doi.org/10.1128/mSystems.01128-22>
- Grossin, M., Mazel, F., Sanders, J. G., Smillie, C. S., Lavergne, S., Thuiller, W., & Alm, E. J. (2017). Unraveling the processes shaping mammalian gut microbiomes over evolutionary time. *Nature Communications*, 8(1), 14319. <https://doi.org/10.1038/ncomms14319>
- Hajjo, H., & Geva-Zatorsky, N. (2021). Strain-level immunomodulatory variation of gut bacteria. *FEBS Letters*, 595(9), 1322–1327. <https://doi.org/10.1002/1873-3468.14057>
- Hale, V. L., Tan, C. L., Knight, R., & Amato, K. R. (2015). Effect of preservation method on spider monkey (*Ateles geoffroyi*) fecal microbiota over 8 weeks. *Journal of Microbiological Methods*, 113, 16–26. <https://doi.org/10.1016/j.mimet.2015.03.021>
- Hall, A. B., Yassour, M., Sauk, J., Garner, A., Jiang, X., Arthur, T., Lagoudas, G. K., Vatanen, T., Fornelos, N., Wilson, R., Bertha, M., Cohen, M., Garber, J., Khalili, H., Gevers, D., Ananthakrishnan, A. N., Kugathasan, S., Lander, E. S., Blainey, P., ... Huttenhower, C. (2017). A novel *Ruminococcus gnavus* clade enriched in inflammatory bowel disease patients. *Genome Medicine*, 9(1), 103. <https://doi.org/10.1186/s13073-017-0490-5>
- Hammer, T. J., Janzen, D. H., Hallwachs, W., Jaffe, S. P., & Fierer, N. (2017). Caterpillars lack a resident gut microbiome. *Proceedings of the National Academy of Sciences of the United States of America*, 114(36), 9641–9646. <https://doi.org/10.1073/pnas.1707186114>
- Hammer, T. J., Sanders, J. G., & Fierer, N. (2019). Not all animals need a microbiome. *FEMS Microbiology Letters*, 366(10), fnz117. <https://doi.org/10.1093/femsle/fnz117>
- Harrison, X. A., Price, S. J., Hopkins, K., Leung, W. T. M., Sergeant, C., & Garner, T. W. J. (2019). Diversity-stability dynamics of the amphibian skin microbiome and susceptibility to a lethal viral pathogen. *Frontiers in Microbiology*, 10, 2883. <https://doi.org/10.3389/fmicb.2019.02883>
- Harrison, X. A., Sewell, T., Fisher, M., & Antwis, R. E. (2020). Designing probiotic therapies with broad-spectrum activity against a wildlife pathogen. *Frontiers in Microbiology*, 10, 3134. <https://doi.org/10.3389/fmicb.2019.03134>
- Heintz-Buschart, A., May, P., Laczny, C. C., Lebrun, L. A., Bellora, C., Krishna, A., Wampach, L., Schneider, J. G., Hogan, A., De Beaufort, C., & Wilmes, P. (2016). Integrated multi-omics of the human gut microbiome in a case study of familial type 1 diabetes. *Nature Microbiology*, 2(1), 16180. <https://doi.org/10.1038/nmicrobiol.2016.180>
- Heintz-Buschart, A., & Wilmes, P. (2018). Human gut microbiome: Function matters. *Trends in Microbiology*, 26(7), 563–574. <https://doi.org/10.1016/j.tim.2017.11.002>
- Heravi, F. S., Zakrzewski, M., Vickery, K., & Hu, H. (2020). Host DNA depletion efficiency of microbiome DNA enrichment methods in infected tissue samples. *Journal of Microbiological Methods*, 170, 105856. <https://doi.org/10.1016/j.mimet.2020.105856>
- Hillmann, B., Al-Ghalith, G. A., Shields-Cutler, R. R., Zhu, Q., Gohl, D. M., Beckman, K. B., Knight, R., & Knights, D. (2018). Evaluating the information content of shallow shotgun metagenomics. *mSystems*, 3(6), e00069-18. <https://doi.org/10.1128/mSystems.00069-18>
- Holmes, N. A., Innocent, T. M., Heine, D., Bassam, M. A., Worsley, S. F., Trottmann, F., Patrick, E. H., Yu, D. W., Murrell, J. C., Schiøtt, M., Wilkinson, B., Boomsma, J. J., & Hutchings, M. I. (2016). Genome analysis of two *Pseudonocardia* phylotypes associated with *Acromyrmex* leafcutter ants reveals their biosynthetic potential. *Frontiers in Microbiology*, 7, 2073. <https://doi.org/10.3389/fmicb.2016.02073>
- Hu, Y., Sanders, J. G., Łukasik, P., D'Amelio, C. L., Millar, J. S., Vann, D. R., Lan, Y., Newton, J. A., Schotanus, M., Kronauer, D. J. C., Pierce, N. E., Moreau, C. S., Wertz, J. T., Engel, P., & Russell, J. A. (2018). Herbivorous turtle ants obtain essential nutrients from a conserved nitrogen-recycling gut microbiome. *Nature Communications*, 9(1), 964. <https://doi.org/10.1038/s41467-018-03357-y>
- Huan, T., Forsberg, E. M., Rinehart, D., Johnson, C. H., Ivanisevic, J., Benton, H. P., Fang, M., Aisporna, A., Hilmers, B., Poole, F. L., Thorgersen, M. P., Adams, M. W. W., Krantz, G., Fields, M. W., Robbins, P. D., Niedernhofer, L. J., Ideker, T., Majumder, E. L., Wall, J. D., ... Siuzdak, G. (2017). Systems biology guided by XCMS online metabolomics. *Nature Methods*, 14(5), 461–462. <https://doi.org/10.1038/nmeth.4260>
- Huang, G., Wang, L., Li, J., Hou, R., Wang, M., Wang, Z., Qu, Q., Zhou, W., Nie, Y., Hu, Y., Ma, Y., Yan, L., Wei, H., & Wei, F. (2022). Seasonal shift of the gut microbiome synchronizes host peripheral circadian rhythm for physiological adaptation to a low-fat diet in the giant panda. *Cell Reports*, 38(3), 110203. <https://doi.org/10.1016/j.celrep.2021.110203>
- Ingala, M. R., Simmons, N. B., Dunbar, M., Wultsch, C., Krampis, K., & Perkins, S. L. (2021). You are more than what you eat: Potentially adaptive enrichment of microbiome functions across bat dietary niches. *Animal Microbiome*, 3(1), 82. <https://doi.org/10.1186/s4252-3-021-00139-8>
- Jacobs, J. P., Lagishetty, V., Hauer, M. C., Labus, J. S., Dong, T. S., Toma, R., Vuyisich, M., Naliboff, B. D., Lackner, J. M., Gupta, A., Tillisch, K., & Mayer, E. A. (2023). Multi-omics profiles of the intestinal microbiome in irritable bowel syndrome and its bowel habit subtypes.

- Microbiome*, 11(1), 5. <https://doi.org/10.1186/s40168-022-01450-5>
- Jiang, Y., Xiong, X., Danska, J., & Parkinson, J. (2016). Metatranscriptomic analysis of diverse microbial communities reveals core metabolic pathways and microbiome-specific functionality. *Microbiome*, 4(1), 2. <https://doi.org/10.1186/s40168-015-0146-x>
- Jurado-Rueda, F., Alonso-Guirado, L., Perea-Chamblee, T. E., Elliott, O. T., Filip, I., Rabadán, R., & Malats, N. (2023). Benchmarking of microbiome detection tools on RNA-seq synthetic databases according to diverse conditions. *Bioinformatics Advances*, 3(1), vbad014. <https://doi.org/10.1093/bioadv/vbad014>
- Kang, D. D., Li, F., Kirtan, E., Thomas, A., Egan, R., An, H., & Wang, Z. (2019). MetaBAT 2: An adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ*, 7, e7359. <https://doi.org/10.7717/peerj.7359>
- Kearns, P. J., Fischer, S., Fernández-Beakoetxea, S., Gabor, C. R., Bosch, J., Bowen, J. L., Tlustý, M. F., & Woodhams, D. C. (2017). Fight fungi with fungi: Antifungal properties of the amphibian mycobiome. *Frontiers in Microbiology*, 8, 2494. <https://doi.org/10.3389/fmicb.2017.02494>
- Kim, J., Kim, M. S., Koh, A. Y., Xie, Y., & Zhan, X. (2016). FMAP: Functional mapping and analysis pipeline for metagenomics and metatranscriptomics studies. *BMC Bioinformatics*, 17(1), 420. <https://doi.org/10.1186/s12859-016-1278-0>
- Kleiner, M. (2019). Metaproteomics: Much more than measuring gene expression in microbial communities. *mSystems*, 4(3), e00115-19. <https://doi.org/10.1128/mSystems.00115-19>
- Knowles, S. C. L., Eccles, R. M., & Baltrūnaitė, L. (2019). Species identity dominates over environment in shaping the microbiota of small mammals. *Ecology Letters*, 22(5), 826–837. <https://doi.org/10.1111/ele.13240>
- Kueneman, J. G., Woodhams, D. C., Harris, R., Archer, H. M., Knight, R., & McKenzie, V. J. (2016). Probiotic treatment restores protection against lethal fungal infection lost during amphibian captivity. *Proceedings of the Royal Society B: Biological Sciences*, 283(1839), 20161553. <https://doi.org/10.1098/rspb.2016.1553>
- Kwong, W. K., Engel, P., Koch, H., & Moran, N. A. (2014). Genomics and host specialization of honey bee and bumble bee gut symbionts. *Proceedings of the National Academy of Sciences of the United States of America*, 111(31), 11509–11514. <https://doi.org/10.1073/pnas.1405838111>
- La Reau, A. J., Strom, N. B., Filvaroff, E., Mavrommatis, K., Ward, T. L., & Knights, D. (2023). Shallow shotgun sequencing reduces technical variation in microbiome analysis. *Scientific Reports*, 13(1), 7668. <https://doi.org/10.1038/s41598-023-33489-1>
- Lagier, J.-C., Dubourg, G., Million, M., Cadoret, F., Bilen, M., Fenollar, F., Lévassieur, A., Rolain, J.-M., Fournier, P.-E., & Raoult, D. (2018). Culturing the human microbiota and culturomics. *Nature Reviews Microbiology*, 16(9), 540–550. <https://doi.org/10.1038/s41579-018-0041-0>
- Lasken, R. S. (2009). Genomic DNA amplification by the multiple displacement amplification (MDA) method. *Biochemical Society Transactions*, 37(2), 450–453. <https://doi.org/10.1042/BST0370450>
- Leclaire, S., Pineaux, M., Blanchard, P., White, J., & Hatch, S. A. (2022). Microbiota composition and diversity of multiple body sites vary according to reproductive performance in a seabird. *Molecular Ecology*, 32, 2115–2133. <https://doi.org/10.1111/mec.16398>
- Lee, J. W. J., Plichta, D., Hogstrom, L., Borren, N. Z., Lau, H., Gregory, S. M., Tan, W., Khalili, H., Clish, C., Vlamakis, H., Xavier, R. J., & Ananthakrishnan, A. N. (2021). Multi-omics reveal microbial determinants impacting responses to biologic therapies in inflammatory bowel disease. *Cell Host & Microbe*, 29(8), 1294–1304.e4. <https://doi.org/10.1016/j.chom.2021.06.019>
- Levin, D., Raab, N., Pinto, Y., Rothschild, D., Zanir, G., Godneva, A., Mellul, N., Futorian, D., Gal, D., Leviatan, S., Zeevi, D., Bachelet, I., & Segal, E. (2021). Diversity and functional landscapes in the microbiota of animals in the wild. *Science*, 372(6539), eabb5352. <https://doi.org/10.1126/science.abb5352>
- Li, D., Liu, C.-M., Luo, R., Sadakane, K., & Lam, T.-W. (2015). MEGAHIT: An ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*, 31(10), 1674–1676. <https://doi.org/10.1093/bioinformatics/btv033>
- Lim, M. Y., Hong, S., Kim, B.-M., Ahn, Y., Kim, H.-J., & Nam, Y.-D. (2020). Changes in microbiome and metabolomic profiles of fecal samples stored with stabilizing solution at room temperature: A pilot study. *Scientific Reports*, 10(1), 1789. <https://doi.org/10.1038/s41598-020-58719-8>
- Lloyd-Price, J., Arze, C., Ananthakrishnan, A. N., Schirmer, M., Avila-Pacheco, J., Poon, T. W., Andrews, E., Ajami, N. J., Bonham, K. S., Brislawn, C. J., Casero, D., Courtney, H., Gonzalez, A., Graeber, T. G., Hall, A. B., Lake, K., Landers, C. J., Mallick, H., Plichta, D. R., ... IBDMDB Investigators. (2019). Multi-omics of the gut microbial ecosystem in inflammatory bowel diseases. *Nature*, 569(7758), 655–662. <https://doi.org/10.1038/s41586-019-1237-9>
- Louca, S., Polz, M. F., Mazel, F., Albright, M. B. N., Huber, J. A., O'Connor, M. I., Ackermann, M., Hahn, A. S., Srivastava, D. S., Crowe, S. A., Doebeli, M., & Parfrey, L. W. (2018). Function and functional redundancy in microbial systems. *Nature Ecology & Evolution*, 2(6), 936–943. <https://doi.org/10.1038/s41559-018-0519-1>
- Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K., & Knight, R. (2012). Diversity, stability and resilience of the human gut microbiota. *Nature*, 489(7415), 220–230. <https://doi.org/10.1038/nature11550>
- Mallott, E. K., Skovmand, L. H., Garber, P. A., & Amato, K. R. (2022). The faecal metabolome of black howler monkeys (*Alouatta pigra*) varies in response to seasonal dietary changes. *Molecular Ecology*, 31(15), 4146–4161. <https://doi.org/10.1111/mec.16559>
- Marotz, C. A., Sanders, J. G., Zuniga, C., Zaramela, L. S., Knight, R., & Zengler, K. (2018). Improving saliva shotgun metagenomics by chemical host DNA depletion. *Microbiome*, 6(1), 42. <https://doi.org/10.1186/s40168-018-0426-3>
- Martinez, X., Pozuelo, M., Pascal, V., Campos, D., Gut, I., Gut, M., Azpiroz, F., Guarner, F., & Manichanh, C. (2016). MetaTrans: An open-source pipeline for metatranscriptomics. *Scientific Reports*, 6(1), 26447. <https://doi.org/10.1038/srep26447>
- McIver, L. J., Abu-Ali, G., Franzosa, E. A., Schwager, R., Morgan, X. C., Waldron, L., Segata, N., & Huttenhower, C. (2018). bioBakery: A meta-omic analysis environment. *Bioinformatics*, 34(7), 1235–1237. <https://doi.org/10.1093/bioinformatics/btx754>
- Menke, S., Meier, M., & Sommer, S. (2015). Shifts in the gut microbiome observed in wildlife faecal samples exposed to natural weather conditions: Lessons from time-series analyses using next-generation sequencing for application in field studies. *Methods in Ecology and Evolution*, 6(9), 1080–1087. <https://doi.org/10.1111/2041-210X.12394>
- Meyer, F., Fritz, A., Deng, Z.-L., Koslicki, D., Lesker, T. R., Gurevich, A., Robertson, G., Alser, M., Antipov, D., Beghini, F., Bertrand, D., Brito, J. J., Brown, C. T., Buchmann, J., Buluç, A., Chen, B., Chikhi, R., Clausen, P. T. L. C., Cristian, A., ... McHardy, A. C. (2022). Critical assessment of metagenome interpretation: The second round of challenges. *Nature Methods*, 19(4), 429–440. <https://doi.org/10.1038/s41592-022-01431-4>
- Michel, A. J., Ward, L. M., Goffredi, S. K., Dawson, K. S., Baldassarre, D. T., Brenner, A., Gotanda, K. M., McCormack, J. E., Mullin, S. W., O'Neill, A., Tender, G. S., Uy, J. A. C., Yu, K., Orphan, V. J., & Chaves, J. A. (2018). The gut of the finch: Uniqueness of the gut microbiome of the Galápagos vampire finch. *Microbiome*, 6(1), 167. <https://doi.org/10.1186/s40168-018-0555-8>
- Milani, C., Alessandri, G., Mancabelli, L., Mangifesta, M., Lugli, G. A., Viappiani, A., Longhi, G., Anzalone, R., Duranti, S., Turrioni, F., Ossiprandi, M. C., Van Sinderen, D., & Ventura, M. (2020).

- Multi-omics approaches to decipher the impact of diet and host physiology on the mammalian gut microbiome. *Applied and Environmental Microbiology*, 86(23), e01864-20. <https://doi.org/10.1128/AEM.01864-20>
- Mizrahi, I., Wallace, R. J., & Moraïs, S. (2021). The rumen microbiome: Balancing food security and environmental impacts. *Nature Reviews Microbiology*, 19(9), 553–566. <https://doi.org/10.1038/s41579-021-00543-6>
- Mongad, D. S., Chavan, N. S., Narwade, N. P., Dixit, K., Shouche, Y. S., & Dhotre, D. P. (2021). MicFunPred: A conserved approach to predict functional profiles from 16S rRNA gene sequence data. *Genomics*, 113(6), 3635–3643. <https://doi.org/10.1016/j.ygeno.2021.08.016>
- Mordant, A., & Kleiner, M. (2021). Evaluation of sample preservation and storage methods for metaproteomics analysis of intestinal microbiomes. *Microbiology Spectrum*, 9(3), e01877-21. <https://doi.org/10.1128/Spectrum.01877-21>
- Motta, E. V. S., & Moran, N. A. (2023). The honeybee microbiota and its impact on health and disease. *Nature Reviews Microbiology*, 22, 122–137. <https://doi.org/10.1038/s41579-023-00990-3>
- Muth, T., Behne, A., Heyer, R., Kohrs, F., Benndorf, D., Hoffmann, M., Lehtevä, M., Reichl, U., Martens, L., & Rapp, E. (2015). The MetaProteomeAnalyzer: A powerful open-source software suite for metaproteomics data analysis and interpretation. *Journal of Proteome Research*, 14(3), 1557–1565. <https://doi.org/10.1021/pr501246w>
- Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., Schilling, J. S., & Kennedy, P. G. (2016). FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecology*, 20, 241–248. <https://doi.org/10.1016/j.funeco.2015.06.006>
- Nicholson, J. K., Holmes, E., Kinross, J., Burcelin, R., Gibson, G., Jia, W., & Pettersson, S. (2012). Host-gut microbiota metabolic interactions. *Science*, 336(6086), 1262–1267. <https://doi.org/10.1126/science.1223813>
- Nishida, A. H., & Ochman, H. (2018). Rates of gut microbiome divergence in mammals. *Molecular Ecology*, 27(8), 1884–1897. <https://doi.org/10.1111/mec.14473>
- Niu, S.-Y., Yang, J., McDermaid, A., Zhao, J., Kang, Y., & Ma, Q. (2017). Bioinformatics tools for quantitative and functional metagenome and metatranscriptome data analysis in microbes. *Briefings in Bioinformatics*, 19, 1415–1429. <https://doi.org/10.1093/bib/bbx051>
- O'Hara, E., Neves, A. L. A., Song, Y., & Guan, L. L. (2020). The role of the gut microbiome in cattle production and health: Driver or passenger? *Annual Review of Animal Biosciences*, 8(1), 199–220. <https://doi.org/10.1146/annurev-animal-021419-083952>
- Parry, R. H., Slonchak, A., Campbell, L. J., Newton, N. D., Debat, H. J., Gifford, R. J., & Khromykh, A. A. (2023). A novel tamanavirus (*Flaviviridae*) of the European common frog (*Rana temporaria*) from the UK. *The Journal of General Virology*, 104(12), 001927. <https://doi.org/10.1099/jgv.0.001927>
- Pereira-Marques, J., Hout, A., Ferreira, R. M., Weber, M., Pinto-Ribeiro, I., Van Doorn, L.-J., Knetsch, C. W., & Figueiredo, C. (2019). Impact of host DNA and sequencing depth on the taxonomic resolution of whole metagenome sequencing for microbiome analysis. *Frontiers in Microbiology*, 10, 1277. <https://doi.org/10.3389/fmicb.2019.01277>
- Peters, D. L., Wang, W., Zhang, X., Ning, Z., Mayne, J., & Figeys, D. (2019). Metaproteomic and metabolomic approaches for characterizing the gut microbiome. *Proteomics*, 19(16), 1800363. <https://doi.org/10.1002/pmic.201800363>
- Piovia-Scott, J., Rejmanek, D., Woodhams, D. C., Worth, S. J., Kenny, H., McKenzie, V., Lawler, S. P., & Foley, J. E. (2017). Greater species richness of bacterial skin symbionts better suppresses the amphibian fungal pathogen *Batrachochytrium dendrobatidis*. *Microbial Ecology*, 74(1), 217–226. <https://doi.org/10.1007/s0024-8-016-0916-4>
- Pluskal, T., Castillo, S., Villar-Briones, A., & Orešič, M. (2010). MZmine 2: Modular framework for processing, visualizing, and analyzing mass spectrometry-based molecular profile data. *BMC Bioinformatics*, 11(1), 395. <https://doi.org/10.1186/1471-2105-11-395>
- Pölme, S., Abarenkov, K., Henrik Nilsson, R., Lindahl, B. D., Clemmensen, K. E., Kauterud, H., Nguyen, N., Kjølter, R., Bates, S. T., Baldrian, P., Frøslev, T. G., Adojaan, K., Vizzini, A., Suija, A., Pfister, D., Baral, H.-O., Järv, H., Madrid, H., Nordén, J., ... Tedersoo, L. (2020). FungalTraits: A user-friendly traits database of fungi and fungus-like stramenopiles. *Fungal Diversity*, 105(1), 1–16. <https://doi.org/10.1007/s13225-020-00466-2>
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2012). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research*, 41(D1), D590–D596. <https://doi.org/10.1093/nar/gks1219>
- Quince, C., Walker, A. W., Simpson, J. T., Loman, N. J., & Segata, N. (2017). Shotgun metagenomics, from sampling to analysis. *Nature Biotechnology*, 35(9), 833–844. <https://doi.org/10.1038/nbt.3935>
- Ramamoorthy, S., Levy, S., Mohamed, M., Abdelghani, A., Evans, A. M., Miller, L. A. D., Mehta, L., Moore, S., Freinkman, E., & Hourigan, S. K. (2021). An ambient-temperature storage and stabilization device performs comparably to flash-frozen collection for stool metabolomics in infants. *BMC Microbiology*, 21(1), 59. <https://doi.org/10.1186/s12866-021-02104-6>
- Ranjan, R., Rani, A., Metwally, A., McGee, H. S., & Perkins, D. L. (2016). Analysis of the microbiome: Advantages of whole genome shotgun versus 16S amplicon sequencing. *Biochemical and Biophysical Research Communications*, 469(4), 967–977. <https://doi.org/10.1016/j.bbrc.2015.12.083>
- Rasko, D. A., Rosovitz, M. J., Myers, G. S. A., Mongodin, E. F., Fricke, W. F., Gajer, P., Crabtree, J., Sebahia, M., Thomson, N. R., Chaudhuri, R., Henderson, I. R., Sperandio, V., & Ravel, J. (2008). The Pangenome structure of *Escherichia coli*: Comparative genomic analysis of *E. coli* commensal and pathogenic isolates. *Journal of Bacteriology*, 190(20), 6881–6893. <https://doi.org/10.1128/JB.00619-08>
- Raulo, A., Allen, B. E., Troitsky, T., Husby, A., Firth, J. A., Coulson, T., & Knowles, S. C. L. (2021). Social networks strongly predict the gut microbiota of wild mice. *The ISME Journal*, 15(9), 2601–2613. <https://doi.org/10.1038/s41396-021-00949-3>
- Rebollar, E. A., Antwis, R. E., Becker, M. H., Belden, L. K., Bletz, M. C., Brucker, R. M., Harrison, X. A., Hughey, M. C., Kueneman, J. G., Loudon, A. H., McKenzie, V., Medina, D., Minbiole, K. P. C., Rollins-Smith, L. A., Walke, J. B., Weiss, S., Woodhams, D. C., & Harris, R. N. (2016). Using “omics” and integrated multi-omics approaches to guide probiotic selection to mitigate chytridiomycosis and other emerging infectious diseases. *Frontiers in Microbiology*, 7, 68. <https://doi.org/10.3389/fmicb.2016.00068>
- Rebollar, E. A., Gutiérrez-Preciado, A., Noecker, C., Eng, A., Hughey, M. C., Medina, D., Walke, J. B., Borenstein, E., Jensen, R. V., Belden, L. K., & Harris, R. N. (2018). The skin microbiome of the neotropical frog *Craugastor fitzingeri*: Inferring potential bacterial-host-pathogen interactions from metagenomic data. *Frontiers in Microbiology*, 9, 466. <https://doi.org/10.3389/fmicb.2018.00466>
- Rebollar, E. A., Martínez-Ugalde, E., & Orta, A. H. (2020). The amphibian skin microbiome and its protective role against chytridiomycosis. *Herpetologica*, 76(2), 167. <https://doi.org/10.1655/0018-0831-76.2.167>
- Reese, A. T., Phillips, S. R., Owens, L. A., Venable, E. M., Langergraber, K. E., Machanda, Z. P., Mitani, J. C., Muller, M. N., Watts, D. P., Wrangham, R. W., Goldberg, T. L., Emery Thompson, M., & Carmody, R. N. (2021). Age patterning in wild chimpanzee gut microbiota diversity reveals differences from humans in early life. *Current Biology*, 31(3), 613–620.e3. <https://doi.org/10.1016/j.cub.2020.10.075>

- Ren, T., Boutin, S., Humphries, M. M., Dantzer, B., Gorrell, J. C., Coltman, D. W., McAdam, A. G., & Wu, M. (2017). Seasonal, spatial, and maternal effects on gut microbiome in wild red squirrels. *Microbiome*, 5(1), 163. <https://doi.org/10.1186/s40168-017-0382-3>
- Rennison, D. J., Rudman, S. M., & Schluter, D. (2019). Parallel changes in gut microbiome composition and function during colonization, local adaptation and ecological speciation. *Proceedings of the Royal Society B: Biological Sciences*, 286(1916), 20191911. <https://doi.org/10.1098/rspb.2019.1911>
- Renwick, S., Ganobis, C. M., Elder, R. A., Gianetto-Hill, C., Higgins, G., Robinson, A. V., Vancuren, S. J., Wilde, J., & Allen-Vercoe, E. (2021). Culturing human gut microbiomes in the laboratory. *Annual Review of Microbiology*, 75(1), 49–69. <https://doi.org/10.1146/annurev-micro-031021-084116>
- Risely, A., Müller-Klein, N., Schmid, D. W., Wilhelm, K., Clutton-Brock, T. H., Manser, M. B., & Sommer, S. (2023). Climate change drives loss of bacterial gut mutualists at the expense of host survival in wild meerkats. *Global Change Biology*, 29(20), 5816–5828. <https://doi.org/10.1111/gcb.16877>
- Risely, A., Schmid, D. W., Müller-Klein, N., Wilhelm, K., Clutton-Brock, T. H., Manser, M. B., & Sommer, S. (2022). Gut microbiota individuality is contingent on temporal scale and age in wild meerkats. *Proceedings of the Royal Society B: Biological Sciences*, 289(1981), 20220609. <https://doi.org/10.1098/rspb.2022.0609>
- Risely, A., Wilhelm, K., Clutton-Brock, T., Manser, M. B., & Sommer, S. (2021). Diurnal oscillations in gut bacterial load and composition eclipse seasonal and lifetime dynamics in wild meerkats. *Nature Communications*, 12(1), 6017. <https://doi.org/10.1038/s41467-021-26298-5>
- Rojas, C. A., Holekamp, K. E., Viladomat Jasso, M., Souza, V., Eisen, J. A., & Theis, K. R. (2023). Taxonomic, genomic, and functional variation in the Gut microbiomes of wild spotted hyenas across 2 decades of study. *mSystems*, 8(1), e00965-22. <https://doi.org/10.1128/msystems.00965-22>
- Royo, D., Méndez-García, C., Raczkowska, B. A., Bargiela, R., Moya, A., Ferrer, M., & Barbas, C. (2017). Exploring the human microbiome from multiple perspectives: Factors altering its composition and function. *FEMS Microbiology Reviews*, 41(4), 453–478. <https://doi.org/10.1093/femsre/fuw046>
- Roodgar, M., Good, B. H., Garud, N. R., Martis, S., Avula, M., Zhou, W., Lancaster, S. M., Lee, H., Babveyh, A., Nesamoney, S., Pollard, K. S., & Snyder, M. P. (2021). Longitudinal linked-read sequencing reveals ecological and evolutionary responses of a human gut microbiome during antibiotic treatment. *Genome Research*, 31(8), 1433–1446. <https://doi.org/10.1101/gr.265058.120>
- Round, J. L., & Mazmanian, S. K. (2009). The gut microbiota shapes intestinal immune responses during health and disease. *Nature Reviews Immunology*, 9(5), 313–323. <https://doi.org/10.1038/nri2515>
- Rühlemann, M. C., Bang, C., Gogarten, J. F., Hermes, B. M., Groussin, M., Waschina, S., Poyet, M., Ulrich, M., Akoua-Koffi, C., Deschner, T., Muyembe-Tamfum, J. J., Robbins, M. M., Surbeck, M., Wittig, R. M., Zuberbühler, K., Baines, J. F., Leendertz, F. H., & Franke, A. (2024). Functional host-specific adaptation of the intestinal microbiome in hominids. *Nature Communications*, 15(1), 326. <https://doi.org/10.1038/s41467-023-44636-7>
- Sadoughi, B., Schneider, D., Daniel, R., Schülke, O., & Ostner, J. (2022). Aging gut microbiota of wild macaques are equally diverse, less stable, but progressively personalized. *Microbiome*, 10(1), 95. <https://doi.org/10.1186/s40168-022-01283-2>
- Sajulga, R., Easterly, C., Riffle, M., Mesuere, B., Muth, T., Mehta, S., Kumar, P., Johnson, J., Gruening, B. A., Schiebenhoefer, H., Kolmeder, C. A., Fuchs, S., Nunn, B. L., Rudney, J., Griffin, T. J., & Jagtap, P. D. (2020). Survey of metaproteomics software tools for functional microbiome analysis. *PLoS One*, 15(11), e0241503. <https://doi.org/10.1371/journal.pone.0241503>
- Sanders, J. G., Beichman, A. C., Roman, J., Scott, J. J., Emerson, D., McCarthy, J. J., & Girguis, P. R. (2015). Baleen whales host a unique gut microbiome with similarities to both carnivores and herbivores. *Nature Communications*, 6(1), 8285. <https://doi.org/10.1038/ncomms9285>
- Sanders, J. G., Sprockett, D. D., Li, Y., Mjungu, D., Lonsdorf, E. V., Ndjango, J.-B. N., Georgiev, A. V., Hart, J. A., Sanz, C. M., Morgan, D. B., Peeters, M., Hahn, B. H., & Moeller, A. H. (2023). Widespread extinctions of co-diversified primate gut bacterial symbionts from humans. *Nature Microbiology*, 8(6), 1039–1050. <https://doi.org/10.1038/s41564-023-01388-w>
- Scharf, M. E., & Peterson, B. F. (2021). A century of synergy in termite symbiosis research: Linking the past with new genomic insights. *Annual Review of Entomology*, 66(1), 23–43. <https://doi.org/10.1146/annurev-ento-022420-074746>
- Schiebenhoefer, H., Schallert, K., Renard, B. Y., Trappe, K., Schmid, E., Benndorf, D., Riedel, K., Muth, T., & Fuchs, S. (2020). A complete and flexible workflow for metaproteomics data analysis based on MetaProteomeAnalyzer and Prophan. *Nature Protocols*, 15(10), 3212–3239. <https://doi.org/10.1038/s41596-020-0368-7>
- Schmid, D. W., Capilla-Lasheras, P., Dominoni, D. M., Müller-Klein, N., Sommer, S., & Risely, A. (2023). Circadian rhythms of hosts and their gut microbiomes: Implications for animal physiology and ecology. *Functional Ecology*, 37(3), 476–487. <https://doi.org/10.1111/1365-2435.14255>
- Schneider, T., Schmid, E., De Castro, J. V., Cardinale, M., Eberl, L., Grube, M., Berg, G., & Riedel, K. (2011). Structure and function of the symbiosis partners of the lung lichen (*Lobaria pulmonaria* L. Hoffm.) analyzed by metaproteomics. *Proteomics*, 11(13), 2752–2756. <https://doi.org/10.1002/pmic.201000679>
- Sczyrba, A., Hofmann, P., Belmann, P., Koslicki, D., Janssen, S., Dröge, J., Gregor, I., Majda, S., Fiedler, J., Dahms, E., Bremges, A., Fritz, A., Garrido-Oter, R., Jørgensen, T. S., Shapiro, N., Blood, P. D., Gurevich, A., Bai, Y., Turavaev, D., ... McHardy, A. C. (2017). Critical assessment of metagenome interpretation—A benchmark of metagenomics software. *Nature Methods*, 14(11), 1063–1071. <https://doi.org/10.1038/nmeth.4458>
- Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation. *Bioinformatics*, 30(14), 2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>
- Shakya, M., Lo, C.-C., & Chain, P. S. G. (2019). Advances and challenges in metatranscriptomic analysis. *Frontiers in Genetics*, 10, 904. <https://doi.org/10.3389/fgene.2019.00904>
- Sommer, F., & Bäckhed, F. (2013). The gut microbiota—Masters of host development and physiology. *Nature Reviews Microbiology*, 11(4), 227–238. <https://doi.org/10.1038/nrmicro2974>
- Song, S. J., Amir, A., Metcalf, J. L., Amato, K. R., Xu, Z. Z., Humphrey, G., & Knight, R. (2016). Preservation methods differ in fecal microbiome stability, affecting suitability for field studies. *mSystems*, 1(3), e00021-16. <https://doi.org/10.1128/mSystems.00021-16>
- Song, S. J., Sanders, J. G., Delsuc, F., Metcalf, J., Amato, K., Taylor, M. W., Mazel, F., Lutz, H. L., Winker, K., Graves, G. R., Humphrey, G., Gilbert, J. A., Hackett, S. J., White, K. P., Skeen, H. R., Kurtis, S. M., Withrow, J., Braile, T., Miller, M., ... Knight, R. (2020). Comparative analyses of vertebrate gut microbiomes reveal convergence between birds and bats. *MBio*, 11(1), e02901-19. <https://doi.org/10.1128/mBio.02901-19>
- Stoffel, M. A., Acevedo-Whitehouse, K., Morales-Durán, N., Grosser, S., Chakarov, N., Krüger, O., Nichols, H. J., Elorriaga-Verplancken, F. R., & Hoffman, J. I. (2020). Early sexual dimorphism in the developing gut microbiome of northern elephant seals. *Molecular Ecology*, 29(11), 2109–2122. <https://doi.org/10.1111/mec.15385>
- Stothart, M. R., McLoughlin, P. D., & Poissant, J. (2022). Shallow shotgun sequencing of the microbiome recapitulates 16S amplicon results and provides functional insights. *Molecular Ecology*

- Resources, 23, 549–564. <https://doi.org/10.1111/1755-0998.13713>
- Sun, S., Jones, R. B., & Fodor, A. A. (2020). Inference-based accuracy of metagenome prediction tools varies across sample types and functional categories. *Microbiome*, 8(1), 46. <https://doi.org/10.1186/s40168-020-00815-y>
- Suzuki, T. A. (2017). Links between natural variation in the microbiome and host fitness in wild mammals. *Integrative and Comparative Biology*, 57(4), 756–769. <https://doi.org/10.1093/icb/ix104>
- Tamames, J., Cobo-Simón, M., & Puente-Sánchez, F. (2019). Assessing the performance of different approaches for functional and taxonomic annotation of metagenomes. *BMC Genomics*, 20(1), 960. <https://doi.org/10.1186/s12864-019-6289-6>
- Taxis, T. M., Wolff, S., Gregg, S. J., Minton, N. O., Zhang, C., Dai, J., Schnabel, R. D., Taylor, J. F., Kerley, M. S., Pires, J. C., Lamberson, W. R., & Conant, G. C. (2015). The players may change but the game remains: Network analyses of ruminal microbiomes suggest taxonomic differences mask functional similarity. *Nucleic Acids Research*, 43, 9600–9612. <https://doi.org/10.1093/nar/gkv973>
- Teullet, S., Tilak, M.-K., Magdeleine, A., Schaub, R., Weyer, N. M., Panaino, W., Fuller, A., Loughry, W. J., Avenant, N. L., De Thoisy, B., Borrel, G., & Delsuc, F. (2023). Metagenomics uncovers dietary adaptations for chitin digestion in the gut microbiota of convergent myrmecophilous mammals. *mSystems*, 8, e00388-23. <https://doi.org/10.1128/msystems.00388-23>
- Thomas, T., Moitinho-Silva, L., Lurgi, M., Björk, J. R., Easson, C., Astudillo-García, C., Olson, J. B., Erwin, P. M., López-Legentil, S., Luter, H., Chaves-Fonnegra, A., Costa, R., Schupp, P. J., Steindler, L., Erpenbeck, D., Gilbert, J., Knight, R., Ackermann, G., Victor Lopez, J., ... Webster, N. S. (2016). Diversity, structure and convergent evolution of the global sponge microbiome. *Nature Communications*, 7(1), 11870. <https://doi.org/10.1038/ncomms11870>
- Tung, J., Barreiro, L. B., Burns, M. B., Grenier, J.-C., Lynch, J., Grieneisen, L. E., Altmann, J., Alberts, S. C., Blekhman, R., & Archie, E. A. (2015). Social networks predict gut microbiome composition in wild baboons. *eLife*, 4, e05224. <https://doi.org/10.7554/eLife.05224>
- Turnbaugh, P. J., Ridaura, V. K., Faith, J. J., Rey, F. E., Knight, R., & Gordon, J. I. (2009). The effect of diet on the human gut microbiome: A metagenomic analysis in humanized gnotobiotic mice. *Science Translational Medicine*, 1(6), 6ra14. <https://doi.org/10.1126/scitranslmed.3000322>
- Uehara, M., Inoue, T., Kominato, M., Hase, S., Sasaki, E., Toyoda, A., & Sakakibara, Y. (2022). Intra-intestinal analysis of the functional activity of microbiomes and its application to the common marmoset intestine. *mSystems*, 7(5), e00520-22. <https://doi.org/10.1128/msystems.00520-22>
- Uren Webster, T. M., Consuegra, S., Hitchings, M., & Garcia de Leaniz, C. (2018). Interpopulation variation in the atlantic salmon microbiome reflects environmental and genetic diversity. *Applied and Environmental Microbiology*, 84(16), e00691-18. <https://doi.org/10.1128/AEM.00691-18>
- Varga, J. F. A., Bui-Marinos, M. P., & Katzenback, B. A. (2019). Frog skin innate immune defences: Sensing and surviving pathogens. *Frontiers in Immunology*, 9, 3128. <https://doi.org/10.3389/fimmu.2018.03128>
- Vargas-Pellicer, P., Watrobska, C., Knowles, S., Schroeder, J., & Banks-Leite, C. (2019). How should we store avian faecal samples for microbiota analyses? Comparing efficacy and cost-effectiveness. *Journal of Microbiological Methods*, 165, 105689. <https://doi.org/10.1016/j.mimet.2019.105689>
- Verschaffelt, P., Van Den Bossche, T., Martens, L., Dawyndt, P., & Mesuere, B. (2021). Unipept desktop: A faster, more powerful metaproteomics results analysis tool. *Journal of Proteome Research*, 20(4), 2005–2009. <https://doi.org/10.1021/acs.jproteome.0c00855>
- Vieira-Silva, S., Falony, G., Darzi, Y., Lima-Mendez, G., Garcia Yunta, R., Okuda, S., Vandeputte, D., Valles-Colomer, M., Hildebrand, F., Chaffron, S., & Raes, J. (2016). Species–function relationships shape ecological properties of the human gut microbiome. *Nature Microbiology*, 1(8), 16088. <https://doi.org/10.1038/nmicrobiol.2016.88>
- Vlíčková, K., Mrázek, J., Kopečný, J., & Petrželková, K. J. (2012). Evaluation of different storage methods to characterize the fecal bacterial communities of captive western lowland gorillas (*Gorilla gorilla gorilla*). *Journal of Microbiological Methods*, 91(1), 45–51. <https://doi.org/10.1016/j.mimet.2012.07.015>
- Wanelik, K. M., Raulo, A., Troitsky, T., Husby, A., & Knowles, S. C. L. (2023). Maternal transmission gives way to social transmission during gut microbiota assembly in wild mice. *Animal Microbiome*, 5(1), 29. <https://doi.org/10.1186/s42523-023-00247-7>
- Wang, C., Zhang, L., Jiang, X., Ma, W., Geng, H., Wang, X., & Li, M. (2022). Toward efficient and high-fidelity metagenomic data from sub-nanogram DNA: Evaluation of library preparation and decontamination methods. *BMC Biology*, 20(1), 225. <https://doi.org/10.1186/s12915-022-01418-9>
- Wang, Q., Ye, J., Fang, D., Lv, L., Wu, W., Shi, D., Li, Y., Yang, L., Bian, X., Wu, J., Jiang, X., Wang, K., Wang, Q., Hodson, M. P., Thibaut, L. M., Ho, J. W. K., Giannoulitou, E., & Li, L. (2020). Multi-omic profiling reveals associations between the gut mucosal microbiome, the metabolome, and host DNA methylation associated gene expression in patients with colorectal cancer. *BMC Microbiology*, 20(S1), 83. <https://doi.org/10.1186/s12866-020-01762-2>
- Wang, Z., Zolnik, C. P., Qiu, Y., Usyk, M., Wang, T., Strickler, H. D., Isasi, C. R., Kaplan, R. C., Kurland, I. J., Qi, Q., & Burk, R. D. (2018). Comparison of fecal collection methods for microbiome and metabolomics studies. *Frontiers in Cellular and Infection Microbiology*, 8, 301. <https://doi.org/10.3389/fcimb.2018.00301>
- Ward, D. V., Scholz, M., Zolfo, M., Taft, D. H., Schibler, K. R., Tett, A., Segata, N., & Morrow, A. L. (2016). Metagenomic sequencing with strain-level resolution implicates uropathogenic *E. coli* in necrotizing enterocolitis and mortality in preterm infants. *Cell Reports*, 14(12), 2912–2924. <https://doi.org/10.1016/j.celrep.2016.03.015>
- Warne, R. W., Kirschman, L., & Zeglin, L. (2019). Manipulation of gut microbiota during critical developmental windows affects host physiological performance and disease susceptibility across ontogeny. *Journal of Animal Ecology*, 88, 845–856. <https://doi.org/10.1111/1365-2656.12973>
- Wemheuer, F., Taylor, J. A., Daniel, R., Johnston, E., Meinicke, P., Thomas, T., & Wemheuer, B. (2020). Tax4Fun2: Prediction of habitat-specific functional profiles and functional redundancy based on 16S rRNA gene sequences. *Environmental Microbiomes*, 15(1), 11. <https://doi.org/10.1186/s40793-020-00358-7>
- Wilmes, P., & Bond, P. L. (2006). Metaproteomics: Studying functional gene expression in microbial ecosystems. *Trends in Microbiology*, 14(2), 92–97. <https://doi.org/10.1016/j.tim.2005.12.006>
- Wood, D. E., Lu, J., & Langmead, B. (2019). Improved metagenomic analysis with kraken 2. *Genome Biology*, 20(1), 257. <https://doi.org/10.1186/s13059-019-1891-0>
- Woodhams, D. C., McCartney, J., Walke, J. B., & Whetstone, R. (2023). The adaptive microbiome hypothesis and immune interactions in amphibian mucus. *Developmental & Comparative Immunology*, 145, 104690. <https://doi.org/10.1016/j.dci.2023.104690>
- Worsley, S. F., Davies, C. S., Mannarelli, M.-E., Hutchings, M. I., Komdeur, J., Burke, T., Dugdale, H. L., & Richardson, D. S. (2021). Gut microbiome composition, not alpha diversity, is associated with survival in a natural vertebrate population. *Animal Microbiome*, 3(1), 84. <https://doi.org/10.1186/s42523-021-00149-6>
- Worsley, S. F., Davies, C. S., Mannarelli, M.-E., Komdeur, J., Dugdale, H. L., & Richardson, D. S. (2022). Assessing the causes and consequences of gut mycobiome variation in a wild population of the Seychelles

- warbler. *Microbiome*, 10(1), 242. <https://doi.org/10.1186/s40168-022-01432-7>
- Xie, F., Jin, W., Si, H., Yuan, Y., Tao, Y., Liu, J., Wang, X., Yang, C., Li, Q., Yan, X., Lin, L., Jiang, Q., Zhang, L., Guo, C., Greening, C., Heller, R., Guan, L. L., Pope, P. B., Tan, Z., ... Mao, S. (2021). An integrated gene catalog and over 10,000 metagenome-assembled genomes from the gastrointestinal microbiome of ruminants. *Microbiome*, 9(1), 137. <https://doi.org/10.1186/s40168-021-01078-x>
- Yang, C., Chowdhury, D., Zhang, Z., Cheung, W. K., Lu, A., Bian, Z., & Zhang, L. (2021). A review of computational tools for generating metagenome-assembled genomes from metagenomic sequencing data. *Computational and Structural Biotechnology Journal*, 19, 6301–6314. <https://doi.org/10.1016/j.csbj.2021.11.028>
- Yasuda, S., Okahashi, N., Tsugawa, H., Ogata, Y., Ikeda, K., Suda, W., Arai, H., Hattori, M., & Arita, M. (2020). Elucidation of gut microbiota-associated lipids using LC-MS/MS and 16S rRNA sequence analyses. *iScience*, 23(12), 101841. <https://doi.org/10.1016/j.isci.2020.101841>
- Ye, S. H., Siddle, K. J., Park, D. J., & Sabeti, P. C. (2019). Benchmarking metagenomics tools for taxonomic classification. *Cell*, 178(4), 779–794. <https://doi.org/10.1016/j.cell.2019.07.010>
- Youngblut, N. D., De La Cuesta-Zuluaga, J., Reischer, G. H., Dauser, S., Schuster, N., Walzer, C., Stalder, G., Farnleitner, A. H., & Ley, R. E. (2020). Large-scale metagenome assembly reveals novel animal-associated microbial genomes, biosynthetic gene clusters, and other genetic diversity. *mSystems*, 5(6), e01045-20. <https://doi.org/10.1128/mSystems.01045-20>
- Youngblut, N. D., Reischer, G. H., Walters, W., Schuster, N., Walzer, C., Stalder, G., Ley, R. E., & Farnleitner, A. H. (2019). Host diet and evolutionary history explain different aspects of gut microbiome diversity among vertebrate clades. *Nature Communications*, 10(1), 2200. <https://doi.org/10.1038/s41467-019-10191-3>
- Zepeda Mendoza, M. L., Xiong, Z., Escalera-Zamudio, M., Runge, A. K., Thézé, J., Streicker, D., Frank, H. K., Loza-Rubio, E., Liu, S., Ryder, O. A., Samaniego Castruita, J. A., Katzourakis, A., Pacheco, G., Taboada, B., Löber, U., Pybus, O. G., Li, Y., Rojas-Anaya, E., Bohmann, K., ... Gilbert, M. P. T. (2018). Hologenomic adaptations underlying the evolution of sanguivory in the common vampire bat. *Nature Ecology & Evolution*, 2(4), 659–668. <https://doi.org/10.1038/s41559-018-0476-8>
- Zhang, M., Wang, X., Wang, Z., Mao, S., Zhang, J., Li, M., & Pan, H. (2023). Metatranscriptomic analyses reveal important roles of the gut microbiome in primate dietary adaptation. *Genes*, 14(1), 228. <https://doi.org/10.3390/genes14010228>
- Zhang, X., Li, L., Butcher, J., Stintzi, A., & Figeys, D. (2019). Advancing functional and translational microbiome research using meta-omics approaches. *Microbiome*, 7(1), 154. <https://doi.org/10.1186/s40168-019-0767-6>
- Zhang, Y., Thompson, K. N., Branck, T., Yan, Y., Nguyen, L. H., Franzosa, E. A., & Huttenhower, C. (2021). Metatranscriptomics for the human microbiome and microbial community functional profiling. *Annual Review of Biomedical Data Science*, 4(1), 279–311. <https://doi.org/10.1146/annurev-biodatasci-031121-103035>
- Zhou, D., Zhu, W., Sun, T., Wang, Y., Chi, Y., Chen, T., & Lin, J. (2021). iMAP: A web server for metabolomics data integrative analysis. *Frontiers in Chemistry*, 9, 659656. <https://doi.org/10.3389/fchem.2021.659656>
- Zhu, Q., Huang, S., Gonzalez, A., McGrath, I., McDonald, D., Haiminen, N., Armstrong, G., Vázquez-Baeza, Y., Yu, J., Kuczynski, J., Sepich-Poore, G. D., Swafford, A. D., Das, P., Shaffer, J. P., Lejzerowicz, F., Belda-Ferre, P., Havulinna, A. S., Méric, G., Niiranen, T., ... Knight, R. (2022). Phylogeny-aware analysis of metagenome community ecology based on matched reference genomes while bypassing taxonomy. *mSystems*, 7(2), e00167-22. <https://doi.org/10.1128/msystems.00167-22>

How to cite this article: Worsley, S. F., Videvall, E., Harrison, X. A., Björk, J. R., Mazel, F., & Wanelik, K. M. (2024). Probing the functional significance of wild animal microbiomes using omics data. *Functional Ecology*, 38, 2329–2349. <https://doi.org/10.1111/1365-2435.14650>