


Impact of gaming disorder on first episode psychosis patients' evolution: Protocol for a multicentered prospective study

Maxime Huot-Lavoie^{1,2,3}  | Charles Desmeules^{1,2,4} | Olivier Corbeil^{2,3,5} |
 Laurent Bécharé^{2,3,5} | Sébastien Brodeur^{1,2,3} | Anne-Marie Essiambre^{2,3,6} |
 Chantale Thériault^{2,3} | Elizabeth Anderson^{2,3,6} | Lauryann Bachand⁶ |
 Zain Al-Aabideen Haider¹ | Amal Abdel-Baki^{7,8} | Yasser Khazaal^{9,10} |
 Isabelle Giroux^{6,11} | Marie-France Demers^{2,3,5} | Marc-André Roy^{1,2,3} 

¹Faculty of Medicine, Université Laval, Quebec City, Quebec, Canada

²CERVO Brain Research Centre, Université Laval, Quebec City, Quebec, Canada

³Clinique Notre-Dame des Victoires, Institut universitaire en santé mentale de Québec, Centre intégré universitaire en santé et services sociaux de la capitale nationale, Montreal, Quebec, Canada

⁴Service de médecine des addictions, Lausanne University Hospital Research Center, Lausanne, Switzerland

⁵CHU de Québec-Université Laval Research Center, Université Laval, Quebec City, Quebec, Canada

⁶Faculty of Pharmacy, Université Laval, Quebec City, Quebec, Canada

⁷Centre Québécois d'Excellence pour la Prévention et le Traitement du Jeu, Université Laval, Quebec City, Quebec, Canada

⁸School of Psychology, Université Laval, Quebec City, Quebec, Canada

⁹Faculty of Medicine, Department of Psychiatry, Université de Montréal, Montreal, Quebec, Canada

¹⁰Clinique JAP, Montreal University Hospital Research Center, Montreal, Quebec, Canada

¹¹Department of Psychiatry, Lausanne University, Lausanne, Switzerland

Correspondence

Maxime Huot-Lavoie, Faculty of Medicine, Université Laval, 1050, Av. de la Médecine, Quebec City, QC G1V 0A6, Canada.
 Email: maxime.huot-lavoie.1@ulaval.ca

Funding information

Canadian Institutes of Health Research, Grant/Award Number: FID-172598; Mitacs, Grant/Award Number: IT34510

Abstract

Aims: The objective of this study is to underline the impact of Gaming Disorder on the clinical evolution of patients with First Episode Psychosis. The specific aims of the study are to determine the prevalence of gaming disorder among those patients and assess the consequences of gaming on their clinical trajectory.

Methods: This is a prospective multicenter cohort study that will enrol 800 patients diagnosed with a first episode psychosis, with a follow-up period of up to 3 years. Using a systematic screening procedure for gaming disorder, the clinical staff will assess patients gaming habits at admission and every 6 months thereafter. Information from patients' medical records will also be extracted using the same timeframe.

Results: The patients' characteristics at admission and during follow-up will be presented in the form of descriptive statistics and compared between different subgroups of patients using uni- and multivariate logistic regression models. Repeated

Abbreviations: CGI, clinical global impression scale; DSM-5, diagnostic and statistical manual of mental disorders, fifth edition; FEP, First-episode psychosis; GD, Gaming disorder; ICD-11, International classification of diseases, 11th edition; PANSS, positive and negative syndrome scale; REDCap, research electronic data capture; SOFAS, social and occupational functioning assessment scale.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Early Intervention in Psychiatry* published by John Wiley & Sons Australia, Ltd.

measures ANCOVA will also be performed to analyse the impact of gaming disorders on patients' clinical path as assessed by the Positive and Negative Syndrome Scale and the Clinical Global Impression scale, considering covariates such as psychiatric diagnosis, pharmacological treatment, age, sex/gender, and duration of untreated psychosis.

Conclusion: These findings will guide the development of prevention, detection, and treatment strategies for the comorbidity between gaming disorder and first episode psychosis, ultimately improving the patients' recovery.

KEYWORDS

early interventions, gaming disorder, psychotic disorder, schizophrenia, video games

1 | BACKGROUND

Psychotic disorders (PD), are major mental disorders that affect 3% of the population (Perälä et al., 2007). They occur during a critical period of life, specifically in late adolescence and early adulthood (Birchwood, 2014). Their consequences include an increased rate of suicide and acts of violence (Fu et al., 2021; Sicotte et al., 2021; Whiting et al., 2021), low employment rates (Plana-Ripoll et al., 2023), reduced life expectancy by 15 years (Laursen et al., 2014; Romain et al., 2021), as well as homelessness and experiences of stigmatization (Deschênes et al., 2021; Roy et al., 2022). In Canada, the economic burden of these conditions was estimated to be \$7 billion in 2004 while the global estimates indicate a range between 0.02% and 1.65% of the gross domestic product in 2016 (Chong et al., 2016; Goeree et al., 2005). While antipsychotic medications are effective in alleviating certain symptoms, such as delusions and hallucinations, they are not sufficient for achieving full recovery (Bécharde et al., 2021). In fact, PD are frequently associated with psychiatric comorbidities, including substance use disorders, personality disorders, and attention-deficit/hyperactivity disorder (ADHD), which can further exacerbate the consequences associated with PD (Corbeil et al., 2021). Therefore, treating comorbidities is crucial for achieving recovery, particularly in patients with first episode psychosis (FEP), where early intervention has demonstrated the greatest impact on patients' long-term outcomes (Correll et al., 2018). Among other comorbidities, excessive video gaming has become a cause for concerns for clinicians due to its impact on patients' lives, in particular among young adults, but a literature review conducted by our team (Huot-Lavoie et al., 2022) has highlighted a significant paucity of data on video gaming comorbid with FEP, despite recent interest and developments on this topic.

In recent decades, there has been a growing interest about the consequences of excessive video gaming on mental health. This attention has been prompted by the inclusion of Internet Gaming Disorder (IGD) as a 'Condition for Further Study' in the Diagnostic and Statistical Manual of Mental Disorders, fifth edition (DMS-5), in 2013 (American Psychiatric Association, 2013). The recognition of this disorder as an official diagnosis in the 11th edition of the International

Classification of Diseases (ICD-11) under the term Gaming Disorder (GD) has reinforced the importance and urgency to better understand the clinical impacts of this condition (WHO, 2019). GD is defined as a persistent and recurrent pattern of gaming behaviour, involving digital or video gaming, characterized by an impaired control over gaming as well as an increased priority placed on gaming over other activities to the extent that it takes precedence over other interests and daily activities. Individuals with GD continue or escalate their gaming behaviour despite experiencing negative consequences such as reduced occupational or academic functioning (WHO, 2019).

The reported prevalence of GD in the general population varies from 0.2% to 20% across studies (WHO, 2019). This variability can be explained by several factors, including the age of the participants studied, the country where the study was conducted, or the definition of GD used in the study. There are three main categories of risk factors associated with GD: (1) those related to gaming and its practice (e.g., duration and frequency of gaming, online and multiplayer games), (2) individual factors (e.g., isolation, low self-esteem, impulsivity, difficulties in emotion regulation, feelings of stigmatization, sex, gender, young age, low education level), and (3) environmental factors (e.g., family dysfunction, harassment and childhood neglect) (Ji et al., 2022; Richard et al., 2020). GD has also been associated with other symptoms, mental disorders, and impairments in functioning, such as depression, social anxiety, ADHD, impaired academic performance, and deterioration in interpersonal relationships, but a causal relationship has not been clearly established (Andreassen et al., 2016; Brunborg et al., 2014; González-Bueso et al., 2018; Ha et al., 2007; Kuss, 2013; Lee et al., 2021; Lo et al., 2005; Sussman et al., 2018; Weinstein & Weizman, 2012; Yu et al., 2020). In terms of treatment, promising results have been achieved with various approaches, including cognitive-behavioural therapy. However, there is currently insufficient data to draw solid conclusions regarding the effectiveness of existing approaches (Gorowska et al., 2022).

Regarding the comorbidity between GD and PD, the available literature is limited to a few case reports describing the occurrence of brief psychotic episodes following excessive video game use or abrupt cessation (Huot-Lavoie et al., 2022). Research on this comorbidity is therefore scarce and lacks prospective data, which hinders the

improvement of prevention, detection, and treatment strategies. However, individuals with PD present several common risk factors with GD: the male gender, social isolation, low self-esteem, difficulties in emotion regulation, impulsivity or behavioural inhibition, and onset of disorders during adolescence or early adulthood (Arango et al., 2021; Ji et al., 2022; Kahn, 2020; Richard et al., 2020; van Os et al., 2010). Additionally, little is known about the patterns of video game use among individuals with PD (e.g., gaming time, game content, interface [console, mobile, computer, online or offline]). For the affected youth, the consequences of GD added to those of PD may further hinder recovery. Studying GD among young adults with PD is then particularly relevant, as the risk factors common to GD and PD are even more prevalent in this population compared to the overall population with PD alone (e.g., substance use disorders, male predominance, younger age) (Hartz et al., 2014; Khokhar et al., 2018). Furthermore, it is well established that clinical interventions in FEP significantly influence clinical outcomes of affected individuals (Correll et al., 2018).

In line with prevalent societal beliefs, video games have often been associated with a negative perception that links them to aggressive and criminal behaviour (Calvert et al., 2017). However, an increasing body of evidence suggests that video games can have beneficial effects and even serve as a therapeutic tool. The term 'serious games' refers to software that incorporates playful elements for educational, training, and therapeutic purposes. Such games have been employed in the treatment of individuals with PD, demonstrating a significant reduction in psychotic symptoms (Leff et al., 2013; Moritz et al., 2014). Additionally, they have shown improvements in cognitive functions, social cognitions, and occupational status (Amado et al., 2016; Chan et al., 2010; Rus-Calafell et al., 2014; Wiederhold & Riva, 2013). Although the understanding of the impact of commercial video games on PD is limited, a recent literature review highlights several advantages of these types of games, with seven studies reporting positive outcomes (Quiles & Verdoux, 2023). Among them, four studies revealed enhanced processing speed, memory, and executive functions through the use of commercial video games, while three studies demonstrated improvements in aerobic fitness and walking speed in patients who engaged in active video games utilizing the Kinect for Xbox game system (Bang-Kittilsen et al., 2021; Roberts et al., 2021; Shimizu et al., 2017; Välimäki et al., 2021). These potential benefits in individuals with PD underscore the unique nature of gaming addiction, where the objective is to promote healthy gaming habits rather than complete avoidance, as seen in other forms of addiction. In order to help patients to reach these healthy habits, it is imperative to enhance our comprehension of the influence of gaming on the clinical trajectory of individuals diagnosed with PD.

Using an innovative screening and assessment procedure for GD tailored to young adults with FEP, the aim of this study is to better understand the impact of gaming in people with FEP by: (1) determining the prevalence of GD among FEP patients; (2) determining the consequences of GD on the clinical trajectory of FEP patients; and (3) assessing individual factors that differentiate patients who developed GD among the overall study population.

2 | METHODS

2.1 | Study design

This is a prospective multicenter cohort study, akin to our previous work (Corbeil et al., 2023), aiming to enrol a total of 800 patients who have been diagnosed with FEP and with follow-up periods of up to 3 years. The recruitment process began on 1 November 2019, and will extend until 1 November 2023. Throughout this period, all individuals admitted at the two designated sites will undergo comprehensive GD assessments by clinicians upon admission and subsequently every 6 months, using a specialized screening procedure developed exclusively for this study. During this follow-up period, that will conclude on 1 May 2024, the research team will extract independent variables from the patients' medical records.

2.2 | Study settings

This study is being conducted within 2 specialized programs for individuals experiencing FEP in the province of Quebec, Canada. These programs, which operate at multidisciplinary clinics, admit annually approximately 200 patients presenting a FEP. The patients are accompanied for up to 3 years on a case management approach that involves close collaboration between a certified psychiatrist, both centres having a combined total of 11 psychiatrists, and a designated case manager, who assumes the responsibility of overseeing and coordinating the patient's care and services, and could include either psychologist, social works, occupational therapists, nurses and specialized educators. In this framework, an individual case manager typically oversees the care of 16 patients, as suggested in the guidelines for implementing community mental health support measures (MSSS, 2002).

Patients are also offered family intervention, individual psychotherapy and community outreach services during their follow-up period. The case managers offer an extensive systematic clinical follow-up, employing standardized questionnaires and involving input from the patients' family members. All relevant clinical data derived from these activities are extensively documented in the patients' medical records. No consent to participate in the study is required given the lack of any contact between the patients and the research staff.

2.3 | Participants

During the recruitment period, all patients admitted at the two study sites will be included in the study. Clinic admission prerequisites necessitate individuals to be within the age range of 18 to 35 years, hold a primary diagnosis of FEP according to DSM-5 criteria (encompassing both affective and nonaffective psychoses, along with substance-induced psychotic disorder), and exhibit limited exposure to continuous antipsychotic treatment (i.e., 6 months or less). The only exclusions criteria are related to admittance criteria in the clinics,

specifically, organic psychosis and moderate/severe intellectual disability as defined by the DSM-5.

2.4 | Variables

The primary outcome is the prevalence of a GD diagnosis based on ICD-11 criteria established by the treating psychiatrist. As mentioned earlier, the evaluation of this outcome is conducted by the case managers and treating psychiatrists utilizing a specialized, GD screening procedure designed explicitly for this study in collaboration with experts in GD and FEP. It is a 3-step procedure that includes: (1) a questionnaire for assessing video gaming habits (frequency and duration of gaming session) (supplementary material 1), (2) an evaluation of the severity of the gaming habits using the *Internet Gaming Disorder Scale - Short Form (IGDS9-SF)* (supplementary material 1) (Poon et al., 2021), and (3) a diagnostic interview assessing the ICD-11 gaming addiction diagnosis criteria. The patients' case managers conduct the initial two steps, while the treating psychiatrist oversees the final step. All participating professionals received training on gaming disorders, encompassing the systematic screening procedure and the diagnostic criteria from the ICD-11. Additionally, instructional videos have been made available as a reference for all clinicians.

The results of this procedure are documented in the patients' medical records and subsequently extracted by the research staff. Variables that are related to the clinical manifestations of the FEP are documented using three standardized scales rated by treating psychiatrists every 6 months as part of the patients' systematic follow-up. These variables are recorded in the patients' medical files and extracted by the research staff. The severity of psychotic symptoms is measured using the 'Positive and Negative Syndrome Scale' (PANSS), which is a routinely used measure that allows clinicians to monitor the progression of symptoms in patients with psychotic disorders (Kay et al., 1987). The PANSS-6, a shortened version that has excellent convergent validity with the original PANSS, will be used in this study (Østergaard et al., 2016). The severity of the FEP will also be assessed using the 'Clinical Global Impression - Severity' (CGI-S) scale, which quantifies the clinical presentation of the disease using simple score ranging between 0 and 7 (Busner & Targum, 2007; Haro et al., 2003). CGI-S scores have good correlation with PANSS scores (Samara et al., 2014). Finally, the individual's functioning level will be evaluated using the 'Social and Occupational Functioning Assessment Scale' (SOFAS), a graded 0–100 scale quantifying patients' social and occupational functioning (Rybarczyk, 2011). All psychiatrists underwent prior training on these scales (PANSS, CGI-S, SOFAS) since these assessments are integral to the systematic follow-up within the participating clinics. All evaluation being systematically part of the clinical follow-ups, the evaluators are blind to the analysis performed by the research staff.

The research staff will extract various independent variables from the patients' medical records, encompassing socio-demographic factors (such as sex at birth, gender identity, ethnicity, employment status, education level, relationship status, living arrangements, and

criminal history) and other clinical variables (including the primary DSM-5 psychiatric diagnosis, comorbid DSM-5 psychiatric diagnoses, psychiatric hospitalizations, community treatment order, and medication treatment, which comprises antipsychotics and other psychotropic drugs like antidepressants, benzodiazepines, hypnotics, mood stabilizers, and psychostimulants). Regarding medication information, both current (during the follow-up period) and past (prior to admission in the clinics) medication details will be collected, encompassing dosages, frequency, and route of administration. A patient will be considered exposed to a medication after receiving it for more than 1 month (or one injection in the case of long-acting injectable antipsychotics). Medications administered on an as-needed basis will not be included in the analysis.

3 | STATISTICAL ANALYSES

Statistical analysis will be conducted in partnership with expert statisticians from the CERVO Research Centre, Quebec, Canada, to guarantee rigorous data interpretation and maximize the validity of our findings. To meet the first two objectives, the patients' characteristics at admission and during follow-up will be presented in the form of descriptive statistics and compared between different subgroups of patients using uni- and multivariate logistic regression models. All tests will be performed with a bilateral significance level set at 5%.

To meet the third objective, repeated measures ANCOVA will be performed. In this analysis, the independent variable will be the presence or absence of GD as defined by ICD-11 criteria, while the dependent variables will be the clinical ratings (PANSS-6, CGI-S, SOFAS) measured at admission and repeated every 6 months during the patients' follow-up. The covariates of the analysis will be: (1) the main psychiatric diagnosis; (2) the pharmacological treatment used; (3) the age and sex/gender of the patients; (4) the duration of untreated psychosis, defined as the interval between onset of first psychotic symptoms and admission to treatment centre; (5) impact of previous diagnoses and current comorbidities; these factors being determinant in the clinical evolution of the patients living with FEP. The results from these analyses will be performed with a bilateral significance level set at 5%.

4 | DISCUSSION

This aims to improve our understanding of the relationship between GD and PD. It will be the first empirical study that explores this comorbidity, which has only been previously discussed in a few case reports. The results will not only confirm or refute the association between GD and PD, but also provide a more detailed description of the factors linking these two conditions and their consequences.

A notable strength of this study is the systematic screening procedure of GD that has been developed, which eliminates any detection bias and improves sensibility. All patients newly admitted to

either of the two participating FEP programs will undergo the same screening process, irrespective of their disease progression and treatment history. This systematic approach is particularly crucial given the limited knowledge about the consequences of GD, which may be underestimated by both patients and clinicians. On completion of the study, the screening procedure for is expected to be implemented across all FEP programs in Quebec, and potentially in other regions in Canada as well.

The study's large sample size and extended follow-up duration are also significant strengths, ensuring sufficient statistical power to generate meaningful findings. The lack of a consent requirement, combined with the diverse areas served by the two participating FEP programs, including urban and rural areas, ensures that the sample is representative of the FEP population in Quebec. Consequently, the findings are expected to be generalizable to all FEP programs in the province.

By collecting variables that are routinely documented by clinicians as part of their systematic clinical follow-up, the ecological validity of this study is further strengthened. The close collaboration between our research team and clinicians facilitates knowledge transfer and improves the quality of patient care.

The primary limitation of this research project lies in its dependence on clinician cooperation in adhering to the GD screening procedure, as we have no control over their clinical practices and prescribing patterns. However, we have implemented a monitoring committee to assess clinicians' adherence to the GD screening process on a regular basis. Within each participating center, designated research professionals rigorously track patient screenings, ensuring biannual procedures by case managers and psychiatrists; lapses prompt immediate reminders to uphold consistent clinical evaluations. Furthermore, all new clinicians joining the study receive comprehensive training in the utilization of this procedure. By considering these methodological aspects collectively, we anticipate that the results will substantially enhance our understanding of this understudied comorbidity.

ACKNOWLEDGEMENTS

The authors are thankful to the clinical staff working at the 2 study sites for their precious collaboration and their unwavering dedication to patients' recovery.

FUNDING INFORMATION

This project is supported by a Mitacs Grant (IT34510). MHL is also supported by the Frederick Banting and Charles Best Canada Graduate Scholarship Doctoral Awards from the Canadian Institute of Health Research (FID-172598). The funders played no role in this study, and we have no restrictions regarding the submission of this protocol for publication.

CONFLICT OF INTEREST STATEMENT

OC has received an honorarium from Janssen and Otsuka-Lundbeck Alliance. MFD has received grants from Janssen and Otsuka-Lundbeck Alliance. MAR has received grants from Otsuka-Lundbeck

Alliance, Janssen, Sunovion and Mylan. All other authors have no conflicts of interest that are directly relevant to this study.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

ORCID

Maxime Huot-Lavoie  <https://orcid.org/0000-0001-7378-9776>

Marc-André Roy  <https://orcid.org/0000-0002-0123-6651>

REFERENCES

- Amado, I., Brénugat-Herné, L., Orriols, E., Desombre, C., Santos, M. D., Prost, Z., Krebs, M.-O., & Piolino, P. (2016). A serious game to improve cognitive functions in schizophrenia: A pilot study. *Frontiers in Psychiatry, 7*, 64.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders: DSM-5* (5th ed.). American Psychiatric Association.
- Andreassen, C. S., Billieux, J., Griffiths, M. D., Kuss, D. J., Demetrovics, Z., Mazzoni, E., & Pallesen, S. (2016). The relationship between addictive use of social media and video games and symptoms of psychiatric disorders: A large-scale cross-sectional study. *Psychology of Addictive Behaviors, 30*(2), 252–262.
- Arango, C., Dragioti, E., Solmi, M., Cortese, S., Domschke, K., Murray, R. M., Jones, P. B., Uher, R., Carvalho, A. F., Reichenberg, A., Shin, J. I. I., Andreassen, O. A., Correll, C. U., & Fusar-Poli, P. (2021). Risk and protective factors for mental disorders beyond genetics: An evidence-based atlas. *World Psychiatry, 20*(3), 417–436.
- Bang-Kittilsen, G., Egeland, J., Holmen, T. L., Bigseth, T. T., Andersen, E., Mordal, J., Ulleberg, P., & Engh, J. A. (2021). High-intensity interval training and active video gaming improve neurocognition in schizophrenia: A randomized controlled trial. *European Archives of Psychiatry and Clinical Neuroscience, 271*(2), 339–353.
- Béchar, L., Corbeil, O., Malenfant, E., Lehoux, C., Stip, E., Roy, M. A., & Demers, M. F. (2021). Psychopharmacology of first episode psychosis: An approach based on recovery. *Sante Mentale au Quebec, 46*(2), 113–137.
- Birchwood, M. (2014). Early intervention in psychosis services: The next generation. *Early Intervention in Psychiatry, 8*(1), 1–2.
- Brunborg, G. S., Mentzoni, R. A., & Frøyland, L. R. (2014). Is video gaming, or video game addiction, associated with depression, academic achievement, heavy episodic drinking, or conduct problems? *Journal of Behavioral Addictions, 3*(1), 27–32.
- Busner, J., & Targum, S. D. (2007). The clinical global impressions scale: Applying a research tool in clinical practice. *Psychiatry (Edgmont), 4*(7), 28.
- Calvert, S. L., Appelbaum, M., Dodge, K. A., Graham, S., Nagayama Hall, G. C., Hamby, S., Fasig-Caldwell, L. G., Citkowitz, M., Galloway, D. P., & Hedges, L. V. (2017). The American Psychological Association task force assessment of violent video games: Science in the service of public interest. *American Psychologist, 72*, 126–143.
- Chan, C. L., Ngai, E. K. Y., Leung, P. K. H., & Wong, S. (2010). Effect of the adapted virtual reality cognitive training program among Chinese older adults with chronic schizophrenia: A pilot study. *International Journal of Geriatric Psychiatry, 25*(6), 643–649.
- Chong, H. Y., Teoh, S. L., Wu, D. B., Kotirum, S., Chiou, C. F., & Chaikunapruk, N. (2016). Global economic burden of schizophrenia: A systematic review. *Neuropsychiatric Disease and Treatment, 12*, 357–373.
- Corbeil, O., Bérubé, F. A., Artaud, L., & Roy, M. A. (2021). Detecting and treating comorbid disorders in first-episode psychosis: A lever for recovery. *Sante Mentale au Quebec, 46*(2), 307–330.

- Corbeil, O., Soulard, M., Huot-Lavoie, M., Béchar, L., Fournier, É., Brodeur, S., Essiambre, A. M., Desmeules, C., Thériault, C., Abdel-Baki, A., Jacques, C., Giroux, I., Dorval, M., Roy, M. A., & Demers, M. F. (2023). Problem gambling among people with first-episode psychosis: Protocol for a prospective multicenter cohort study. *BMC Psychiatry*, 23(1), 287.
- Correll, C. U., Galling, B., Pawar, A., Krivko, A., Bonetto, C., Ruggeri, M., Craig, T. J., Nordentoft, M., Srihari, V. H., Guloksuz, S., Hui, C. L. M., Chen, E. Y. H., Valencia, M., Juarez, F., Robinson, D. G., Schooler, N. R., Brunette, M. F., Mueser, K. T., Rosenheck, R. A., et al. (2018). Comparison of early intervention services vs treatment as usual for early-phase psychosis: A systematic review, meta-analysis, and meta-regression. *JAMA Psychiatry*, 75(6), 555–565.
- Deschênes, J. M., Roy, L., Girard, N., & Abdel-Baki, A. (2021). How to help young people with early psychosis avoid homelessness? *Sante Mentale au Quebec*, 46(2), 189–216.
- Fu, X.-L., Qian, Y., Jin, X. H., Yu, H. R., Wu, H., du, L., Chen, H. L., & Shi, Y. Q. (2021). Suicide rates among people with serious mental illness: A systematic review and meta-analysis. *Psychological Medicine*, 53, 1–11.
- Goeree, R., Farahati, F., Burke, N., Blackhouse, G., O'Reilly, D., Pyne, J., & Tarride, J. E. (2005). The economic burden of schizophrenia in Canada in 2004. *Current Medical Research and Opinion*, 21(12), 2017–2028.
- González-Bueso, V., Santamaría, J., Fernández, D., Merino, L., Montero, E., & Ribas, J. (2018). Association between internet gaming disorder or pathological video-game use and comorbid psychopathology: A comprehensive review. *International Journal of Environmental Research and Public Health*, 15(4), 668.
- Gorowska, M., Tokarska, K., Zhou, X. Y., Gola, M. K., & Li, Y. (2022). Novel approaches for treating internet gaming disorder: A review of technology-based interventions. *Comprehensive Psychiatry*, 115, 152312.
- Ha, J. H., Kim, S. Y., Bae, S. C., Bae, S., Kim, H., Sim, M., Lyoo, I. K., & Cho, S. C. (2007). Depression and internet addiction in adolescents. *Psychopathology*, 40(6), 424–430.
- Haro, J. M., Kamath, S. A., Ochoa, S., Novick, D., Rele, K., Fargas, A., Rodríguez, M. J., Rele, R., Orta, J., Kharbeng, A., Araya, S., Gervin, M., Alonso, J., Mavreas, V., Lavrentzou, E., Lontos, N., Gregor, K., Jones, P. B., & SOHO Study Group. (2003). The clinical global impression-schizophrenia scale: A simple instrument to measure the diversity of symptoms present in schizophrenia. *Acta Psychiatrica Scandinavica*, 107(s416), 16–23.
- Hartz, S. M., Pato, C. N., Medeiros, H., Cavazos-Rehg, P., Sobell, J. L., Knowles, J. A., Bierut, L. J., Pato, M. T., & Genomic Psychiatry Cohort Consortium. (2014). Comorbidity of severe psychotic disorders with measures of substance use. *JAMA Psychiatry*, 71(3), 248–254.
- Huot-Lavoie, M., Gabriel-Courval, M., Béchar, L., Corbeil, O., Brodeur, S., Massé, C., Fournier, É., Essiambre, A. M., Anderson, E., Cayouette, A., Giroux, I., Khazaal, Y., Demers, M.-F., & Roy, M.-A. (2022). Gaming disorder and psychotic disorders: A scoping review. *Psychopathology*, 56, 315–323.
- Ji, Y., Yin, M. X. C., Zhang, A. Y., & Wong, D. F. K. (2022). Risk and protective factors of internet gaming disorder among Chinese people: A meta-analysis. *The Australian and New Zealand Journal of Psychiatry*, 56(4), 332–346.
- Kahn, R. S. (2020). On the origins of schizophrenia. *The American Journal of Psychiatry*, 177(4), 291–297.
- Kay, S. R., Fiszbein, A., & Opler, L. A. (1987). The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin*, 13(2), 261–276.
- Khokhar, J. Y., Dwi, L. L., Henricks, A. M., Doucette, W. T., & Green, A. I. (2018). The link between schizophrenia and substance use disorder: A unifying hypothesis. *Schizophrenia Research*, 194, 78–85.
- Kuss, D. J. (2013). Internet gaming addiction: Current perspectives. *Psychology Research and Behavior Management*, 6, 125–137.
- Laursen, T. M., Nordentoft, M., & Mortensen, P. B. (2014). Excess early mortality in schizophrenia. *Annual Review of Clinical Psychology*, 10, 425–448.
- Lee, J., Bae, S., Kim, B. N., & Han, D. H. (2021). Impact of attention-deficit/hyperactivity disorder comorbidity on longitudinal course in internet gaming disorder: A 3-year clinical cohort study. *Journal of Child Psychology and Psychiatry*, 62(9), 1110–1119.
- Leff, J., Williams, G., Huckvale, M. A., Arbuthnot, M., & Leff, A. P. (2013). Computer-assisted therapy for medication-resistant auditory hallucinations: Proof-of-concept study. *The British Journal of Psychiatry*, 202(6), 428–433.
- Lo, S.-K., Wang, C.-C., & Fang, W. (2005). Physical interpersonal relationships and social anxiety among online game players. *Cyberpsychology & Behavior*, 8(1), 15–20.
- Moritz, S., Voigt, M., Köther, U., Leighton, L., Kjahili, B., Babur, Z., Jungclaussen, D., Veckenstedt, R., & Grzella, K. (2014). Can virtual reality reduce reality distortion? Impact of performance feedback on symptom change in schizophrenia patients. *Journal of Behavior Therapy and Experimental Psychiatry*, 45(2), 267–271.
- MSSS. (2002). *Lignes directrices pour l'implantation de mesures de soutien dans la communauté en santé mentale*. Gouvernement du Québec.
- Østergaard, S. D., Lemming, O. M., Mors, O., Correll, C. U., & Bech, P. (2016). PANSS-6: A brief rating scale for the measurement of severity in schizophrenia. *Acta Psychiatrica Scandinavica*, 133(6), 436–444.
- Perälä, J., Suvisaari, J., Saarni, S. I., Kuoppasalmi, K., Isometsä, E., Pirkola, S., Partonen, T., Tuulio-Henriksson, A., Hintikka, J., Kieseppä, T., Härkönen, T., Koskinen, S., & Lönnqvist, J. (2007). Lifetime prevalence of psychotic and bipolar I disorders in a general population. *Archives of General Psychiatry*, 64(1), 19–28.
- Plana-Ripoll, O., Weyer, N., Knudsen, A. K., Hakulinen, C., Madsen, K. B., Christensen, M. K., Agerbo, E., Laursen, T. M., Nordentoft, M., Timmermann, A., Whiteford, H., Øverland, S., Iburg, K. M., & McGrath, J. J. (2023). The association between mental disorders and subsequent years of working life: A Danish population-based cohort study. *The Lancet Psychiatry*, 10(1), 30–39.
- Poon, L. Y., Tsang, H. W. H., Chan, T. Y. J., Man, S. W. T., Ng, L. Y., Wong, Y. L. E., Lin, C.-Y., Chien, C.-W., Griffiths, M. D., Pontes, H. M., & Pakpour, A. H. (2021). Psychometric properties of the internet gaming disorder scale-short-form (IGDS9-SF): Systematic review. *Journal of Medical Internet Research*, 23(10), e26821.
- Quiles, C., & Verdoux, H. (2023). Benefits of video games for people with schizophrenia: A literature review. *Current Opinion in Psychiatry*, 36(3), 184–193.
- Richard, J., Temcheff, C. E., & Derevensky, J. L. (2020). Gaming disorder across the lifespan: A scoping review of longitudinal studies. *Current Addiction Reports*, 7(4), 561–587.
- Roberts, M. T., Lloyd, J., Välimäki, M., Ho, G. W., Freemantle, M., & Békefi, A. Z. (2021). Video games for people with schizophrenia. *Cochrane Database of Systematic Reviews*, 2, CD012844.
- Romain, A. J., Bernard, P., Piché, F., Kern, L., Ouellet-Plamondon, C., Abdel-Baki, A., & Roy, M. A. (2021). A healthy mind in a healthy body: The value of physical activity for youth with first episode psychosis. *Sante Mentale au Quebec*, 46(2), 249–276.
- Roy, L., Abdel-Baki, A., Bérubé, F. A., Crocker, A., de Benedictis, L., Dostie, M., Latimer, E., & Roy, M. A. (2022). Housing trajectories and the risk of homelessness among new mental health service users: Protocol for the AMONT study. *Early Intervention in Psychiatry*, 17, 495–501.
- Rus-Calafell, M., Gutiérrez-Maldonado, J., & Ribas-Sabaté, J. (2014). A virtual reality-integrated program for improving social skills in patients with schizophrenia: A pilot study. *Journal of Behavior Therapy and Experimental Psychiatry*, 45(1), 81–89.
- Rybarczyk, B. (2011). Social and occupational functioning assessment scale (SOFAS). In J. S. Kreutzer, J. DeLuca, & B. Caplan (Eds.), *Encyclopedia of clinical neuropsychology* (p. 2313). Springer.

- Samara, M. T., Engel, R. R., Millier, A., Kandenwein, J., Toumi, M., & Leucht, S. (2014). Equipercentile linking of scales measuring functioning and symptoms: Examining the GAF, SOFAS, CGI-S, and PANSS. *European Neuropsychopharmacology*, *24*(11), 1767–1772.
- Shimizu, N., Umemura, T., Matsunaga, M., & Hirai, T. (2017). An interactive sports video game as an intervention for rehabilitation of community-living patients with schizophrenia: A controlled, single-blind, crossover study. *PLoS One*, *12*(11), e0187480.
- Sicotte, R., Iyer, S. N., Kiepura, B., & Abdel-Baki, A. (2021). A systematic review of longitudinal studies of suicidal thoughts and behaviors in first-episode psychosis: Course and associated factors. *Social Psychiatry and Psychiatric Epidemiology*, *56*(12), 2117–2154.
- Sussman, C. J., Harper, J. M., Stahl, J. L., & Weigle, P. (2018). Internet and video game addictions: Diagnosis, epidemiology, and neurobiology. *Child and Adolescent Psychiatric Clinics of North America*, *27*(2), 307–326.
- Välimäki, M., Yang, M., Lam, Y. T. J., Lantta, T., Palva, M., Palva, S., Yee, B., Yip, S. H., Yu, K. S. D., Chang, H. C. C., Cheng, P. Y. I., & Bressington, D. (2021). The impact of video gaming on cognitive functioning of people with schizophrenia (GAME-S): Study protocol of a randomised controlled trial. *BMC Psychiatry*, *21*(1), 1–12.
- van Os, J., Kenis, G., & Rutten, B. P. F. (2010). The environment and schizophrenia. *Nature*, *468*(7321), 203–212.
- Weinstein, A., & Weizman, A. (2012). Emerging association between addictive gaming and attention-deficit/hyperactivity disorder. *Current Psychiatry Reports*, *14*, 590–597.
- Whiting, D., Lichtenstein, P., & Fazel, S. (2021). Violence and mental disorders: A structured review of associations by individual diagnoses, risk factors, and risk assessment. *The Lancet Psychiatry*, *8*(2), 150–161.
- WHO. (2019). *International statistical classification of diseases and related health problems*. WHO.
- Wiederhold, B., & Riva, G. (2013). Cognitive rehabilitation of schizophrenia through NeuroVr training. *Annual Review of Cybertherapy and Telemedicine*, *2013*, 158.
- Yu, Y., Yang, X., Wang, S., Wang, H., Chang, R., Tsamag, L., Zhang, S., Xu, C., Yu, X., Cai, Y., & Lau, J. T. F. (2020). Serial multiple mediation of the association between internet gaming disorder and suicidal ideation by insomnia and depression in adolescents in Shanghai, China. *BMC Psychiatry*, *20*(1), 1–9.

How to cite this article: Huot-Lavoie, M., Desmeules, C., Corbeil, O., Béchar, L., Brodeur, S., Essiambre, A.-M., Thériault, C., Anderson, E., Bachand, L., Haider, Z. A.-A., Abdel-Baki, A., Khazaal, Y., Giroux, I., Demers, M.-F., & Roy, M.-A. (2023). Impact of gaming disorder on first episode psychosis patients' evolution: Protocol for a multicentered prospective study. *Early Intervention in Psychiatry*, 1–7. <https://doi.org/10.1111/eip.13482>