



Scientific Evidence for the Updated Guidelines on Indications for Metabolic and Bariatric Surgery (IFSO/ASMBS)

Maurizio De Luca¹ · Scott Shikora² · Dan Eisenberg³ · Luigi Angrisani⁴ · Chetan Parmar⁵ · Aayed Alqahtani⁶ · Ali Aminian⁷ · Edo Aarts⁸ · Wendy Brown⁹ · Ricardo V. Cohen¹⁰ · Nicola Di Lorenzo¹¹ · Silvia L. Faria¹² · Kasey P. S. Goodpaster¹³ · Ashraf Haddad¹⁴ · Miguel Herrera¹⁵ · Raul Rosenthal¹⁶ · Jacques Himpens¹⁷ · Angelo Iossa¹⁸ · Mohammad Kermansaravi¹⁹ · Lilian Kow²⁰ · Marina Kurian²¹ · Sonja Chiappetta²² · Teresa LaMasters²³ · Kamal Mahawar²⁴ · Giovanni Merola²⁵ · Abdelrahman Nimeri² · Mary O’Kane²⁶ · Pavlos Papasavas²⁷ · Giacomo Piatto²⁸ · Jaime Ponce²⁹ · Gerhard Prager³⁰ · Janey S. A. Pratt³ · Ann M. Rogers³¹ · Paulina Salminen³² · Kimberley E. Steele³³ · Michel Suter³⁴ · Salvatore Tolone³⁵ · Antonio Vitiello³⁶ · Marco Zappa³⁷ · Shanu N. Kothari³⁸

Received: 14 May 2024 / Accepted: 21 May 2024 / Published online: 25 September 2024

© 2024 The Authors. Published by Elsevier Inc on behalf of American Society for Metabolic & Bariatric Surgery (ASMBS) and Springer Nature on behalf of International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO). This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>) 2024

Abstract

The 2022 American Society of Metabolic and Bariatric Surgery (ASMBS) and International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO) updated the indications for Metabolic and Bariatric Surgery (MBS), replacing the previous guidelines established by the NIH over 30 years ago. The evidence supporting these updated guidelines has been strengthened to assist metabolic and bariatric surgeons, nutritionists, and other members of multidisciplinary teams, as well as patients. This study aims to assess the level of evidence and the strength of recommendations compared to the previously published criteria.

Keywords Obesity · Metabolic and bariatric surgery · IFSO · ASMBS · Guidelines · Indications

Abbreviations

AAHKS	American Association for Hip and Knee Surgeons
ACS-NSQIP	American College of Surgeons National Surgical Quality Improvement Program
AGB	Adjustable gastric banding
ASMBS	American Society for Metabolic and Bariatric Surgery
BMI	Body mass index
BPD	Bilio-pancreatic diversion
EAES	European Association for Endoscopic Surgery
EASO	European Association for the Study of Obesity
EBMIL	Excess of BMI loss
EWL	Excess weight loss

GI	Gastrointestinal
GRADE	Grading of Recommendations, Assessment, Development and Evaluations
HF	Heart failure
HTN	Hypertension
IFSO	International Federation for the Surgery of Obesity and Metabolic Disorders
LOS	Length of stay
LVAD	Left ventricular assist device
LVEF	Left ventricular ejection fraction
MACE	Major adverse cardiovascular event
MAFDL	Metabolic dysfunction–associated liver disease
MBS	Metabolic bariatric surgery
MBSAQIP	Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program
MDT	Multidisciplinary team
NIH	National Institute of Health
OAGB	One anastomosis gastric bypass
OSA	Obstructive sleep apnea
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses

This paper was jointly developed by the American Society for Metabolic and Bariatric Surgery and International Federation for the Surgery of Obesity and Metabolic Disorders and jointly published in *Surgery for Obesity and Related Diseases* and *Obesity Surgery*. The articles are identical except for minor stylistic and spelling differences in keeping with each journal's style. Either citation can be used when citing this article.

Extended author information available on the last page of the article

PWS	Prader Willi syndrome
RCT	Randomized controlled trial
RWG	Recurrent weight gain
RYGB	Roux en Y gastric bypass
SADI-S	Single anastomosis duodeno-ileal bypass with sleeve gastrectomy
SBO	Small bowel obstruction
SG	Sleeve gastrectomy
SOT	Solid organ transplantation
T2DM	Type 2 diabetes mellitus
TKA	Total knee arthroplasty
TWL	Total weight loss
VHR	Ventral hernia repair
WHO	World Health Organization

Introduction

Since its inception in the mid-1950s, gastrointestinal surgery to treat excess adiposity and associated medical conditions has significantly changed [1]. Weight loss surgery, in its early history, lacked regulation, was associated with a high risk of adverse events, and had a high rate of recurrent weight gain (RWG). There were no uniform guidelines for patient selection, preoperative work-up, procedure selection, and long-term follow-up. These deficiencies contributed to the high rates of suboptimal outcomes.

To create uniform criteria for bariatric surgery, in 1991, the National Institute of Health (NIH) in the USA held a consensus conference on gastrointestinal (GI) surgery for the treatment of severe obesity [2]. A multidisciplinary panel of “experts” reviewed the available peer-reviewed literature and patient experience and created the first criteria for the practice of metabolic bariatric surgery (MBS). However, since 1991, there have been dramatic changes in the field of MBS, including fellowship training, accreditation of MBS centers of excellence, development of MBS registries, the introduction of minimally invasive surgery, and new procedures such as sleeve gastrectomy (SG), and dramatic improvements in perioperative and long-term patient care and safety. Despite these improvements in surgical techniques and perioperative care for patients undergoing MBS, the reliance on the 1991 NIH criteria for determining patient candidacy for surgery remained unchanged, and surprisingly, it is still in wide use more than 33 years later.

In 2022, a group of metabolic bariatric surgeons and other clinicians caring for people with obesity recognized that the 33-year-old guidelines based on expert opinion in the era of open surgery did not reflect the current published literature or state of the field. There was a growing interest in revisiting the 1991 NIH criteria and revising it to reflect MBS’s current practice.

The two largest MBS organizations in the world, the American Society for Metabolic and Bariatric Surgery (ASMBS) and the International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO), agreed to partner to create new guidelines

that would be evidence-based and rely on the most up to date high quality published literature along with current expert global practice. The group searched the literature for high-level evidence using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [3]. Without supportive literature, a Delphi survey of experts in the field was performed [4]. Systematic reviews were performed on 13 topics highlighted in the recently published MBS guidelines. This study aimed to determine the level of evidence and the grade of the recommendations of these 13 previously published criteria [5, 6]. Tables 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 and 13

Methods

In order to methodologically support the previously published ASMBS/IFSO guidelines, two international teams of writers were created.

One team of seven researchers (MDL, GM, AI, GP, ST, SC, AV) performed a systematic search of high-level evidence for different items, according to the PRISMA (see *PRISMA Prospect*). Two independent researchers analyzed each article, first by title and abstract and subsequently by the full text, and extracted the relevant data. In case of disagreement, a third researcher (MDL) was consulted.

Eventually, 12 different systematic reviews from the 13 PRISMA were carried out. PRISMA on item 2 (BMI 35–40 kg/m² without obesity-associated medical problems) produced no studies.

The level of evidence and grade of recommendation are categorized in Table 14.

The second team (MDL, MK, ST) was tasked to resolve any issues not answered by the systematic reviews. For these situations, a Delphi survey was constructed and consisted of two consecutive rounds. Forty-nine recognized MBS experts from 18 countries participated in this Delphi survey to address nine statements that did not have strong backing from the literature search (Table 2 and Table 15). Consensus was reached when the agreement/disagreement rate was equal to or greater than 70%. An online platform (Survey Monkey on <https://www.surveymonkey.com/r/MBS-Criteria>) was used. Seven statements reached consensus in the first round, and two reached consensus in the second round of voting (Table 2 and Table 15). Statements 1 to 5 referred to item 2 (body mass index [BMI] 35–40 kg/m² without comorbidities), and statements 6 to 9 referred to item 6 (joint arthroplasty).

Results

BMI criteria for MBS

1. MBS for BMI 30–34.9 kg/m² [7–35]
PRISMA Appendix 1 [PubMed, Cochrane, Embase]
Systematic Review Table 1

Table 1 MBS Indications for Individuals with BMI 30–34.9

First author (year)	Study design	Quality assessment (NOS)	Asian/Non-Asian	Number of patients	BMI	Comparison to surgical treatment	Number of non-surgical patients	Intervention	Operative time (min)	Length of stay (days)	Weight loss	Complication 1–2	Complication 3–4	Complication 5 (surgical related mortality)	Nutritional complications
Billiter et al. (2022) [7]	Prospective	Good quality	Non-Asian	20	25 < BMI < 35	NO	N/A	RYGB	Not reported	Not reported	8.3 Δ BMI	5%	5%	0%	Not reported
Chaturvedi et al. (2022) [8]	Retrospective/simulation	N/A	Non-Asian	347	30 < BMI < 35	NO	N/A	RYGB, SG	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported	Not reported
Altieri et al. (2022) [9]	Retrospective	Fair quality	Non-Asian	1296	30 < BMI < 35	NO	N/A	RYGB, SG	Not reported	Not reported	30% BMI loss	Not reported	Not reported	Not reported	Not reported
Singh et al. (2022) [10]	Retrospective	Fair quality	Non-Asian	20	30 < BMI < 35	NO	N/A	SG	Not reported	Not reported	18% TWL; 70.3% EWL	Not reported	0%	0%	0%
Baldwin et al. (2021) [11]	Retrospective	Fair quality	Non-Asian	30	BMI < 35	NO	N/A	RYGB, SG	Not reported	Not reported	20–21% TWL; 83–94% EWL	Not reported	Not reported	Not reported	Not reported
Gupta et al. (2020) [12]	Retrospective	Fair quality	Non-Asian	132	30 < BMI < 35	NO	N/A	LAGB to RYGB	Not reported	Not reported	44% EWL	7.8%	23.4% < 30 days; 50% > 30 days	0%	Not reported
Varban et al. (2020) [13]	Retrospective	Fair quality	Non-Asian	1073	BMI < 35	NO	N/A	SG	Not reported	3	22% TWL; 71% EWL	0.7%	3.4%	0%	Not reported
Navarette et al. (2020) [14]	Prospective	Fair quality	Non-Asian	16	30 < BMI < 35	NO	N/A	OAGB	70	2	87.6% EWL	0%	0%	Not reported	Not reported
Gamme et al. (2019) [15]	Retrospective	Good quality	Non-Asian	9094	30 < BMI < 35	Comparison to Class II	9094	RYGB, SG	82	1.6	Not reported	0.9%	3.9%	0%	Not reported
Feng et al. (2019) [16]	Retrospective	Good quality	Non-Asian	8628	30 < BMI < 35	NO	N/A	RYGB, SG	80	1.6	Not reported	0.6%	0.7%	0%	Not reported

Table 1 (continued)

First author (year)	Study design	Quality assessment (NOS)	Asian/non-Asian	Number of surgical patients	BMI	Comparison to surgical treatment	Number of non-surgical patients	Intervention	Operative time (min)	Length of stay (days)	Weight loss	Complications 1–2	Complications 3–4	Complications 5 (surgical related mortality)	Nutritional complications
Vitello et al. (2019) [17]	Retrospective	Fair quality	Non-Asian	56	30 < BMI < 35	YES	20	L AGB, R YGB, SG	Not reported	Not reported	69% BMI loss	0%	7%	7%	Not reported
Noun et al. (2016) [18]	Prospective	Fair quality	Non-Asian	541	30 < BMI < 35	NO	N/A	SG	74	1.7	24% TWL	1.8%	0%	0%	0%
Maiz et al. (2015) [19]	Retrospective	Fair quality	Non-Asian	1119	BMI < 35	NO	N/A	R YGB, SG	70	3	107% EWL	3.8%	0.7%	0%	Not reported
Kaska et al. (2014) [20]	Retrospective	Poor quality	Non-Asian	30	30 < BMI < 35	Comparison to Class II	82	R YGB	Not reported	Not reported	5 Δ BMI	20%	3%	0%	3.6%
Walker et al. (2014) [21]	Prospective	Fair quality	Non-Asian	52	30 < BMI < 35	NO	N/A	L AGB to R YGB	105	3	3 Δ BMI	5%	20%	0%	Not reported
Boza et al. (2014) [22]	Prospective	Fair quality	Non-Asian	100	BMI < 35	NO	N/A	R YGB	110	3	93% EWL	5%	9%	0%	Not reported
Scopinaro et al. (2014) [23]	Retrospective	Fair quality	Non-Asian	10	30 < BMI < 35	NO	N/A	BPD	Not reported	Not reported	6 Δ BMI	40%	40%	0%	Not reported
Serrot et al. (2011) [24]	Retrospective	Fair quality	Non-Asian	17	30 < BMI < 35	YES	17	R YGB	Not reported	Not reported	70% EWL	11.7%	11.7%	0%	0%
Gianos et al. (2011) [25]	Retrospective	Fair quality	Non-Asian	42	30 < BMI < 35	YES	17	L AGB, R YGB, SG	Not reported	Not reported	7–8 Δ BMI	Not reported	Not reported	Not reported	Not reported
Choi et al. (2010) [26]	Retrospective	Fair quality	Non-Asian	66	30 < BMI < 35	Comparison to Class II	438	AGB	Not reported	Not reported	40% EWL	4.5%	1.5%	0%	Not reported
Varela et al. (2011) [27]	Retrospective	Fair quality	Non-Asian	10	30 < BMI < 35	Comparison to Class II	20	AGB	118	1.3	20% TWL	0%	0%	0%	Not reported

Table 1 (continued)

First author (year)	Study design	Quality assessment (NOS)	Asian/non-Asian	Number of surgical patients	BMI	Comparison to surgical treatment	Number of non-surgical patients	Intervention	Operative time (min)	Length of stay (days)	Weight loss (Δ BMI)	Complications 1–2	Complications 3–4	Complications 5 (surgical related mortality)	Nutritional complications	
Scopinaro et al. (2011) [28]	Retrospective	Fair quality	Non-Asian	40	25 < BMI < 35	NO	N/A	BPD	Not reported	Not reported	5 Δ BMI	0%	2.5%	0%	2.5%	
De Maria et al. (2010) [29]	Retrospective	Fair quality	Non-Asian	235	BMI < 35	NO	N/A	AGB, RYGB	Not reported	Not reported	4 Δ BMI	10%	1.3%	0%	Not reported	
Parikh et al. (2010) [30]	Prospective	Fair quality	Non-Asian	93	30 < BMI < 35	NO	N/A	AGB	Not reported	Not reported	54% EWL	1%	3.2%	0%	Not reported	
Sultan et al. (2009) [31]	Prospective	Fair quality	Non-Asian	53	30 < BMI < 35	NO	N/A	AGB	Not reported	Not reported	69.7% EWL	7.6%	1.9%	0%	Not reported	
Cohen et al. (2006) [32]	Retrospective	Fair quality	Non-Asian	33	30 < BMI < 35	NO	N/A	RYGB	56	3	81%	0%	0%	0%	0%	
Angrisani et al. (2004) [33]	Retrospective	Fair quality	Non-Asian	225	BMI < 35	NO	N/A	AGB	Not reported	Not reported	5.2%	2.8%	0%	0%	Not reported	
Cevallos (2021) [34]	Prospective	Fair quality	Non-Asian (Latinos)	51	30 < BMI < 35	NO	N/A	RYGB	Not reported	Not reported	25% TWL; 74% EWL	0%	1.9% < 30 days; 7.8% > 30 days	0%	0%	
Espinosa (2018) [35]	Prospective	Fair quality	Non-Asian (Latinos)	23	30 < BMI < 35	NO	N/A	RYGB	168	3.2	24% TWL	13%	0%	0%	Not reported	
				23,452												

Forty-three articles were included in the present review, 29 (69%) were conducted on non-Asian patients [7–35] and 13 (31%) on Asian patients.

Nine retrospective (31%) and 20 (69%) prospective studies reported MBS results. All articles had a good/fair quality. Two articles investigated the effects of surgery on patients with BMI <30 kg/m², four papers compared outcomes in low BMI with results in patients with severe obesity, and three other studies made a comparison with lifestyle intervention.

Seventeen articles reported results after Roux-en-Y gastric bypass (RYGB), 11 after SG, 1 after one anastomosis gastric bypass (OAGB), 2 after biliopancreatic diversion (BPD), 7 after adjustable gastric banding (AGB), and 2 after revisional surgery from AGB to RYGB with an overall medium follow-up of 29.3 [12–120] months.

Operative time and length of stay (LOS) appeared comparable to available data in the literature for MBS in BMI ≥35 kg/m². All articles reported satisfactory weight loss with no mortality. Clavien–Dindo complications grades 3–4 ranged from 0 to 40% (40% in a paper on BPD complications). A higher complication rate was reported after revisional surgery. Remission from type 2 diabetes mellitus (T2DM) and hypertension (HTN) ranged from 33 to 100% and from 28 to 100%, respectively.

Recommendation

- *MBS is recommended for patients with T2DM and a BMI of 30–34.9 kg/m².*
- *MBS is recommended for patients with a BMI of 30–34.9 kg/m² and one obesity-associated medical problem.*
- *MBS should be considered in patients with a BMI of 30–34.9 kg/m² who do not achieve substantial or durable weight loss or co-morbidity improvement using nonsurgical methods.*

Level of Evidence 2a

Grade of recommendation B

2. MBS for BMI 35–40 kg/m² without obesity-associated medical problems

PRISMA Appendix 2 [PubMed, Cochrane, Embase] not enough studies

No Systematic Review Delphi Table 2

Although previous studies support the superiority of MBS compared to non-surgical therapy in patients with BMI ≥35 kg/m² with no obesity-associated complications, there is a lack of high-grade evidence to support this item. Considering the lack of data from the literature, the leaderships of IFSO and ASMBS have convened a Delphi survey. According to the survey results of 49 experts, MBS is indicated in patients with class II obesity, a BMI of 35–40 kg/m², with no associated

medical problems in all groups of ages following a comprehensive multidisciplinary team (MDT) assessment. The consensus also supported the fact that MBS is cost-effective in patients with class II obesity when compared to non-surgical therapy.

Recommendation

- MBS is recommended for patients with a BMI ≥35 kg/m² regardless of the presence, absence, or severity of obesity-related complications.

Level of Evidence 5

Grade of recommendation D

3. BMI thresholds in the Asian population [36–54]

PRISMA Appendix 3

Systematic Review Table 3

Seven retrospective (54%, 2 multicenter) and 6 (46%) prospective studies reported the results of MBS on the Asian population. All articles have a good/fair quality. The articles investigated the effects of surgery on patients with BMI <30 kg/m².

Eight articles reported results after RYGB, 5 after SG, 2 after OAGB, and one study after SADI-S with an overall medium follow-up of 33.4 [12–84] months. Operative time and LOS appeared comparable to data already published in the literature for MBS in patients with a BMI >35 kg/m².

All articles reported satisfactory weight loss with no mortality. Clavien–Dindo complications grades 3–4 ranged from 0 to 7.3%. Higher long-term nutritional complications were recorded after hypoabsorptive procedures. Remission from T2DM and HTN ranged from 38 to 100% and 30 to 83%, respectively.

Recommendations

- Clinical obesity in the Asian population is recognized in patients with BMI ≥25 kg/m². Access to MBS should not be denied solely based on the traditional BMI criteria.

Level of Evidence 2a

Grade of recommendation B

Extreme of Age

4. MBS in the older population [55–72]

PRISMA Appendix 4 [PubMed, Cochrane, Embase]

Systematic Review Table 4

Eighteen papers have been retrieved [55–72] for qualitative analysis. One RCT [56] and one prospective multicenter paper [62] have been found. Papers were categorized as comparative [55, 56, 62–65, 68, 69, 72] and non-comparative [57–61, 66, 67, 70, 71]. In the comparative group, two

Table 2 IFSO/ASMBS Delphi Results of MBS indications individuals with class II obesity with no associated medical problems

Statement	Round 1	Round 2	Final result
1. Metabolic and bariatric surgery (MBS) is indicated in 18–65-year-old individuals with Class II obesity with no associated medical problems (body mass index of ≥ 35 kg/m ²)	95.7% Agree	–	CONSENSUS (AGREE)
2. MBS is indicated in <18-year-old individuals with Class II obesity with no associated medical problems (body mass index of ≥ 35 kg/m ²)	76.6% Agree	–	CONSENSUS (AGREE)
3. MBS is indicated in >65-year-old individuals with Class II obesity with no associated medical problems (body mass index of ≥ 35 kg/m ²)	85.1% Agree	–	CONSENSUS (AGREE)
4. MBS is indicated for individuals with Class II obesity and have no associated medical problems following comprehensive multidisciplinary team (MDT) assessment (body mass index of ≥ 35 kg/m ²)	54.1% Agree	100% Agree	CONSENSUS (AGREE)
5. MBS is cost-effective in individuals with Class II obesity compared to non-surgical therapy	97.8% Agree	–	CONSENSUS (AGREE)

Table 3 BMI thresholds in the Asian population for MBS

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)
Mazidi, 2017 [36]	Prospective	5	152 type II diabetic obese; all RYGB
Osman, 2019 [37]	Prospective	5	17 type II diabetic obese, all SAGB
Ma, 2022 [38]	Retrospective	5	49 T2DM, all SG
Park, 2021 [39]	Prospective non-randomized controlled	5	17 T2DM BMI 30–35 vs 115 medical therapy; 7 RYGB, 10 SG
Luo, 2020 [40]	Retrospective	5	87 T2DM patients, 25 SG, 62 RYGB
Nautiyal, 2019 [41]	Retrospective	5	113 T2DM
Huang, 2020 [42]	Retrospective multicenter db	5	1199 patients BMI >25
Malapan, 2014 [43]	Prospective	5	29 T2DM RYGB
Zuo, 2020 [44]	Retrospective	5	17 RYGB, 3 SG
Zhao, 2018 [45]	Retrospective	5	78 T2DM, RYGB
Fan, 2014 [46]	Retrospective	5	19 BMI <35 lap band
Kim, 2014 [47]	Retrospective	5	107 BMI <30, SAGB
Zhang, 2017 [48]	Retrospective	5	25 T2DM BMI <30 and 28 T2DM BMI <28
Kwon, 2017 [49]	Prospective non-randomized	6	15 T2DM BMI 23–30 RYGB
Du, 2018 [50]	Retrospective	5	58 T2DM BMI 27.5–32.5, RYGB
Liang, 2018 [51]	Retrospective	5	54 BMI <30 T2DM, RYGB
Widjaja, 2020 [52]	Retrospective	5	18 BMI 27.5–30, SG
Yu, 2021 [53]	Retrospective	5	90 RYGB, 22 SG BMI <32.5 T2DM
Mazidi, 2017 [54]	Retrospective	5	209 RYGB

subgroups have been identified: older [age ≥ 65 years old] versus younger age [55, 65, 68, 69, 72]. SG, RYGB, and AGB were more representative surgical operations in these studies [56, 62–64]. The other studies were not comparative.

Five studies compared the intra- and post-operative complications of MBS between the elderly and non-elderly populations [55, 65, 68, 69, 72]. Despite the high-risk populations evaluated, the studies found no differences in postoperative complications, weight loss, and comorbidities resolutions.

Five studies evaluated the efficacy of AGB [57, 60, 61, 65, 71]. Despite its low peri-operative complication rates, all studies concluded that other procedures, such as SG or

the RYGB, have better post-operative outcomes regarding weight loss and comorbidity resolution or improvement.

According to Gondal et al. [73], rather than age alone, frailty is independently associated with higher rates of postoperative complications following MBS. Furthermore, when considering MBS in older patients, the risk of surgery should be evaluated against the morbidity risk of obesity-related problems. Thus, there is no evidence to support an age limit on patients seeking MBS, but a careful selection that includes an assessment of frailty is recommended.

Two systematic reviews that included studies with elderly groups aged more than 60 years were found in the literature

Table 4 MBS in the older population

Author, year [ref]	Study design	Quality assessment	Number of patients	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidity-resolution rates	Comorbidities	Intra-operative complications	Post-operative mortality	Long-term complications
Athanasidis DI, 2021 [55]	Retrospective comparative	9 (MINORS)	29 (LRYGB72.4%/LSG) VS 997 (LRYGB74.6%/LSG)	72±1.7 (44.5±11.5) VS 48	48	(41.1±6.8) VS NR (46.2±8)	BMI loss - 9.6+7.5 VS 10.2+6.5	DM (50%) VS NR - HT (41.38%) (92.9%) - HT - HC (82.8%) (84.6%) - HC - OSAS (69%) - (62.5%) OSAS (44.8%) VS DM (37.5%) VS DM - HT (20%) (66.2%) - HT - HC (44.9%) (41.7%) - HC - OSAS (23.4%) (44.9%) - OSAS (12%)	DM (50%) VS NR (41.38%) - HT (92.9%) - HT - HC (82.8%) (84.6%) - HC - OSAS (69%) - (62.5%) OSAS (44.8%) VS DM (37.5%) VS DM - HT (20%) (66.2%) - HT - HC (44.9%) (41.7%) - HC - OSAS (23.4%) (44.9%) - OSAS (12%)	NR	8% VS 6% NR	38% VS 23% reoperation during 4-year FU	
Pajacki D, 2021 [56]	RCT	3 (JADAD)	18 (LSG) VS 18 (LRYGB)	67 (65–69) VS 12 67 (65–68)		41.9 (38–47.1) NR VS 47.6 (43.1–51.3)	EWL 29.4% - TWL 31.4% VS EWL 60% - TWL 68%	DM (72%) VS NR - HT (46.15%) (100%) - HT VS DM (-1 pts) (77.7%) VS DM - HT (85.7%) (88.9%) - HT (-2 pts)	DM (72%) VS NR (46.15%) - HT (100%) - HT VS DM (-1 pts) (77.7%) VS DM - HT (85.7%) (88.9%) - HT (-2 pts)	Reported in previous paper	Reported in previous paper	NR	
Nor Hanipah Z, 2018 [57]	Retrospective non-comparative	8 (MINORS)	19 (LRYGB 2/LSG 11/LAGB 4/LGP 2)	76 (75–81)	48 (12–120)	41.4 (35.8–57.5)	EWL 5 years 32.8 (3–5 years) VS 31.4%	DM (58%) VS NR - HT (89%) (89%) - heart disease (58%) - HC (74%) - COPD (26%) - OSAS (58%)	DM (58%) VS NR (6/8) - 0 HT (7/8) - HC (0/7) disease (58%) - HC (74%) - COPD (26%) - OSAS (58%)	3 FA	0	1-year FU (2 FA and 1 poor intake)	

Table 4 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidity-resolution rates	Comorbidity-resolution rates	Intra-operative complications	Post-operative complications	Long-term complications
Hammond JB, 2020 [58]	Retrospective non-comparative	7 (MINORS)	23 (LRYGB)	72 (70–80)	12	43.3 (37.3–56)	NR	%TWL 29(13–46) - EWL 60 (21–105)	DM (17/23) - HT (22/23) - HC (17/23)	DM (16/17) - HT (27%) - HC (16/17)	NR	1 (aspiration pneumonia)	1 poor intake
Gholizadeh B, 2021 [59]	Retrospective non-comparative	7 (MINORS)	61 (OAGB)	67.62±2.03	(12–60)	46.42±5.46	NR	%TWL 60 months (29.56+0.54) - %EWL (70.53+11.87)	DM (60.65%) - HT (72.13%) - HC (45.9%) - GERD (19.67%) - OSAS (18.03%)	DM (70%) - HT (63.41%) - HC (56%) - GERD (100%) - OSAS (100%)	12 months 0	4 (2 bleeding - 1 leak - wound infection) - 4 readmission	7 (3 hypoalbuminemia - 2 marginal ulcer - 2 perforation of marginal ulcer)
Ramirez A, 2012 [60]	Retrospective non-comparative	8 (MINORS)	42 (LRYGB (19%)/LSG (28.6%)/LAGB (52.4%))	73.5 (71–80)	12	44 (34–81)	NR	%EWL (LRYGB - 63.6+32.2) - (SG - 34.9+15.4) - (LAGB - 32.8+17)	DM (38.5%) - HT (66%) - HC (57%) - MI (19%) - OSAS (26%)	DM (53%) - HT (56%) - HC (54%) - OSAS (33%)	2 (accidental colotomy)	0	NR

Table 4 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications
Daigle 2015 [61]	Retrospective non-comparative	7 (MINORS)	30 (LRYGB 16/LSG 6/LAGB 8)	67.1±2.7	37 (6–95)	55.9±3.9	42.3±6.7	%EWL (45.1+21.2)	DM (30%) - HT (90%) - HC (53.3%) - OSAS (46.7%)	DM (40%) - HC (50%)	NR	16.7% (Overall 5 - 1 anastomotic leak after LRYGB - 1 FA after LSG- 1 delayed gastric emptying after LAGB - 1 lower limb deep vein thrombosis after LSG - 1 pulmonary embolism after LRYGB)	NR	0
Casillas RA, 2017 [62]	Prospective multicenter	11 (MINORS)	177 (LRYGB) VS 252 (LSG)	67.5 (67.6)	48	(42.6) VS (42.5)	NR	%EWL (66.1%) VS %EWL (42.3%)	DM (57%) - HT (86%) VS DM (53%) - HT (85%)	DM (57%) NR	NR	6 VS 11 (major complications) VS 12 VS 11 (minor complications)	0 (30 days) VS 3 (major LRYGB complications) VS 0 VS 11 (minor LSG days) VS 11 (minor complications)	10 VS 20 (major complications) VS 0 VS 11 (minor complications)

Table 4 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications
Frieder JS, 2021 [63]	Retrospective comparative	12 (MINORS)	244 (LSG) VS 321 (LRYGB)	(71.1±4) VS (71.7±4.54)	24	(40.5±5.5) VS (43.7±7.22)	VS 34.3±6.6	%EBMIL (43.9+32.2) VS %EBMIL (77.4+26.1)	DM (67.2%) - HT (61.1%) VS DM (38.6%) - HT (31.8%)	NR	NR	9.4% VS 27.7%	0	NR
Quebbemann B, 2005 [64]	Retrospective comparative	11 (MINORS)	13 (LRYGB) VS 14 (LAGB)	68 (66–73) VS 68 (66–73)	12	(45.8±8.6) VS (48.78±5.7)	VS NR	%EWL (75) VS %EWL (46)	4.1% VS 3.9%	2.2% left VS 2.8% left	NR	1 (Anastomotic stricture) VS 2 (conversion to LRYGB; port fracture)	0	NR
Quirante FP, 2017 [65]	Retrospective comparative	10 (MINORS)	1220 (252 LRYGB/726 LS/34 LAGB) VS 393 (130 LRYGB/164 LS/43 LAGB)	(<65) VS (>65)	NR	42.2 (IQR 37.8–47.6) VS 41.6 (IQR 37.8–46.1)	NR	NR	DM (59%) - HT (59%) - HC (33%) - GERD (52%) - OSAS (43%) VS DM (59%) - HT (85%) - HC (44%) - GERD (42%) - OSAS (56%)	NR	NR	448 VS 42	NR	NR

Table 4 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/ TWL/EBMIL	Pre-operative comorbidity-resolution rates	Comorbidities	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications
Lainas P, 2018 [66]	Retrospective non-comparative	8 (MINORS)	54 (LSG)	68 (65–75)	36 (14–62)	43 (35.1–59)	30.7 (23.6–39)	30.7 (23.6–39) ± 5.7 %EWL (76.3 ± 16.5)	DM (70.3%) - HT (90.7%) - HC (42.6%) - OSAS (75.9%)	DM (17.8%) left) - HT (10.7%) left) - HC (10.7%) - OSAS (10.7%) left) - OSAS (21.4%) left)	0	1 (Intra-abdominal hematoma) - 2 (staple line leak) - 2 (other)	0	NR
Danan M, 2019 [67]	Retrospective non-comparative	7 (MINORS)	93 (LSG)	68.8 (65–78)	12	43.6±5.4	30.2 ± 5.7 %EWL (67.1)	DM (20) - HT (51) - HC (17) - OSAS (33)	DM (65%) - HT (72.5%) - HC (47.1%) - OSAS (65%)	DM (65%) left)	NR	1	0	NR

Table 4 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/ TWL/EBMIL	Pre-operative comorbidity-resolution rates	Comorbidity-resolution rates	Intra-operative complications	Post-operative mortality	Long-term complications
van Rutte PW, 2013 [68]	Retrospective comparative	12 (MINORS)	73 (LSG) VS 50 (LSG) VS 12 (LSG)	57 (55–59) VS 62 (60–64) VS 66 (65–70)	VS 14.6	43.8 (29.8–65.1) VS 44.2 (29.8–64.8) VS 45.8 (31.9–65.1)	33.6±6.2 VS 35±5.2 VS 34.2±5	%EWL (55.2%) VS (52.2%) VS (59.9%)	DM (62.3%) - HT (49%) - HC (38.4%) - GERD (26%) - OSAS (30.1%) VS DM (54%) - HT (82%) - HC (42%) - GERD (18%) - OSAS (38%) VS DM (58.3%) - HT (100%) - HC (58.3%) - GERD (8.3%) - OSAS (41.7%)	DM (62.3%) - HT (49%) - HC (35.7%) - GERD (36.2%) - OSAS (59.1%) VS DM (25.9%) - HT (53.7%) - HC (47.6%) - GERD (66.7%) - OSAS (60%) VS DM (42.9%) - HT (33.3%) - HC (28.6%) - GERD (0%) - OSAS (60%)	NR	6 (3 Bleeding - 1 leakage - 1 dehydration - 1 cardiac complication) VS 6 (1 Bleeding - 2 Dysphagia - 2 Cardiac complication - 1 pulmonary) VS 3 (1 complication)	2 (1 Dehydration - 1 incisional hernia) VS 3 (1 dehydration - 2 dysphagia) VS 1 (dysphagia)

Table 4 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidity-resolution	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications
O'Keefe KL, 2010 [69]	Retrospective comparative	II (MINORS)	157 (LRYGB) VS 37 (LAGB) VS 6 (LSG)	67.3±2.3	12	48.5±6.6 (125 PTS) VS 45.9±6.9 (34 PTS) VS 50.0±12.0 (6 PTS)	32.7±5.9 (125 PTS) VS 39.4±7.0 (22 PTS) VS 34.3±19.5 (4 PTS)	%EWL (59.8+14.6) VS (33.5+15.8) VS (34.3+19.5)	NR	NR	9 (1 Obstruction - 4 gastrointestinal bleeding - 1 post-operative respiratory failure - 1 pneumonia - 2 myocardial infarction) VS 2 (2 obstruction) VS 2 (1 obstruction - 1 pneumonia)	2 VS 0 VS 0	NR
Garofalo F, 2017 [70]	Retrospective non-comparative	7 (MINORS)	30 (LSG)	67.2 (65–74)	>24	45.1±5.6	NR	% EWL (52.9±21.8) - %TWL (24±9.9)	DM (56.7%) - HT (86.7%) - HC (56.7%) - OSAS (56.7%)	NR	5 (3 minor complications - 2 major complications)	NR	NR

Table 4 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidity-resolution rates	Comorbidity-resolution rates	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications
Loy JJ, 2014 [71]	Retrospective non-comparative	8 (MINORS)	55 (LAGB)	72.4±2.5	>36	45±6.2	36±6.7 at 2 years	Global mean BMI at 3 years 35±5.6	DM (7%) - HT (89%) - HC (73%) - IMA (29%) - OSAS (56%)	DM (35%) - HT (27%) - HC (28%)	NR	0	0	3 (1 slip-page - 1 band intolerance - 1 incisional hernia in port site)
Nevo N, 2019 [72]	Retrospective comparative	11 (MINORS)	66 (LSG) VS 65 (LSG)	67.6±2.6 (38.4±11)	VS 21	(44.2±7) VS (42.7±5.4)	34.2±5.4 VS 29.3±4.7	%EBMIL (77.3) VS (53.5)	DM (63.6%) - HT (75.8%) - HC (53%) - OSAS (31.8%) VS DM (66.6%) - HT (56.2%) - HC (33.3%) - OSAS (31.8%) VS DM (31.8%) VS DM (23.1%) - HT (24.6%) - HC (23.1%) - OSAS (16.9%)	DM (66.6%) - HT (56.2%) - HC (33.3%) VS DM (59.5) - HT (44%) - HC (74.2%)	0	3 (2 Bleeding - 1 other) VS 3 (1 bleeding - 1 leak - 1 other)	0	8 (5 GERD - 3 stricture) VS 6 (3 GERD - 2 stricture - 1 incisional hernia)

Table 5 MBS for the pediatrics and adolescents

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Paul E. O'Brien, 2015 [76]	RCT	3 (JADAD)	25 (LAGB) VS 25 (lifestyle modification)	(16.5+1.4) VS (16.6+1.2)	VS24	(42.3+6.1) vs (40.4+3.1)	VS NR	%EWL 78.8% VS 13.2%	63 total (27HT - 35 solved) - HC - 1 glucose intolerance)	0	0	8 LAGB (6 proximal gastric enlargement - 2 needle stick injury to tubing)	In this study, gastric banding proved to be an effective intervention leading to a substantial and durable reduction in obesity and to better health. The adolescent and parents must understand the importance of careful adherence to recommended eating behaviors and of seeking early consultation if symptoms of reflux, heartburn, or vomiting occur. As importantly, they should be in a setting in which they can maintain contact with health professionals who understand the process of care. This study indicates that, in such a setting, the laparoscopic adjustable gastric banding process can achieve important improvements in weight, health, and quality of life in severely obese adolescents

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications	Long-term complications	Conclusion	
Carbajo M, 2019 [77]	Retrospective non-comparative	8 (MINORS)	39 (OAGB)	17.8 ± 2 (13–19)	60	48.2 ± 5.7	25.9 ± 5.3	%EWL 94.7 ± 17.9 - %TWL 38.8 ± 12.7	DM (7.9%) - HT (10.3%) - HC (100%) - joint pain (23.1%) - OSAS (30.8%) - GERD (38.5%)	DM (100%) - HT (100%) - HC (100%) - GERD (100%)	0	0	1 severe anemia due to iron deficiency	OAGB is a valid alternative for long-term weight loss and remission of comorbidities in childhood and adolescence. There are no cases of malnutrition and the scarce vitamin or mineral deficiencies can be controlled with specific supplementation, without affecting normal growth. The measurement of the total bowel length is essential to perform a safe technique and to achieve adequate long-term results

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications	Long-term complications	Conclusion	
Dobritoiu D, 2019 [78]	Retrospective non-comparative	7 (MINORS)	64 (2 RYGB - 62 LSG)	Range 12–18	12	39/5	24/92	WL% 36.38	DM (6) - glucose intolerance (6) - HT (6) - OSAS (3) - GERD (14) - liver steatosis (12)	DM (66.6%) - glucose intolerance (100%) - HT (100%) - HC (100%) - OSAS (88.88%) - GERD (100%) - OSAS (64.28%) - liver steatosis (100%)	0	1 prolonged dysphagia	Constipation 22 - iron deficiency 6 - folate deficiency 4 - temporary hair loss 12	Bariatric surgery is already an important part of the management of adolescents with obesity. Among techniques available, laparoscopic sleeve gastrectomy proves to be, again and again, a procedure with lower risks and multiple advantages. Like with every surgical procedure, even lower risks appear when patients are managed in centers of excellence, where multidisciplinary team, with pediatrician and pediatric surgeon included, addresses the patient as a whole and collaborates for best results, in the best interest of the patient

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications	Long-term complications	Conclusion
Elhag W, 2017 [79]	Retrospective non-comparative	8 (MINORS)	102 (LSG)	15.99 ± 1.07 (13–17)	24	46.04 ± 5.99	28.3 ± 6.07	%EWL 81.08 ± 19.65	DM (10) - glucose intolerance (17)	DM (80%) - glucose intolerance (100%)	NR	NR	Our findings confirm that, among obese adolescents, LSG achieves significant weight loss and improves anthropometric parameters and 10 cardiovascular metabolic risk factors, without the development of trace element deficiency after surgery. Conversely, the preoperative nutritional deficiencies (vitamin D, anemia, and hypalbuminemia) persisted or worsened postoperatively
Elhag W, 2021 [80]	Retrospective non-comparative	8 (MINORS)	146 (LSG) - 12 (REDO)	16.51 ± 1.29	108	46.95 ± 7.28	30.20 ± 3.92	%EWL 136.78 ± 38.69	DM (8) - glucose intolerance (38) - HT (1) - HC (53) - OSAS (0) - GERD (2) - liver steatosis (39) - asthma (13) - depression (1) - hypothyroidism (3)	DM (100%) - glucose intolerance (100%)	NR	NR	The findings of current study represent the first contribution to start an evidence base of the long-term outcomes of LSG among adolescents. LSG resulted in marked and durable weight loss and cardiovascular risk reduction, e.g., amelioration of prediabetes, T2DM, hypertension, dyslipidemia, elevated liver enzymes, and hyperuricemia

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Raziel A, 2014 [81]	Prospective non-comparative	8 (MINORS)	32 (LSG)	16.7 (14–18)	60	43.2 (35–54)	NR	%EWL 101.6	34 [DM (2) - HT (7) - HC (8) - joint pain (10) - asthma (5) - OSAS (2)]	28 [DM (2) - HT (5) - HC (6) - joint pain (10) - asthma (3) - OSAS (2)]	0	1 leak 1 cholecystitis	Research evaluating the treatment of morbidly obese adolescents has improved in terms of quality and quantity in the past several years; however, it is still questionable. LSG seems to be an excellent, safe, and efficient bariatric procedure for the treatment of morbid obesity in adolescents
Yitzhak A, 2006 [82]	Retrospective non-comparative	6 (MINORS)	60 (LAGB)	16 (9–18)	39.5	43 (35–61)	30 (20–39)	NR	18 [DM (2) - HT (3) - Asthma (3) - OSAS (10)]	All solved	0	0	8 Band slip-page

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Holterman A-X, 2012 [83]	Prospective comparative	10 (MINORS)	18 (11 Morbidly obese VS 7 Superobese) LAGB	16 ± 1 VS 16.6 ± 0.8	18	44 ± 4 VS 61 ± 7.5	NR	%EBMIL 20.5% ± 10.3% VS 22.8% ± 11.6%	Insulin resist- ance 82% - HC 82% % - NAFLD 91% VS insulin resistance 29% - HC 100% - HC 86% - NAFLD 86%	Insulin resist- ance 78% - HC 67% NAFLD NR VS insulin resistance 17% - NAFLD NR	NR	NR	These short-term outcome data of bariatric adolescent patients demonstrating differential and delayed metabolic response of SO adolescents in a gastric banding-based weight loss program despite weight loss provide the rationale for early referral for timely intervention of obesity in adolescents before they progress to the SO state to optimize their metabolic response to bariatric interventions
Holterman A-X, 2010 [84]	Prospective non-comparative	9 (MINORS)	26 (LAGB)	16 ± 1	29 ± 9	50 ± 10	40.6 ± 4.6	%EWL 41 ± 27%	DM 90% - HTDM 39% - HC 45% - HC 80% - fatty liver 88%	DM 39% - HTDM 33% - HC 27%	0	0	3 wound infection - 2 laparoscopic revisions for 1 band malfunction and 1 hiatal hernia repair

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications	Long-term complications	Conclusion
Inge TH, 2004 [85]	Prospective non-comparative	6 (MINORS)	20 (RYGB laparoscopic 2 open)	18NR	24	NR	NR	%EWL NR	NR	Reported as all pre-operative comorbidities solved	NR	1 Roux limb obstruction	Carefully designed clinical studies should be performed to better define the safety of RYGBP in this population and identify which patients will benefit most from gastric bypass, what factors might predict postoperative complications, and what postoperative management approaches are most helpful for adolescents
Jaramillo JD, 2017 [86]	Retrospective non-comparative	8 (MINORS)	38 LSG	16.8 ± 1.3	12	46.7 ± 5.7	35/5	%EWL 47.7	DM 11 - insulin resistance 9 - HT 13 - HC 22 - fatty liver 5 - OSAS 24	DM 10 - insulin resistance 4 - HT 13 - HC 18 - fatty liver 5 - OSAS 22	0	0	0
Conroy R, 2011 [87]	Prospective non-comparative non-randomized	9 (MINORS)	88 LAGB	16.8 ± 0.1	12	45.9 ± 1	NR	%EWL 17.1 ± 2.2%	NR	NR	NR	NR	NR

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Riquin E, 2018 [88]	Retrospective comparative	6 (MINORS)	16 LAGB vs 35 lifestyle modification	15.3 ± 1.2	24	40.6±4.4 VS 39.9±4.5	NR	%EWL 42.8% vs 51%	DM 8 - HT 3 NR - joint pain 7 - OSAS 2 - NAFLD 5 VS endocrine and metabolic 20 - muscular/skeletal 17 - respiratory 19	0	0	0	In conclusion, this study highlights the usefulness of laparoscopic adjustable gastric band (LAGB) for weight loss and also shows the significant psychiatric vulnerability of severely obese adolescents. Given the vulnerability of the patients in this cohort, we argue that lengthy and close psychiatric follow-up before surgery is necessary
Loy JJ, 2015 [89]	Retrospective non-comparative	8 (MINORS)	52 LAGB	16.1 (14–17.8)	24	48.8±7	36.8±8.2	%EWL 45.7±23.1%	NR	0	0	4 slippage	LAGB in suitably selected and counseled adolescents is safe and effective. The excess weight loss observed and the resolution of the components of the metabolic syndrome were significant and seen at 1 year postoperatively. These improvements were sustained at 2 years

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Manco M, 2016 [90]	Retrospective comparative	9 (MINORS)	20 LSG - 20 BIB - 53 lifestyle modification	16.71 (1.44) VS 14.13 (2.12) VS 14.67 (1.89)	12	48.56 (4.15) VS 40.24 (5.02) VS 40.40 (3.55)	38.54 (3.51) VS 38.93 (4.67) VS 39.61 (3.71)	nr	DM 25% - HT60% - HC 60% - OSAS 45% VS DM 20% - HT45% - HC 40% - OSAS 30% VS DM 15 - HT 39.6 - HC 36.3 - OSAS22.6	Left: DM 0% - HT 0% - HC 5% - OSAS 15% VS DM10 - HT 40% - HC 25% - OSAS 20% VS DM 13.6% - HT 22.7 - HC -36.3 - OSAS 27.27	0	1 Pneumo-0 nia after LSG	LSG reverted steatohepatitis and reduced hepatic fibrosis in morbidly obese adolescents with NAFLD 1 year after surgery. It was also beneficial for resolving hypertension and ameliorating dyslipidemia and OSAS. In contrast, lifestyle intervention, alone or in combination with IGWLD, was not able to induce a sustained weight loss and therefore was less effective in reverting liver histology and metabolic abnormalities
Nadler E, 2009 [91]	Prospective non-comparative	9 (MINORS)	54 LAGB	16.1 ± 1.2	24	48.1 ± 6.4	35.8 ± 7.9	%EWL 46.8 ± 21.9	DM 9 - HT 11 - HC 17 - OSAS 5 - GERD 2	DM 9 - HT 11 - HC 12 - OSAS 1 - GERD 2	NR	NR	In summary, we found that the LAGB provides an excellent option for weight loss in the morbidly obese adolescent population. The EWL after LAGB in morbidly obese adolescents is approximately 45% at 1- and 2-year follow-up, with the majority of weight loss consisting of android fat

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Zeller MH, 2009 [92]	Retrospective non-comparative	8 (MINORS)	31 LRYGB	16.4 (1.4)	12	63.5 (10.6)	39.4±7.8	NR	NR	NR	NR	NR	
Xiao N, 2015 [93]	Retrospective comparative	8 (MINORS)	22 Severe obese VS 44 lean	16.5 (15.0, 17.4) VS 16.5 (14.5, 17.0)	12	48.4 (42.0, 51.7)	32.2 (28.6, 37.1)	NR	DM 1 - HT 4 All solved	NR	NR	NR	This study demonstrated that adolescents with severe obesity have sub-clinical kidney injury in the presence of normal kidney function and absence of microalbuminuria. The study also determined that sub-clinical kidney injury persists 1 year after significant weight loss induced by bariatric surgery
Sarwer D, 2017 [94]	Prospective comparative	9 (MINORS)	119 Surgery VS 169 lifestyle modification	17 ± 1.5 vs 15.24 ± 1.4	52 ± 8.9 VS 37 ± 5.2	37.7±8.4 VS 35.8±4.8	%TWL 31.6±1.1 VS 0.3±0.9	NR	NR	NR	NR	NR	The study had several limitations. Our two groups differed in age, race, gender, and BMI. Since the start of this study, a number of investigations have raised concerns about the accuracy and validity of information that adolescents provide in these interviews, as noted above. (48–51). We also experienced some attrition. While this did not appear to be systematic, it may have biased the results

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications	Conclusion
Ryder J, 2016 [95]	Prospective non-comparative	8 (MINORS)	206 (139 LRYGB - 56 LSG - 11 LAGB)	17.1 (1.57)	24	51.7 (8.5)	NR	%EBMIL 40.3%	NR	NR	NR	NR	NR	Bariatric surgery in adolescents with severe obesity significantly improves resting HR, completion time of a standardized 400-m walk test, and immediate post-test HR response, and reduces walking-related musculoskeletal pain complaints at 6 months post-surgery. These meaningful improvements were maintained up to 2 years post-surgery. Whether these positive changes in functional mobility and musculoskeletal pain persist over the long-term and lead to further improvements in cardiometabolic risk requires evaluation

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Alqatani AR, 2016 [96]	Retrospective comparative	9 (MINORS)	24 LSG in PW vs 72 LSG non-PW	NR	60	46.4 ± 12.0 VS 46.4 ± 11.7	35.9 ± 12.5 VS 25.1 ± 7.0	%EWL 38.4 ± 25.6 VS 75.4 ± 28.3	Only in PW (DM 6 - HT 10 - HC 15 - OSAS 24)	DM 6 - HT 7 - HC 9 - OSAS 21	0	0	In our experience, LSG is a well-tolerated, effective treatment option for severely obese PWS patients. The surgical procedure similarly resulted in significant weight loss and maintained resolution of comorbidities in both study participant groups (PWS and non-syndromic obese patients), particularly with limited other options for treatment of marked obesity. Nevertheless, long-term studies are needed to confirm the durability of this weight loss, the co-morbidity resolution, and long-term complications. The inclusion of patient-reported outcomes such as QOL is vital to fully understand the "success" of bariatric surgery, with promotion of QOL identified as a primary goal of Healthy People 202
Reiter-Purtill J, 2020 [97]	Retrospective comparative	9 (MINORS)	139 Surgery (88 LRYGB - 50 LSG - 3 LAGB) VS 83 lifestyle modification	16.86 ± 1.39 VS 16.11 ± 1.40	24	51.52 ± 8.32 VS 46.85 ± 6.12	36.01 ± 8.55 VS 48.65 ± 8.37	NR	NR	NR	NR	NR	

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications	Conclusion
Xanthakos S. 2020 [98]	Prospective comparative	10 (MINORS)	226 (161 LRYGB - 65 LSG)	16.6 (1.6) VS 16.4 (1.6)	60	53.7 (9.6) VS 50.2 (8.3)	39.0 (32.0, 48.2) VS 37.0 (32.1, 40.8)	NR	(DM)16% - HT 46% - HC 79% - GERD 12% - liver steatosis 44% VS (DM)11% - HT 36% - HC 70% - GERD 16% - liver steatosis 28%)	NR	NR	NR	NR	In a prospective study of adolescents who underwent RYGB or VSG, we observed nutritional deficiencies by 5 years after the procedures—particularly in iron and vitamin B ₁₂ after RYGB

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications	Conclusion
Nehus EI, 2017 [99]	Prospective non-comparative	8 (MINORS)	242 (140 LRYGB - 52 LSG - 14 LAGB)	17/1	36	50.5 (45.2, 58.3)	36.2 (30.2, 44.9)	NR	DM 30 - HT 104 - HC 179	DN 29 - HT 75 - HC 125	NR	NR	NR	In conclusion, we report 3-year kidney outcomes of the largest study to date of adolescents undergoing bariatric surgery. Kidney function and albuminuria improved following weight loss surgery in those participants with evidence of pre-operative kidney disease. Furthermore, BMI levels of greater than 40 kg/m ² at follow-up were associated with a progressive decline in kidney function. These data support the addition of kidney dysfunction as a selection criterion for bariatric surgery in adolescents who reach a BMI of 40 kg/m ² to optimize chances for reversal of severe obesity and kidney risks

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Kaar J, 2021 [100]	Retrospective non-comparative	8 (MINORS)	81 (LSG)	16.9 ± 2.0	36	47.9 ± 7.3	37.2 ± 5.1 in 15 pts with OSAS remission	DM 16 - HT 20 - OSAS 44 TWL -27% in patient with OSAS remission and -25% in patient without OSAS remission	DM 16 - HT 20 - OSAS 15	NR	NR	NR	Over half of our adolescents with severe obesity undergoing MBS had a diagnosis of OSA. Of those adolescents who had completed a post-operative PSG, almost two-thirds had remission of OSA following surgery. Those with higher baseline BMIs may be less likely to achieve remission. Obtaining both pre- and post-MBS PSGs is recommended for routine clinical care to accurately identify patients with OSA to promote positive health outcomes
Derderian SC, 2020 [101]	Prospective comparative	9 (MINORS)	192 (140 LRYGB - 52 LSG) (114 greater weight loss VS 78 small weight loss)	16.7 (1.56) VS 16.1 (1.60)	60	52.6 (9.40) VS 53.2 (10.0)	NR	%TWL 34.1 ± 9.5% VS 25.6 ± 7.9% (at 1-year follow-up) 33.8 ± 9.2% VS 8.6 ± 9.5% (at 5 years)	DM 12 - HT 37 - HC 27 - DM 8 - VS (DM 13 - HT 17 - HC 28)	DM 10 - HT 27 - HC 62	NR	NR	These results emphasize the metabolic benefit of bariatric surgery. Whether modest weight loss (5–10% total body weight loss) achieves similar metabolic benefits is an area of future research

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Nobili V, 2017 [102]	Prospective non-comparative	8 (MINORS)	20 (LSG)	16.71 ± 1.44	12	49.3 ± 4.8	37.5 ± 5.3	NR	20 NAFLD				In conclusion, our findings suggest a mechanistic explanation of the beneficial effect of LSG-associated weight loss on liver histology in adolescent NAFLD and highlight that the histologic improvement induced after LSG is associated with the activation state of local cellular compartment and their crosstalk
Katra M, 2005 [103]	Retrospective non-comparative	7 (MINORS)	35 (LRYGB)	17.57 ± 1.82	12	60.8 ± 11.07	41.6 ± 9.5	%EBMIL	OSAS 19	NR	NR	NR	In summary, our study indicated that OSA was highly prevalent in extremely overweight adolescents meeting eligibility criteria for bariatric surgery. The significant weight loss after bariatric surgery was associated with either the resolution of OSA in a majority or a significant reduction in OSA severity

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications	Conclusion
Watanabe Y, 2022 [104]	Retrospective comparative	8 (MINORS)	122 Childhood obesity VS 183 post-puberty obesity (LSG)	<13years VS >13years	36	43.8 (38.6–50.6) VS 40.3 (36.7–45.7)	-13.9±0.6 VS -12.9±0.5	(30.1±11.7) VS (29.7±10.6)	(DM 60.7% VS 68.9%) HT 73.7% VS 81.8% HC 98.4% VS 95.1% OSAS 47.4% VS 32.1% VS 65.6% VS 75.4%	(DM 83.6% VS 58.4%) HT 47.4% VS 32.1% VS 65.6% VS 75.4%	NR	NR	NR	In conclusion, this study suggests that severely obese patients with childhood onset tend to have more severe and subcutaneous fat-dominant obesity compared to those with post-puberty onset, and subcutaneous fat predominance may be associated with lower HbA1c in those with childhood onset
Hjeltnesæth J, 2020 [105]	Prospective comparative	10 (MINORS)	39 (LRYGB) VS 96 (lifestyle intervention)	16.7 (1.0) vs 15.6 (1.3)	12	45.6 (4.4) VS 43.3 (4.1)	31.5 (4.4) VS 42.7 (5.2)	%TWL -30 (-33 to -27) VS 1 (-1 to 3)	(DM 9 VS 8) HT 22 VS 17 NAFLD 15 VS 19 HC 65 VS 43	(DM 9) VS (DM 8)	2 MINOR SURGICAL COMPLICATION	0	Hypovitaminosis (10% anemia - 21% iron deficiency - 10% vitamin B ₁₂ deficiency)	

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Tashiro J, 2022 [106]	Retrospective comparative	8 (MINORS)	76 (LSG with parent history of bariatric surgery) 184 (with-out)	16.4 (2.07) VS 16.8 (1.81)	12	50.3 (10.9) VS 47.8 (8.47)	-21.0 (18.1) VS -24.5 (14.7)	%EBMIL 53.4 (19.6) VS 59.6 (25.8)	NR	NR	NR	NR	Programs offering adolescent bariatric surgery, as well as insurance companies, should consider eliminating their minimum age requirement for surgery. Although there were no differences in weight loss outcomes for those patients whose parents have or have not had bariatric surgery themselves, given their heavier size at age of surgery, there may be benefit in ensuring even earlier access to care for these children and adolescents
Lee DY, 2012 [107]	Retrospective comparative	10 (MINORS)	32 (LRYGB) VS 23 (LAGB)	18.6±0.6 VS 17.2±1.5	24	50.6±7.0 VS 47.0±7.4	NR	%EWL 83.4±20.5 VS 29.7±18.9	(DM 3 - HT 2 - VS (HC OSAS 5) 1) (DM 3 - HC 2) VS (HC 1)	0	0	1 Gastro-gastric fistula and 1 dehydration in LRYGB 5 iron deficiency vitamin D deficiency VS 2 band removal for slip-page 2 IRON DEF- CENCY 2 VIT D DEF- CENCY	LRYGB achieved superior weight loss in adolescents compared to LAGB in a short-term follow-up in our series. Randomized studies with long-term follow-ups will be needed before definitive recommendations can be made on the appropriate operation for this age group

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Cozacov Y, 2014 [108]	Retrospective non-comparative	8 (MINORS)	18 (LRYGB and LSG)	17.5	54	47.2	30.1	(DM 1 - HT 2 - HC 1 - GERD 4 - OSAS 6)	(DM 1 - HT 2 - HC 1 - GERD 3 - OSAS 3)	0	0	1 (Hartburn LSG) (LSG)	LSG and RYGB are safe and effective, as the long-term success of this study shows, in treating the morbidly obese adolescent population. We have shown in the longest prospective post-LAGB study in Australian adolescents with severe obesity that LAGB improves BMI in the majority of adolescents with minimal complications. LAGB may be the preferred 'initial' operation for adolescents with severe obesity as part of a "stepped up" surgical management plan
Peña AS, 2017 [109]	Prospective cohort study	9 (MINORS)	21 (LAGB)	17.4 [16.5–17.7]	45.5 [32–50]	47.3 ± 8.4	35 [32–51]	%BMIL 15 [7.24] - %TWL 16 [7.24]	NR	NR	1 Hematoma - 2 food intolerance - 1 infection	2 Food intolerance - 7 band removal	We have shown in the longest prospective post-LAGB study in Australian adolescents with severe obesity that LAGB improves BMI in the majority of adolescents with minimal complications. LAGB may be the preferred 'initial' operation for adolescents with severe obesity as part of a "stepped up" surgical management plan

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications	Conclusion
Dumont PN, 2017 [110]	Prospective cohort study	9 (MINORS)	97 (LAGB)	17.2 ± 0.7	56.0 ± 22.0	44.9 ± 6.1	NR	%EWL 46.6 ± 39.5	DM 1 - HT 8 - HC 1 - OSAS 5	DM 1 - HT 8 - HC 1 - OSAS 3	NR	NR	19 Band removal	The effectiveness of LAGB appears to improve over the long term in adolescents. LAGB is a reliable, reversible technique that may be an appropriate and ethical first-line surgical option for obese youth. In the current study of adolescents undergoing LAGB, the procedure was safe and there was a significant positive correlation between excess weight loss and follow-up duration (mean 56.0 ± 22.0 months); overall excess weight loss was 46.6%

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative mortality	Long-term complications	Conclusion
Lainas P, 2019 [111]	Retrospective non-comparative	8 (MINORS)	84 (LSG)	17 [15–17]	24	43.7 [31.5–74.9]	28.8 [21–38.6]	%EWL 79.1 ± 15.5 %TWL 29.1 ± 5.4	DM 88 - HT 12 - HC 5 - GERD 4 - OSAS 22	DM 85 - HT 6 - HC 1 - GERD 2 - OSAS 8	1 (Pulmonary complication)	NR	This study suggests that LSG is safe and effective for patients under 18 years old, resulting in significant weight loss, comorbidity remission, and QoL improvement. Weight loss is associated with a marked metabolic improvement, depicted in this series by the disappearance of insulin resistance in all but one patient
Mohammed MR, 2021 [112]	Retrospective non-comparative	8 (MINORS)	72 (LSG)	17.4 ± 1.85	12	47.9 ± 7.1	31.76 ± 6.15	%EWL 73.5 ± 22.1% - %TWL 34.9 ± 9.35	DM 6 - HT 5 HC 1 - OSAS 3	DM 6 - HT 4 - HC 1 - OSAS 3	0	0	1 Severe dys-SG is an effective and safe procedure in adolescent patients with severe obesity. SG achieved satisfactory weight loss in the majority of patients along with excellent remission in comorbidities, namely, DM. The complication rate of SG in adolescents is acceptably low, yet nutritional deficiencies warrant strict follow-up

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications	Long-term complications	Conclusion
Teepie EA, 2012 [113]	Retrospective non-comparative	7 (MINORS)	15 (LRYGB)	NR	24	58.8 ± 10.7	34.9 ± 5.6	%EWL 62.2 ± 14.6	Non-extractable data	0	1 Ileus - 1 anastomotic leak - 1 anastomotic bleeding	1 port site hernia - 1 gastrojejunal stricture	In conclusion, bariatric surgery is a safe and efficacious treatment option for morbidly obese adolescents with significant obesity-related comorbid conditions
Tuna T, 2020 [114]	Retrospective non-comparative	8 (MINORS)	16 (LSG)	15–19	36	48.6 (42–56.4)	29.6 (24.5–33.2)	%EWL 79.3 ± 20.5	DM 13 - HT 3 - HC 5 - OSAS 8 - NAFLD 13	DM 13 - HT 2 - HC 3 - OSAS 8 - NAFLD 8	0	0	Laparoscopic sleeve gastrectomy results in significant weight loss and leads to resolution of comorbidities in most patients, with a low rate of complications or re-intervention. Although current evidence points to MBS as a safe and effective option for treatment of morbidly obese adolescents, our limited sample and the short FU period do not allow definitive conclusions to be drawn, particularly concerning long-term safety

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative mortality	Long-term complications	Conclusion
White B, 2017 [115]	Prospective comparative	10 (MINORS)	29 (18LSG VS 21 LRYGB) VS lifestyle change	16.1 (1.3) VS 15.9 (1.3)	48	51.3 (7.8) VS 51.1 (7.4)	VS Non-extractable data	Non-extractable data	DM 3 - HT 10 - HC 6 VS (DM 2 - HT 7 - HC4)	Non-extractable data	1 Gastric perforation (LRYGB)	0	1 Gastric perforation - 2 gas-trojejunal stricture - 7 GERD	Adolescent bariatric surgery in an NHS service compares favorably to international cohorts and shows promise as an effective treatment for severe obesity. Further work is needed to improve patient selection, reduce age at surgery, and reduce attrition
Messiah SE, 2013 [116]	Retrospective non-comparative	8 (MINORS)	890 (454 LRYGB - 436 LAGB)	18.5	12	48.5 (48.0-49.0)	37.0 (36.3-37.8)	%EWL 23.5 (22.4-24.7)	NR	NR	NR	NR	38 Gastrointestinal complication - 26 nutritional deficiency	1 Death 5 months after LRYGB

Table 5 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean age (years)	Mean follow-up (months)	BMI basal	BMI follow-up	BMI loss/EWL/TWL/EBMIL	Pre-operative comorbidities resolution	Intra-operative complications	Post-operative complications mortality	Long-term complications	Conclusion
Kauffman AM, 2010 [117]	Retrospective non-comparative	8 (MINORS)	61 (LRYGB)	17.31.9	24	54.46.8	NR	TWL 49.22.1	DM 11% - HT 30% - NALFD 18% - OSAS 63%	NR	NR	NR	In our study, a 7.4% decline in WB BMC was shown in adolescents who underwent bariatric surgery. The decrease in bone mass was associated, to a small extent (14% for BMC), with weight loss in the first year. Future studies are needed to elucidate the types of weight loss (lean versus fat mass) and molecular pathways that influence changes in bone mass, and the clinical consequences of these changes

Table 6 MBS prior to joint arthroplasty (as a bridge)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Choi, 2020 [120]	Retrospective	5	1327 bariatric with hip repair vs 2127 only hip repair	10 years	NA	NA	NA	NA	NA	NA	NA	NA	Hip fractur	Survival rates lower in bariatric (87.2% vs 91.8% <i>P</i> 0.048), no differences for complications rates at 30 dd, greater readmission rates for bariatric patients (OR 1.46, 95%)
Inacio, 2014 [121]	Retrospective	5	69 patients with bariatric surgery >2 years and 102 within 2 years of total joint arthroplasty	1 year	NA	NA	NA	NA	NA	NA	NA	NA	171 (hip 21%, knee 79%)	Similar post-operative complications than non-operated obese
Werner, 2015 [122]	Retrospective	5	219 patients with previous bariatric surgery vs 11,294 obese	2 years	NA	NA	NA	NA	NA	NA	NA	NA	219 total knee arthroplasty	Lower 90-day complication rate than non-operated obese, but increased than lean control

Table 6 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Nickel, 2016 [123]	Retrospective	5	5918 bariatric vs 26616 non-operated obese BMI >40 vs 6480 lean	2 years	NA	NA	NA	NA	NA	NA	NA	NA	5918 total knee arthroplasty	Higher 30-day complication rates for bariatric patients vs non-operated obese vs lean
McLawhorn, 2018 [124]	Propensity score matched analysis	5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	2636 bariatric vs 2636 non-operated TKA; 792 bariatric vs 792 non-operated obese THA	Lower in-hospital complications rate in bariatric patients; similar rates of revision
Lee, 2018 [125]	Retrospective	5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	35 bariatric THA, 70 bariatric TKA	Prior to THA, bariatric surgery patients were at increased risk for post-operative infections; prior to TKA were at increased risk of revision but lower risk for infections

Table 6 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Wang, 2019 [126]	Propensity score matched analysis	5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	2540 bariatric vs 2540 non-operated THA; 9803 bariatric vs 9803 non-operated TKA	Prior to THA, bariatric surgery patients were at increased risk for blood transfusion and anemia; prior to TKA were at increased risk of for blood transfusion and anemia but lower risk for pulmonary embolism
Sax, 2022 [127]	Retrospective	5	NA	6 to 12 months	NA	NA	NA	NA	NA	NA	NA	NA	1901 BS 6 months before TKA; 14,022 BS 1 year before TKA; vs 121,934 lean and 87,449 BMI >40	Lower complications than BMI >40; similar rates BS at 6months vs BS at 1 year

Table 6 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Ryan, 2022 [128]	Retrospective	5	64 RYGB, 9 LAGB, 8 SG, 7 NA bypass	NA	NA	NA	NA	NA	NA	NA	NA	NA	THA	BS were more likely to have reoperation and periprosthetic joint infection than lean control; 90-day complications were lower in BS than BMI >40
Martin, 2015 [129]	Retrospective	5	NA	5 years	51.1	36.5	NA	NA	NA	NA	NA	NA	91 TKA after BS vs 91 TKA high BMI and 182 low BMI	increased risk of reoperation in BS group vs high-BMI group and vs lean
Watts, 2016 [130]	Retrospective	5	47	10 years	49.7	35.3	NA	NA	NA	NA	NA	NA	47 THA after BS vs 94 THA in non-operated obese	Lower rate of reoperation and revision after BS

Table 6 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Liu, 2021 [131]	Retrospective	5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1894 BS before TKA/THA vs 1000 obese TKA/THA	After 6 months from the primary surgery, BS patients had less complications than THA/TKA alone
Nickel, 2018 [132]	Retrospective	5	NA	NA	NA	36.5	NA	NA	NA	NA	NA	NA	1545 BS before THA vs 6918 BMI >40 vs 3697 lean	Dislocation and revision rate increased after BS; lower rate at 90-day complications BS vs BMI >40
Dowsey, 2022 [133]	RCT	8	39 LAGB	12 months	43.8	36.5	20%	NA	NA	NA	NA	NA	29 BS before TKA vs 39 obese TKA	Peri-operative complications were lower in BS group
Ighani Arani, 2021 [134]	Retrospective	5	SG and RYGB	12 months	43	31	NA	NA	NA	NA	NA	NA	441 BS before TKA vs 95,948 TKA same age and same BMI	Similar risk for revision and infection

Table 6 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Nearing II, 2016 [135]	Retrospective	5	92 RYGB and 10 SG	4.9	NA	37.6	NA	4 T2DM, 8 OSAS	NA	NA	NA	NA	49 TKA and 17 THA after BS vs 23 TKA and 13 THA before BS	Lower operative time and length of stay, similar rate of 30-day complications and reinterventions over long-term FU
Liu, 2020 [136]	Retrospective	5	NA	2 years	NA	NA	NA	NA	NA	NA	NA	NA	1478 BS before TJA, 60,259 obese before TJA, 281,973 lean before TJA	BS before TJA had increased complications, especially blood transfusion, but lower pneumonia than obese before TJA

Table 6 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Meller, 2019 [137]	Retrospective	5	2044 LAGB, 1671 RYGB, 1025 SG, 21,112 non-specified BS		NA	NA	NA	NA	NA	NA	NA	NA	25,852 BS before TKA vs 2,675,575 no previous BS TKA	Readmission in 90-day post-TKA was increased in BS, a low increased trend for 90-day complications was observed in BS than in non-operated BS can reduce knee symptoms and the necessity to undergo knee surgery. After BS complications were lower than those who underwent first to TKA
Purcell, 2022 [138]	Retrospective	5	355 SG	5 years	51.4	41.3	66.3 lb	6 resolution of knee pain	NA	NA	NA	NA	27 SG before TKA vs 24 TKA/arthroscopy before SG	BS can reduce knee symptoms and the necessity to undergo knee surgery. After BS complications were lower than those who underwent first to TKA

Table 6 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidities resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Schwarzkopf, 2018 [139]	Retrospective	5	1347	NA	NA	NA	NA	NA	NA	NA	NA	NA	330 BS before THA and 1017 BS before TKA	No association between the time of BS and arthroplasty and 90-day complication, whereas those who underwent BS less than 6 months are at increased risk for readmission
Ighani Arani, 2022 [140]	Retrospective	5	NA	1.1 years	30.6	33.6% TWL	NA	NA	NA	NA	NA	NA	44 BS before TKA vs 3524 no BS TKA	No clinical differences in 1-year post-operative score and function score
Severson, 2012 [141]	Retrospective	5	NA	NA	37.9	NA	NA	NA	NA	NA	NA	NA	61 BS >2 years before TKA, vs 25 BS within 2 years TKA vs 39 TKA before BS	Reduced post-operative time, similar 90-day complication

Table 7 MBS and abdominal wall hernia repair

Author, year [ref]	Study design	Quality assessment	Number of patients	Intervention	Length of stay	Concomitant hernia repair (yes/no)	Type of concomitant hernia repair	Patients with concomitant VHR	BMI at VHR	EWL/TWL	OR time (min)	Complications MBS
Morrell, David J et al. [143]	Retrospective	3	20	LSG	1.6 ± 0.8	No	No	0	45.6 ± 6.1	20.7 ± 12.3% TBMIL	121.2	<i>n</i> = 1 bleeding
Moolla, Muhammad et al. [144]	MBSAQIP	2	4690 concomitant VHR vs 4648 matched	RYGB/LSG	NR	Yes <i>n</i> = 4690; no <i>n</i> = 4648	Epigastric, incisional, umbilical, Spigelian hernias	4690	46.1 ± 8.4	NR	112.6 ± 57.9 vs 93.7 ± 50, <i>p</i> < 0.001	
Krivan, Sylvia et al. [145]	Miss Retrospective	3	106	RYGB/LSG	4.1 ± 1.5	Yes laparoscopic and open	Open/lps, primary closure/mesh	106	53 ± 9	25.4% at 24 month	NR	6 of 106 (omental bleedings, 2 anastomotic leak, gastrojejunostomy stenosis)
Sharma, G et al. [146]	Retrospective	3	159	105 RYGB, 50 SG, 4 GB	3 (2–4)	Yes	Umbilical, (99), incisional (52), others (16) - 144 lps, 15 open	159	48.2	56.4% (primary) and 55.9% (mesh) at 1 year	210 min (156.5–243.5)	4 major complications, 16 minor complications
Olimi, Stefano et al. [147]	Retrospective	3	Group A <i>n</i> = 30 before MBS, <i>n</i> = 170 after weight loss; Group B <i>n</i> = 170	SG (group B), RYGB 2, GB 2	Group A: 2 ± 2.7 vs group B: 2.8 ± 1.9 (<i>p</i> < 0.5)	No	All lps Parietex composite	No	Group A: 37.8 ± 5.7; Group B: 24.6 ± 4.5	OR hernia	NR	NR
Raziel, Asnat et al. [148]	Retrospective	3	54	SG 48, lps RYGB 2, open RYGB 2, GB 2	NR	Yes	Dual mesh during MBS	54	44.2 ± 4.5	EWL 57.7% ± 9.2 at 12 months	NR	3 leaks, 1 pulmonary embolism
Chandeze, Marie-Maëlle et al. [149]	Retrospective	3	Group A after MBS <i>n</i> = 30, during MBS, group B <i>n</i> = 60	SG 17, RYGB 18, GB 4	Group A: 6.2 ± 2.6, group B: 10.7 ± 9.3 (<i>p</i> = 0.002)	Yes	Continuous suture with slowly absorbable monofilament	14	46.7 ± 6.4 vs 34.1 ± 6.5 vs 42.4 ± 7.2	NR	NR	NR

Table 7 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients	Intervention	Length of stay	Concomitant hernia repair (yes/no)	Type of concomitant hernia repair	Patients with concomitant VHR	BMI at VHR	EWL/TWL	OR time (min)	Complications
Morrell, David J et al. [143]	Retrospective	3	20	LSG	1.6 ± 0.8	No	No	0	45.6 ± 6.1	34.9	20.7 ± 12.3% TBMIL	<i>n</i> = 1 bleeding
Praveen Raj, Palanivelu et al. [150]	Retrospective	3	156	SG 120, RYGB 36	3 (2–5)	Yes	IPOM	156	43.64 ± 6.8 and 42.49 ± 8.57	NR	SG 118.34 ± 29.88; RYGB 154.69 ± 35.54	NR
Clapp, Benjamin et al. [151]	MBSAQIP	2	5463	1908 RYGB; 3555 SG	2.4 RYGB and 1.9 SG	Yes	Mostly laparoscopic	5463	46.2 RYGB, 44.1 SG	NR	142 RYGB + hernia; 105 SG + hernia	Death < 1%, reoperation 3.8% RYGB, 2.1% SG
Chan, Daniel Leonard et al. [152]	Retrospective	3	45	36 RYGB and SG, 9 GB	3.3 (0–13)	Yes	Laparoscopically	45	40.3 (31.4–57)	NR	151 (60–360)	NR
Eid, G M et al. [153]	Retrospective	3	85	RYGB	3.2 (2–5)	Yes <i>n</i> = 73	Primary or small intestinal submucosa	73	50.9 (36–70.9)	NR	NR	NR
Moszkowicz, David et al. [154]	National discharge summary	2	2039 VHR 2 in the 2 years before, 3388 concomitant, 6260 VHR within 2 years after MBS	Bypass and sleeve	BS first 12.1 ± 14.5 days, VHR first 9.3 ± 8 and concomitant 7.6 ± 8.2	Yes <i>n</i> = 3388	NR	3388	NR	NR	NR	Major complications 11.1 (VHR first), 7.8% (concomitant), 16.9 BS first
Newcomb, W L et al. [155]	Retrospective	3	27	Gastric bypass (<i>n</i> = 22 open, <i>n</i> = 5 lps)	NR	No	No	0	51 (39–69)	33 (25–37)	0	0
Datta, Tejwant et al. [156]	Retrospective	3	26	Gastric bypass	2.7 (VHR)	Yes <i>n</i> = 18	8 primary, 10 prosthetic mesh	10 18	NR	NR	NR	NR
Spaniolas, Konstantinos et al. [157]	ACS-NSQIP	2	503	SG (<i>n</i> = 70) or RYGB (<i>n</i> = 433)	NR	Yes	All concomitant	503	47.2	NR	NR	NR

Table 7 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients	Intervention	Length of stay	Concomitant hernia repair (yes/no)	Type of concomitant hernia repair	BMI at VHR	EWL/TWL	OR time (min)	Complications
Morrell, David J et al. [143]	Retrospective	3	20	LSG	1.6 ± 0.8	No	No	45.6 ± 6.1	20.7 ± 12.3% TBMIL	121.2	<i>n</i> = 1 bleeding
Eid, George M et al. [158]	Retrospective	3	28	RYGB and SG	NR	Yes <i>n</i> = 20	NR	NR	NR	NR	NR
Khorgami, Zhamak et al. [159]	ACS-NSQIP	2	988	concomitant VHR matched 1:1 (988)	2.3 ± 2.7	Yes <i>n</i> = 988	Open VHR (<i>n</i> = 322), lps VHR (<i>n</i> = 666)	48 ± 9	NR	134.2 ± 52.8	Adverse events 4.8%, 30-day reoperation 3.3%
Praveen Raj, P et al. [160]	Retrospective	3	36	RYGB (<i>n</i> = 11), SG (25)	4.2 and 3.9 days	Yes	Concomitant mesh repair for VH	41.1 (RYGB) and 43.8 (SG)	NR	149 (120–210)	None
Praveenraj, Palanivelu et al. [161]	Retrospective	3	23	SG (<i>n</i> = 22), RYGB (<i>n</i> = 1)	3.3 (2–8)	Yes	IPOM	43.24	NR	112 (65–220)	NR
Borbely, Yves et al. [162]	Retrospective	3	15	SG	NR	No	No	45.3 (36.3–65.8)	33.6 (20.9–45.7)	62.5 (41–120)	1 trocar site hematoma, 1 SSI, 1 readmission for dehydration
Kaminski, D L [163]	Retrospective	3	70	Gastric restrictive procedures	NR	No	No	NR	NR	NR	NR
Marzouk, Ahmed M S M et al. [164]	Retrospective	3	15	SG	2	Yes	Laparoscopic intraperitoneal only mesh	45.2 (38.7–56.2)	NR	162.1	NR
Vitiello A. et al. [165]	Retrospective	3	20 (concomitant vs 20 (delayed))	SG	3.6 ± 0.4	Yes	Intraperitoneal mesh after fascial closing	42.7 ± 4.9 and 43 ± 5.1	NR	96 ± 17.4 versus 91.7 ± 15.2	NR

Table 7 (continued)

Author, year [ref]	Compl-ications between MBS and hernia repair	Complica-tions hernia repair	Early mor-tality	Late mortal-ity	Long-term complications	Weight loss prior to hernia repairno (kg)	Bridge to surgery yes/no	Time between hernia consultation and ultimate repair (months)	Time between MBS and hernia repair (months)	Type of hernia	Hernia repair with mesh (yes/no)	Outcomes of hernia repair	Mean follow-up after VHR (months)
Morrell, David J et al. [143]	<i>n</i> = 2 (SBO, 1 con-servative, 1 surgery)	NR	0	0	NR	Mean BMI loss 20.7 ± 12.3%	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	494.9 ± 221.2 cm ³	NR	No recur-rence	20.9 ± 16.5 LSG in patients with com-plex AWH successfully
Muolla, Muhammad et al. [144]	NR	Concurrent VHR 5.8 vs 3.8%, <i>p</i> < 0.001, greater incidence of reopera-tion 2.3% vs 1.1%, <i>p</i> < 0.001, deep surgi-cal site infection 0.7 vs 0.3%, <i>p</i> = 0.025 and sepsis 0.3 vs 0%, <i>p</i> < 0.004	0.3 vs 0.2%	0	Repeat VHR within 30 days 0.45 vs 0.006%	Mean BMI loss 20.7 ± 12.3%	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	494.9 ± 221.2 cm ³	NR	No recur-rence	20.9 ± 16.5 LSG in patients with com-plex AWH successfully
													Elevated risk of major complica-tions

Table 7 (continued)

Author, year [ref]	Complications between MBS and hernia repair	Complications hernia repair	Early mortality	Late mortality	Long-term complications	Weight loss prior to hernia repair (kg)	Bridge to surgery	Time between hernia consultation and ultimate repair (months)	Time between MBS and hernia repair (months)	Type of hernia	Hernia repair with mesh (yes/no)	Outcomes of hernia repair	Mean follow-up after VHR (months)
Morrell, David J et al. [143]	<i>n</i> = 2 (SBO, 1 conservative, 1 surgery)	NR	0	0	NR	Mean BMI loss 20.7 ± 12.3%	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	494.9 ± 221.2 cm ³	NR	No recurrence	20.9 ± 16.5 LSG in patients with complex AWH successfully
Krivan, Miss Sylvia et al. [145]	<i>n</i> = 1 adhesions between mesh and small bowel with reoperation, 6 wound infection, 6 hematoma, 4 seroma (3.8%) 5 mesh repairs	<i>n</i> = 1 adhesions between mesh and small bowel with reoperation, 6 wound infection, 6 hematoma, 4 seroma (3.8%) 5 mesh repairs	0	0	No difference between BS and BS + VHR	NR	NR	NR	NR	NR	NR	5 mesh repairs, recurrence: 12	VHR + BS feasible with low recurrence rate
Sharma, G et al. [146]	5 reoperations: SBO, sepsis, mesh dehiscence	0	0	0	Hernia recurrence 31% at 5 years primary repair, 34.8% at 5 years with mesh	NR	No	NR	NR	7.1 cm ² primary repair (<i>n</i> = 115), 12.6 cm ² mesh repair (<i>n</i> = 115)	44 with mesh, 115 primary	Hernia recurrence at 5 years primary repair, 34.8% at 5 years with mesh	5 years Primary and mesh repair acceptable
Olmi, Stefano et al. [147]	Group A 13.3% vs group B 5.9% seroma <i>p</i> < 0.05	Group A 13.3% vs group B 5.9% seroma <i>p</i> < 0.05	0	0	Recurrence 3.3% vs 2.4% and bulging 10% vs 2.4% <i>p</i> = 0.23	NR	Yes	NR	8 months–2 years	All types	Yes	Recurrence and bulging not statistically different	1 year

Table 7 (continued)

Author, year [ref]	Compl-ications between MBS and hernia repair	Complica-tions hernia repair	Early mor-tality	Late mortal-ity	Long-term compli-cations	Weight loss prior to hernia repairno (kg)	Bridge to surgery yes/ no	Time between hernia consultation and ultimate repair (months)	Time between MBS and hernia repair (months)	Type of hernia	Hernia repair with mesh (yes/ no)	Outcomes of hernia repair	Mean follow-up after VHR (months)
Morrell, David J et al. [143]	<i>n</i> = 2 (SBO, NR)	NR	0	0	NR	Mean BMI loss 20.7 ± 12.3%	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	494.9 ± 221.2 cm ³	NR	No recur-rence	20.9 ± 16.5 LSG in patients with com-plex AWH successfully
Raziel, Asnat et al. [148]	NR	Two abdominal wall hema-tomas	0	0	1 recurrence (1.8%)	NR	No	NR	NR	23 ventral, 29 umbilli-cal, 2 ventral and umbilical	Dual mesh	1 recur-rence	12 months
Chandeze, Marie-Maëlle et al. [149]	No stran-gulations, only 30 of 41 patients underwent VHR (<i>n</i> = not lose enough weight)	Chronic sepsis 7% vs 2%, SBO 0 vs 5%, wound abscess 7% vs 3%	0	0	Recurrence 6.7% vs 24% <i>p</i> = 0.048	28 ± 11%	Yes	NR	21.5 months (range 7–87)	<7 and >7 cm	Open mesh in sublay (87%) and underlay (13%)	Post-operative morbidity 37% vs 44% (<i>p</i> = 0.45)	4.6 ± 4.1 years
Praveen Raj, Palanivelu et al. [150]	NR	Seroma 3.2%, par-alytic ileus 3.84%	0	0	Recurrence 0.91%	NR	No	NR	NR	Umbilical/ perium-bilical, intraum-bilical, supraum-bilical, Swiss cheese defect	Yes	<1% recur-rence	12 months

Table 7 (continued)

Author, year [ref]	Complications between MBS and hernia repair	Complications hernia repair	Early mortality	Late mortality	Long-term complications	Weight loss prior to hernia repair (kg)	Bridge to surgery yes/no	Time between hernia consultation and ultimate repair (months)	Time between MBS and hernia repair (months)	Type of hernia	Hernia repair with mesh (yes/no)	Outcomes of hernia repair	Mean follow-up after VHR (months)
Morrell, David J et al. [143]	<i>n</i> = 2 (SBO, 1 conservative, 1 surgery)	NR	0	0	NR	Mean BMI loss 20.7 ± 12.3%	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	494.9 ± 221.2 cm ³	NR	No recurrence	20.9 ± 16.5 LSG in patients with complex AWH successfully
Clapp, Benjamin et al. [151]	NR	NR	<1%	NR	Readmission, reoperations and reoperations increased in concomitant VHR	NR	No	NR	NR	Ventral, epigastric, incisional, inguinal	NR	Readmission, reoperations and reoperations increased in concomitant VHR	30 days
Chan, Daniel Leonard et al. [152]	NR	Mesh infection 5.56%, no recurrence	0	0	Mesh infection 5.56%, no recurrence	NR	No	NR	NR	Abdominal wall	Yes	No recurrence, total rate of mesh infection 4.44%	13 months
Eid, G M et al. [153]	NR	Wound cellulitis and seroma	NR	NR	Primary repair (59), biomaterial mesh (12), deferred treatment (14)	NR	NR	NR	NR	Primary, umbilical, incisional, recurrent	Primary repair (59), biomaterial mesh (12), deferred treatment (14)	22% recurrence in primary group, no recurrence, biomaterial mesh, SBO in deferred treatment 37.5%	All pt with SBO had originally incisional VH with omental incarceration and had undergone adhesiolysis with reduction but without defect repair!

Table 7 (continued)

Author, year [ref]	Complications between MBS and hernia repair	Complications hernia repair	Early mortality	Late mortality	Long-term complications	Weight loss prior to hernia repair (kg)	Bridge to surgery yes/no	Time between hernia repair and ultimate repair (months)	Time between MBS and hernia repair (months)	Type of hernia	Hernia repair with mesh (yes/no)	Outcomes of hernia repair	Mean follow-up after VHR (months)	
Morrell, David J et al. [143]	$n = 2$ (SBO, NR)	NR	0	0	NR	Mean BMI loss $20.7 \pm 12.3\%$	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	$494.9 \pm 221.2 \text{ cm}^3$	NR	No recurrence	20.9 ± 16.5 LSG in patients with complex AWH successfully	
Moszkowicz, David et al. [154]	NR	Mesh infection 1% in VHR first, 4.3% in concomitant, 1.9% BS first	NR	NR	Overall 10-year reoperation rate for VHI: 23.3%, reoperation rate highest in VHR-first group 36.2%	NR	NR	NR	NR	Mixed	Mixed	Overall 10-year reoperation rate for VHI: 23.3%, reoperation rate highest in VHR-first group 36.2%		
Newcomb, W L et al. [155]	3.7 previous failed hernia repairs prior staged repair	$n = 7$ concomitant with 1 dehiscence of biologic mesh	NR	NR	NR	NR	Yes	NR	1.3 years (0.9–3.1)	NR	Open modified Rives–Stoppa polypropylene, lps with mesh	One patients SBO between MBS and VHR	20 months	Complex ventral hernias!
Datta, Tejwant et al. [156]	NR	2 primary with post-operative SBO, 1 chronic pain relieved by lps transection	0	0	2 primary with post-operative SBO, 1 chronic pain relieved by lps transection	NR	Yes	NR	NR	NR	Mixed	2 primary with post-operative SBO, 1 chronic pain relieved by lps transection	14 months (4–30)	

Table 7 (continued)

Author, year [ref]	Complications between MBS and hernia repair	Complications hernia repair	Early mortality	Late mortality	Long-term complications	Weight loss prior to hernia repair (kg)	Bridge to surgery	Time between hernia consultation and ultimate repair (months)	Time between MBS and hernia repair (months)	Type of hernia	Hernia repair with mesh (yes/no)	Outcomes of hernia repair	Mean follow-up after VHR (months)
Morrell, David J et al. [143]	$n = 2$ (SBO, 1 conservative, 1 surgery)	NR	0	0	NR	Mean BMI loss 20.7 ± 12.3%	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	494.9 ± 221.2 cm ³	NR	No recurrence	20.9 ± 16.5 LSG in patients with complex AWH successfully
Spaniolas, Konstantinos et al. [157]	NR	Overall morbidity 8.3%, SSI 4.6%, reoperation 3.4%	0.2%	NR	NR	NR	No	NR	NR	NR	NR	Increase in SSI during chronic VHR	30 days
Eid, George M et al. [158]	NR	NR	NR	NR	NR	NR	Mixed	NR	6 months in MBS first 2 with 9 and 18 months delayed for insufficient WL	VHR first $n = 3$; $n = 3$	2 of 28 recurrences (7%)	2 years	4 groups: favorable anatomy symptomatic/asymptomatic—unfavorable anatomy sym/asympt

Table 7 (continued)

Author, year [ref]	Compli- cations between MBS and hernia repair	Complica- tions hernia repair	Early mor- tality	Late mortal- ity	Long-term complica- tions	Weight loss prior to hernia repair (kg)	Bridge to surgery yes/no	Time between hernia consultation and ultimate repair (months)	Time between MBS and hernia repair (months)	Type of hernia	Hernia repair with mesh (yes/ no)	Outcomes of hernia repair	Mean follow-up after VHR (months)
Morrell, David J et al. [143]	<i>n</i> = 2 (SBO, 1 con- servative, 1 surgery)	NR	0	0	NR	Mean BMI loss 20.7 ± 12.3%	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	494.9 ± 221.2 cm ³	NR	No recur- rence	20.9 ± 16.5 LSG in patients with com- plex AWH successfully
Khorgami, Zhamak et al. [159]	NR	30-day reop- eration for ventral hernia repair 21.2%	0.1%	NR	NR	NR	NR	NR	NR	167 strangu- lated, 821 reducible hernia	CPT codes all together	More composite adverse event (2.7% vs 4.8%, <i>p</i> = 0.01), more return to OR in 30 days (3.3% vs 0.6%, <i>p</i> < 0.01), readmis- sion (3.2 vs 5.9%, <i>p</i> = 0.01)	30 days
Praveen Raj, P et al. [160]	NR	No recur- rence, no mesh infection	NR	NR	NR	NR	No	NR	NR	NR	YES	No recur- rence, no mesh infection	18 months RYGB, 11 months SG
Praveenraj, Palamivelu et al. [161]	NR	4 seromas	NR	NR	No mesh infection, no recur- rence	NR	No	NR	NR	2 previous repairs (1–5)	IPOM in recurrent ventral hernias	No recur- rence, no mesh infection	3.3 years (9 months–5.5 years)

Table 7 (continued)

Author, year [ref]	Complications between MBS and hernia repair	Complications hernia repair	Early mortality	Late mortality	Long-term complications	Weight loss prior to hernia repair (kg)	Bridge to surgery yes/no	Time between hernia consultation and ultimate repair (months)	Time between MBS and hernia repair (months)	Type of hernia	Hernia repair with mesh (yes/no)	Outcomes of hernia repair	Mean follow-up after VHR (months)
Morrell, David J et al. [143]	<i>n</i> = 2 (SBO, 1 conservative, 1 surgery)	NR	0	0	NR	Mean BMI loss 20.7 ± 12.3%	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	494.9 ± 221.2 cm ³	NR	No recurrence	20.9 ± 16.5 LSG in patients with complex AWH successfully
Borbely, Yves et al. [162]	None	2 pneumonia (13%), 2 infected seroma, 3 SSI	0	0	3 reoperations: small recurrence (7%), infected seroma (7%), infected mesh (7%)	TWL 25.7% (12.7–50.8)	Yes	NR	185 days (32–640)	Repair of hernia with loss of domain in 2-step approach; recurrent incisional hernia with >2 failed repairs	Mesh intraperitoneally before fascial closure	1 reoperation for recurrence, 1 for infected seroma, 1 for infected mesh	24 months (6–68)
Kaminski, D L [163]							Yes	NR				Body weight should be < 200 lb to decrease recurrent hernia formation (90 kg)	79 ± 18 months When >250 lb (>113 kg) ventral hernia recurrence of 33%
Marzouk, Ahmed M S M et al. [164]	NR	1 recurrence, 4 seroma	NR	NR	1 recurrence	NR	No	NR	NR	Mean hernia defect size 2.6 cm (1.3–4.2)	Yes	4 patients with self-limited seroma, 1 recurrence at 14 months	12 months

Table 7 (continued)

Author, year [ref]	Compl-ications between MBS and hernia repair	Complica-tions hernia repair	Early mor-tality	Late mortal-ity	Long-term complications	Weight loss prior to hernia repairno (kg)	Bridge to surgery yes/ no	Time between hernia consultation and ultimate repair (months)	Time between MBS and hernia repair (months)	Type of hernia	Hernia repair with mesh (yes/ no)	Outcomes of hernia repair	Mean follow-up after VHR (months)
Morrell, David J et al. [143]	n = 2 (SBO, 1 con-servative, 1 hernia repair surgery)	0	0	0	NR	Mean BMI loss 20.7 ± 12.3%	Yes	22.6 ± 12.5 (months)	13.5 ± 11.7 months	494.9 ± 221.2 cm ³	NR	No recur-rence	20.9 ± 16.5 LSG in patients with com-plex AWH successfully
Vitello A. et al. [165]	NR	2 recur-rences in delayed group, 1 trocar site hematoma in syn-chronous group, and 1 in delayed group	NR	NR	2 recur-rences	NR	Mixed	NR	NR	Periumbili-cal hernia	Yes onlay mesh open in delayed group	2 recur-rences	19,8 ± 5.6

[74, 75]. Both supported MBS in the elderly with a careful selection of patients.

According to the literature, although there was only one RCT, we could state that MBS is a safe and effective treatment of the elderly in carefully selected cases. In this patient population, attention must be paid to patient selection and procedure selection, considering the chance of comorbidity resolution and post-operative follow-up compliance.

Recommendation

- *MBS has been performed successfully in increasingly older patients, including patients ≥ 70 years of age. In septuagenarians, compared with a younger population, MBS is associated with slightly higher rates of postoperative complications but still provides substantial benefits of weight loss and co-morbid disease remission.*
- *Frailty, cognitive capacity, smoking status, and end-organ function have an important role in the indications for MBS.*
- *There is no evidence to support an age limit for older patients seeking MBS, but a careful patient selection that includes a frailty assessment is recommended.*

Level of Evidence 2a

Grade of recommendation B

5. MBS for the pediatrics and adolescents [76–117]
PRISMA Appendix 5 [PubMed, Cochrane, Embase]
Systematic Review Table 5

Forty-two papers have been retrieved for qualitative analysis [76–117]. One RCT [76] and 14 comparative papers [83, 88, 90, 93, 94, 96–98, 101, 104–107, 115] were found.

Seven studies about MBS versus lifestyle modifications [76, 88, 90, 94, 97, 105, 115] were evaluated. The surgical approach was more effective and durable than lifestyle modification regarding excess weight loss (%EWL), total weight loss (%TWL), and comorbidity resolutions.

Ten papers used the teen-LABS database [92–101], comparing different laparoscopic MBS procedures (AGB, RYGB, SG) to assess many aspects of MBS in pediatric and adolescent patients. All papers demonstrated an acceptable lasting %EWL with a good resolution of obesity-related complications.

Sixteen papers evaluated the efficacy of RYGB in adolescent patients [78, 85, 92, 95, 97–99, 101, 103, 105, 107, 108, 113, 115–117]; only six of them were comparative [97, 98, 101, 105, 107, 115]. All studies concluded that RYGB achieved good weight loss, improvement, and/or resolution of comorbidities in the pediatric and adolescent population with an acceptable complication rate.

A matched-control study evaluated the outcomes of MBS in Prader Willi syndrome (PWS) compared with a non-PWS group of patients and concluded that the SG is a well-tolerated, effective treatment option for PWS patients with

obesity. In both groups, the weight loss and the resolution of the comorbidities were similar [96].

Alqathani et al., in a retrospective study with 10 years of follow-up, suggested that MBS would not negatively impact pubertal development or linear growth, and therefore, a specific Tanner stage and bone age should not be considered a requirement for surgery [96].

According to a literature review, the AGB seems to be a safer procedure. However, it achieved a lower weight loss, which was less durable than the RYGB or the SG.

Recommendation

- *MBS does not negatively impact pubertal development or linear growth.*
- *MBS is safe in the population younger than 18 years and produces durable weight loss and improvement in comorbid conditions.*

Level of Evidence 1b

Grade of recommendation A

Bridge to other Treatments

6. MBS prior to joint arthroplasty [120–141]
PRISMA Appendix 6 [PubMed, Cochrane, Embase]
Systematic Review Table 6 (some studies not in favor)
Delphi Table 15

Twenty-two articles were chosen to be included in the present review.

Several studies have shown that patients with severe obesity ($\text{BMI} \geq 40 \text{ kg/m}^2$) were at increased risk of major and minor complications after joint surgery. The American Association for Hip and Knee Surgeons (AAHKS) provided a consensus opinion recommending delaying elective surgery when the BMI exceeds 40 kg/m^2 , and in 2023, adherence to these recommendations was evaluated [142]. Pre-operative health optimization programs, including weight loss with MBS before joint surgery, have been implemented to reduce postoperative complications.

However, the current literature is unclear whether persons undergoing MBS have a lower risk of postoperative complications and need for revisions after joint surgery when compared to people with obesity who have not had MBS. This systematic review (Table 6) demonstrated that only one RCT was available. Additional results were obtained from cohort studies. Some studies have demonstrated the benefits of preoperative MBS, while others have highlighted the risks of prior MBS. In addition, it seems that MBS should be performed within 2 years before joint arthroplasty to decrease the negative impact of metabolic bone disease. Furthermore,

Table 8 MBS prior to organ transplantation

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Type of surgery	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidity resolution	Post-transplant survival
J. Wesley Alexander, Hope Goodman [166]	Retrospective		32 pts with CRF-GBP without KT	GBP		48 (BMI of first group out KT)		68% EBMIL	51% T2DM	92% at 1 year
J. Wesley Alexander, Hope Goodman			9 pts GBP + KT (after)	GBP						92% at 1 year
J. Wesley Alexander, Hope Goodman			10 pts KT + GBP (later)	GBP	13 years			70.5% EBMIL		92% at 1 year
Modanlou, Muthyala, Xiao et al. [167]	Retrospective		72 before listing 188 total	GBP, VBG, restrictive procedures, BPD-DS	36.6 M (18.6, 55.7)	38.1 ± 12.4	35.1 ± 6.2	60.6 (41.5, 72.4)		
Modanlou, Muthyala, Xiao et al.	Retrospective		72 were performed pre-listing, 29 on waitlist 188 total	GBP, VBG, restrictive procedures, BPD-DS	22.6 M (11.1, 47.2)	40.1 ± 9.2	35.1 ± 10.8	60.2 (0, 68.2)		
Modanlou, Muthyala, Xiao et al.	Retrospective		87 after transplant 188 total	GBP, VBG, restrictive procedures, BPD-DS	12 M	46.6 ± 4.6	40.2 ± 7.8	30.8 (8.7, 48.3)		
Mohammad H. Jamal, Ricard Corcelles, Christopher R. Daigle et al. [168]	Retrospective		21	18 GBP/2 SG/1 AGB	27.6 M	47.1 ± 5.5	35.3 ± 8.4	60.5% ± 35.4% (%EWL)	18% T2DM remission	2/11
Anthony B. Mozer, John R. Pender, William H. H. Chapman et al. [169]	Retrospective		138	33.8% (n = 47) AGB, 48.9% (n=68) GBP, and 16.5% (n=23) LSG		45.5±8.1				
C. M. Freeman, E. S. Woodle, J. Shi et al. [170]	Prospective		52	SG	220 ± 152 days	43.0 ± 5.4	36.4 ± 5.4	29.8 ± 18.7%	37.8% (18) hypertension resolution, 30% T2DM resolution	

Table 8 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Type of surgery	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidity resolution	Post-transplant survival
Shadi Al-Bahri, Tannous K. Fakhry, John Paul Gonzalvo et al. [171]	Retrospective		16	12 GBP, 3 AGB, 1 SG	2.8 years median (1–10 years)	48 ± 8	31 ± 7	62 ± 24 (EBWL)		
Gheith, Al-Otaibi, Halim et al. [172]	Retrospective		22 after KT	Unspecified procedure	6 M	38.49 ± 9.1	34.34 ± 7.6			
Y. Kim, A. D. Jung, V. K. Dhar et al. [173]	Retrospective		20	SG before KT		41.5 ± 4.4	32.3 ± 3.1 (BMI at time of KT)		HTN 85% resolution rate; T2DM 42% resolution rate	100%
Ian A. Thomas, Jeffrey J. Gaynor, Tameka Joseph [174]	Retrospective		31	GBP	72 M post-KT	43.5 ± 0.7	28.1 ± 0.8	72.8% ± 3.0%	38.7% HTN resolution; 42% T2DM resolution	DGF was experienced by 21.4% (3/14) of RYGB (similar to control group); complication rate 35.7% (5/14) of RYGB patients vs 42.1% (control group 8/19). Death 0% vs 16 ± 8% mortality 2.3%
Cohen JB, Lim MA, Tewksbury CM et al. [175]	Retrospective		43 pre-KT BS	GBP 27/63%, SG 5/12%, AGB 5/12%, unspecified 1/2%	43 M (inter-quartile range 20–89)	43 (38–48)	32 (28–36)	69%/22% 2-year FU		
Cohen JB, Lim MA, Tewksbury CM et al.	Retrospective		21 post-KT BS	GBP 6/28%, SG 10/48%, AGB 4/19%, VBG 1/5%	130 M (inter-quartile range 74–194)	41 (39–44)	34 (33–37)	71%/59% 2-year FU		0
Renana Yemini, Eviatar Neshet, Idan Carmeli et al. [176]	Retrospective		24 pre-KT	LSG 17/GBP 7	47 months (range 0.5–5 years)	41 kg/m ² (range 35–51)	28 kg/m ² (range 19–36)	66%/29%	T2DM 44%; HTN 45%	dyslipidemia 22%

Table 8 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Type of surgery	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidity resolution	Post-transplant survival
Renana Yemini, Eviatar Neshet, Janos Winkler et al. [177]			34 after transplant	19 SG; 15 GBP; 26 KT, 1 KT + LT, 2 KT + PT, 4 LT, 1 HT		41 (range 35–48) kg/m ²		%EWL 72% (1 year); 84% (3 years)	8/30 HTN complete remission (27%); 67% improvement; 4/21 T2DM complete remission (19%); 67% improvement	
Philippe Bouchard, Jean Tchervenkov, Sebastian Demyttenaere et al. [178]	Retrospective		32	LSG	14 months (± 41)	42.3 (± 5.2) kg/m ²		56% at 1 year	T2DM remission 31%	
Jordana B. Cohen, Colleen M. Tewksbury, Samuel Torres Landa et al. [179]	Retrospective		925 with end-stage kidney disease (ESKD)	775/84% SG, 150/16% GBP		44.7 (40.7–49.6) IQR				
Hilla Schindel BSc, Janos Winkler MD, Renana Yemini MD et al. [180]	Retrospective case-control		30 after KT	19/63% SG; 10/33% GBP, 1/3.3% BPD-DS	2.5 ± 1.3	41.3 ± 3.7	29.5 ± 4.7	71.6 ± 28.7%	The total number of obesity-related comorbidities (T2D, HTN, dyslipidemia, etc.) and number of prescribed medications decreased in the bariatric surgery group (−0.7 and −2, respectively) and increased in the control group (+0.3 and +1.1, respectively) (<i>P</i> < .001).	
Al-Faraaz Kas-sam, Ahmad Mirza, Young Kim et al. [181]	Prospective		243; 198 with ESRD/45 with CKD	SG	2.3 ± 1.5 years	44.0 ± 6.3 kg/m ² (range 35.0–69.7 kg/m ²)	36.7 ± 6.6 kg/m ² (range 20.8–61.7 kg/m ²)	38.2 ± 20.3%	39% HTN resolution. 54% T2DM resolution	

Table 8 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Type of surgery	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidity resolution	Post-transplant survival
Kyle H. Sheetz, Kenneth J. Woodside, Yahakn B. Shahinian [182]	Retrospective		Surgical trends 2006–2008 N=275 (10), 2009–2013 N=1094 (41), 2014–2016 N=1329 (49)	The use of SG increased significantly from 1% in 2006 to 84% in 2016 (10% per year increase; $P=0.001$). AGB declined precipitously 1% of patients by 2016 (4% per year decrease; $P=0.004$). GBP also declined from 58% in 2006 to 13% in 2016 (4% per year decrease; $P=0.02$)						
Małgorzata Dobrzycka, Monika Proczko-Stepaniak, Lukasz Kaska et al. [183]	Retrospective		20 with ESKD	9 OAGB, 9 GBP, 1 year 2 SG	38.7		90% EBML vs 68% of non ESKD group (P n.s.)			
Loubna Outmani, Hendrikus J. A. N. Kimenai, Joke I. Roodhat et al. [184]	Retrospective		23 before KT	11 SG, 9 GBP, 3 AGB	42.3 (41.3–47.8) IQR; 33.8 (31.6–34.1) BMI at time of KT		54 (45–64) 1 year; 67 (31–76) 2 years	8.7% T2DM remission		
Basem G. Soliman, Nabil Tariq, Yi Ying Law et al. [185]	Retrospective		38	24 GBP, 14 SG	44.5 (33.4; 57.0)	35.3±3.8		51.56 ± 32.24 (–56.27; 87.76) 24 M		
Elaine Ku, Charles E. McCulloch, Garrett R. Roll et al. [186]	Retrospective		503 prior to KT	38.6% GBP, 37.8% SG; 23.7% AGB					33.3 ± 4.8	

Table 8 (continued)

Author, year [ref]	Study design	Quality assessment	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions		
Roxaneh Zaminyan, Matias Claus, Steven Paraskevas et al. [187]	Retrospective		32	SG	53 (58 months)	42.3 (5.2) kg/m ²	BMI basal	BMI follow-up	EWL/TWL	Comorbidity resolution	Post-transplant survival
Mark C. Takata, Guilherme M. Campos, and Ruxandra Cioveica [188]	Retrospective		7 on 15	GBP	15.4 M (3–24)	50			EWL 61 (range 41–75)	33% T2DM resolution rate	
Matthew Y.C. Lin, M. Mehdi Tavakol, Ankit Sarin et al. [189]	Retrospective		6 on 26	SG		48.3 kg/m ² (range 38–60.4 kg/m ²)			%EWL 50 (1 year); 66% (2 year)	7/13 54% T2DM resolution	
Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions			
J. Wesley Alexander, Hope Goodman [166]	0	0	0	0	5 experienced (9.8%). Mortality between 112 and 2869 days 10% (causes not related to surgery)		Weight loss after GBP also seems to stabilize or improve renal disease in some patients, suggesting additional benefits for GBP early in CRF. Early GBP may also allow preemptive transplantation				
J. Wesley Alexander, Hope Goodman	0	0	0	0	9 pts successfully transplanted						
J. Wesley Alexander, Hope Goodman	0	0	0	0							
Modanlou, Muthyala, Xiao et al. [167]	0										
Modanlou, Muthyala, Xiao et al.	3.5%						Given the known contributions of obesity to excess morbidity and mortality in this population, BS warrants prospective study as a strategy for improving outcomes before and after kidney transplantation				
Modanlou, Muthyala, Xiao et al.	3.5%										

Table 8 (continued)

Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions
Mohammad H. Jamal, Ricard Corcelles, Christopher R. Daigle et al. [168]	1/4.8% (GBP related to BS)	group, death unre-	14%	19%			16 patients (76%) had lost sufficient weight and were placed on the transplant list, and 2 patients (9.5%) underwent kidney transplantation; 1 patient who had a LAGB was denied because of failure of weight loss, and 2 patients were denied because of their high cardiopulmonary risk	Chronic renal failure requiring dialysis should not be considered a contraindication to bariatric surgery. Our experience with this patient population has shown excellent medium-term weight loss and an acceptable (albeit increased) risk/benefit ratio
Anthony B. Mozer, John R. Pender, William H. H. Chapman et al. [169]	0.7% (1)		5.8%					Bariatric surgery in selected patients with DDRF is safe with a low rate of 30-day morbidity and mortality
C. M. Freeman, E. S. Woodle, J. Shi et al. [170]	37.8% (18)	hypertension resolution, 30% T2DM resolution	1.9% (1)			55.8% of patients were eligible for waitlisting or living donor transplant, 6 (11.5%) were successfully transplanted	1-year mortality 3.8% comparing with 7% of overall waitlist mortality	LSG is a safe and effective means for addressing obesity in kidney transplant candidates in the context of a multidisciplinary approach

Table 8 (continued)

Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions
Shadi Al-Bahri, Tammous K. Fakhry, John Paul Gonzalezvo et al. [171]	0	0	0	0	2 deaths (12.5%) unrelated to surgery	Four patients renal transplantation 2.5–5 years post-BS. No rejection; 5 listed (31%); 2 lost in FU (31%); 2 lost in FU	25% underwent	Bariatric surgery is effective in patients with ESRD and improves access to renal transplantation. Bariatric surgery offers a safe approach to weight loss and improvement in comorbidities in the majority of patients. Referrals of transplant candidates with obesity for bariatric surgery should be considered early in the course of ESRD
Gheith, Al-Otaibi, Halim et al. [172]								<p>Graft function 95% vs 88% in obese non-BS group (44 pts). Survival rate 100% vs 95%. Statin therapy 47% vs 71%. New-onset T2DM 13.6% vs 31.8%. Mean tacrolimus dose lower in the non-bariatric group with significantly lower targeted level ($P = .02$). No significant differences in the dose or targeted cyclosporine levels</p> <p>Bariatric surgeries are feasible, safe procedures for selected obese renal transplant recipients</p>
Y. Kim, A. D. Jung, V. K. Dhar et al. [173]	0	0	0	0	15% readmission 30 days; 40% readmission 1st year	5% Hemodialysis 24 h after SG, survival rate 100%, 10% graft loss at 1 year. All values inferior compared to control group		Post-operative complications frequently experienced among obese recipients, such as DGF, NODAT, and obesity-associated conditions, are favorably impacted by surgical weight loss

Table 8 (continued)

Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions
Ian A. Thomas, Jeffrey J. Gaynor, Tameka Joseph [174]	0		25.8% (2 severe—6.4%)	—outcomes BS related	80.6% (25/31) of patients had been waitlisted for a transplant, and 14 of 31 (45%) patients received a kidney transplant at the center		Outcomes 72 M median post-KT	
Cohen JB, Lim MA, Tewksbury CM et al. [175]	1/43 died/mortality	2.3%						Increased risk of acute rejection at 1 year (aOR 1.19, 95% CI 1.07–1.33) and a decreased risk of delayed graft function (aOR .53, 95% CI .42–.68), all-cause allograft failure (aHR .31, 95% CI .29–.33), and mortality (aHR .57, 95% CI .53–.61)
Cohen JB, Lim MA, Tewksbury CM et al.			3/21 14%					Was associated with a decreased risk of all-cause and allograft failure (aHR .85, 95% CI .85–.86) and mortality (aHR .80, 95% CI .79–.82)
Renana Yemini, Eviatar Nesher, Idan Carmeli et al. [176]	2/8%							Bariatric surgery appears to be a safe and reasonable approach to weight loss, both before and after transplantation. The issues should be discussed in detail with all obese transplant candidates when they are seen for transplant evaluation. Having a kidney transplant should not be a contraindication to bariatric surgery; however, because of the possibility of kidney-related complications, close monitoring of allograft function and calcineurin-inhibitor levels is recommended. Pretransplant bariatric surgery does not seem to significantly delay access to transplantation based on the results of the present study and should be considered for this group of patients
								While the surgical risk is probably higher than that of the regular bariatric surgery population, we believe that the net advantages of the resulting weight reduction before kidney transplantation provide a convincing argument in favor of bariatric surgery for morbidly obese patients who require kidney transplantation (1/16–6.2%)

Table 8 (continued)

Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions
Renana Yemini, Eviatar Neshner, Janos Winkler et al. [177]	1/3%			3%		KT in 14 (44%) patients at a median of 8 (12) months after SG; 20 (63%) listed for KT	2/14 (14%) acute rejections. No graft loss	The acceptable safety and efficacy of SG in this high-risk population makes it an optimal choice as a bridging procedure. Furthermore, SG may allow for kidney transplantation in patients who may otherwise not be considered transplant candidates at some institutions due to their weight and body habitus
Jordana B. Cohen, Colleen M. Tewksbury, Samuel Torres Landa et al. [179]	13 (1.4%)			Reoperation (12.2%); acute kidney injury 0			113 Patients with CKD and ESKD experience higher relative risk of post-bariatric surgery complications compared to patients without CKD. Although patients with ESKD have higher mortality risk compared to patients without CKD, this is comparable to the expected increased mortality rates in the non-surgical ESKD population, and substantially lower than postoperative ESKD mortality rates previously reported in other non-emergent surgeries. Our results reinforce that CKD and ESKD should not be contraindications to undergoing bariatric surgery	

Table 8 (continued)

Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions
	Mortality 3.22/0.1%; patients with CKD had no significant difference in 30-day mortality compared to patients without CKD (adjusted OR [aOR] 1.26, 95% CI 0.59–2.72); however, patients with ESKD had a significantly increased 30-day mortality risk (aOR 8.65, 95% CI 4.78–15.68) compared to patients without CKD	Patients with both CKD and ESKD had a significantly increased risk of 30-day reoperation (CKD aOR 1.72, 95% CI 1.33–2.22; ESKD aOR 2.73, 95% CI 2.01–3.72), intervention (CKD aOR 2.49, 95% CI 2.02–3.07; ESKD aOR 2.28, 95% CI 1.67–3.10), and readmission (CKD aOR 1.78, 95% CI 1.51–2.08; ESKD aOR 2.23, 95% CI 1.82–2.74)	Reoperation 5389 (1.7%); intervention 6266 (1.9%); readmission 15,677 (4.9%); acute kidney injury 552 (0.2%)	There was no effect modification of the association between CKD status and age ≥ 60 years, hypertension, and type of bariatric surgery ($P > .05$)				
	7/0.4%			Reoperation 64 (3.8%); intervention 102 (6%); readmission 182 (10.7%); acute kidney injury 63 (3.7%)				

Table 8 (continued)

Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions
Hilla Schindel BSc, Janos Winkler MD, Renana Yemini MD et al. [180]	2/6.7% vs 16.7% in control group	2/6.7% vs 16.7% in control group	2/6.7% vs 16.7% in control group	2/6.7% vs 16.7% in control group		4.6±3.1 years between BS and KT	CKD classification improved in 30% of the bariatric surgery group (9 patients) and in 12% of the control group (6 patients) ($P = .083$). The eGFR increased in the bariatric surgery group (+13.4) and decreased in the control group (−3.9) ($P < .001$)	The results of this analysis of kidney recipients demonstrated an improvement in renal function among the patients in the bariatric surgery group, while the patients in the control group had worsening of their renal function
Al-Faraaz Kasam, Ahmad Mirza, Young Kim et al. [181]	0		1.2% readmission rate			71 achieved a BMI ≤ 40 kg/m ² and were wait-listed for KT, with 45 (63.4%) of these patients receiving KT and 10 (14.1%) remaining on the waitlist	Mortality rate among wait-listed patients after SG was 1.8 deaths per 100 patient-years, compared to 7.3 deaths per 100 patient-years on the waitlist and 5.5 nationally	SG has significant, sustainable effects on weight loss and medical comorbidities in this high-risk population and improves transplant candidacy safely and effectively, without increasing patient morbidity and mortality while on dialysis. It may also decrease the time from initial referral to KT
Kyle H. Sheetz, Kenneth J. Woodside, Vahagn B. Shahinian [182]			Readmission rate 8.6% vs 5.4% (pts without ESKD)	3.4% vs 3.6% (pts without ESKD)			GBP was associated with more complications, longer hospital stays, and more readmissions for all patients	This study suggests that laparoscopic sleeve gastrectomy has replaced Roux-en-Y gastric bypass as the most common bariatric surgical procedure in patients with ESKD. The data also demonstrate a favorable complication profile in patients with sleeve gastrectomy

Table 8 (continued)

Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions
Małgorzata Dobrzycka, Monika Proczko-Stepaniak, Łukasz Kaska et al. [183]			1/20 5% (leak after OAGB 1st day p.o. reoperated)					Morbidly obese kidney transplantation candidates benefit from bariatric surgery and can be eagerly included in bariatric surgery weight loss programs. Bariatric surgery allows efficient pretransplantation weight loss results, and the procedures in ESKD patients seem as safe as previously published
Loubna Outmani, Hendrikus J. A. N. Kimenai, Joke I. Roodnat et al. [184]	0			39% Grade IIIa; 4.3% grade IIIb; 4.3% grade IV, 0 grade V	Weight gain 1 year post-transplantation was comparable among both groups, 2 kg (−5.5 to 8.5) in the TO and 0.8 kg (−10.1 to 5.5) in the BSG	Time between BS and KT 6.87 years (5.37–9.35)	5-Year survival rate 80% vs 81% of TO group. Graft failure 26. vs 37.6% of TO group	Patients who became eligible for KTx after BS after initial rejection due to obesity have similar results of KTx as matched kidney transplant patient with obesity class II and III who were eligible while being obese. Kidney transplantation after BS does not negatively affect the outcome of KTx compared with transplanting patients with obesity class II or higher
Basem G. Soliman, Nabil Tariq, Yi Ying Law et al. [185]	0	0	Readmission 2/5.3%; reoperation 2/5.3%; bleeding 2/5.3%			8/21% transplanted; 18/47% waitlisted	No reported peri-operative morbidity or mortality, no delayed graft function, or allograft failure over follow-up period of 1–23 months	Bariatric surgery has significant, sustainable effects on weight loss and improves transplant candidacy effectively and can successfully move patients through the care pathway to transplantation

Table 8 (continued)

Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions
Elaine Ku, Charles E. McCulloch, Garrett R. Roll et al. [186]							<p>34% re-admission vs 30% of non-BS pts; 10.5% of graft failure ($N = 53$) vs 14.6% in non-BS. No association between BS and risk of death; odds of acute rejection within the first year after KT tended to be higher (OR 1.14; 95% CI 0.87–1.48) among BS not statistically significant. There was no statistically significant association between a history of bariatric surgery and risk of graft failure. There was a tendency toward higher risk of death in those with a history of bariatric surgery compared with those without, although this finding also did not achieve statistical significance</p>	

Table 8 (continued)

Author, year [ref]	Mortality	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery	Conclusions
Roxaneh Zaminpeyma, Matias Claus, Steven Paraskavas et al. [187]						27 (84%) listed for transplantation and 23 (72%) patients successfully transplanted	0% mortality, 0% SSI, readmission 90-day 4/17%; reintervention 3/13%, reoperation 2/9%; acute graft rejection 2/8.7%; chronic graft rejection 3/13%; return to dialysis 1/4.3%	
Mark C. Takata, Guilherme M. Campos, and Ruxandra Ciovea [188]	0	0	0	0	0	13 patients for whom follow-up data of 3 months were available reached our institution's BMI limits for transplantation and were undergoing their pretransplant evaluation		
Matthew Y.C. Lin, M. Mehdi Tavakol, Ankit Sarin et al. [189]	7/13 54% T2DM resolution	1/17%	0	0	0	1 pts successfully transplanted; another combined KT+LT	LSG in patients with end-organ dysfunction awaiting transplant is well tolerated, is technically feasible, provides excellent weight loss, and improves candidacy for transplantation	

Table 9 MBS for BMI ≥ 60 kg/m²

Author, year	Study design	Population (comparator)	Age (comparator)	Initial BMI (comparator)	Presence of weight-related comorbidities (comparator)	Intervention (comparator)	Length of hospital stay (comparator)
Howell 2021 [192]	Retrospective cohort study	208 people with BMI ≥ 60 kg/m ² (super obese SSO)	43	65.9 \pm 6.0	(Obstructive sleep apnea) OSA 153 (73.6%), HTN (hypertension) 123 (59.1%), gastroesophageal reflux disease) GERD 90 (43.3%), (type 2 diabetes) T2DM 62 (29.8%)	97 Roux-en-Y gastric bypasses RYGB (46%), 88 laparoscopic sleeve gastrectomies SG (42%), 23 adjustable gastric bands LAGB (11%)	2.3 \pm 0.9
Banks 2021 [193]	Prospective cohort study	21 SSO (super obese) people treated with two-stage procedure (21 SSO people treated with single-stage procedure). Overall 42	45 (44)	66.1 \pm 4 (63.7 \pm 3.9)	HTN 10 (47.6%) (9 (42.8%), T2DM 3 (14%), cardiovascular disease 3 (14%) (0), OSA 4 (19%) (8 (38%))	21 two-staged procedure (intragastric balloon for 6 months + sleeve gastrectomy) (21 single stage: 5 RYGB, 16 sleeve gastrectomy)	After definitive procedure 3.3 \pm 1.9 (2.2 \pm 0.6) <i>P</i> =.005
Nasser 2021 [194]	Retrospective cohort study	2505 LRYGB. (3510 LSG). Overall 6015 SSO	43.4 \pm 11.3 (42.8 \pm 11.3)	66.6 \pm 6.2 (66.7 \pm 6.6)	HTN 1580 (63.1%), (2131 (60.7%)) (dyslipidemia) DL 1137 (45.4%), (1349 (38.4%)) T2DM 1037 (41.4%), (1215 (34.6%)), GERD 1180 (47.1%) (1483 (42.3%)), OSA 1657 (66.2%) (2206 (62.9%))	2505 LRYGB. (3510 LSG)	NR
Mahmoud 2021 [195]	Prospective cohort study	52 SSO people. Overall 39.4 \pm 11.6 morbidly obese (MO) people	Overall 39.4 \pm 11.6	>60 for 52 patients	NR	NR	NR
Dupree 2018 [196]	Retrospective cohort study	381 Obese, 225 superobese, 109 super obese. Overall 715	40.7	67.4	T2DM 41 (37.6%), HTN 68 (62.3%), DL 10 (9.1%), OSA 31 (28.4%), GERD 20 (18.3%)	86 SG, 21 RYGB	NR
Samuel 2020 [197]	Retrospective cohort study	168 Obese, 182 super obese, 23 super super obese. Overall 353	40.9	65.03	T2DM 9 (39.1%), HTN 4 (26%), DL 6 (30.4%), OSA 2 (8.6%), GERD 4 (17.3%)	HTN 4 LAGB, 6 SG, 13 RYGB	NR

Table 9 (continued)

Author, year	Study design	Population (comparator)	Age (comparator)	Initial BMI (comparator)	Presence of weight-related comorbidities (comparator)	Intervention (comparator)	Length of hospital stay (comparator)
Stephens 2008 [198]	Retrospective cohort study	291 SSO, (3401 obese or super obese)	41	67 (46)	HTN 88 (30.2%) (1192), T2DM 52 (17.8%) (623), DL 24 (8.2%) (645), OSA 81 (27.8%) (500)	Vertical banded gastropasty–Roux-en-Y gastric bypass VBG-RYGB 130 (1073), RYGB 116 (1356), laparoscopic adjustable gastric banding LAGB 45 (972)	3 (2)
Romero-Velez 2020 [199]	Retrospective database analysis	2322 people with BMI >70 (161,091 obese BMI < 70)	40.18 ± 10.489 (44.68 ± 11.73)	76.55 ± 8.772 (45.55 ± 7.298)	HTN 58.5% (50.6%) $P = .01$, T2DM 29.4% (27.7%), OSA 34.1% (16.3%) $P = .001$	954 SG, 1368 RYGB	2.66 ± 2.66 (2.18 ± 2.231) $P = .0001$
Schwartz 2013 [200]	Prospective cohort study	20 SSO	37.5	63	HTN 13 (65%), OSA 4 (35%), T2DM 7 (20%)	RYGB	6 days
Gonzalez-Heredia 2016 [201]	Retrospective cohort study	12 RYGB (68 SG). Overall 89 SSO	44.4 (38.1)	64.2 (64.9)	HTN 9 (75%) (38 (49.4%), T2DM 6 (50%) (24 (31.2%)), DL 2 (16.7%) (23 (29.9%)), OSA 8 (66.7%) 34 (44.2%)	RYGB (SG)	3 (3.7)
Ochner 2013 [202]	Retrospective cohort study	BMI 35–39.9; $n = 232$; BMI 40–49.9; $n = 1166$; BMI 50–59.9; $n = 429$; BMI ≥ 60; $n = 166$	NR	NR	NR	Not relevant for the study	NR
Arapis 2019 [203]	Retrospective cohort study	210 SSO; 91 SG (RYGB)	44.9 ± 11.4 (39.7 ± 9.9) $P = .0032$	68.2 ± 7.1 (65.1 ± 4.3) $P = .0003$	T2DM 27 (29.6%) (38 (31.9%)), HTN 39 (42.8%) (50 (42.1%)), OSA 65 (68%) (75 (61.4%)), GERD 20 (21%) (38 (31.1%))	SG (RYGB)	7 (12) $P = .003$
Serrano 2016 [204]	Retrospective cohort study	135 SSO; 93 RYGB (LSG)	42.33 ± 11.5 (38.2 ± 11.3)	66.3 ± 5.4 (68.4 ± 7.9)	T2DM 26 (28.0%) (9 (21.4%)), DL 28 (32.2%) (10 (23.8%)), GERD 17 (18.3%) (10 (23.8%)), HTN 60 (64.5%) (20 (50.0)), OSA 43 (46.2%) (27 (64.3%))	RYGB (SG)	3.02 ± 2.8 (3.40 ± 2.7)

Table 9 (continued)

Author, year	Study design	Population (comparator)	Age (comparator)	Initial BMI (comparator)	Presence of weight-related comorbidities (comparator)	Intervention (comparator)	Length of hospital stay (comparator)
Mehaffey 2015 [205]	Retrospective database analysis	328 SSO (1681 non-SSO). Overall 2009.	41.10 ± 9.92 (42.88 ± 9.98)	67.0 ± 6.53 (50.9 ± 9.21)	GERD 28.66% (31.11%), OSA 45.73% (28.47%), HTN 60.67% (55.30%), T2DM 31.10% (32.80%)	RYGB on SSO (RYGB on non-SSO)	3.20 ± 3.27 (2.48 ± 2.02) <i>P</i> =.0341
Nasser 2019 [206]	Retrospective database analysis	18,861 SSO (65,565 SO, 272,195 MO). Overall 356,621	41.3 ± 11.1 (SO 42.7 ± 11.8; MO 45.2 ± 11.9)	66.7 ± 7.44 (SO 53.9 ± 2.75; MO 42.4 ± 3.87)	GERD: 5265 (27.9%) (MO: 85,696 (31.5%); SO: 19,052 (29.1%)). HTN: 10,909 (57.8%) (MO: 128,184 (47.1%); SO: 34,228 (52.2%)). DL: 3775 (20.0%) (MO: 67,448 (24.8%); SO: 14,115 (21.5%)). T2DM: 5481 (29.1%) (MO: 69,697 (25.6%); SO: 18,241 (27.8%)). OSA: 10,460 (55.5%) (MO: 96,745 (35.5%); SO: 30,440 (46.4%)). <i>P</i> <.001	LSG: 12,725 (67.5%) (MO: 198,656 (73.0%); SO: 44,078 (67.2%)) RYGB: 6136 ± 1.19; (MO: 73,539 ± 1.66; SO: 2,06 ± 2.47 (MO: 1.97 (27.0%); SO: 21,487 (32.8%))	LSG: 1.79 ± 2.28 (MO: 1.57 ± 1.28; SO: 1.62 ± 1.19). RYGB: 2.27 ± 2.47 (MO: 1.97 ± 1.66; SO: 2.06 ± 2.47). <i>P</i> <.001
Thereaux 2015 [207]	Prospective cohort study	30 SSO (60 MO)	42.0±12.4 (41.8±11.5)	64.1±4.1 (46.3±5.6)	HTN: 76.7% (53.3%) T2DM: 40% (40%) OSA: 76.7% (56.7%) DL: 30% (45%)	RYGB	NR
Peraglie 2008 [208]	Retrospective cohort study	16 SSO	40	62.4	NR	One anastomosis gastric bypass (OAGB)	NR
Madhok 2016 [209]	Retrospective cohort study	19 SSO OAGB (56 SSO SG). Overall 75 SSO	45 (51)	67 (65)	T2DM: 6 (17); HTN: 8 (28); OSA: 3(12)	OAGB (LSG)	2 (2)
Gegner 2008 [210]	Retrospective cohort study	63 SSO	43	68	NR	LSG	5
Taylor 2006 [211]	Retrospective cohort study	60 SSO (444 MO)	40 (37)	≥60 (<60)	HTN: 3 (5%) (20 (4.5%)); T2DM: 15 (25%) (58 (13%)); OSA: 5 (8.3%) (26 (5.8%))	Open RYGB	5.1 (4.8)

Table 9 (continued)

Author, year	Study design	Population (comparator)	Age (comparator)	Initial BMI (comparator)	Presence of weight-related comorbidities (comparator)	Intervention (comparator)	Length of hospital stay (comparator)
Wilkinson 2019 [212]	Retrospective database analysis	5723 SSO (24,940 SO)	41.65 ± 11.88 (42.63 ± 11.22)	63.67 ± 2.75 (53.91 ± 2.7)	GERD: 1597 (27.9%) (7380 (29.6%)); HTN: 3230 (56.4) (12,893 (51.7%)); DL: 1159 (20.3%) (5309 (21.3%)); T2DM: 1687 (29.5%) (6856 (27.5%)); OSA: 3184 (55.6%) (11,620 (46.6%))	RYGB and LSG	NR
Abeles 2009 [213]	Retrospective cohort study	95 SSO (1311 MO)	42.8 ± 11.8 (42.6 ± 11.2)	64.9 ± 5.3 (45.8 ± 5.7)	NR	RYGB	3.1 ± 1.6 (3.1 ± 2.9)
Parmar 2017 [214]	Retrospective cohort study	19 SSO OAGB, 47 SSO RYGB	45 (47)	67 (64.4)	T2DM: 7 (36.8%) (22 (46.8%)); HTN: 8 (42.1%) (24 (51.1%))	OAGB (RYGB)	2 (2)
Artuso 2004 [215]	Retrospective cohort study	21 SSO (61 SO)	42 ± 10 (43 ± 9)	≥ 60 (< 60)	NR	RYGB	6.6 ± 6 (5.3 ± 3) P < .05
Ece 2018 [216]	Retrospective cohort study	28 SSO (52 SO, 83 MO)	42.5 ± 10.4 (SO: 40.1 ± 11.2; MO: 39.2 ± 14.6)	64.1 ± 3.0 (SO: 55.3 ± 2.7; MO: 44.8 ± 2.4)	T2DM: 15 (53.5%) (SO: 22 (42.3%); MO: 33 (39.7%)); HTN: 19 (67.8%) (SO: 23 (44.2%); MO: 32 (38.5%)); DL: 11 (39.2%) (SO: 17 (32.6%); MO: 25 (30.1%)). P < .05	SG	NR
Sachenz-Santos 2006 [217]	Prospective cohort study	70 SSO (184 SO, 483 MO)	42.2 ± 10.8 (SO: 42.1 ± 11.5; 40 ± 2.6)	66 (SO: 54, MO: 44) Extrapolated from figures	HTN: 32.9% (SO: 34.8%; MO: 29%); OSA: 64.3% (SO: 33.1%; MO: 18.7%); T2DM: 30.4% (SO: 30.9%; MO: 16.9%); DL: 45.7% (SO: 38.7%; MO: 30.5%) P < .05 except for HTN	RYGB	6.3 ± 4.7 (SO: 5.8 ± 7.5; MO: 4.6 ± 2.6)

Table 9 (continued)

Author, year	Study design	Population (comparator)	Age (comparator)	Initial BMI (comparator)	Presence of weight-related comorbidities (comparator)	Intervention (comparator)	Length of hospital stay (comparator)
Schmitz 2022 [218]	Retrospective cohort study	243 SSO, 93 SG, 150 OAGB	39.1 OAGB, 41.5 SG	65.2 (SG: 66.91; OAGB: 64.14)	OSA: SG: 69 (74.2%); OAGB: 113 (75.3%); T2DM: SG: 42 (45.7%); OAGB: 51 (34.0%); HTN: SG: 69 (74.2%); OAGB: 98 (65.3%)	OAGB (SG)	OAGB: 3.4 (SG: 4.5)
Moon 2016 [219]	Retrospective cohort study	230 SSO (661 SO, 1555 MO)	40.1±10.3 (SO: 42.1±10.8; MO: 42.3±11.0)	66.2±6.6 (SO: 54.1±2.7; MO: 44.6±7.7)	Number of comorbidities: 1.4±1.2 (SO: 1.5±1.2; MO: 1.5±1.2)	RYGB	NR
Farkas 2005 [220]	Prospective cohort study	46 SSO (167 MO)	40 (37)	67 (48)	T2DM: 10 (22%) (31 (19%)); HTN: 22 (48%) (52 (31%)); OSA: 9 (20%) (15 (9%))	RYGB	NR
Gould 2006 [221]	Retrospective cohort study	28 SSO (260 MO)	43.8 (45.2)	62.0 ± 2.3 (48.3 ± 5.4)	HTN: 18 (64%) (139 (54%)); T2DM: 8 (29%) (60 (23%)); DL: 6 (21%) (86 (33%)); OSA: 11 (39%) (83 (32%))	RYGB	2.3 (2.2)
Kushmir 2010 [222]	Retrospective cohort study	21 SSO (147 MO)	46.7 (44)	68.6 (45.1)	Mean number of comorbidities: 4 (4)	RYGB	NR
Oliak 2002 [223]	Retrospective cohort study	39 SSO (261 MO)	41 (41)	66 (48)	HTN: 10 (26%) (101 (39%)); T2DM: 4 (10%) (43 (16%)); OSA: 11 (28%) (57 (22%)); DL: 3 (8%) (30 (11%))	RYGB	NR
Fielding 2003 [224]	Retrospective cohort study	76 SSO	39	69 ± 6.2	HTN: 16 (21%); T2DM: 4 (5.2%); OSA: 6 (7.9%)	LAGB	3
Hering 2022 [225]	Retrospective cohort study	26 SSO with balloon (52 SSO without balloon)	48.24 ± 10.2 (47.87 ± 10.1)	69.26 (64.07) P<.01	T2DM: 15 (57.7%) (21 (41.2%)); HTN: 25 (96.2%) (49 (94.2%))	10 Balloon + RYGB (17); 16 balloon + sleeve (35)	7.15 ± (7.67 ± 2.72)
Wang 2014 [226]	Retrospective cohort study	26 SSO	32.1	65.8	DL 25 (96.2%); OSA 14 (53.8%); T2DM 11 (42.3%); HTN 6 (23.1%); GERD 4 (15.4%)	RYGB	8.9

Table 9 (continued)

Author, year	Study design	Population (comparator)	Age (comparator)	Initial BMI (comparator)	Presence of weight-related comorbidities (comparator)	Intervention (comparator)	Length of hospital stay (comparator)
Singhal 2022 [227]	Multicenter observational cohort study	155 SSO (905 SO; 6024 MO)	39.3 ± 11.7 (SO: 41.2 ± 12.5; MO: 40.3 ± 11.8)	>60	<i>P</i> < .001 for all comorbidities among people with BMI >60. T2DM: 48 (31%); HTN: 77 (49.7%); OSA 51 (32.9%); DL: 42 (27.1%)	LSG 111 (71.6%); OAGB 23 (14.8%); RYGB 12 (7.7%); others 9 (5.8%)	NR
Myers 2006 [228]	Retrospective cohort study	53	39.7	66	OSA 64%; HTN 42%; GERD 28%; DL: 23%; T2DM 19%	LAGB	2.7
Tichansky 2005 [229]	Retrospective cohort study	45 SSO (640 MO)	41.6 ± 10.1	65.6 ± 5.3 (47.8 ± 5.2)	HTN 66.7% (49.7%); OSA 44.4% (27.5%); GERD 55.6% (26.7%)	RYGB	6.4
Zerrweck 2012 [230]	Retrospective case-control	23 SSO balloon + RYGB (37 SSO RYGB)	44 ± 10.8 (44.9 ± 10.6)	65 ± 3.8 (66.6 ± 6.7)	T2DM: 8 (34%) (10 (73%)); HTN: 17 (73%) (22 (59%)); OSA: 8 (34%) (16 (43%))	Intra-gastric balloon prior to RYGB (RYGB alone)	5.4 ± 2.4 (7.3 ± 6.6)
Catheline 2012 [231]	Prospective cohort study	30	35	66	OSA 17 (58%); HTN 15 (50%); T2DM 13 (42%); DL 13 (42%)	LSG	7.5
Torchia 2009 [232]	Prospective cohort study	95	38.5 ± 13.5	62.5 ± 4.2	HTN: 44 (49.3%); T2DM: 11 (12.6%); OSA: 35 (39.4%); DL: 22 (29.4%)	LAGB	NR
Date 2013 [233]	Retrospective cohort study	28 SSO (23 MO)	44 (48)	67 (42)	OSA: 14 (7%); HTN: 16 (18%); DL: 6 (13%)	RYGB	NR
Di Betta 2008 [234]	Retrospective cohort study	32	37 ± 9.3	70.1 ± 5.3	T2DM 9 (28.1%); HTN 12 (37.5%); DL 17 (53.1%); OSA 9 (28.2%)	Duodenal switch associated with transitory vertical gastroplasty (DS-TVG)	12
Spyropoulos 2007 [235]	Retrospective cohort study	26	40.8 ± 8.1	65.3 ± 9.8	OSA 21 (81%); T2DM 18 (69%); HTN 7 (27%). Mean comorbidities per patient: 4.33 ± 1.12	Intra-gastric balloon BIB	2

Table 9 (continued)

Author, year	Study design	Population (comparator)	Age (comparator)	Initial BMI (comparator)	Presence of weight-related comorbidities (comparator)	Intervention (comparator)	Length of hospital stay (comparator)
Gottig 2009 [236]	Retrospective cohort study	109	39.1±8.4	68.8±8.9	HTN 74 (67.8%); T2DM 53 (48.6%); OSA 39 (35.7%)	Intra-gastric balloon BIB	NR
Shuhabier 2004 [237]	Retrospective cohort study	25 SSO (25 SO)	38 (38.7)	78 (55)	OSA: 16 (18); HTN: 11 (12); GERD: 3 (3); T2DM: 6 (7)	11 Open RYGB	4.56 (3.04)
Fazylov 2005 [238]	Retrospective cohort study	102 SSO (283 SO)	38.75 (39.35)	68.45 (51)	HTN 55% (44.25%); T2DM 22.5% (24.3%); OSA 80.3% (49.8%); DL 29.4% (35.3%)	Open duodenal switch	NR
Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator)	Long-term complications	Summary of findings	Additional notes
Howell 2021 [192]	30 days	NR	NR	11 complications (5.2%). 1 anastomotic leak (0.48%). No mortality. Complications occurred in 14.8% of conversion/revision cases and only 3.9% in primary cases ($P = .0395$)	NR	Bariatric surgery is feasible in patients with SSO (super super obesity). Revision procedures may increase risk of operative complications	
Banks 2021 [193]	12 months after definitive procedure	BMI 48.4 ± 1.9 (43.5 ± 2) $P=0.57$; %EWL 45 (41.5) $P=47$. %EBMIL 43 (51.3) Δ BMI 17.7	NR	7. 1 port site hernia (1) NR	NR	Routine use of an intra-gastric balloon in super-super obese patients is not required and may be associated with poorer peri-operative outcomes and delayed weight loss	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Nasser 2021 [194]	12 months	%EWL 55.5 ± 14.2 (47.5 ± 14.3) $P < .01$	T2DM 74.1% (58.2%) $P < .01$, HTN 52.9% (48.3%) $P = .10$, DL 62.7% (53.3%) $P = .01$, OSA 52.3% (43.5%) $P < .01$, RGE 78.4% (68.9%) $P = .04$	Any complications 14.5% (6.8%) $P < .01$, leak or perforation 1.0% (0.3%) $P < .01$, bowel obstruction 2.9% (0.4%) $P < .01$, bleeding 2.4% (1.1%) $P < .01$, surgical site infection 5.7% (1.4%) $P < .01$, rate of reoperation 2.8% (1.2%) $P < .01$	NR	<p>LRYG was associated with better weight loss and medication discontinuation 1 year following surgery at the expense of an increase in perioperative complications and resource utilization compared to LSG</p>	
Mahmoud 2021 [195]	NR	NR	NR	NR	NR	<p>Neither super obesity nor super super obesity is associated with difficult intubation or difficult mask ventilation. High STOP-Bang and Mallampati scores are the independent factors of possible difficult intubation in patients undergoing bariatric surgery</p>	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Dupree 2018 [196]	30 days	NR	NR	2 major complications among super obese. 1 leak 1 death	NR	BMI was significantly lower in patients with complications, ($P < .05$), whereas patients' age was significantly higher ($P < .05$) in complication cohort. BMI showed an inverse correlation to the patients' age at surgery ($P < .05$). Conclusion: Super super obesity should not be considered as a limiting factor for bariatric surgery outcome	
Samuel 2020 [197]	24 months	BMI 50.8. %EBMIL 35.54; Δ BMI 14.23	NR	0	NR	The mid-term results for weight loss and resolution of obesity-related comorbidities are best achieved in super-obese patients undergoing LRYGB, without any significant increase in complications with this procedure as compared with LAGB and LSG	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Stephens 2008 [198]	30 days	NR	NR	1 death (4 deaths) no significance	NR	Super-super obese patients required longer total operating room times, a longer hospital length of stay, and were more likely to be discharged to chronic care facilities than were patients with a BMI 60 kg/m ² ; however, the in-hospital mortality was similar for both groups	Longer operating time for SSO. SSO people more likely to be discharged to chronic facilities
Romero-Velez 2020 [199]	30 days	NR	NR	9 Mortality 0.4% (0.1%) NR P=0001, 25 sepsis 1.1% (0.4%) P=.001, 15 septic shock 0.6% (0.2%) P=0001, 67 reoperation 2.9% (2%) P=.003	NR	BMI > 70 is associated with higher morbidity and mortality, still relatively low	
Schwartz 2013 [200]	12 months	BMI 42.9, %EBMIL 52.89; ΔBMI 20.1	HTN 69%, T2DM 75%, OSA 43%	2 minor complications (10%), 1 death (5%)	NR	LGBP appears feasible and effective for SSO, both in terms of weight loss and improvement of comorbidities. Rate of complications is considered low	
Gonzalez-Heredia 2016 [201]	24 months	%EWL 68.5 (45.8) P=.014	NR	0 (2) no deaths	NR	SG and RYGB appear to be viable procedures for SSO patients. RYGB provides a significantly higher %EWL and %WL at 12 and 24 months compared to SG	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Ochner 2013 [202]	36 months	%EWL (extrapolated): BMI 35–39.9: 88%; BMI 40–49.9: 65%; BMI 50–59.9: 46%; BMI ≥60: 38%. <i>P</i> <.0005	NR	NR	NR	A dosage effect of preoperative BMI was apparent, with heavier individuals showing lower percentages of initial and excess weight loss, regardless of BMI above or below 60 kg/m ²	
Arapis 2019 [203]	60 months	BMI 46.07 (42.15), %EBMIL 51.22 (57.23); ΔBMI 22.13 (22.95) <i>P</i> not significant	Extrapolated: T2DM 32% (68%), HTN 38% (22%), GERD 82% (52%) <i>P</i> =.05	6 (6.6%) (14(11.7%), 1(5), 1 death (1). <i>P</i> =.02	26% of patients of the RYGB group. gallstone formation (6.7%), marginal ulcers (5.8%), internal hernias (5.4%), and anastomotic strictures (3.3%). The SG group had a 16.1% complication rate (<i>P</i> = .04). The most common was newly acquired GERD syndrome (7.6%)	SG as a primary procedure for SSO patients remains effective even though RYGB achieves better midterm-outcomes. RYGB associated with higher complications rate, both short and long term	
Serrano 2016 [204]	12 months	BMI 43.6 ± 5.4 (46.9 ± 6.8), %EBMIL 54.96 (49.53); ΔBMI 22.7 (21.5)		RYGB: 15.1% of patients. Occlusion (5.4%), wound infection (4.3%), leak (2.2%), stricture (2.2%), bleeding (1.1%) and pulmonary embolism (2.2%). 1 death. SG: 4.8% of patients. Occlusion (2.4%) and pulmonary embolism (2.4%)	NR	Bariatric surgery is feasible in the SSO patients with comparable EWL outcomes and postoperative complications to historical non-SSO patients. Higher complication rate among RYGB patients	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Mehaffey 2015 [205]	48 months	%EBMIL 61.6 (69.36) P=.018	NR	11.4% (7.97%), 7.93% (7.72%), Mor- tality 0.61% (0.4%)	NR	RYGB is a safe operation for super-obese patients with BMI >60 kg/m ² in experienced centers. SSO patients still have significant reduction in excess BMI despite being less than non-SSO patients undergoing RYGB	
Nasser 2019 [206]	30 days	NR	NR	<u>LSG</u> : 582 (4.57%) (MO: 5060 (2.55%); SO: 1482 (3.36%)). <u>RYGB</u> : 532 (8.67%) (MO: 4,355 (5.92%); SO: 1419 (6.60%)). Mortality: 23 (.18%) (MO: 82 (.04%); SO: 35 (.08%))	NR	SO and SSO patients are at increased risk of 30-day morbidity and mortality compared with MO patients. Despite this elevated perioperative risk, the overall risk of these procedures remains low and acceptable especially as bariatric surgery is the durable treatment option for obesity	
Thereaux 2015 [207]	60 months	BMI: 46.6±8.6 (BMI: 32.6±5.9); %EBMIL 44.7 (64.3); ΔBMI 17.5 (13.7)	HTN: 56.5% (65.6%) T2DM: 83.3% (75.0%) OSA: 60.9% (82.4%) DL: 66.7% (70.4%)	Major adverse events 0% (7.1%)	NR	RYGB is associated with similar and beneficial long-term effects for SSO as for non SSO patients with regard to percentage of weight loss, diabetes and hypertension risks at 5 years	
Peraglie 2008 [208]	24 months	%EWL 65	NR	No complications	NR	OAGB seems safe and effective among SSO people	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Madhok 2016 [209]	24 months	%EWL 66 (38)	T2DM 66% (52%) HTN0 (3) 12.5% (14.2%) OSA 66% (58%)	4 (5.8%)	2 (5)	OAG yields superior weight loss in comparison with SG in obese patients with BMI ≥ 60 kg/m ² without an increase in early complication rate	
Gegner 2008 [210]	12 months	BMI 50%EBMIL41.8; Δ BMI 18	NR	4 (5.8%)	NR	SG represents an excellent procedure for SSO to achieve good weight loss	
Taylor 2006 [211]	30 months	NR	NR	2 (3.33%) (38 (7.65%)) (2 deaths in the control group)	Anemia 7 (11.7%) (37 (8.3%))	Super-super-obese patients should not be excluded from RYGBP because of a perceived increased risk based upon BMI	
Wilkinson 2019 [212]	30 days	NR	NR	Unplanned ICU admission: 63 (1.1%) (185 (0.7%)) $P=$.006; re-admission: 276 (4.8%) (1038 (4.2%)) $P=$.03; mortality: 10 (0.2%) (33 (0.1%)) $P=$.44. Among SSO patients who underwent RYGB experienced more complications compared to those who underwent LSG: Re-admission 116 (6.4) 160 (4.1) $<$.0005 Intervention 53 (2.9) 47 (1.2) $<$.0005 Re-operation 41 (2.3) 33 (0.8) $<$.0005		Patients with SSO undergoing LRYGB or LSG have an increased risk of post-operative 30-day complications compared to patients with SO. For patients with SSO, LSG may be the preferred procedure of choice to counter the increased peri-operative risk associated with multiple pre-operative co-morbidities affording a lower 30-day post-operative complication profile compared to LRYGB	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Abetes 2009 [213]	30 days	NR	NR	Overall complications: 13 (13.7%) (167 (12.7%)) $P=.8$; Death: 0 (3 (0.2)) $P=.08$	NR	RYGB is safe for SSO. Overall, there were no increased risks of intraoperative complications or postoperative morbidity or mortality as compared with MO patients	
Parmar 2017 [214]	24 months	%EWL 70.4 (54.7–87) (57.1 (16–104)) $P=.01$	T2DM 3/7 (42.9%) (13/22 (59.1%)) $P=.45$ Resolution of hypertension 2/8 (25.0%) 11/24 (45.8%) $P=.30$	1 (0)	6 (2)	OAGB/MGB yields superior weight loss at 18- and 24-month follow-ups in comparison with the gold standard RYGB in patients with BMI ≥ 60 kg/m ²	
Artuso 2004 [215]	12 months	%EWL 47 (61) $P<.05$	NR	1 minor, 2 major, 1 death (6 minor, 1 major)	NR	BMI >60 kg/m ² have a higher risk of major complications after LRYGBP than patients with BMI <60. BMI >60 has acceptable morbidity and should not be considered contraindicated to laparoscopic RYGB	
Ece 2018 [216]	41 months	BMI 45.0 \pm 2.8 (SO: 37.0 \pm 1.9; MO: 31.7 \pm 2.1). %EBMIL 48.8. Δ BMI 19.0 \pm 3.1 (SO: 18.2 \pm 2.8; MO: 13.0 \pm 2.7). $P<.012$	T2DM: 11 (73.3%) (SO: 19 (79.1%); MO: 27 (81.8%)) HTN: 14 (73.6%) (SO: 20 (80.0%); MO: 25 (78.1%)) DL: 9 (81.8%) (SO: 15 (83.3%); MO: 20 (87.0%))	5 (17.8%) (SO: 7 (13.4%); MO: 9 (10.8%))	NR	LSG is a safe standalone bariatric surgical procedure to use for the resolution of comorbidities in MO, SO, and SSO patients	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Sachenz-Santos 2006 [217]	60 months	BMI 36 (SO: 30, MO: 29) Extrapolated from figures, %EBMIL 46.8, ΔBMI 30	OSA: 96% (MO: 88%)	Leak: 3.2% (SO: 4.7%; MO: 3.7%); reoperation: 0 (SO: 4.1%; MO: 2.2%); mortality: 1.6% (SO: 1.4%; MO: 0) P =.005	Incisional hernia among laparoscopic RYGB	among RYGB achieved significant durable weight loss and effectively treated co-morbid conditions in SO and SSO patients with acceptable post-operative morbidity and slightly greater mortality than in MO patients	
Schmitz 2022 [218]	36 months	%BMIL: OAGB: 27.4 (SG: 22).	OSA: 50%; T2DM: 30%; OAGB: 22%; HTN: SG:22%; OAGB: 50%	Clavien-Dindo >3: 2.7% (11.8%)	Insufficient weight loss/weight regain: OAGB: 14 (33.3%) (SG: 28 (66.7%)) P<.001; Ulcer: OAGB: 11 (7.3%) (SG: 1 (1%)) P=.033	OAGB can be considered a safe and effective option in the treatment of SSO patients and can possibly even be considered superior to LSG in these patients	
Moon 2016 [219]	21.6 months	%EBMIL 59.1 (SO: 65.9; MO: 80.9)	Number of comorbidities: 1.1	Readmission rate: 11.7% (SO: 9.2%; MO: 10.7%) P=.07. Reoperation rate: 6.1% (SO: 5%; MO:7.3%) P<0.04. Mortality rate: 0.4% (SO: 0.2%; MO: 0.1%) P=0.56	NR	Readmission and reoperation rates were similar in BMI 40–50, 50–60, and ≥60 kg/m ² group. Super-obese and super-super obese patients are not at greater risk for surgical complications when compared to those with lower BMIs	
Farkas 2005 [220]	24 months	%EWL 57 ± 7 (67 ± 18)	NR	Major complications: 5 (9); minor complications: 4 (10)	NR	LR YGBP can be performed safely and effectively in super-obese patients (BMI >60). Although these patients have less %EWL than lighter patients, they still end up with a good result	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Goold 2006 [221]	24 months	%EWL 60.8 ± 10.1 (70.9 ± 15.4)	T2DM: 100% (80%); HTN: 63% (89%); DL: 80% (92%); OSA 91% (92%)	1 (3.6%) leak; 1 (3.6%) wound infection	8 (28%) anastomotic stenosis (24 (9.2%)) $P < .01$; 3 (10.7%) marginal ulcers	Laparoscopic Roux-en-Y gastric bypass can be accomplished safely even in extremely obese patients. Although excess weight loss in the super obese is diminished postoperatively when compared with less obese patients, health is improved and quality of life is good regardless of a patient's preoperative BMI	
Kushmir 2010 [222]	30 days	NR	NR	Major complications: 0 (2(1.4%)); minor complications: 1 (4.8%) (18 (11.8%))	NR	Patients with BMI ≥ 60 kg/m ² do not have a higher postoperative morbidity compared with other patients undergoing laparoscopic Roux-en-Y gastric bypass	
Oliak 2002 [223]	30 days	NR	NR	Major complications: 4 (10%) (17 (6%)); 2 deaths (5%). Minor complications: 8 (21%) (34 (13%))	NR	Laparoscopic RYGBP is feasible for patients with BMI ≥ 60 . Our data suggest that these patients are at a higher risk for GI leak, postoperative infection, and death	
Fielding 2003 [224]	60 months	BMI 35.09 ± 5.37 %EBMIL 77.27; ΔBMI 34	HTN: 9 (56%); T2DM: 4 (100%); OSA: 4 (66.6%)	2 minor (2.63%)	NR	Laparoscopic adjustable gastric banding is a valid surgical approach for mega obese patients	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Hering 2022 [225]	24 months	BMI 51.15 ± 6.99 (42.25 ± 6.62) %EBMIL 40.9; ΔBMI 18 (%EMBIL 56.4; ΔBMI 22)	NR	1 (3.8%) (1 (1.9%))	NR	No differences among complications, lower %EBMIL and ΔBMI in the balloon + surgery group	
Wang 2014 [226]	30 months	BMI 35.5±4.1 %EBMIL 74.2; ΔBMI 30.3	DL 20 (80%); OSA 8 (57%); T2DM 8 (72%); HTN 4 (66%); GERD 0 (100%)	2 minor (7.7%)	6 minor (23%)	LRYG is feasible for Chinese super-obese patients, with significant short-term results	
Singhal 2022 [227]	30 days	NR	NR	12 (7.7%) <i>P</i> not significant, 3 deaths (1.9%) (SO: 0.1%; MO: 0.1%), <i>P</i> =0.001	NR	The 30-day mortality rate was significantly higher in patients with BMI >60 kg/m ² . There was, however, no significant difference in complications rates in different BMI groups	
Myers 2006 [228]	18 months	BMI 47.3 %EBMIL 45.6; ΔBMI 18.7	NR	One band removal for chronic obstruction (1.9%), one band revision for slip (1.9%), and one nonfatal pulmonary embolism (1.9%)	NR	LAGB is an appropriate surgical option for treatment of massive super-obesity. The procedure can be performed with minimal morbidity and mortality and leads to promising medium-term weight loss	
Tichansky 2005 [229]	12 months	%EWL 58 ± 13 (70 ± 16)	NR	Marginal ulcer 2 (4.4%) (25 (3.9%)); stomal stenosis 3 (6.7%) (18 (2.8%)); SBO requiring operation 3 (6.7%) (12(1.9%)) 0.07. Anastomotic leak 1 (2.2%) (26 (4.1%)); internal hernia 1 (2.2%) (5 (0.8))	NR	The complication and mortality rates are not increased in super-obese patients who undergo RYGB, with acceptable weight loss	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Zerrweck 2012 [230]	12 months	%EBMIL 52.4±17.3 (50.3±12.7)	NR	2 (8%) (4 (11%))	NR	IGB prior to LGBP in super-super obese patients significantly reduced excess BMI. It was associated with a shorter operative time and a lower overall risk of significant adverse outcomes	
Catheline 2012 [231]	24 months	BMI 45 %EBMIL 51.2 ΔBMI 21	NR	4 (14%): two subphrenic hematoma, one leak, one pulmonary embolism	2 (7%) port site hernia, 6 (20%) reintervention for insufficient weight loss	Laparoscopic sleeve gastrectomy is a safe and efficient procedure for treating super obesity. In the case of insufficient weight loss, a second-stage operation like resleeve gastrectomy or gastric bypass can be proposed	
Torchia 2009 [232]	48 months	BMI 28.9; %EBMIL 89.6 ΔBMI 33.6	HTN: 86%; T2DM: 90.9%; OSA: 100%; DL: 54.5%	2 minor	0	LAGB can be considered an appropriate bariatric surgical option in super-super obese patients, both for low morbidity rate and weight loss	
Date 2013 [233]	12 months	BMI 45 %EBMIL 52.3 ΔBMI 22 (28 %EBMIL 82.35 ΔBMI 14)	86% (79%)	2 (2)	NR	Higher %EBMIL but lower ΔBMI for MO compared to SSO	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Di Betta 2008 [234]	48 months	BMI 37.5±7.5 %EBMIL 72.3 ΔBMI 32.6	Improvement 100%	Major: Gastrointestinal bleeding 1 (3.1%) Abdominal rupture after laparotomy 1 (3.1%) Acute pancreatitis 1 (3.1%) Pulmonary embolism 2 (6.2%) Minor: Urinary infection 1 (3.1%) Wound infection 2 (6.2%)	NR	In persons with BMI >60 DSTVG should be considered as a valid surgical option with two-staged laparoscopic procedures	
Spyropoulos 2007 [255]	6 months	BMI 54.3 ± 9.9 %EBMIL 27.29 ΔBMI 11	OSA 89%; T2DM 81.5%; HTN 85.7%	Pneumonia 1 (3.1%) 1 vomiting treated with 0 medical therapy for 5 days		BIB placement can be considered an effective first-stage treatment of high-risk mega-obese patients in need of surgical intervention	
Gottig 2009 [236]	6 months	%EBMIL 19.7±10.2 ΔBMI 8.7±5.1		3 minor	0	The study shows safety and efficacy of intragastric balloon in extremely obese patients particularly as a first step before a definitive anti-obesity operation	
Shuhabier 2004 [237]	30 days	NR	NR	Intensive care unit days: 17 (0); days on ventilator: 7 (0). P<.05	NR	Gastric bypass in mega-obese persons can be performed safely. A longer LOS, need for ICU stay and mechanical ventilation should be anticipated	

Table 9 (continued)

Author, year	Follow-up duration	Weight loss achieved (comparator)	Comorbidity resolution (comparator)	Early complications requiring reoperation (comparator requiring reoperation)	Long-term complications	Summary of findings	Additional notes
Fazylov 2005 [238]	30 days	NR	NR	Mortality: 8 (7.8%) vs 0 in SO group, $P < 0.01$; among male patients (7/8) 16.7% vs 0. Other complication similar between SSO and SO	0 Incisional hernia and small bowel obstruction similar between two groups	SSO patients (males especially) experience significantly higher mortality compared with SO patients	

given the setting of these studies, there is the possibility of bias due to the selection of patients.

In an RCT on 82 patients with obesity and osteoarthritis, 39 were randomized to AGB 12 months prior to total knee arthroplasty (TKA), and 41 patients were randomized to receive the usual nonoperative weight management prior to TKA. In a median follow-up of 2 years after TKA, 14.6% of patients in the MBS group incurred the primary outcome of composite complications, compared with 36.6% in the control group (difference 22.0%, $P = .02$). The incident TKA decreased by 29.3% in the MBS group because of symptom improvement following weight loss, compared with only 4.9% in the control group [133].

MBS can be performed safely before joint arthroplasty. However, further data are needed with specifically designed trials to clarify the causal role of MBS on the outcomes of subsequent joint arthroplasties.

Considering the conflicting data obtained from the literature, the IFSO and ASMBS decided to conduct a Delphi analysis on the topic of joint arthroplasty in patients with obesity (Table 15). This included the role of MBS as a bridge to joint arthroplasty and the proper time to arthroplasty after MBS.

During the two rounds of the Delphi analysis, the 49 experts reached a consensus on 5 statements concluding that MBS is indicated in patients with class II and III obesity (BMI ≥ 35 kg/m²) even with no other medical conditions and in all age groups following a comprehensive multidisciplinary team assessment. In this survey, consensus was reached in four statements. First, MBS can be considered a bridge to joint arthroplasty in patients with a BMI ≥ 30 kg/m². Second, MBS can decrease the operating time, risk of readmission, and short-term complications of subsequent joint arthroplasty in patients with a BMI ≥ 30 kg/m². Third, MBS can decrease the need for joint arthroplasty in patients with BMI ≥ 30 kg/m². Fourth, the experts also reached a consensus that joint arthroplasty in patients with a BMI ≥ 30 kg/m² should be done 6 months to 1 year after MBS, depending on the severity of the joint disease, if there is weight loss stabilization and if the patient has good muscle mass and nutritional status.

Recommendation

- Obesity is associated with poor outcomes after total joint arthroplasty. Orthopedic surgical societies discourage hip and knee replacement in patients with BMI ≥ 40 kg/m², mainly due to the increased risk of readmission and surgical complications, such as wound infection and deep vein thrombosis.
- Before total knee and hip arthroplasty, MBS has decreased operative time, hospital LOS, and early post-operative complications.
- According to experts, MBS can be considered a bridge to joint arthroplasty in patients with BMI ≥ 30 kg/m².

Table 10 MBS in patients with liver cirrhosis

PMID	Author, year [ref]	Publication	Evidence	Number of patients	Intervention	Length of stay	Mean follow-up (months)	Weight (kg) follow-up	BMI basal	BMI follow-up	EWL/TWL %	Comorbidity resolution
33900587	Kaul A et al. [239]	<i>Obes Surg.</i> 2020	Level 3	22	Cirrhosis SG 20, RYGB 2	4/8	27 (6–51)	86.4 ± 14.8	48.8 ± 7.5	33.1 ± 5.5	62.4 ± 15.8 (EBMIL)	Amelioration DM2
30109667	Hanipah ZN et al. [240]	<i>Obes Surg.</i> 2018	Level 3	13 (+ portal hypertension)	Cirrhosis SG 10, RYGB 3	3 (2–4)	24/00	97.1 (71.7–132.9)	48	36.7 (28.3–43.8)	49/25	100% diabetes, 100% hypertension, 50% dyslipidemia
31853866	Younus H et al. [241]	<i>Obes Surg.</i> 2020	Level 2	26	Cirrhosis GB 1, SG 7, RYGB 17, open RYGB 1,	5	54 (24–132)	NR	46	NR	NR	NR
32808168	Quezada N et al. [242]	<i>Obes Surg.</i> 2020	Level 2	16 (including 3 with portal hypertension)	Cirrhosis SG 11, RYGB 5	3 ± 1	24/00	NR	39 ± 6.8	NR	84.9%/28%	Hypertension 14%, DM2 50%, dyslipidemia 85%
27881858	Singh T et al. [243]	<i>Int J Obes (Lond).</i> 2017	Level 2	99 (stage 3 and 4)	(fibrosis GB5, SG 19, RYGB 75)	4	12/00	NR	46.4 ± 9.6	33.5 ± 7.7	NR	NR
32152677	Salman MA et al. [244]	<i>Surg Endosc.</i> 2021	Level 2	71 (related cirrhosis)	(NASH-SG 71)	NR	30/00	122.5 ± 14.5	44.1 ± 4.3	95.1	NR	DM 50%, hypertension 61.5%, dyslipidemia 70.6%
32285332	Mumtaz K et al. [245]	<i>Obes Surg.</i> 2020	Level 2	3086 (compensated cirrhosis, 103 decompensated cirrhosis)	RYGB 2367, SG 719, RYGB 78, SG 25	1.7 (1.2–2.6) and 3.4 (2.0–7.6)	NR	NR	NR	NR	NR	NR
33156104	Are VS et al. [246]	<i>Am J Gastroenterol.</i> 2020	Level 2	9802 (9356 CC, 446 DC)	46% restrictive procedure	3.4 ± 10.5%	NR	NR	NR	NR	NR	NR
34351576	Miller A et al. [247]	<i>Updates Surg.</i> 2021	Level 2	3032	RYGB 1864, SG 1168	3, 4 RYGB vs 3 SG	NR	NR	NR	NR	NR	NR
30794300	Klebanoff et al. [248]	<i>MJAMA Netw Open.</i> 2019	Level 2	161	SG, RYGB	NR	NR	NR	NR	NR	NR	NR

Table 10 (continued)

PMID	Author, year [ref]	Publication	Evidence	Number of patients	Intervention	Length of stay	Mean follow-up (months)	Weight (kg) follow-up	BMI basal	BMI follow-up	EWL/TWL %	Comorbidity resolution
23201210	Shimizu H et al. [249]	<i>Surg Obes Relat Dis.</i> 2013	Level 3	23	RYGB 14, SG 8, GB 1	4.3 ± 2.7	37 ± 11.1	124.8 ± 33.7	48.2 ± 8.6	33.7 ± 7.7	67.7 ± 24.8	Lipid profile improved in majority of pt, 86.7% improved glucose control in dm2, clinical improve-ment dyslipidemia 66.7 and hypertension 68.7
14980033	Dallal RM et al. [250]	<i>Obes Surg.</i> 2004	Level 3	30	Cirrhosis 27 RYGB, 3 SG	4 ± 4	16/00	NR	52.6 ± 8.3	NR	63 ± 15	NR
34129090	Mittal T et al. [251]	<i>Surg Endosc.</i> 2022	Level 3	15	Portal hypertension	2.73 ± 0.59	12/00	86.05 ± 14.4	43.7 ± 5.79	31.16 ± 3.82	26.5 ± 5.42	Diabetes resolution 80%, hypertension resolution 72.72%
30397876	Minambres I et al. [252]	<i>Obes Surg.</i> 2019	Level 3	41	SG 28, GB 11, BPD 2	NR	38 ± 20	NR	45 ± 8.3	NR	21.16 ± 15.32	DM2 remission 53.6%, TWL 29.1 ± 10.9
32925171	Vuppalanchi R et al. [253]	<i>Ann Surg.</i> 2022 (full text missing)	Level 3	106		3.7 ± 4						
PMID	Author, year [ref]	Intra-operative complications	Peri-operative complications	Post-operative complications	Early mortality	Late mortality	Long-term complications	Child–Pugh	Diagnosis of cirrhosis	Liver outcome	TIPS	Transplantation list
33900587	Kaul A et al. [239]	0	0	n = 1 operative liver decompensation, n = 1 postoperative flank echymoses (both after SG)	0	1 (after 6 months)	0	A	Transient elastography, intraoperative and percutaneous biopsy	n = 3 worsening on transient elastography	0	0

Table 10 (continued)

PMID	Author, year [ref]	Intra-operative complications	Peri-operative complications	Post-operative complications	Early mortality	Late mortality	Long-term complications	Child–Pugh	Diagnosis of cirrhosis	Liver outcome	TIPS	Transplantation list	
30109667	Hanipah ZN et al. [240]	0	0	Infection (1), intra-abdominal hematoma (1), subcutaneous hematoma (1)	0	1 (8 months)	GERD 1, dysphagia 1, cholecystitis 1	NR	Endoscopy, imaging studies, routine intra-operative liver biopsy, mean MELD prior to MBS 9 (7–17)	MELD post-operative 10 (7–13), no changes in the liver function	6 (4 prior, 2 after MBS)	4—but nobody transplanted	
31853866	Younus H et al. [241]	0	(10/26)	$n = 3$ CD III (12%)	0	0	0	A	Intra-operative biopsies	MELD score from 7 to 6	NR	NR	
32808168	Quezada N et al. [242]	0	(5/16)	$n = 2$; CD III (12%)	0	1 (traffic accident)	$n = 1$ hepatoma at 6 years—LT	A	Intra-operative biopsies	MELD pre 7.38±1, 2-year FU: 8.63±3 ($P = .52$)	NR	NR	
27881858	Singh T et al. [243]	0	(13/99)	17 (17.2%)	0	0	36.4%	A	Intra-operative biopsies	MELD pre 7 (6–7) to post 6 (6–7)	0	0	
32152677	Salman MA et al. [244]	0	(9/71)	Bleeding 4, leakage 2, ascites 1, HE 1, ARDS 1	0	0	0	A	Wedge biopsy	Fibrosis score at 30 months: F2 26.8%, F3 40.8%, F4 32.4%	NR	NR	
32285332	Mumtaz K et al. [245]	NR	NR	NR	20 (0.6% with CC, 20 (19.4% with DC)	NR	NR	NR	NR	NR	NR	NR	Higher mortality in DC, undergoing RYGB, low-volume centers (<50 BS)

Table 10 (continued)

PMID	Author, year [ref]	Intra-operative complications	Peri-operative complications	Post-operative complications	Early mortality	Late mortality	Long-term complications	Child–Pugh cirrhosis	Diagnosis of Liver outcome	TIPS	Transplantation list
33156104	Are VS et al. [246]	NR	2% (1.4 CC, 14.1 DC)	Bleeding 2.9%, sepsis 1.8%, urinary tract infection 1.9%, acute kidney injury 4.5%	9356: 0.8% in CC and 446 22.1% in DC	NR	NR	NR	NR	NR	Higher inpatient mortality in non-strictive surgery and low-volume centers (<50 BS/year)
34351576	Miller A et al. [247]	NR	See post-operative	Ascites 1.7 vs 1, variceal bleeding <1 vs 1.1, encephalopathy 0.6 vs <1	21 (1.1%) RYGB vs <10 SG	42 (2.2%) RYGB vs <10 SG	Ascites 3.5 vs 1.4, variceal bleeding <1 vs 1.1, encephalopathy 2.3 vs <1	NR	NR	NR	RYGB increased healthcare utilization in CC
30794300	Klebanoff et al. [248]	MJNR	NR	NR	NR	NR	NR	NR	NR	NR	BS highly cost-effective in patients with NASH and compensated cirrhosis, with SG being the most cost-effective option across all weight classes

Table 10 (continued)

PMID	Author, year [ref]	Intra-operative complications	Peri-operative complications	Post-operative complications	Early mortality	Late mortality (9 months)	Long-term complications	Child–Pugh	Diagnosis of cirrhosis	Liver outcome	TIPS	Transplantation list
23201210	Shimizu H et al. [249]	NR	RYGB changed in SG in 3 pt	8 (34.8%): 1 gastrodujejunal leak, 2 anastomotic strictures, 1 infected hematoma, 1 leak, 1 stricture, 1 pneumonia, 1 bleeding	0	1 (9 months)	NR	A (22), B (1)	21 Liver biopsy during surgery	Follow-up biopsies in 3 patients: marked improvement in the degree of fatty change and inflammation, others not evaluated	2 prior	NR
14980033	Dallal RM et al. [250]	Greater average blood loss 290 ml (vs 115 ml in general)	RYGB changed in SG in 3 pt	10/30 4 tubular necrosis, 1 anastomotic leak, 2 transfusions, 1 prolonged ileus, 2 intubation	0	1	9/30 1 prolonged and malnutrition, 3 abdominal pain, 1 malnutrition, 1 calculi, 2 acute	nausea and malnutrition, 3 abdominal pain, 1 malnutrition, 1 renal marginal ulcer, 1 acute cholecystitis	Intra-operatively in 90%	No liver-related complications	NR	NR
34129090	Mittal T et al. [251]	0	0	0	0	0	3 with GERDA	Endoscopy with dilated esophageal varices	NR	NR	NR	NR

Table 10 (continued)

PMID	Author, year [ref]	Intra-operative complications	Peri-operative complications	Post-operative complications	Early mortality	Late mortality	Long-term complications	Child–Pugh	Diagnosis of cirrhosis	Liver outcome	TIPS	Transplantation list
30397876	Minambres I et al. [252]	NR	7/41 patients: fistula, hemoperitoneum, upper GI bleeding in ulcers, portal thrombosis, wound infection, 2 ascites	2 liver decompensation, 3 late liver decompensation	0	0	6 patients: HCC post-surgically	A 40, B 1	Biopsy during surgery, 17 prior to surgery	MELD 7.2 ± NR 1.9, 7.8± 2.2 1 year, 8.4±3.1 3 years, 9.8 ± 4.6 5 years		1 of 41
32925171	Vuppalaanchi R et al. [253]				0	3					46% prior to surgery	97%

Level of Evidence 2b

Grade of recommendation B

7. MBS and abdominal wall hernia repair [143–165]
PRISMA Appendix [PubMed, Cochrane, Embase]
Systematic Review Table 7

Twenty-three studies were included [143–165]. Five studies were extracted from national registers, including the Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program (MBSAQIP) [144, 151], the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) [159], and the French hospital discharge summaries database system [154]. The other 18 studies were single-cohort retrospective studies. The studies were heterogeneous regarding timing and technique. Timing is mainly divided into ventral hernia repair (VHR) before MBS (symptomatic, low- or high-grade intestinal obstruction), concomitant VHR, and VHR after MBS. Some authors presented treatment algorithms regarding timing in their studies [147, 158, 161]. Of the 23 studies, 18 studies included a concomitant VHR and 5 studies a staged procedure [143, 147, 155, 162, 163]. Ventral hernias included epigastric, incisional, umbilical, paraumbilical, and Spigelian hernias, and one study reported the multistep approach in complex hernias with loss of domain [162]. Studies included primary and recurrent incisional hernia repair. VHR included open and laparoscopic techniques, with and without mesh. MBS included AGB, SG, and open and laparoscopic RYGB.

The studies analyzed postoperative morbidity and mortality, long-term complications, and recurrence rates. Abdominal wall hematoma, seroma, and surgical site infections were the most reported complications associated with all types of VHR. Small bowel obstructions (SBOs) and mesh dehiscence were reported in some studies, with the highest incidence of SBO at 37.5% in one study with deferred treatment [153].

Early mortality was reported in four register studies that analyzed concomitant VHR and was reported to be 0.3% [144], <1% [151], 0.2% [157], and 0.1% [159].

The literature presents a large amount of heterogeneous data regarding VHR in patients with obesity, and only five studies evaluated a staged approach. Due to the higher risk of reoperation for recurrence, VHR would be avoided in bariatric patients before MBS.

Recommendation

- Obesity is a risk factor for the development of ventral hernias.
- In persons with obesity and an abdominal wall hernia, MBS-induced weight loss is suggested before ventral hernia repair in order to reduce the rate of postoperative complications.

Level of Evidence 2b

Table 11 MBS in patients with heart failure

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidity resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Blumer 2020 [254]	Retrospective	5	5216 (2636 RYGB, 2272 SG, 308 LAGB)	NA	Range from <35 to >60	NA	NA	NA	Myocardial infarction 9% vs 0.1%; AF 28.7% vs 1.9%; pneumonia 12.7% vs 0.82%; respiratory failure 16.8% vs 1%; AKI 27.9% vs 1.9%	None	NA	NA	788,195 BS vs 5216 BS with HF	Higher in-hospital mortality than BS without HF (10.3 vs 0.4)
Sun, 2022 [255]	Retrospective	5	318 SG (72%), 110 RYGB (25%), 15 others (14%)	NA	72% BI >40	NA	NA	None	None	10/433 worsening of HF	NA	NA	433 with hypertrophic cardiomyopathy	Safe surgery with few post-operative complications
Balakura-man, 2022 [256]	Retrospective	5	10 SG, 5 RYGB	2	50	38.3	35.77 TWL	NA	NA	NA	NA	NA	NA	12/15 had recovery for LVEF
Mentias, 2022 [257]	Retrospective	5	94,885 BS vs 94,885 matched cohort	4 years	44.7	NA	NA	NA	NA	NA	NA	NA	NA	54% lower risk of new onset HF; lower rate of readmission for HF
Val, 2020 [258]	Retrospective	5	23 SG, 3 RYGB	12 mm	42.8	31.7	35 lb	NA	None	2 DVT, 4 AKI, 3 SSI	NA	1 death	48 LVAD inserted	Complication rates remain in an acceptable range

Table 11 (continued)

Author, year [ref]	Study design Quality assessment	Number of patients (interven- tion)	Mean follow-up (months)	BMI basal	BMI follow- up	EWL/TWL	Comorbidity resolution	Intra-opera- tive compli- cations	Peri-opera- tive compli- cations	Post-opera- tive compli- cations	Long-term complica- tions	Bridge to surgery	Outcomes of second- ary surgery
Alsbrook, 2006 [259]	Retrospec- tive	32 RYGB	2 yy	56.5	38.9	48 kg	15/16 DM, 17/23 HTN, 20/23 OSAS	None	May-32	1 death	NA	32 CHF	Acceptable morbidity and mortality
Vest, 2016 [260]	Retrospec- tive	8 LAGB (19%), 23 RYGB (55%), 11 SG (26%)	12 mm	48.2	35.5	22.6%	NA	None	HF/pul- monary edema 10%, MI 2% AKI 2%, ICU 5%, read- mission 12%	NA	NA	42 LSVD	Improve- ment of LVEF; acceptable morbidity
Naslund, 2021 [261]	Retrospec- tive	465 RYGB, 44 SG	4.6 yy	40.6	28	29% TWL	51% DM, 24.7% HTN 66.1% OSAS	NA	8.4%, seri- ous 3.8%	NA	1 death 0.2%	53 HF	Lower risk of new- onset HF and MI
Doumouras, 2021 [262]	Retrospec- tive propensity score match	1319 BS (79.5% RYGB, 20.5% SG) vs 1319 non-BS	NA	NA	NA	NA	NA	NA	NA	NA	Overall 7.7% and 0.15% mortality	274 HF	Bariatric surgery was associated with a significant lower incidence of MACE in patients with CVD and severe obesity. These observed results apply to both patients with IHD and HF

Table 11 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidity resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Aminian 2021 [263]	Retrospective observational matched	5	1362 RYGB; 693 SG	5 yy	45.3	NA	NA	NA	NA	NA	NA	NA	204 HF	Lower incidence of MACE; reduction in incidence of HF
Lundberg, 2022 [264]	Retrospective population-based cohort	5	27,882 RYGB	10 yy	NA	NA	NA	NA	NA	NA	NA	NA	1 HF	Lower risk of developing HF than non-operated obese match, higher risk than controls (but slightly)
Yuan, 2021 [265]	Retrospective	5	308 RYGB	1 y	>35	NA	NA	NA	NA	NA	NA	NA	1 HF	Lower rates of developing MACE
Ng, 2022 [266]	Retrospective	5	22 SG	18 mm	43.3	31.5	26.8% TWL	NA	NA	NA	NA	NA	22 LVAD positioned before SG	2 LVAD explained, 10 HT, 5 listed for HT
Benotti, 2017 [267]	Retrospective longitudinal cohort	5	1724 RYGB and 1724 matched control	8 yy	46.5	NA	NA	70% DM	NA	NA	NA	NA	79 HF	Reduced risk of relapse
Hoskuldottir, 2021 [268]	Cohort study retrospective	5	5321 RYGB and 5321 matched	4.5 yy	42	32	NA	NA	NA	NA	NA	NA	142 HF	Lower rates of rehospitalization for HF and reduction in mortality

Table 11 (continued)

Author, year [ref]	Study design	Quality assessment	Number of patients (intervention)	Mean follow-up (months)	BMI basal	BMI follow-up	EWL/TWL	Comorbidity resolution	Intra-operative complications	Peri-operative complications	Post-operative complications	Long-term complications	Bridge to surgery	Outcomes of secondary surgery
Yang, 2020 [269]	Retrospective	5	18 LAGB, 2SG, 1 BPD	1 y	46.2	38	43.8 % EWL	NA	NA	NA	4 Clavien–Dindo 1, 3 CD 2, 2 CD 3b	1 Infected band	3 Heart transplant	1 had successful HT, 2 were removed from HT list because of increased LVEF; 76,2% had an increase of at least 10% of LVEF
Strzelczyk, 2021 [270]	Retrospective	5	65	1 y	43.7	30	31.2 % EWL	NA	NA	NA	NA	NA	NA	Left atrial function increased after BS, with overall cardiac improvement
Stenberg, 2022 [271]	Observational matched cohort database	6	974 RYGB and 191 SG with HD vs 4870 RYGB and 955 SG without HD	2 yy	NA	NA	28.7 % TWL	NA	2.80%	9.40%	NA	0.70%	NA	Same complication rate in HD obese patients. Higher but acceptable cardio-vascular complication rate for HF 1.2% vs 0.2%

Table 12 Multidisciplinary care

Practical Recommendations of the Obesity Management Task Force of the European Association for the Study of Obesity for the Post-Bariatric Surgery Medical Management [272]	<i>Obes Facts</i>	Eng	2017	MEDLINE
<p>Busetto L, Dicker D, Azran C, Batterham RL, Farpour-Lambert N, Fried M, Hjeltnesæth J, Kinz J, Leitner DR, Makaronidis JM, Schindler K, Toplak H, Yumuk V</p>	<p><i>Obes Facts</i></p>	<p>Eng</p>	<p>2017</p>	<p>Metabolic/obesity surgery is today the most effective long-term therapy for the management of patients with severe obesity, and its use is recommended by the relevant guidelines of the management of obesity in adults. Bariatric surgery is in general safe and effective, but it can cause new clinical problems and is associated with specific diagnostic, preventive, and therapeutic needs. For clinicians, the acquisition of special knowledge and skills is required in order to deliver appropriate and effective care to the post-bariatric patient. In the present recommendations, the basic notions needed to provide first-level adequate medical care to post-bariatric patients are summarized. Basic information about nutrition, management of comorbidities, pregnancy, psychological issues as well as weight regain prevention and management is derived from current evidences and existing guidelines. A short list of clinical practical recommendations is included for each item. It remains clear that referral to a bariatric multidisciplinary center, preferably the one performing the original procedure, should be considered in case of more complex clinical situations</p>

Table 12 (continued)

Optimizing Bariatric Surgery Multidisciplinary Follow-up: A Focus on Patient-Centered Care [273]	<i>Obes Surg</i>	Eng	2017	MEDLINE
<p>Aarts MA, Sivapalan N, Nikzad SE, Serodio K, Sockalingam S, Conn LG</p>		<p>Background: Failure to follow-up post-bariatric surgery has been associated with higher postoperative complications, lower percentage weight loss, and poorer nutrition</p> <p>Objective: This study aimed to understand the patient follow-up experience in order to optimize follow-up care within a comprehensive bariatric surgery program</p> <p>Methods: Qualitative telephone interviews were conducted in patients who underwent surgery through a publicly funded multidisciplinary bariatric surgery program in 2011, in Ontario, Canada. Inductive thematic analysis was used</p> <p>Results: Of the 46 patients interviewed, 76.1% were female, mean age was 50, and 10 were lost to follow-up within 1 year post-surgery. Therapeutic continuity was the most important element of follow-up care identified by patients and was most frequently established with the dietician, as this team member was highly sought and accessible. Patients who attended regularly (1) appreciated the specialized care, (2) favored ongoing monitoring and support, (3) were committed to the program, and (4) felt their family doctor had insufficient experience/knowledge to manage their follow-up care. Of the 36 people who attended the clinic regularly, 8 were not planning to return after 2 years due to (1) perceived diminishing usefulness, (2) system issues, (3) confidence that their family physician could continue their care, or (4) higher priority personal/health issues. Patients lost to follow-up stated similar barriers</p> <p>Conclusion: Patients believe the follow-up post-bariatric surgery is essential in providing the support required to maintain their diet and health. More personalized care focusing on continuity and relationships catering to individual patient needs balanced with local health-care resources may redefine and reduce attrition rates</p>		

Table 12 (continued)

	<i>Transplantation</i>	2011	MEDLINE
Panel report: best practices for the surgical treatment of obesity [274]	Gould J, Ellsmere J, Fanelli R, Hutter M, Jones S, Pratt J, Schauer P, Schirmer B, Schwartzberg S, Jones DB		<p>The multidisciplinary bariatric patient care team and the Eng bariatric program accreditation process are key factors in best outcomes. The multidisciplinary WLS team should include trained surgeon(s), a WLS program coordinator, nutritionist, primary care physician, medical subspecialists, and the operating room team. Optimal perioperative care of the WLS patient involves the use of multiple medical disciplines and the multidisciplinary team. For this reason, WLS should be focused at centers where these resources are readily available. The multidisciplinary WLS team is an important component of any bariatric surgery program for a variety of reasons. First of all, bariatric surgery patients have needs that are very different from patients undergoing other types of surgery. Education and behavior modification are important for WLS to succeed. These complex needs, coupled with an extremely low tolerance for poor outcomes (public scrutiny), the essentially elective nature of these operations, and a lack of sympathy for and bias against obesity, create an environment where multidisciplinary programs and accreditation of these programs is essential. There are currently two systems of accreditation for WLS programs not run by individual insurance companies. The American College of Surgeons Bariatric Surgery Centers Network and the Surgical Review Corporation Bariatric Surgery Center of Excellence Program (affiliated with the American Society of Metabolic and Bariatric Surgery) are similar in many ways [22, 23]. Both programs require specific resources (facilities and specialized equipment), and evaluate key personnel, the bariatric surgeon(s), the patient selection process, and patient education as well as outcomes and follow-up. There are some minor differences in terms of the data collection process, fees, and the fact that the ACS only accredits centers where the Surgical Review Corporation accredits both surgeons and centers. As outcomes data from these accredited centers have accumulated over the years, it has become apparent that the morbidity and mortality rates for these centers are lower than expected based on published data [24]. Future steps in the accreditation process include developing a risk-adjusted system where outcomes can replace surgical volume as a surrogate for excellence. It is likely that the future of bariatric surgery accreditation and reimbursement will take into account these outcomes</p>

Table 12 (continued)

	Warman JL	<i>Crit Care Nurs Q</i>	2013	MEDLINE
<p>The application of laparoscopic bariatric surgery for treatment of severe obesity in adolescents using a multidisciplinary adolescent bariatric program [275]</p>		<p>Q The evolution of laparoscopic surgery has made bariatric surgery acceptable for weight loss; however, much controversy exists about its appropriateness for adolescents. Despite the controversial issues, the growing epidemic in adolescent obesity has resulted in rising numbers of applications for bariatric surgery. There are few bariatric surgical programs designed for adolescents. Pediatric settings face high start-up costs and poor reimbursement and lack established bariatric surgeons. Even so, bariatric surgery is increasingly being performed on adolescents in alarming numbers. To avoid adverse physical and psychosocial outcomes, the application of the principles of growth and development is essential. The program should be established as a multidisciplinary approach to management of adolescents and should be in institutions capable of meeting the guidelines for surgical treatment outlined by the American Society of Bariatric Surgery. To prevent postoperative complication, a multidisciplinary team of experienced medical and surgical specialists is needed for optimal preoperative decision making and postoperative management and long-term follow-up. Laparoscopic Roux-en-Y gastric bypass is a safe procedure and an effective means to treat obesity-related morbidity in the adolescent. Results have been excellent and justify a clinical trial to confirm the safety and efficacy of bariatric surgery in the adolescent population</p>	Eng	
<p>Collazo-Clavell ML, Clark MM, McAlpine DE, et al. Assessment and preparation of patients for bariatric surgery. <i>Mayo Clin Proc.</i> 2006;81:S11–7.39. 2012report2/downloads/BS_full-report.pdf [276]</p>	Collazo-Clavell ML, Clark MM, McAlpine DE, et al.			Eng

Table 12 (continued)

<p>Proposal for a multidisciplinary approach to the patient with morbid obesity: the St. Francisus Hospital morbid obesity program [277]</p>	<p>Elte JW, Castro Cabezas M, Vrijland WW, Ruseler CH, Groen M, Mannaerts GH</p>	<p><i>Eur J Intern Med</i> Morbid obesity is a serious disease as it is accompanied Eng by substantial comorbidity and mortality. The prevalence is increasing to an alarming extent, in Europe as well as in the USA. In the past few decades, bariatric surgery has developed and gained importance. It currently represents the only long-lasting therapy for this group of patients, resulting in an efficient reduction in body weight and obesity-related medical conditions, mostly cardiovascular in nature. The importance of a standardized protocol, the use of selection criteria, and a multidisciplinary approach have been stressed but not yet described in detail. Therefore, in this article, the multidisciplinary approach and the treatment protocol that have been applied in our hospital for more than 20 years are set out in a detailed manner. The application of a strict protocol may help to select and follow up motivated patients and to organize multidisciplinary research activities</p>	<p>2008</p>
--	--	--	-------------

Table 12 (continued)

Apovian CM, Cummings S, Anderson W, et al. Best practice updates for multidisciplinary care in weight loss surgery. <i>Obesity</i> . 2009;17:871–89. https://doi.org/10.1038/oby.2008.58 [278]	The objective of this study is to update evidence-based best practice guidelines for multidisciplinary care of weight loss surgery (WLS) patients. We performed systematic search of English-language literature on WLS, patient selection, and medical, multidisciplinary, and nutritional care published between April 2004 and May 2007 in MEDLINE and the Cochrane Library. Key words were used to narrow the search for a selective review of abstracts, retrieval of full articles, and grading of evidence according to systems used in established evidence-based models. A total of 150 papers were retrieved from the literature search and 112 were reviewed in detail. We made evidence-based best practice recommendations from the most recent literature on multidisciplinary care of WLS patients. New recommendations were developed in the areas of patient selection, medical evaluation, and treatment. Regular updates of evidence-based recommendations for best practices in multidisciplinary care are required to address changes in patient demographics and levels of obesity. Key factors in patient safety include comprehensive preoperative medical evaluation, patient education, appropriate perioperative care, and long-term follow-up	Eng	2009	MEDLINE	Santry HP, Chin MH, Cagney KA, Alverdy JC, Lauerderdale DS. The use of multidisciplinary teams to evaluate bariatric surgery patients: results from a national survey in the USA. <i>Obes Surg</i> 2006;16:59–66. [PubMed: 16417760]
<p>Apovian CM, Cummings S, Anderson W, Boyer K, Day K, Hatchigian E, Hodges B, Patti ME, Pettus M, Perma F, Rooks D, Saltzman E, Skoropowski J, Tantillo MB, Thomason P</p> <p>Best practice updates for multidisciplinary care in weight loss surgery</p>	<p>The objective of this study is to update evidence-based best practice guidelines for multidisciplinary care of weight loss surgery (WLS) patients. We performed systematic search of English-language literature on WLS, patient selection, and medical, multidisciplinary, and nutritional care published between April 2004 and May 2007 in MEDLINE and the Cochrane Library. Key words were used to narrow the search for a selective review of abstracts, retrieval of full articles, and grading of evidence according to systems used in established evidence-based models. A total of 150 papers were retrieved from the literature search and 112 were reviewed in detail. We made evidence-based best practice recommendations from the most recent literature on multidisciplinary care of WLS patients. New recommendations were developed in the areas of patient selection, medical evaluation, and treatment. Regular updates of evidence-based recommendations for best practices in multidisciplinary care are required to address changes in patient demographics and levels of obesity. Key factors in patient safety include comprehensive preoperative medical evaluation, patient education, appropriate perioperative care, and long-term follow-up</p>	Eng	2009	MEDLINE	<p>Santry HP, Chin MH, Cagney KA, Alverdy JC, Lauerderdale DS. The use of multidisciplinary teams to evaluate bariatric surgery patients: results from a national survey in the USA. <i>Obes Surg</i> 2006;16:59–66. [PubMed: 16417760]</p>
					<p>8. Giusti V, De Lucia A, Di Vetta V, et al. Impact of preoperative teaching on surgical option of patients qualifying for bariatric surgery. <i>Obes Surg</i> 2004;14:1241–1246. [PubMed: 15527642]</p>
					<p>9. Cunningham E. What is the registered dietitian's role in the preoperative assessment of a client contemplating bariatric surgery? <i>J Am Diet Assoc</i> 2006;106:163. [PubMed: 16390679]</p>

Table 12 (continued)

Clinical practice guidelines of the European Association for Endoscopic Surgery (EAES) on bariatric surgery: update 2020 endorsed by IFSO-EC, EASO and ESPCOP [279]	Di Lorenzo N, Antoniou SA, Batterham RL, Busetto L, Godoroja D, Iossa A, Carrano FM, Agresta F, Alarçon I, Azran C, Bouvy N, Balaguè Ponz C, Buza M, Copăescu C, De Luca M, Dickler D, Di Vincenzo A, Felsenreich DM, Francis NK, Fried M, Gonzalo Prats B, Goitein D, Halford JCG, Herlesova J, Kalogridaki M, Ket H, Morales-Conde S, Piatto G, Prager G, Pruijssers S, Pucci A, Rayman S, Romano E, Sanchez-Cordero S, Vilallonga R, Silecchia G	<i>Surg Endoscop</i>	Preoperative dietitian consultation should be considered Eng	2020	Paper for justification Antoniou SA, Anastasiadou A, Antoniou GA, Granderath F-A, Kafatos A (2017) Preoperative nutritional counseling versus standard care prior to bariatric surgery: effects on postoperative weight loss. <i>Eur Surg Acta Chir Aust</i> . https://doi.org/10.1007/s10353-016-0459-4
			<p>for patients undergoing bariatric surgery</p> <p>Strong recommendation</p> <p>Justification</p> <p>A meta-analysis reporting 3 RCTs was found on this topic [40]. Analyses were re-performed due to error in the primary meta-analysis (calculation of WMD instead of standardized MD, SMD). The overall quality of evidence was very low for weight loss and low for postoperative complications due to risk of bias across RCTs, inconsistency (conceptual and statistical heterogeneity due to variety of preoperative interventions for weight loss, and heterogeneity in the duration of follow-up) and indirectness (follow-up duration for weight loss insufficient for generalizability of findings). Postoperative weight loss was more pronounced in the preoperative diet consultation group (SMD 0.4, 95% CI 0.03 to 0.78 higher). No difference in the odds of postoperative complications was found (risk ratio, RR, 0.80, 95% CI 0.22 to 2.86), although interval estimates were wide. Confidence in the evidence was generally low (Supplementary Table S3); however, the panel favored a strong recommendation after consulting with the patient representative who expressed a strong preference for a holistic approach of the bariatric patient with continuous preoperative and postoperative consultation. The panel considered this practice feasible, requiring moderate human and financial resources, and being acceptable to stakeholders. There was no evidence of any risk for the intervention according to the panel's judgement</p>		
van Hout GC, Vreeswijk CM, van Heck GL. Bariatric surgery and bariatric psychology: evolution of the Dutch approach. <i>Obes Surg</i> . 2008 Mar;18(3):321-5. https://doi.org/10.1007/s11695-007-9271-3 . Epub 2008 Jan 17. PMID: 18202896. [280]					0

Table 12 (continued)
 It's Time for Multi-
 disciplinary Obe-
 sity Management
 Centers Comment
 Obesity (Silver
 Spring). 2019
 Apr;27(4):534.
<https://doi.org/10.1002/oby.22450>.
 COMMENTARY
 [281]

Walter J Pories I, Louis J Aronne 2 US problems

Predictors of attrition in a multidisciplinary adult weight management clinic. Gill RS, Karmali S, Hadi G, Al-Adra DP, Shi X, Birch DW. *Can J Surg*. 2012 Aug;55(4):239-43. <https://doi.org/10.1503/cjs.035710>. PMID: 22617538; PMCID: PMC3404143 [282]

Background: Worldwide, more than 1.7 billion individuals may be classified as overweight and are in need of appropriate medical and surgical treatments. The primary goal of a comprehensive weight management program is to produce sustainable weight loss. However, for such a program to be effective, the patient must complete it. We analyzed attrition rates and predictors of attrition within a publicly funded, multidisciplinary adult weight management program

Methods: We retrospectively reviewed charts from an urban multidisciplinary adult weight management clinic program database. Patients received medical or surgical treatment with appropriate follow-up. We collected information on demographics and comorbidities. Patients in the surgical clinics received either laparoscopic gastric band insertion or gastric bypass. We conducted univariate analysis and multivariate analyses on predictors of attrition

Results: A total of 1205 patients were treated in the weight management program: 887 in the medical clinic and 318 with surgery and follow-up in a surgical clinic. Overall, 516 patients left the program or were lost to follow-up (attrition rate 42.8%). The attrition rate was 53.9% in the medical clinic and 11.9% in the surgical clinic. Multivariate analyses identified participation in the medical clinic, younger patient age, and lower body mass index as predictors of attrition

Conclusion: We found lower attrition rates among surgically than medically treated patients in a multidisciplinary weight management clinic. Further research is needed to understand those variables that lead to improved attrition rates

Table 12 (continued)

Andalib A, Bouchard P, Bougie A, Loïselle SE, Demyttenaere S, Court O. Variability in bariatric surgical care among various centers: a survey of all bariatric surgeons in the Province of Quebec, Canada. *Obes Surg*. 2018 Aug;28(8):2327–2332. <https://doi.org/10.1007/s11695-018-3157-4>. PMID: 29492752. [283]

Aboueid S, Jasinska M, Bourgeault I, Giroux I. Current Canada situation

weight management approaches used by primary care providers in six multidisciplinary healthcare settings in Ontario. *Can J Nurs Res*. 2018 Dec;50(4):169–178. <https://doi.org/10.1177/0844562118769229>. Epub 2018 Apr 17. PMID: 29665702. [284]

Table 13 Revisional surgery

First author (year)	Study design	Quality assessment (NOS)	Number of patients	BMI	Reason for conversion/revision	Conversion/revision	Laparoscopic/robotic/open	Intervention	Operative time (min)	Length of stay (days)	Weight loss	Complications	Complications	Complications	Nutritional complications	Follow-up (months)	Other outcomes
Vahibe (2023) [285]	Retrospective	Fair quality	53	Not available	Malnutrition	Revision	Laparoscopic	Different types	Not available	Not available	Not available	45.2%	Not available	3.8%	5.7%	24	Improvement of nutritional complications
Vanetta (2022) [286]	Retrospective	Good quality	20,387	39.5–47.2	Weight regain, GERD, complications	Conversion	Laparoscopic/robotic	Different types (especially from AGB and SG)	103–196,913	2.9	Not available	3.8%	9%	0.2%	Not available	30 days	
Major (2022) [287]	Retrospective	Fair quality	799	48	Weight regain, complications	Conversion	Laparoscopic	Different types (especially from AGB and VGB to RYGB and OAGB)	Not available	3.5	33.4%	9.52%	4.76%	0%	4.76%	22.7	43% Remission from TD2M; 31% remission from hypertension
Xie (2022) [288]	Retrospective	Good quality	221	45.6	Weight regain, GERD, complications	Conversion	Laparoscopic/robotic	Different types (especially from AGB and SG)	149.2	2	17.3%	WL7.7%	3.1%	0.4%	0.9%	24	
Hernandez (2021) [289]	Retrospective	Fair quality	54	41.7	Weight regain, GERD, complications	Revision	Laparoscopic	Revisional (RYGB, AGB, SG)	Not available	4.1	Not available	Not available	0.9% early and 1.8% late	0%	Not available	Not available	Not available

Table 13 (continued)

First author (year)	Study design	Quality assessment (NOS)	Number of patients	BMI	Reason for conversion/revision	Conversion/revision	Laparoscopic/robotic/open	Intervention	Operative time (min)	Length of stay (days)	Weight loss	Complication	Complications	Complications	Nutritional complications	Follow-up (months)	Other outcomes
Gero (2021) [290]	Retrospective	Good quality	3143	35.2	Weight regain, GERD, complications	Revision, conversion	Laparoscopic	Different types	93	not available	17.7% WL	Not available	23.8%	0.06%	Not available	12	Secondary BS is safe, although postoperative morbidity exceeds the established benchmarks for primary BS
Dreifuss (2021) [291]	Retrospective	Good quality	76	45.7	Weight regain, GERD, complications	Revision, conversion	Robotic	Different types (especially from AGB and SG to RYGB)	182	2.1	22.4% WL	Not available	3.9% early and 5.2% late	1%	Not available	24	
King (2020) [292]	Retrospective	Good quality	167	37–39.5	Complications, weight regain	Revision	laparoscopic/robotic	Revisional	Not available	5.2–5.8%	Not available	5.2–5.8%	1.9–5.2%	0%	Not available	30 days	Comparable results between laparoscopic and robotic revisional surgery

Table 13 (continued)

First author (year)	Study design	Quality assessment (NOS)	Number of patients	BMI	Reason for conversion/revision	Conversion/revision	Laparoscopic/robotic/open	Intervention	Operative time (min)	Length of stay (days)	Weight loss	Complication Dindo 1–2	Complication Clavien–Dindo 3–4	Complication Clavien–Dindo 5 (surgical related mortality)	Nutritional complications	Follow-up (months)	Other outcomes
Cheema (2021) [293]	Retrospective	Fair quality	266	39.8–45	Weight regain, GERD, complications	Revision/conversion	Laparoscopic	Revisional RYGB, conversion from AGB and SG	Not available	2	10–30% WL	Not available	2.6%	0%	Not available	24 months	Improvement of HbA1c and CV risk
El Chaar (2021) [294]	Retrospective	Good quality	440	42.4	Not available	Revision	Laparoscopic/robotic	Revisional RYGB, revisional SG	145.5	Not available	Not available	3%	0%	0%	Not available	30 days	
Mora Oliver (2020) [295]	Retrospective	Fair quality	112	41.9	Weight regain	Conversion	Laparoscopic	Different types (especially from AGB, VBG, and SG to OAGB)	135.8	4.9	27.5% WL	3%	2.7%	0%	Not available	20.8	Improvement of TD2M and HTN
Keren (2019) [296]	Retrospective	Good quality	266	41.3	Weight regain (90%), complications	Revision/conversion	Laparoscopic/open	Different types (especially from AGB and SG)	Not available	3.2	30.5% WL	4.8%	2.4	2%	Not available	12	
Accedo (2020) [297]	Retrospective	Good quality	2288	40.9	Not available	Revision/conversion	Laparoscopic/robotic	Revisional RYGB, revisional SG	125.4	2.2	Not available	Not available	3.2%	0.2%	Not available	30 days	

Table 13 (continued)

First author (year)	Study design	Quality assessment (NOS)	Number of patients	BMI	Reason for conversion/revision	Conversion/revision	Laparoscopic/robotic/open	Intervention	Operative time (min)	Length of stay (days)	Weight loss	Complications	Complications	Complications	Nutritional complications	Follow-up (months)	Other outcomes
Clapp (2019) [298]	Retrospective	Good quality	37,916	41.6	Not available	Revision/robotic	Laparoscopic/robotic	Revisional RYGB, conversion from AGB and SG	103–167	1.7–2.3	10 Δ BMI	Not available	Not available	0.1%	Not available	12	
Aleassa (2019) [299]	Retrospective	Fair quality	81	41.2–47.2	Weight regain, complications	Revision/conversion	Laparoscopic	Revisional RYGB, conversion of VBG, AGB, and SG to RYGB	Not available	Not available	20.5% WL	Not available	Not available	Not available	Not available	22	23.1–35% Remission from TD2M
Qiu (2018) [300]	Retrospective	Good quality	84	38–42	Weight regain, complications	Revision/conversion	Laparoscopic	Revisional RYGB, conversion of VBG, AGB, and SG to RYGB	133–175	2	7.7–30.2% WL	8.3%	6%	0%	Not available	12	
Gray (2018) [301]	Retrospective	Good quality	84	39–45	Weight regain, complications	Revision/conversion	Laparoscopic/robotic	Revisional RYGB, conversion from AGB and SG	177–238	3.7–5.8	Not available	Not available	5.9%	0%	Not available	12	
Souto (2018) [302]	Retrospective	Fair quality	67	36.9	Malnutrition, weight regain	Revision/conversion	Laparoscopic	Revisional JIB, revisional BPD-DF	Not available	Not available	28.7–77% EWL	11.9%	11.9%	9.2%	Over 29	years	

Table 13 (continued)

First author (year)	Study design	Quality assessment (NOS)	Number of patients	BMI	Reason for conversion/revision	Conversion/revision	Laparoscopic/robotic/open	Intervention	Operative time (min)	Length of stay (days)	Weight loss	Complications	Complications	Nutritional complications	Follow-up (months)	Other outcomes	
Fulton (2017) [303]	Retrospective	Fair quality	117	44.7	Weight regain, malnutrition	Revision/conversion	Laparoscopic/open	Revisional RYGB, conversion from AGB and SG	168	4	61.2% EWL	Not available	10.8%	0%	Not available	12	
Daigle (2016) [304]	Retrospective	Fair quality	121	47.5	Weight regain	Revision/conversion	Laparoscopic	Revisional RYGB, conversion from AGB, SG, and VSG	Not available	6	59.4% EWL	17%	3.3%	0%	Not available	40	Revisable bariatric surgery is capable of treating both inadequate weight loss and refractory metabolic disease
Shimizu (2013) [305]	Retrospective	Fair quality	154	44	Weight regain, complications	Revision/conversion	Laparoscopic/open	Different types	268–280	5.4–9.5	37.6% EWL	10.3%	12.9%	0.6%	Not available	12	
Kuesters (2011) [306]	Retrospective	Fair quality	100	28–62	Weight regain, complications	Revision/conversion	Laparoscopic/open	Different types	Not available	Not available	56% EWL	Not available	Not available	0%	Not available	12	
Fronza (2010) [307]	Retrospective	Fair quality	63	38–41	Weight regain, malnutrition	Revision/conversion	Laparoscopic/open	Different types	Not available	Not available	>50% EWL	19%	11%	0%	Not available	12	

Table 13 (continued)

First author (year)	Study design	Quality assessment (NOS)	Number of patients	BMI	Reason for conversion/revision	Conversion/revision	Laparoscopic/robotic/open	Intervention	Operative time (min)	Length of stay (days)	Weight loss	Complications Clavien-Dindo 1–2	Complications Clavien-Dindo 3–4	Complications Clavien-Dindo 5 (surgical related mortality)	Nutritional complications	Follow-up (months)	Other outcomes
Spyropoulos (2010) [308]	Retrospective	Fair quality	56	46.9	Weight regain, malnutrition	Revision/conversion	Open	Revisional RYGB, revisional BPD-DS	210	16.5	68.9% EWL	20.8%	13.1%	0%	3.6%	102	
Lim (2009) [309]	Retrospective	Fair quality	75	46.3	Weight regain, malnutrition	Revision/conversion	Laparoscopic/open	Revisional RYGB, conversion from AGB and SG	152–231	2–5.8	47.8% EWL	17.3%	4.0%	0%	Not available	6	
Nesset (2009) [310]	Retrospective	Fair quality	218	42	Weight regain, complication, malnutrition	Revision/conversion	Open/laparoscopic	Revisional RYGB, revisional JIB, revisional VBG	298	9	13 Δ BMI	Not available	26%	0.9%	Not available	84	
			67,408														

Table 14 Grade of recommendation and level of evidence

Grade of recommendation	Level of evidence	Type of study
A	1a	Systematic review of [homogeneous] randomized controlled trials
A	1b	Individual randomized controlled trials [with narrow confidence intervals]
B	2a	Systematic review of [homogeneous] cohort studies of “exposed” and “unexposed” subjects
B	2b	Individual cohort study/low-quality randomized control studies
B	3a	Systematic review of [homogeneous] case–control studies
B	3b	Individual case–control studies
C	4	Case series, low-quality cohort, or case–control studies
D	5	Expert opinions based on non-systematic reviews of results or mechanistic studies

Evidence-Based Medicine, Stony Brook University Libraries, 14 March 2023

Table 15 IFSO/ASMBS delphi results on MBS in individuals need joint arthroplasty

1. MBS can be considered a bridge to joint arthroplasty in patients with a Body Mass Index of ≥ 30 kg/m ²	84.7% Agree	-	CONSENSUS (AGREE)
2. MBS can decrease the operating time, risk of readmission, and short-term complications of subsequent joint arthroplasty in individuals with a Body Mass Index of ≥ 30 kg/m ²	82.9% Agree	-	CONSENSUS (AGREE)
3. MBS can decrease the need for Joint arthroplasty in patients with a Body Mass Index of ≥ 30 kg/m ² .	84.7% Agree	-	CONSENSUS (AGREE)
4. Joint arthroplasty in patients with a Body Mass Index of ≥ 30 kg/m ² should be done 6 months to 1 year after MBS depending on the severity of their arthritis or if their weight loss stabilizes and they have sufficient muscle mass and good nutritional status.	50.0% Agree	88.3% Agree	CONSENSUS (AGREE)

Grade of recommendation B

8. MBS prior to organ transplantation [166–189]
PRISMA Appendix 8 [PubMed, Cochrane, Embase]
Systematic Review Table 8

Generally, extremely high or low BMI is considered a contraindication to solid organ transplantation (SOT) due to poor outcomes. Class III obesity may prevent access to transplantation since it is considered a relative contraindication and poses specific technical challenges during surgery [190, 191]. MBS, despite worldwide recognition as the most effective treatment for obesity, may be overlooked as an option in patients with severe end-stage organ disease. Nonetheless, MBS has been described in patients with end-stage organ disease to improve their candidacy for transplantation.

A systematic review of 2241 papers identified 24 thoroughly analyzed studies. The studies included different SOT summarized as heart/lung, kidney, and liver.

The literature search considered several variables, such as surgical procedures, disease status, patient age, and follow-up time. In many studies, specific data points such as weight loss, operative time, and complication rates

were missing. In addition, there were differences between patients and studies, including different transplant timing and surgical techniques.

Recommendation

- *Obesity is associated with end-stage organ disease and may limit access to transplantation. Obesity is also a relative contraindication for solid organ transplantation and poses unique technical challenges during surgery.*
- *Published data supports considering patients with end-stage renal disease and obesity grade 3 being able to be listed for kidney transplant after MBS.*
- *MBS is shown to be safe and effective as a bridge to liver transplantation in selected patients who would otherwise be ineligible.*
- *MBS can also improve heart transplants outcomes.*
- *Limited data suggest that MBS could improve eligibility for lung transplantation.*
- *MBS can be performed post-SOT or concomitantly to reduce complication rates and mortality.*

Level of Evidence 2b Grade of recommendation B

MBS in the High-Risk Patients

9. MBS for BMI ≥ 60 kg/m² [192–238]

PRISMA Appendix 9 [PubMed, Cochrane, Embase]
Systematic Review Table 9

Forty-seven papers have been retrieved for qualitative analysis [192–238].

Twelve studies were focused on the safety and feasibility of MBS among patients with severe obesity at 30 days of follow-up after surgery with no reported data on weight loss or obesity-related comorbidities. Thirty-five studies analyzed MBS's safety, feasibility, and medium to long-term results in patients with obesity and BMI ≥ 60 kg/m².

Concerning weight loss, the mean initial BMI was ≥ 66.64 kg/m² (SD ± 3.05). After a mean follow-up of 28 months, the mean %EBMIL was 51.5 (SD ± 16) with a mean Δ BMI of 21.64 kg/m² (SD ± 7.16). Improvement or resolution of the obesity-related complications were reported in 17 studies, including patients with BMI ≥ 60 kg/m². The mean percentage of improvement/resolution of T2DM was 67.35% (SD ± 24.79). The mean percentage of improvement/resolution of HTN was 54.01% (SD $\pm .93$). The mean percentage of improvement/resolution of obstructive sleep apnea (OSA) was 63.61% (SD ± 21.51), while the mean percentage of improvement/resolution of dyslipidemia was 70.95% (SD ± 10.31).

Early complications (within 30 days from surgery) were reported in 45 studies.

The overall mean percentage of early complications was 7.57% (SD ± 6.28), and the mean percentage of early complications requiring reoperation was 4.9% (SD ± 3.48). The overall mean mortality was 1.61% (SD ± 2.29).

Long-term complications were reported in 13 studies. The mean percentage of long-term complications was 13.56% (SD ± 10.93).

Recommendation

- MBS is safe and effective in patients with BMI ≥ 60 kg/m².
- Evidence suggests a higher rate of perioperative complications after MBS in patients with BMI ≥ 60 kg/m².
- According to the literature, MBS appears safe in patients with initial BMI ≥ 70 kg/m².

Level of Evidence 2a Grade of recommendation B

10 MBS in patients with liver cirrhosis [239–253]

PRISMA Appendix 10 [PubMed, Cochrane, Embase]
Systematic Review Table 10

Fifteen studies were included in this systematic review. Some studies differed between compensated and decompensated liver cirrhosis.

The early mortality was reported as 0.6 and 0.8% in the Metabolic Dysfunction-associated Liver Disease (MAFLD) or compensated liver cirrhosis, and 19.4 and 22.1% in decompensated liver cirrhosis. Mumatz et al. and Are et al. [245, 246] underlined the higher mortality of patients in low-volume centers (<50/year). Miller et al. analyzed 3032 patients undergoing SG ($n = 1168$) and RYGB ($n = 1864$) with compensated liver cirrhosis and reported early mortality in 21 (1.1%) of patients after RYGB and 10 patients after SG (<1%). Late mortality occurred in 42 patients after RYGB (2.2%) and under 10 patients after SG (<0.8%) [247].

Based on the current systematic review, patients with MAFLD or compensated liver cirrhosis have acceptable perioperative morbidity and mortality. However, patients with obesity and decompensated liver cirrhosis are at much higher risk for perioperative complications and perioperative mortality following MBS. Those patients should only be considered for surgery on a selective basis after a comprehensive risk assessment and only in high-volume centers. The risk of postoperative liver decompensation is low but should not be underestimated. Weight loss and remission of comorbidities are similar to the general bariatric surgical population. Careful patient selection and consideration of the choice of surgical procedure are important to ensure the best outcomes.

Recommendation

- Obesity is a significant risk factor for MAFLD and liver cirrhosis.
- MBS has been associated with histologic improvement of MAFLD and regression of liver fibrosis.
- MBS is associated with a risk reduction of progression of MAFLD to liver cirrhosis.
- MBS in patients with “decompensated” cirrhosis is associated with high perioperative mortality.
- Careful patient selection and consideration of the choice of surgical procedure are important to ensure the best outcomes.

Level of Evidence 2b**Grade of recommendation B**

11. MBS in patients with heart failure [254–271]

PRISMA Appendix 11 [PubMed, Cochrane, Embase]

Systematic Review Table 11

Thirty-one full-text articles were assessed for eligibility. Eighteen studies are included in the qualitative synthesis [254–271].

MBS is associated with a lower risk of major adverse cardiovascular events (MACE), including myocardial infarction, ischemic heart disease, or heart failure (HF) in patients with severe obesity [255–257].

The overall risk for early (less than 30 days) and late (30 days or more) complications was similar for patients with cardiovascular disease and the matched group that did not have cardiovascular disease [258–262]. Some studies reported an increased risk for early cardiovascular complications as well as a higher 90-day mortality rate (still within an acceptable range) for patients with heart disease, such as HF [263–266].

Current data suggest that MBS can be a useful adjunct to treatment in patients with obesity and HF before heart transplantation or placement of a left ventricular assistance device (LVAD) [266–268]. Patients who underwent MBS were observed to have improvement in cardiac function [269, 270]. This had several beneficial effects, such as a reduction in re-hospitalization for HF, and improvement in their left ventricular ejection fraction (LVEF). MBS could increase the patient's likelihood of receiving a heart transplant. On the other hand, some patients had enough improvement in their cardiac function to no longer require a heart transplant [269, 270].

Recommendation

- *MBS in patients with obesity and HF is associated with improvement of LVEF, improvement of functional capacity, and higher chances for receiving heart transplantation.*
- *In patients with obesity and HF, MBS has low morbidity and mortality and can be a useful adjunct before heart transplantation or placement of LVAD.*

Level of Evidence 2b**Grade of recommendation B****Patient Evaluation**

12. Multidisciplinary care [272–284]

PRISMA Appendix 12 [PubMed, Cochrane, Embase]

Systematic Review Table 12

The search screened 95 papers, but only 6 were thoroughly analyzed. There were guidelines or consensus statements, including those from the European Association for the Study of Obesity (EASO) and the European Association for Endoscopic Surgery (EAES) [272, 279]. Standardized pre-operative multidisciplinary evaluations have been reported to reduce major complications and reoperation rates.

The studies of this systematic review support the protective role of the multidisciplinary team (MDT) to ensure patient safety.

Registered experts in nutrition in MBS can assist in the management of post-operative patients who may experience issues such as food intolerances, malabsorption, micronutrient deficiencies, dumping syndrome, hypoglycemia, and RWG. Licensed mental health providers with specialty knowledge and experience in MBS behavioral health are necessary to assess patients for psychopathology and determine the candidate's ability to cope with the adversity of surgery, the changing body image, and the lifestyle changes required after MBS.

Based on the EAES guidelines, scheduled multidisciplinary post-operative follow-up should be provided to every patient undergoing MBS [279].

Recommendation

- MDT has an important role in MBS patients' pre- and post-operative management.

Level of Evidence 2c**Grade of recommendation B****Revisional Surgery**

13. Revisional MBS [285–310]

PRISMA Appendix 13 [PubMed, Cochrane, Embase]

Systematic Review Table 13

Twenty-six studies were selected for this systematic review. All studies were retrospective with a good/fair quality.

Recent articles report conversion from AGB and SG and revision of RYGB and OAGB. Revisional MBS is currently performed laparoscopically and robotically, with a growing trend toward a robotic approach. Operative time and LOS of revisional surgery were reduced with time and experience, which could be comparable to those reported in the literature for primary surgery.

All revisional and conversional interventions lead to additional weight loss. Clavien–Dindo complications 3–4 ranged from 0.9 to 26%. Mortality was lower than 1% for conversions from restrictive procedures, and up to 11.9% was reported after revisional stapling procedures. Revisional surgery appeared to induce further remission from T2DM and HTN.

Recommendation

- *Indication for revisional surgery after MBS varies among patients but may include insufficient weight loss, weight regain, insufficient remission of comorbidities, and management of complications (e.g., gastroesophageal reflux).*
- *Due to its complexity, revisional MBS may be associated with higher rates of perioperative complications. However, revisional MBS induces satisfactory metabolic outcomes with acceptable complications and mortality rates.*

Level of Evidence 2b

Grade of recommendation B

Discussion (see Criteria Table 16)

The indications for MBS have not changed since the NIH proposed them in 1991. In other words, the indications have not kept up with the evolution of surgical technique from open laparotomy to minimally invasive, the changing procedure types, the improved safety of MBS, and the emerging evidence on numerous health benefits of weight loss.

IFSO and ASMBS joined forces to tackle this major problem, and the new MBS guidelines were published in October 2022. Updated guidelines based on current literature and data are vital as access to this life-saving surgery is still

very low despite the available evidence—in most countries, access to MBS is less than 2% of eligible candidates.

This study systematically reviewed the best literature available for the outcomes of MBS for various populations with differing demographics and obesity-related complications. Eleven of the 13 criteria were supported by the literature. Where there was a lack of evidence, a Delphi process was employed to achieve expert consensus. *PRISMA Prospect* summarized the findings.

From these data, MBS impacted positively a range of populations and settings. The majority of examined populations had Grade B recommendations for the indications of surgery. Expert opinion (Grade D) was only relied upon to strengthen the evidence for the role of MBS in a few unique circumstances. This includes patients with a BMI of 35–40 kg/m² who have no comorbidities, patients with a concurrent need for arthroplasty, and the role of the multidisciplinary team. Particularly in the pediatric and adolescent populations, the strength of the available data supported a Grade A recommendation. Improved access to surgery in adolescents was one of the two major new emphases of the new IFSO/ASMBS guidelines.

This systematic review highlights the need for well-designed RCTs or large prospective cohort studies to enable better-informed decision-making for clinicians and patients. Clinicians working in the field innately understand the benefit of multidisciplinary teamwork. However, it has yet to be proven in high-quality studies.

Just as the NIH indications from 1991 became outdated as surgical techniques, with a better understanding of the pathophysiology of obesity and improved perioperative safety, these current guidelines should be regularly revisited when new evidence emerges to inform treatment decisions.

List of Delphi consensus Experts

First Name	Last Name	Country
Edo	Aaarts	Netherland
Ahmad	Aly	Australia
Ali	Aminian	USA
Luigi	Angrisani	Italy
Ahmad Abdallah	Bashir	Jordan
Estuardo	Behrens	Guatemala
Helmuth Thorlakur	Billy	USA
Sonja	Chiappetta	Italy
Jean-Marc	Chevallier	France
Ricardo Vitor	Cohen	Brazil
Maurizio	De Luca	Italy
Pierre Y	Garneau	Canada
Khaled Aly	Gawdat	Egypt
Ashraf	Haddad	Jordan
Jacques M	Himpens	Belgium
Farah Anwari	Husain	USA
Angelo	Iossa	Italy
Mohammad	ermansaravi	Iran
Shanu Nikhil	Kothari	USA
Lilian	Kow	Australia
Marina	Kurian	USA
Teresa LeAnn	LaMasters	USA
Silvia	Leite Faria	Brazil
Ken Wing King	Loi	Australia
Kamal K	Mahawar	UK
Corrigan Lee	McBride	USA
Giovanni	Merola	Italy
Monali	Misra	USA
Abdelrahman Ali	Nimeri	USA
Joe	Northup	USA
Mary	O'Kane	UK
Pavlos	Papasavas	USA
Richard M	Peterson	USA
Giacomo	Piatto	Italy
Luis	Poggi	Peru
Jaime	Ponce	USA
Gerhard	Prager	Austria
Janey Sue Andrews	Pratt	USA
Almino Cardoso	Ramos	Brazil
Ann M	Rogers	USA
Paulina	Salminen	Finland
Nathaniel James	Sann	USA
John David	Scott	USA
Scott Alan	Shikora	USA
Michel	Suter	Switzerland
Salvatore	Tolone	Italy
Antonio	Vitiello	Italy
Cunchuan	Wang	China

Table 16 Summary of recommendations with their grade and level of evidence

Criteria	PRISMA and DELPHI	Appendix/ Table	Level of evidence	Grade of recommen- dation	Recommendation	References
MBS for BMI 30–34.9 kg/m ²	PRISMA	1	2a	B	MBS is recommended for patients with BMI 30–34.9 kg/m ² with T2DM and/or other obesity-associated medical problems	[7–35]
MBS for BMI 35–40 kg/m ² without obesity-associated comorbidities	PRISMA Insufficient data DELPHI	2	5	D	MBS is recommended regardless of the presence, absence, or severity of obesity-associated medical problems	–
BMI thresholds in the Asian population	PRISMA	3	2a	B	Access to MBS should not be denied solely based on the BMI	[36–54]
MBS in the older population	PRISMA	4	2a	B	There is no evidence to support an age limit	[55–72]
MBS for pediatric and adolescents	PRISMA	5	1b	A	MBS is safe in the population younger than 18 years, produces durable weight loss, and improves obesity-associated medical problems	[76–117]
MBS prior to joint arthroplasty	PRISMA Conflicting data DELPHI	6	2b	B	MBS can be considered a bridge to joint arthroplasty in patients with BMI ≥30 kg/m ²	[120–141]
MBS and abdominal wall hernia repair	PRISMA	7	2b	B	In patients with severe obesity and an abdominal wall hernia, MBS-induced weight loss is suggested before hernia repair	[143–165]
MBS prior to organ transplantation	PRISMA	8	2b	B	Published data supports considering patients in need of SOT first to undergo MBS to improve their eligibility for transplantation	[166–189]
MBS for BMI ≥60 kg/m ²	PRISMA	9	2a	B	MBS is safe and effective in patients with BMI ≥60 kg/m ²	[192–238]
MBS in patients with liver cirrhosis	PRISMA	10	2b	B	MBS is associated with a reduction of progression of MAFDL to cirrhosis	[239–253]
MBS in patients with heart failure	PRISMA	11	2b	B	MBS can be a useful treatment adjunct in patients with obesity and heart failure	[254–271]
Multidisciplinary care	PRISMA	12	2c	B	Despite the low evidence level, MDT is at present the unmodifiable core of pre- and post-operative obesity management	[272–284]
Revisional surgery	PRISMA	13	2b	B	Revisional MBS induces satisfactory metabolic outcomes with acceptable rates of complications and mortality	[285–310]

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11695-024-07370-7>.

Data Availability The data that support the findings of this study are available on request from the corresponding author.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and

reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by

statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Mun EC, Blackburn GL, Matthews JB. Current status of medical and surgical therapy for obesity. *Gastroenterology*. 2001;120(3):669–81.
- Gastrointestinal surgery for severe obesity. Consensus Statement. 1991;9(1):1–20.
- Page MJ, McKenzie JE, Bossuyt PM, et al. statement: an updated guideline for reporting systematic reviews. *BMJ*. 2020;372:n71.
- Nasa P, Jain R, Juneja D. Delphi methodology in healthcare research: how to decide its appropriateness. *World J Methodol*. 2021;11(4):116–29.
- Eisenberg D, Shikora SA, Aarts E, et al. 2022 American Society of Metabolic and Bariatric Surgery (ASMBS) and International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO) indications for metabolic and bariatric surgery. *Obes Surg*. 2023;33(1):3–14.
- Eisenberg D, Shikora SA, Aarts E, et al. 2022 American Society for Metabolic and Bariatric Surgery (ASMBS) and International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO): indications for metabolic and bariatric surgery. *Surg Obes Relat Dis*. 2022;18(12):1345–56.
- Billeter AT, Scheurlen KM, Israel B, et al. Gastric bypass resolves metabolic dysfunction-associated fatty liver disease (MAFLD) in low-BMI patients: a prospective cohort study. *Ann Surg*. 2022;276(5):814–21.
- Chaturvedi R, Gracner T, Tysinger B, et al. The long-term value of bariatric surgery interventions for American adults with type 2 diabetes mellitus. *Ann Surg*. 2023;277(5):789–97.
- Altieri MS, DeMaria E, Lensing C, et al. Real-world retrospective analysis of outcomes in patients undergoing bariatric surgery with class 1 obesity. *Surg Obes Relat Dis*. 2022;18(5):569–76.
- Singh D, Baksi A, Ramana P, Singla V, et al. Five-year outcomes of sleeve gastrectomy in patients with class I obesity and type 2 diabetes mellitus. *Obes Surg*. 2022;32(1):96–102.
- Baldwin D, Sanchez-Johnsen L, Bustos R, et al. Metabolic surgery outcomes in U.S. patients with class I obesity. *Bariatric Surg Pract Patient Care*. 2021;16(2):85–91.
- Gupta S, Beitner M, Skinner C, et al. Medium-term outcomes of converting laparoscopic adjustable gastric band to Roux-en-Y gastric bypass in patients with body mass index <35 kg/m(2): a uniquely Australian perspective. *Surg Obes Relat Dis*. 2020;16(4):485–91.
- Varban OA, Bonham AJ, Finks JF, et al. Is it worth it? Determining the health benefits of sleeve gastrectomy in patients with a body mass index <35 kg/m(2). *Surg Obes Relat Dis*. 2020;16(2):248–53.
- Navarrete Aulestia S, LeybaNavarrete Llopis JLS, et al. One anastomosis gastric bypass/minigastric bypass in patients with BMI < 35 kg/m(2) and type 2 diabetes mellitus: preliminary report. *Obes Surg*. 2019;29(12):3987–91.
- Gamme G, Dang JT, Switzer N, et al. Evaluating the safety of bariatric surgery for weight loss in class I obesity: a propensity-matched analysis of North American data. *Surg Obes Relat Dis*. 2019;15(4):629–35.
- Feng X, Andalib A, Brethauer SA, et al. How safe is bariatric surgery in patients with class I obesity (body mass index 30–35 kg/m(2))? *Surg Obes Relat Dis*. 2019;15(2):253–60.
- Vitiello A, Angrisani L, Santonicola A, et al. Bariatric surgery versus lifestyle intervention in class I obesity: 7–10-year results of a retrospective study. *World J Surg*. 2019;43(3):758–62.
- Noun R, Slim R, Nasr M, et al. Results of laparoscopic sleeve gastrectomy in 541 consecutive patients with low baseline body mass index (30–35 kg/m(2)). *Obes Surg*. 2016;26(12):2824–8.
- Maiz C, Alvarado J, Quezada N, et al. Bariatric surgery in 1119 patients with preoperative body mass index <35 (kg/m(2)): results at 1 year. *Surg Obes Relat Dis*. 2015;11(5):1127–32.
- Kaska L, Proczko M, Kobiela J, et al. Dynamics of type 2 diabetes mellitus laboratory remission after Roux-en-Y gastric bypass in patients with body mass index lower than 35 kg/m(2) and higher than 35 kg/m(2) in a 3-year observation period. *Wideochir Inne Tech Maloinwazyjne*. 2014;9(4):523–30.
- Walker DM, Hii MW, Skinner CE, et al. Roux-en-Y gastric bypass after successful weight loss with a laparoscopic adjustable gastric band: rationales and early outcomes in patients of body mass index <35 kg/m(2). *Surg Obes Relat Dis*. 2014;10(6):1104–8.
- Boza C, Valderas P, Daroch DA, et al. Metabolic surgery: roux-en-Y gastric bypass and variables associated with diabetes remission in patients with BMI <35. *Obes Surg*. 2014;24(8):1391–7.
- Scopinaro N, Adami GF, Papadia FS, et al. Effects of gastric bypass on type 2 diabetes in patients with BMI 30 to 35. *Obes Surg*. 2014;24(7):1036–43.
- Serrot FJ, Dorman RB, Miller CJ, et al. Comparative effectiveness of bariatric surgery and nonsurgical therapy in adults with type 2 diabetes mellitus and body mass index <35 kg/m2. *Surgery*. 2011;150(4):684–91.
- Gianos M, Abdemur A, Fendrich I, et al. Outcomes of bariatric surgery in patients with body mass index <35 kg/m2. *Surg Obes Relat Dis*. 2012;8(1):25–30.
- Choi J, Digiorgi M, Milone L, et al. Outcomes of laparoscopic adjustable gastric banding in patients with low body mass index. *Surg Obes Relat Dis*. 2010;6(4):367–71.
- Varela JE. Bariatric surgery: a cure for diabetes? *Curr Opin Clin Nutr Metab Care*. 2011;14(4):396–401.
- Scopinaro N, Adami GF, Papadia FS, et al. The effects of biliopancreatic diversion on type 2 diabetes mellitus in patients with mild obesity (BMI 30–35 kg/m2) and simple overweight (BMI 25–30 kg/m2): a prospective controlled study. *Obes Surg*. 2011;21(7):880–8.
- DeMaria EJ, Pate V, Warthen M, et al. Baseline data from American Society for Metabolic and Bariatric Surgery-designated Bariatric Surgery Centers of Excellence using the Bariatric Outcomes Longitudinal Database. *Surg Obes Relat Dis*. 2010;6(4):347–55.
- Parikh M, Duncombe J, Fielding GA. Laparoscopic adjustable gastric banding for patients with body mass index of or <=35 kg/m2. *Surg Obes Relat Dis*. 2006;2(5):518–22.
- Sultan S, Parikh M, Youn H, et al. Early U.S. outcomes after laparoscopic adjustable gastric banding in patients with a body mass index less than 35 kg/m2. *Surg Endosc*. 2009;23(7):1569–73.
- Cohen RV, Pereira TV, Aboud CM, et al. Microvascular outcomes after metabolic surgery (MOMS) in patients with type 2 diabetes mellitus and class I obesity: rationale and design for a randomised controlled trial. *BMJ Open*. 2017;7(1):e013574.
- Angrisani L, Favretti F, Furbetta F, et al. Italian Group for Lap-Band System: results of multicenter study on patients with BMI < or =35 kg/m2. *Obes Surg*. 2004;14(3):415–8.
- Cevallos L, Rodriguez FM, Herrera A, et al. Metabolic surgery and class 1 obesity (35 kg/m(2)): a prospective study with short-, mid-, and long-term results among Latinos. *Obes Surg*. 2021;31(6):2401–9.
- Espinosa O, Pineda O, Maydon HG, et al. Type 2 diabetes mellitus outcomes after laparoscopic gastric bypass in patients with BMI <35 kg/m(2) using strict remission criteria: early outcomes of a prospective study among Mexicans. *Surg Endosc*. 2018;32(3):1353–9.

36. Mazidi M, Gao HK, Li L, et al. Effects of Roux-en-Y gastric bypass on insulin secretion and sensitivity, glucose homeostasis, and diabetic control: a prospective cohort study in Chinese patients. *Surgery*. 2017;161(5):1423–9.
37. Osman Abouzeid TA, Ain Shoka AA, Atia KS, et al. From diabetes remedy to diabetes remission, could single-anastomosis gastric bypass be a safe bridge to reach target in non-obese patients? *Asian J Surg*. 2019;42(1):307–13.
38. Ma S, Wang L, Chen J, et al. The effect of laparoscopic sleeve gastrectomy on type 2 diabetes remission outcomes in patients with body mass index between 25 kg/m² and 32.5 kg/m². *Asian J Surg*. 2022;45(1):315–9.
39. Park DJ, An S, Park YS, et al. Bariatric surgery versus medical therapy in Korean obese patients: prospective multicenter, non-randomized controlled trial (KOBESS trial). *Ann Surg Treat Res*. 2021;101(4):197–205.
40. Luo Y, Guo Z, He H, et al. Predictive model of type 2 diabetes remission after metabolic surgery in Chinese patients. *Int J Endocrinol*. 2020;2020:2965175.
41. Nautiyal HK, Guan W, Lin S, et al. Preoperative predictors of early relapse/no-remission of type-2 diabetes after metabolic surgery in Chinese patients. *Clin Obes*. 2020;10(2):e12350.
42. Huang YM, Lin YK, Lee WJ, et al. Long-term outcomes of metabolic surgery in overweight and obese patients with type 2 diabetes in Asia. *Diabetes Obes Metab*. 2021;23(3):742–53.
43. Malapan K, Goel R, Tai CM, et al. Laparoscopic Roux-en-Y gastric bypass for nonobese type II diabetes mellitus in Asian patients. *Surg Obes Relat Dis*. 2014;10(5):834–40.
44. Zuo D, Xiao X, Yang S, et al. Effects of bariatric surgery in Chinese with obesity and type 2 diabetes mellitus: a 3-year follow-up. *Medicine (Baltimore)*. 2020;99(34):e21673.
45. Zhao L, Li W, Su Z, et al. Preoperative fasting C-peptide predicts type 2 diabetes mellitus remission in low-BMI Chinese patients after Roux-en-Y gastric bypass. *J Gastrointest Surg*. 2018;22(10):1672–8.
46. Fan J, Xu JH, Wang J, et al. Effects of laparoscopic adjustable gastric banding on weight loss, metabolism, and obesity-related comorbidities: 5-year results in China. *Obes Surg*. 2014;24(6):891–6.
47. Kim MJ, Hur KY. Short-term outcomes of laparoscopic single anastomosis gastric bypass (LSAGB) for the treatment of type 2 diabetes in lower BMI (30 kg/m²) patients. *Obes Surg*. 2014;24(7):1044–51.
48. Zhang H, Han X, Yu H, et al. Effect of Roux-en-Y gastric bypass on remission of T2D: medium-term follow-up in Chinese patients with different BMI obesity class. *Obes Surg*. 2017;27(1):134–42.
49. Kwon O, Lee YJ, Yu JH, et al. The recovery of beta-cell function is critical for antidiabetic outcomes of gastric bypass in Asian subjects with type 2 diabetes and a body mass index below 30. *Obes Surg*. 2017;27(2):541–4.
50. Du X, Fu XH, Shi L, et al. Effects of laparoscopic Roux-en-Y gastric bypass on Chinese type 2 diabetes mellitus patients with different levels of obesity: outcomes after 3 years' follow-up. *Obes Surg*. 2018;28(3):702–11.
51. Liang H, Cao Q, Liu H, et al. The predictive factors for diabetic remission in Chinese patients with BMI > 30 kg/m² and BMI < 30 kg/m² are different. *Obes Surg*. 2018;28(7):1943–9.
52. Widjaja J, Pan H, Dolo PR, et al. Short-term diabetes remission outcomes in patients with BMI ≤/≥ 30 kg/m² following sleeve gastrectomy. *Obes Surg*. 2020;30(1):18–22.
53. Yu Z, Li W, Sun X, et al. Predictors of type 2 diabetes mellitus remission after metabolic surgery in Asian patients with a BMI <32.5 kg/m². *Obes Surg*. 2021;31(9):4125–33.
54. Mazidi M, Gao HK, Hui H, et al. Changes in adiposity and other cardiometabolic risk factors following Roux-en-Y gastric bypass: a 12-month prospective cohort study in Chinese patients. *Indian J Gastroenterol*. 2017;36(4):258–62.
55. Athanasiadis DI, Hernandez E, Dirks RC, et al. Postoperative 4-year outcomes in septuagenarians following bariatric surgery. *Obes Surg*. 2021;31(12):5127–31.
56. Pajceki D, Dantas ACB, Tustumi F, et al. Sleeve gastrectomy versus Roux-en-Y gastric bypass in the elderly: 1-year preliminary outcomes in a randomized trial (BASE Trial). *Obes Surg*. 2021;31(6):2359–63.
57. Nor Hanipah Z, PUNCHAI S, Karas LA, et al. The outcome of bariatric surgery in patients aged 75 years and older. *Obes Surg*. 2018;28(6):1498–503.
58. Hammond JB, Webb CJ, Pulivarthi V, et al. Is there an upper age limit for bariatric surgery? Laparoscopic gastric bypass outcomes in septuagenarians. *Obes Surg*. 2020;30(7):2482–6.
59. Gholizadeh B, Makhosoi BR, Valizadeh R, et al. Safety and efficacy of one anastomosis gastric bypass on patients with severe obesity aged 65 years and above. *Obes Surg*. 2022;32(5):1610–6.
60. Ramirez A, Roy M, Hidalgo JE, et al. Outcomes of bariatric surgery in patients >70 years old. *Surg Obes Relat Dis*. 2012;8(4):458–62.
61. Daigle CR, Andalib A, Corcelles R, et al. Bariatric and metabolic outcomes in the super-obese elderly. *Surg Obes Relat Dis*. 2016;12(1):132–7.
62. Casillas RA, Kim B, Fischer H, et al. Comparative effectiveness of sleeve gastrectomy versus Roux-en-Y gastric bypass for weight loss and safety outcomes in older adults. *Surg Obes Relat Dis*. 2017;13(9):1476–83.
63. Frieder JS, Montorfano L, Gomez CO, et al. Sleeve gastrectomy versus Roux-en-Y gastric bypass in patients aged ≥/≥65 years: a comparison of short-term outcomes. *Surg Obes Relat Dis*. 2021;17(8):1409–15.
64. Quebbemann B, Engstrom D, Siegfried T, et al. Bariatric surgery in patients older than 65 years is safe and effective. *Surg Obes Relat Dis*. 2005;1(4):389–92.
65. Quirante FP, Montorfano L, Rammohan R, et al. Is bariatric surgery safe in the elderly population? *Surg Endosc*. 2017;31(4):1538–43.
66. Lainas P, Dammaro C, Gaillard M, et al. Safety and short-term outcomes of laparoscopic sleeve gastrectomy for patients over 65 years old with severe obesity. *Surg Obes Relat Dis*. 2018;14(7):952–9.
67. Danan M, Nedelcu A, Noel P, et al. Operative morbidity of laparoscopic sleeve gastrectomy in subjects older than age 65. *Surg Obes Relat Dis*. 2019;15(1):8–11.
68. van Rutte PW, Smulders JF, de Zoete JP, et al. Sleeve gastrectomy in older obese patients. *Surg Endosc*. 2013;27(6):2014–9.
69. O'Keefe KL, Kemmeter PR, Kemmeter KD. Bariatric surgery outcomes in patients aged 65 years and older at an American Society for Metabolic and Bariatric Surgery Center of Excellence. *Obes Surg*. 2010;20(9):1199–205.
70. Garofalo F, Denis R, Pescarus R, et al. Long-term outcome after laparoscopic sleeve gastrectomy in patients over 65 years old: a retrospective analysis. *Surg Obes Relat Dis*. 2017;13(1):1–6.
71. Loy JJ, Youn HA, Schwack B, et al. Safety and efficacy of laparoscopic adjustable gastric banding in patients aged seventy and older. *Surg Obes Relat Dis*. 2014;10(2):284–9.
72. Nevo N, Eldar SM, Lessing Y, et al. Sleeve gastrectomy in the elderly. *Obes Facts*. 2019;12(5):502–8.
73. Gondal AB, Hsu CH, Zeeshan M, et al. A frailty index and the impact of frailty on postoperative outcomes in older patients after bariatric surgery. *Surg Obes Relat Dis*. 2019;15(9):1582–8.
74. Giordano S, Victorzon M. Bariatric surgery in elderly patients: a systematic review. *Clin Interv Aging*. 2015;10:1627–35.
75. Shenoy SS, Gilliam A, Mehanna A, et al. Laparoscopic sleeve gastrectomy versus laparoscopic Roux-en-Y gastric bypass in elderly bariatric patients: safety and efficacy—a systematic review and meta-analysis. *Obes Surg*. 2020;30(11):4467–73.

76. O'Brien PE, Sawyer SM, Laurie C, et al. Laparoscopic adjustable gastric banding in severely obese adolescents: a randomized trial. *JAMA*. 2010;303(6):519–26.
77. Carbajo MA, Gonzalez-Ramirez G, Jimenez JM, et al. A 5-year follow-up in children and adolescents undergoing one-anastomosis gastric bypass (OAGB) at a European IFSO Excellence Center (EAC-BS). *Obes Surg*. 2019;29(9):2739–44.
78. Dobritoiu D, Buza M, Catanescu E, et al. Excellent short- and medium-term result of bariatric surgery in adolescence A single center study. *Chirurgia (Bucur)*. 2019;114(6):753–60.
79. Elhag W, El Ansari W, Abdulrazzaq S, et al. Evolution of 29 anthropometric, nutritional, and cardiometabolic parameters among morbidly obese adolescents 2 years post sleeve gastrectomy. *Obes Surg*. 2018;28(2):474–82.
80. Elhag W, El Ansari W. Durability of cardiometabolic outcomes among adolescents after sleeve gastrectomy: first study with 9-year follow-up. *Obes Surg*. 2021;31(7):2869–77.
81. Raziel A, Sakran N, Szold A, et al. Mid-term follow-up after laparoscopic sleeve gastrectomy in obese adolescents. *Isr Med Assoc J*. 2014;16(1):37–41.
82. Yitzhak A, Mizrahi S, Avinoach E. Laparoscopic gastric banding in adolescents. *Obes Surg*. 2006;16(10):1318–22.
83. Holterman AX, Holterman M, Browne A, et al. Patterns of surgical weight loss and resolution of metabolic abnormalities in super-obese bariatric adolescents. *J Pediatr Surg*. 2012;47(9):1633–9.
84. Holterman AX, Browne A, Tussing L, et al. A prospective trial for laparoscopic adjustable gastric banding in morbidly obese adolescents: an interim report of weight loss, metabolic and quality of life outcomes. *J Pediatr Surg*. 2010;45(1):74–8.
85. Inge TH, Garcia V, Daniels S, et al. A multidisciplinary approach to the adolescent bariatric surgical patient. *J Pediatr Surg*. 2004;39(3):442–7.
86. Jaramillo JD, Snyder E, Farrales S, et al. A multidisciplinary approach to laparoscopic sleeve gastrectomy among multiethnic adolescents in the United States. *J Pediatr Surg*. 2017;52(10):1606–9.
87. Conroy R, Lee EJ, Jean A, et al. Effect of laparoscopic adjustable gastric banding on metabolic syndrome and its risk factors in morbidly obese adolescents. *J Obes*. 2011;2011:906384.
88. Riquin E, Schmitt F, Bouhours-Nouet N, et al. Laparoscopic adjustable gastric banding in adolescents with severe obesity: psychological aspects, decision makers of surgery, and 2-year outcomes. A case series *Arch Pediatr*. 2018;25(4):269–73.
89. Loy JJ, Youn HA, Schwab B, et al. Improvement in non-alcoholic fatty liver disease and metabolic syndrome in adolescents undergoing bariatric surgery. *Surg Obes Relat Dis*. 2015;11(2):442–9.
90. Manco M, Mosca A, De Peppo F, et al. The benefit of sleeve gastrectomy in obese adolescents on nonalcoholic steatohepatitis and hepatic fibrosis. *J Pediatr*. 2017;180:31–7.
91. Nadler EP, Reddy S, Isenalumhe A, et al. Laparoscopic adjustable gastric banding for morbidly obese adolescents affects android fat loss, resolution of comorbidities, and improved metabolic status. *J Am Coll Surg*. 2009;209(5):638–44.
92. Zeller MH, Modi AC, Noll JG, et al. Psychosocial functioning improves following adolescent bariatric surgery. *Obesity (Silver Spring)*. 2009;17(5):985–90.
93. Xiao N, Devarajan P, Inge TH, et al. Subclinical kidney injury before and 1 year after bariatric surgery among adolescents with severe obesity. *Obesity (Silver Spring)*. 2015;23(6):1234–8.
94. Sarwer DB, Dilks RJ, Spitzer JC, et al. Changes in dietary intake and eating behavior in adolescents after bariatric surgery: an ancillary study to the Teen-LABS Consortium. *Obes Surg*. 2017;27(12):3082–91.
95. Ryder JR, Edwards NM, Gupta R, et al. Changes in functional mobility and musculoskeletal pain after bariatric surgery in teens with severe obesity: Teen-Longitudinal Assessment of Bariatric Surgery (LABS) Study. *JAMA Pediatr*. 2016;170(9):871–7.
96. Alqahtani AR, Elahmedi MO, Al Qahtani AR, et al. Laparoscopic sleeve gastrectomy in children and adolescents with Prader-Willi syndrome: a matched-control study. *Surg Obes Relat Dis*. 2016;12(1):100–10.
97. Reiter-Purtill J, Ley S, Kidwell KM, et al. Change, predictors and correlates of weight- and health-related quality of life in adolescents 2-years following bariatric surgery. *Int J Obes (Lond)*. 2020;44(7):1467–78.
98. Xanthakos SA, Khoury JC, Inge TH, et al. Nutritional risks in adolescents after bariatric surgery. *Clin Gastroenterol Hepatol*. 2020;18(5):1070-81e5.
99. Nehus EJ, Khoury JC, Inge TH, et al. Kidney outcomes three years after bariatric surgery in severely obese adolescents. *Kidney Int*. 2017;91(2):451–8.
100. Kaar JL, Morelli N, Russell SP, et al. Obstructive sleep apnea and early weight loss among adolescents undergoing bariatric surgery. *Surg Obes Relat Dis*. 2021;17(4):711–7.
101. Derderian SC, Patten L, Kaizer AM, et al. Influence of weight loss on obesity-associated complications after metabolic and bariatric surgery in adolescents. *Obesity (Silver Spring)*. 2020;28(12):2397–404.
102. Nobili V, Carpino G, De Peppo F, et al. Laparoscopic sleeve gastrectomy improves nonalcoholic fatty liver disease-related liver damage in adolescents by reshaping cellular interactions and hepatic adipocytokine production. *J Pediatr*. 2018;194(100–8):e3.
103. Kalra M, Inge T, Garcia V, et al. Obstructive sleep apnea in extremely overweight adolescents undergoing bariatric surgery. *Obes Res*. 2005;13(7):1175–9.
104. Watanabe Y, Yamaguchi T, Tanaka S, et al. Characteristics of childhood onset and post-puberty onset obesity and weight regain after laparoscopic sleeve gastrectomy in Japanese subjects: a subgroup analysis of J-SMART. *Obes Facts*. 2022;15(4):498–507.
105. Hjelmæsæth J, Hertel JK, Holt AH, Benestad B, Seeberg LT, Lindberg M, Halvorsen E, Júlíusson PB, Sandbu R, Lekhal S, Laparoscopic gastric bypass versus lifestyle intervention for adolescents with morbid obesity. *Tidsskr Nor Laegeforen*. English, Norwegian. 2020;140(16). <https://doi.org/10.4045/tidsskr.20.0526>.
106. Tashiro J, McKenna E, Alberto EC, et al. The impact of parental bariatric surgery and patient age on laparoscopic sleeve gastrectomy outcomes in adolescents. *Surg Endosc*. 2022;36(10):7392–8.
107. Lee DY, Guend H, Park K, et al. Outcomes of laparoscopic Roux-en-Y gastric bypass versus laparoscopic adjustable gastric banding in adolescents. *Obes Surg*. 2012;22(12):1859–64.
108. Cozacov Y, Roy M, Moon S, et al. Mid-term results of laparoscopic sleeve gastrectomy and Roux-en-Y gastric bypass in adolescent patients. *Obes Surg*. 2014;24(5):747–52.
109. Pena AS, Delko T, Couper R, et al. Laparoscopic adjustable gastric banding in Australian adolescents: should it be done? *Obes Surg*. 2017;27(7):1667–73.
110. Dumont PN, Blanchet MC, Gignoux B, et al. Medium- to long-term outcomes of gastric banding in adolescents: a single-center study of 97 consecutive patients. *Obes Surg*. 2018;28(1):285–9.
111. Lainas P, De Filippo G, Giuro G, et al. Laparoscopic sleeve gastrectomy for adolescents under 18 years old with severe obesity. *Obes Surg*. 2020;30(1):267–73.
112. Mohammed MR, Mahdy T, Hashem A, et al. Impact of baseline BMI and adherence to follow-up on the outcome of sleeve gastrectomy in treatment of adolescent obesity. *Obes Surg*. 2021;31(6):2567–75.
113. Teeple EA, Teich S, Schuster DP, et al. Early metabolic improvement following bariatric surgery in morbidly obese adolescents. *Pediatr Blood Cancer*. 2012;58(1):112–6.
114. Tuna T, Espinheira MDC, Vasconcelos C, et al. Laparoscopic sleeve gastrectomy in morbidly obese adolescents: initial

- experience of a pediatric multidisciplinary unit. *Arch Pediatr.* 2020;27(6):310–4.
115. White B, Doyle J, Matschull K, Adamo M, Christie D, Nicholls D, Kinra S, Wong IC, Viner RM. Outcomes of 50 patients entering an adolescent bariatric surgery programme. *Archives of disease in childhood.* 2017;102(12):1152–6.
 116. Messiah SE, Lopez-Mitnik G, Winegar D, et al. Effect of ethnicity on weight loss among adolescents 1 year after bariatric surgery. *World J Diabetes.* 2013;4(5):202–9.
 117. Kaulfers AM, Bean JA, Inge TH, et al. Bone loss in adolescents after bariatric surgery. *Pediatrics.* 2011;127(4):e956–61.
 118. Reilly JJ, Kelly J. Long-term impact of overweight and obesity in childhood and adolescence on morbidity and premature mortality in adulthood: systematic review. *Int J Obes (Lond).* 2011;35(7):891–8.
 119. Alqahtani AR, Elahmedi M, Abdurabu HY, et al. Ten-year outcomes of children and adolescents who underwent sleeve gastrectomy: weight loss, comorbidity resolution, adverse events, and growth velocity. *J Am Coll Surg.* 2021;233(6):657–64.
 120. Choi JT, Foster BD, Soraya Heidari K, et al. Effect of bariatric surgery on outcomes in the operative treatment of hip fractures. *Injury.* 2020;51(3):688–93.
 121. Inacio MC, Paxton EW, Fisher D, et al. Bariatric surgery prior to total joint arthroplasty may not provide dramatic improvements in post-arthroplasty surgical outcomes. *J Arthroplasty.* 2014;29(7):1359–64.
 122. Werner BC, Kurkis GM, Gwathmey FW, et al. Bariatric surgery prior to total knee arthroplasty is associated with fewer postoperative complications. *J Arthroplasty.* 2015;30(9 Suppl):81–5.
 123. Nickel BT, Klement MR, Penrose CT, et al. Lingering risk: bariatric surgery before total knee arthroplasty. *J Arthroplasty.* 2016;31(9 Suppl):207–11.
 124. McLawhorn AS, Levack AE, Lee YY, et al. Bariatric surgery improves outcomes after lower extremity arthroplasty in the morbidly obese: a propensity score-matched analysis of a New York statewide database. *J Arthroplasty.* 2018;33(7):2062–9e4.
 125. Lee GC, Ong K, Baykal D, et al. Does prior bariatric surgery affect implant survivorship and complications following primary total hip arthroplasty/total knee arthroplasty? *J Arthroplasty.* 2018;33(7):2070–4e1.
 126. Wang Y, Deng Z, Meng J, et al. Impact of bariatric surgery on inpatient complication, cost, and length of stay following total hip or knee arthroplasty. *J Arthroplasty.* 2019;34(12):2884–9e4.
 127. Sax OC, Chen Z, Bains SS, et al. Timing and type of bariatric surgery preceding total knee arthroplasty leads to similar complications and outcomes. *J Arthroplasty.* 2022;37(8S):S842–8.
 128. Ryan SP, Couch CG, Duong SQ, et al. Frank Stinchfield Award: does bariatric surgery prior to primary total hip arthroplasty really improve outcomes? *J Arthroplasty.* 2022;37(7S):S386–90.
 129. Martin JR, Watts CD, Taunton MJ. Bariatric surgery does not improve outcomes in patients undergoing primary total knee arthroplasty. *Bone Joint J.* 2015;97-B(11):1501–5.
 130. Watts CD, Martin JR, Houdek MT, et al. Prior bariatric surgery may decrease the rate of re-operation and revision following total hip arthroplasty. *Bone Joint J.* 2016;98-B(9):1180–4.
 131. Liu JX, Paoli AR, Mahure SA, et al. Preoperative bariatric surgery utilization is associated with increased 90-day postoperative complication rates after total joint arthroplasty. *J Am Acad Orthop Surg.* 2020;28(5):e206–12.
 132. Nickel BT, Klement MR, Penrose C, et al. Dislocation rate increases with bariatric surgery before total hip arthroplasty. *Hip Int.* 2018;28(5):559–65.
 133. Dowsey MM, Brown WA, Cochrane A, et al. Effect of bariatric surgery on risk of complications after total knee arthroplasty: a randomized clinical trial. *JAMA Netw Open.* 2022;5(4):e226722.
 134. Ighani Arani P, Wretenberg P, Ottosson J, et al. Bariatric surgery prior to total knee arthroplasty is not associated with lower risk of revision: a register-based study of 441 patients. *Acta Orthop.* 2021;92(1):97–101.
 135. Nearing EE 2nd, Santos TM, Topolski MS, et al. Benefits of bariatric surgery before elective total joint arthroplasty: is there a role for weight loss optimization? *Surg Obes Relat Dis.* 2017;13(3):457–62.
 136. Liu J, Zhong H, Poeran J, et al. Bariatric surgery and total knee/hip arthroplasty: an analysis of the impact of sequence and timing on outcomes. *Reg Anesth Pain Med.* 2021;46(11):941–5.
 137. Meller MM, Goodman S, Gonzalez MH, Lau E. Does bariatric surgery normalize risks after total knee arthroplasty? Administrative medicare data. *J Am Acad Orthop Surg Glob Res Rev.* 2019;3(12):e19.00102.
 138. Purcell S, Hossain I, Evans B, et al. Morbid obesity and severe knee osteoarthritis: which should be treated first? *J Gastrointest Surg.* 2022;26(7):1388–93.
 139. Schwarzkopf R, Lavery JA, Hooper J, et al. Bariatric surgery and time to total joint arthroplasty: does it affect readmission and complication rates? *Obes Surg.* 2018;28(5):1395–401.
 140. Ighani Arani P, Wretenberg P, Ottosson J, et al. Pain, function, and satisfaction after total knee arthroplasty, with or without bariatric surgery. *Obes Surg.* 2022;32(4):1164–9.
 141. Severson EP, Singh JA, Browne JA, et al. Total knee arthroplasty in morbidly obese patients treated with bariatric surgery: a comparative study. *J Arthroplasty.* 2012;27(9):1696–700.
 142. Adrados M, Samuel LT, Locklear TM, et al. Institutional adherence to the American Association of Hip and Knee Surgeons body mass index guidelines lowers perioperative emergency department visits in primary total knee arthroplasty. *J Arthroplasty.* 2023;38(6S):S88–93.
 143. Morrell DJ, Pauli EM, Lyn-Sue JR, et al. Laparoscopic sleeve gastrectomy in patients with complex abdominal wall hernias. *Surg Endosc.* 2021;35(7):3881–9.
 144. Moolla M, Dang J, Modasi A, et al. Concurrent laparoscopic ventral hernia repair with bariatric surgery: a propensity-matched analysis. *J Gastrointest Surg.* 2020;24(1):58–66.
 145. Krivan MS, Giorga A, Barreca M, et al. Concomitant ventral hernia repair and bariatric surgery: a retrospective analysis from a UK-based bariatric center. *Surg Endosc.* 2019;33(3):705–10.
 146. Sharma G, Boules M, Punchai S, et al. Outcomes of concomitant ventral hernia repair performed during bariatric surgery. *Surg Endosc.* 2017;31(4):1573–82.
 147. Olmi S, Uccelli M, Cesana GC, et al. Laparoscopic ventral hernia repair in bariatric patients: the role of defect size and deferred repair. *Obes Surg.* 2020;30(10):3905–11.
 148. Raziell A, Sakran N, Szold A, et al. Concomitant bariatric and ventral/incisional hernia surgery in morbidly obese patients. *Surg Endosc.* 2014;28(4):1209–12.
 149. Chandeze MM, Moszkowicz D, Beauchet A, et al. Ventral hernia surgery in morbidly obese patients, immediate or after bariatric surgery preparation: results of a case-matched study. *Surg Obes Relat Dis.* 2019;15(1):83–8.
 150. Praveen Raj P, Senthilnathan P, Kumaravel R, et al. Concomitant laparoscopic ventral hernia mesh repair and bariatric surgery: a retrospective study from a tertiary care center. *Obes Surg.* 2012;22(5):685.
 151. Clapp B, Lee I, Liggett E, et al. Are concomitant operations during bariatric surgery safe? An analysis of the MBSAQIP Database. *Obes Surg.* 2020;30(11):4474–81.
 152. Chan DL, Talbot ML, Chen Z. Simultaneous ventral hernia repair in bariatric surgery. *ANZ J Surg.* 2014;84(7–8):581–3.
 153. Eid GM, Mattar SG, Hamad G, et al. Repair of ventral hernias in morbidly obese patients undergoing laparoscopic gastric bypass should not be deferred. *Surg Endosc.* 2004;18(2):207–10.

154. Moszkowicz D, Jacota M, Nkam L, et al. Ventral hernia repair and obesity: results from a nationwide register study in France according to the timeframes of hernia repair and bariatric surgery. *Obes Surg.* 2021;31(12):5251–9.
155. Newcomb WL, Polhill JL, Chen AY, et al. Staged hernia repair preceded by gastric bypass for the treatment of morbidly obese patients with complex ventral hernias. *Hernia.* 2008;12(5):465–9.
156. Datta T, Eid G, Nahmias N, Dallal RM. Management of ventral hernias during laparoscopic gastric bypass. *Surg Obes Relat Dis.* 2008;4(6):754–7.
157. Spaniolas K, Kasten KR, Mozer AB, et al. Synchronous ventral hernia repair in patients undergoing bariatric surgery. *Obes Surg.* 2015;25(10):1864–8.
158. Eid GM, Wikel KJ, Entabi F, et al. Ventral hernias in morbidly obese patients: a suggested algorithm for operative repair. *Obes Surg.* 2013;23(5):703–9.
159. Khorgami Z, Haskins IN, Aminian A, et al. Concurrent ventral hernia repair in patients undergoing laparoscopic bariatric surgery: a case-matched study using the National Surgical Quality Improvement Program Database. *Surg Obes Relat Dis.* 2017;13(6):997–1002.
160. Praveen Raj P, Bhattacharya S, Saravana Kumar S, et al. Morbid obesity with ventral hernia: is concomitant bariatric surgery with laparoscopic ventral hernia mesh repair the best approach? An experience of over 150 cases. *Surg Obes Relat Dis.* 2019;15(7):1098–103.
161. Praveenraj P, Gomes RM, Kumar S, et al. Concomitant bariatric surgery with laparoscopic intra-peritoneal onlay mesh repair for recurrent ventral hernias in morbidly obese patients: an evolving standard of care. *Obes Surg.* 2016;26(6):1191–4.
162. Borbely Y, Zerkowski J, Altmeier J, et al. Complex hernias with loss of domain in morbidly obese patients: role of laparoscopic sleeve gastrectomy in a multi-step approach. *Surg Obes Relat Dis.* 2017;13(5):768–73.
163. Kaminski DL. The role of gastric restrictive procedures in treating ventral hernias in morbidly obese patients. *Int J Surg Investig.* 2000;2(2):159–64.
164. Marzouk A, Ali HOE. Laparoscopic ventral hernia repair combined with sleeve gastrectomy in morbidly obese patients: early outcomes. *Surg J (N Y).* 2019;5(3):e87–91.
165. Vitiello A, Berardi G, Velotti N, et al. Simultaneous small/medium umbilical hernia repair with laparoscopic sleeve gastrectomy (LSG): results of a retrospective case-matched study. *Surg Laparosc Endosc Percutan Tech.* 2021;31(5):519–22.
166. Alexander JW, Goodman H. Gastric bypass in chronic renal failure and renal transplant. *Nutr Clin Pract.* 2007;22(1):16–21.
167. Modanlou KA, Muthyala U, Xiao H, et al. Bariatric surgery among kidney transplant candidates and recipients: analysis of the United States renal data system and literature review. *Transplantation.* 2009;87(8):1167–73.
168. Jamal MH, Corcelles R, Daigle CR, et al. Safety and effectiveness of bariatric surgery in dialysis patients and kidney transplantation candidates. *Surg Obes Relat Dis.* 2015;11(2):419–23.
169. Mozer AB, Pender JRt, Chapman WH, et al. Bariatric surgery in patients with dialysis-dependent renal failure. *Obes Surg.* 2015;25(11):2088–92.
170. Freeman CM, Woodle ES, Shi J, et al. Addressing morbid obesity as a barrier to renal transplantation with laparoscopic sleeve gastrectomy. *Am J Transplant.* 2015;15(5):1360–8.
171. Al-Bahri S, Fakhry TK, Gonzalvo JP. Bariatric surgery as a bridge to renal transplantation in patients with end-stage renal disease. *Obes Surg.* 2017;27(11):2951–5.
172. Gheith O, Al-Otaibi T, Halim MA, et al. Bariatric surgery in renal transplant patients. *Exp Clin Transplant.* 2017;15(Suppl 1):164–9.
173. Kim Y, Jung AD, Dhar VK, et al. Laparoscopic sleeve gastrectomy improves renal transplant candidacy and posttransplant outcomes in morbidly obese patients. *Am J Transplant.* 2018;18(2):410–6.
174. Thomas IA, Gaynor JJ, Joseph T, et al. Roux-en-Y gastric bypass is an effective bridge to kidney transplantation: results from a single center. *Clin Transplant.* 2018;32(5):e13232.
175. Cohen JB, Lim MA, Tewksbury CM, et al. Bariatric surgery before and after kidney transplantation: long-term weight loss and allograft outcomes. *Surg Obes Relat Dis.* 2019;15(6):935–41.
176. Yemini R, Neshar E, Carmeli I, et al. Bariatric surgery is efficacious and improves access to transplantation for morbidly obese renal transplant candidates. *Obes Surg.* 2019;29(8):2373–80.
177. Yemini R, Neshar E, Winkler J, et al. Bariatric surgery in solid organ transplant patients: long-term follow-up results of outcome, safety, and effect on immunosuppression. *Am J Transplant.* 2018;18(11):2772–80.
178. Bouchard P, Tchervenkov J, Demyttenaere S, et al. Safety and efficacy of the sleeve gastrectomy as a strategy towards kidney transplantation. *Surg Endosc.* 2020;34(6):2657–64.
179. Cohen JB, Tewksbury CM, Torres Landa S, et al. National post-operative bariatric surgery outcomes in patients with chronic kidney disease and end-stage kidney disease. *Obes Surg.* 2019;29(3):975–82.
180. Schindel H, Winkler J, Yemini R, et al. Survival benefit in bariatric surgery kidney recipients may be mediated through effects on kidney graft function and improvement of co-morbidities: a case-control study. *Surg Obes Relat Dis.* 2019;15(4):621–7.
181. Kassam AF, Mirza A, Kim Y, et al. Long-term outcomes in patients with obesity and renal disease after sleeve gastrectomy. *Am J Transplant.* 2020;20(2):422–9.
182. Sheetz KH, Woodside KJ, Shahinian VB, et al. Trends in bariatric surgery procedures among patients with ESKD in the United States. *Clin J Am Soc Nephrol.* 2019;14(8):1193–9.
183. Dobrzycka M, Proczko-Stepaniak M, Kaska L, et al. Weight loss after bariatric surgery in morbidly obese end-stage kidney disease patients as preparation for kidney transplantation. Matched pair analysis in a high-volume bariatric and transplant center. *Obes Surg.* 2020;30(7):2708–14.
184. Outmani L, Kimenai H, Roodnat JJ, et al. Clinical outcome of kidney transplantation after bariatric surgery: a single-center, retrospective cohort study. *Clin Transplant.* 2021;35(3):e14208.
185. Soliman BG, Tariq N, Law YY, et al. Effectiveness of bariatric surgery in increasing kidney transplant eligibility in patients with kidney failure requiring dialysis. *Obes Surg.* 2021;31(8):3436–43.
186. Ku E, McCulloch CE, Roll GR, et al. Bariatric surgery prior to transplantation and risk of early hospital re-admission, graft failure, or death following kidney transplantation. *Am J Transplant.* 2021;21(11):3750–7.
187. Zaminpeyma R, Claus M, Paraskevas S, et al. Outcomes of kidney transplant recipients who underwent pre-transplant bariatric surgery for severe obesity: a long-term follow-up study. *Surg Endosc.* 2023;37(1):494–502.
188. Takata MC, Campos GM, Ciovica R, et al. Laparoscopic bariatric surgery improves candidacy in morbidly obese patients awaiting transplantation. *Surg Obes Relat Dis.* 2008;4(2):159–64.
189. Lin MY, Tavakol MM, Sarin A, et al. Laparoscopic sleeve gastrectomy is safe and efficacious for pretransplant candidates. *Surg Obes Relat Dis.* 2013;9(5):653–8.
190. Diwan TS, Lee TC, Nagai S, et al. Obesity, transplantation, and bariatric surgery: an evolving solution for a growing epidemic. *Am J Transplant.* 2020;20(8):2143–55.
191. Ahmed Z, Khan MA, Vazquez-Montesino LM, et al. Bariatric surgery, obesity and liver transplantation. *Transl Gastroenterol Hepatol.* 2022;7:25.
192. Howell RS, Liu HH, Boinpally H, et al. Outcomes of bariatric surgery: patients with body mass index 60 or greater. *JLSLS.* 2021;25(2):e2020.00089.

193. Banks J, Abouelazayem M, Kaur V, et al. Routine intra-gastric balloon insertion in the management of 'super-super-obese' patients: an obituary? *Obes Surg.* 2021;31(5):2319–23.
194. Nasser H, Ivanics T, Varban OA, et al. Comparison of early outcomes between Roux-en-Y gastric bypass and sleeve gastrectomy among patients with body mass index ≥ 60 kg/m². *Surg Endosc.* 2021;35(6):3115–21.
195. Mahmoud M, Hasanin AM, Mostafa M, et al. Evaluation of super-obesity and super-super-obesity as risk factors for difficult intubation in patients undergoing bariatric surgery. *Surg Obes Relat Dis.* 2021;17(7):1279–85.
196. Dupree A, El Gammal AT, Wolter S, et al. Perioperative short-term outcome in super-super-obese patients undergoing bariatric surgery. *Obes Surg.* 2018;28(7):1895–901.
197. Samuel N, Jalal Q, Gupta A, et al. Mid-term bariatric surgery outcomes for obese patients: does weight matter? *Ann R Coll Surg Engl.* 2020;102(1):54–61.
198. Stephens DJ, Saunders JK, Belsley S, et al. Short-term outcomes for super-super obese (BMI > or =60 kg/m²) patients undergoing weight loss surgery at a high-volume bariatric surgery center: laparoscopic adjustable gastric banding, laparoscopic gastric bypass, and open tubular gastric bypass. *Surg Obes Relat Dis.* 2008;4(3):408–15.
199. Romero-Velez G, Pechman DM, Munoz Flores F, et al. Bariatric surgery in the super-super morbidly obese: outcome analysis of patients with BMI >70 using the ACS-NSQIP database. *Surg Obes Relat Dis.* 2020;16(7):894–9.
200. Schwartz A, Etchehoury L, Collet D. Outcome after laparoscopic gastric bypass for super-super obese patients. *J Visc Surg.* 2013;150(2):145–9.
201. Gonzalez-Heredia R, Sanchez-Johnsen L, Valbuena VS, et al. Surgical management of super-super obese patients: Roux-en-Y gastric bypass versus sleeve gastrectomy. *Surg Endosc.* 2016;30(5):2097–102.
202. Ochner CN, Jochner MC, Caruso EA, et al. Effect of preoperative body mass index on weight loss after obesity surgery. *Surg Obes Relat Dis.* 2013;9(3):423–7.
203. Arapis K, Macrina N, Kadouch D, et al. Outcomes of Roux-en-Y gastric bypass versus sleeve gastrectomy in super-super-obese patients (BMI ≥ 60 kg/m²): 6-year follow-up at a single university. *Surg Obes Relat Dis.* 2019;15(1):23–33.
204. Serrano OK, Tannebaum JE, Cumella L, et al. Weight loss outcomes and complications from bariatric surgery in the super super obese. *Surg Endosc.* 2016;30(6):2505–11.
205. Mehaffey JH, LaPar DJ, Turrentine FE, et al. Outcomes of laparoscopic Roux-en-Y gastric bypass in super-super-obese patients. *Surg Obes Relat Dis.* 2015;11(4):814–9.
206. Nasser H, Ivanics T, Leonard-Murali S, et al. Perioperative outcomes of laparoscopic Roux-en-Y gastric bypass and sleeve gastrectomy in super-obese and super-super-obese patients: a national database analysis. *Surg Obes Relat Dis.* 2019;15(10):1696–703.
207. Thereaux J, Czernichow S, Corigliano N, et al. Five-year outcomes of gastric bypass for super-super-obesity (BMI ≥ 60 kg/m²): a case matched study. *Surg Obes Relat Dis.* 2015;11(1):32–7.
208. Peraglie C. Laparoscopic mini-gastric bypass (LMGB) in the super-super obese: outcomes in 16 patients. *Obes Surg.* 2008;18(9):1126–9.
209. Madhok B, Mahawar KK, Boyle M, et al. Management of super-super obese patients: comparison between mini (one anastomosis) gastric bypass and sleeve gastrectomy. *Obes Surg.* 2016;26(7):1646–9.
210. Gagner M, Gumbs AA, Milone L, et al. Laparoscopic sleeve gastrectomy for the super-super-obese (body mass index >60 kg/m²). *Surg Today.* 2008;38(5):399–403.
211. Taylor JD, Leitman IM, Hon P, et al. Outcome and complications of gastric bypass in super-super obesity versus morbid obesity. *Obes Surg.* 2006;16(1):16–8.
212. Wilkinson KH, Helm M, Lak K, et al. The risk of post-operative complications in super-super obesity compared to super obesity in accredited bariatric surgery centers. *Obes Surg.* 2019;29(9):2964–71.
213. Abeles D, Kim JJ, Tarnoff ME, et al. Primary laparoscopic gastric bypass can be performed safely in patients with BMI > or =60. *J Am Coll Surg.* 2009;208(2):236–40.
214. Parmar C, Abdelhalim MA, Mahawar KK, et al. Management of super-super obese patients: comparison between one anastomosis (mini) gastric bypass and Roux-en-Y gastric bypass. *Surg Endosc.* 2017;31(9):3504–9.
215. Artuso D, Wayne M, Kaul A, et al. Extremely high body mass index is not a contraindication to laparoscopic gastric bypass. *Obes Surg.* 2004;14(6):750–4.
216. Ece I, Yilmaz H, Alptekin H, et al. Comparative effectiveness of laparoscopic sleeve gastrectomy on morbidly obese, super-obese, and super-super obese patients for the treatment of morbid obesity. *Obes Surg.* 2018;28(6):1484–91.
217. Sanchez-Santos R, Vilarrasa N, Pujol J, et al. Is Roux-en-Y gastric bypass adequate in the super-obese? *Obes Surg.* 2006;16(4):478–83.
218. Schmitz SM, Alizai PH, Kroh A, et al. Clinical outcomes after one anastomosis gastric bypass versus sleeve gastrectomy in super-super-obese patients. *Surg Endosc.* 2022;36(6):4401–7.
219. Moon RC, Nelson L, Teixeira AF, et al. Outcomes of Roux-en-Y gastric bypass in the super obese: comparison of body mass index 50–60 kg/m² and ≥ 60 kg/m² with the morbidly obese. *Surg Obes Relat Dis.* 2016;12(2):292–6.
220. Farkas DT, Vemulapalli P, Haider A, et al. Laparoscopic Roux-en-Y gastric bypass is safe and effective in patients with a BMI > or =60. *Obes Surg.* 2005;15(4):486–93.
221. Gould JC, Garren MJ, Boll V, et al. Laparoscopic gastric bypass: risks vs. benefits up to two years following surgery in super-super obese patients. *Surgery.* 2006;140(4):524–9.
222. Kushnir L, Dunnican WJ, Benedetto B, et al. Is BMI greater than 60 kg/m² a predictor of higher morbidity after laparoscopic Roux-en-Y gastric bypass? *Surg Endosc.* 2010;24(1):94–7.
223. Oliak D, Ballantyne GH, Davies RJ, et al. Short-term results of laparoscopic gastric bypass in patients with BMI > or = 60. *Obes Surg.* 2002;12(5):643–7.
224. Fielding GA. Laparoscopic adjustable gastric banding for massive superobesity (> 60 body mass index kg/m²). *Surg Endosc.* 2003;17(10):1541–5.
225. Hering I, Dorries L, Flemming S, et al. Impact of preoperative weight loss achieved by gastric balloon on peri- and postoperative outcomes of bariatric surgery in super-obese patients: a retrospective matched-pair analysis. *Langenbecks Arch Surg.* 2022;407(5):1873–9.
226. Wang C, Yang W, Yang J. Surgical results of laparoscopic Roux-en-Y gastric bypass in super obese patients with BMI ≥ 60 in China. *Surg Laparosc Endosc Percutan Tech.* 2014;24(6):e216-20.
227. Singhal R, Omar I, Madhok B, et al. Effect of BMI on safety of bariatric surgery during the COVID-19 pandemic, procedure choice, and safety protocols - an analysis from the GENEVA Study. *Obes Res Clin Pract.* 2022;16(3):249–53.
228. Myers JA, Sarker S, Shayani V. Treatment of massive super-obesity with laparoscopic adjustable gastric banding. *Surg Obes Relat Dis.* 2006;2(1):37–40.
229. Tichansky DS, DeMaria EJ, Fernandez AZ, et al. Postoperative complications are not increased in super-super obese patients who undergo laparoscopic Roux-en-Y gastric bypass. *Surg Endosc.* 2005;19(7):939–41.

230. Zerrweck C, Maunoury V, Caiazzo R, et al. Preoperative weight loss with intragastric balloon decreases the risk of significant adverse outcomes of laparoscopic gastric bypass in super-super obese patients. *Obes Surg.* 2012;22(5):777–82.
231. Catheline JM, Fysekidis M, Dbouk R, et al. Weight loss after sleeve gastrectomy in super superobesity. *J Obes.* 2012;2012:959260.
232. Torchia F, Mancuso V, Civitelli S, et al. LapBand System in super-superobese patients (>60 kg/m²): 4-year results. *Obes Surg.* 2009;19(9):1211–5.
233. Date RS, Walton SJ, Ryan N, et al. Is selection bias toward super obese patients in the rationing of metabolic surgery justified?—A pilot study from the United Kingdom. *Surg Obes Relat Dis.* 2013;9(6):981–6.
234. Di Betta E, Mittempergher F, Nascimbeni R, et al. Outcome of duodenal switch with a transitory vertical gastroplasty, in super-super-obese patients in an 8-year series. *Obes Surg.* 2008;18(2):182–6.
235. Spyropoulos C, Katsakoulis E, Mead N, et al. Intragastric balloon for high-risk super-obese patients: a prospective analysis of efficacy. *Surg Obes Relat Dis.* 2007;3(1):78–83.
236. Gottig S, Daskalakis M, Weiner S, et al. Analysis of safety and efficacy of intragastric balloon in extremely obese patients. *Obes Surg.* 2009;19(6):677–83.
237. Shuhaiber J, Vitello J. Is gastric bypass associated with more complications in patients weighing >500 lb (>227 kg)? *Obes Surg.* 2004;14(1):43–6.
238. Fazylov RM, Savel RH, Horovitz JH, et al. Association of super-super-obesity and male gender with elevated mortality in patients undergoing the duodenal switch procedure. *Obes Surg.* 2005;15(5):618–23.
239. Kaul A, Singla V, Baksi A, et al. Safety and efficacy of bariatric surgery in advanced liver fibrosis. *Obes Surg.* 2020;30(11):4359–65.
240. Hanipah ZN, Punchai S, McCullough A, et al. Bariatric surgery in patients with cirrhosis and portal hypertension. *Obes Surg.* 2018;28(11):3431–8.
241. Younus H, Sharma A, Miquel R, et al. Bariatric surgery in cirrhotic patients: is it safe? *Obes Surg.* 2020;30(4):1241–8.
242. Quezada N, Maturana G, Irarrazaval MJ, et al. Bariatric surgery in cirrhotic patients: a matched case-control study. *Obes Surg.* 2020;30(12):4724–31.
243. Singh T, Kochhar GS, Goh GB, et al. Safety and efficacy of bariatric surgery in patients with advanced fibrosis. *Int J Obes (Lond).* 2017;41(3):443–9.
244. Salman MA, Mikhail HMS, Nafea MA, et al. Impact of laparoscopic sleeve gastrectomy on fibrosis stage in patients with Child-A NASH-related cirrhosis. *Surg Endosc.* 2021;35(3):1269–77.
245. Mumtaz K, Lipshultz H, Jalil S, et al. Bariatric surgery in patients with cirrhosis: careful patient and surgery-type selection is key to improving outcomes. *Obes Surg.* 2020;30(9):3444–52.
246. Are VS, Knapp SM, Banerjee A, et al. Improving outcomes of bariatric surgery in patients with cirrhosis in the United States: a nationwide assessment. *Am J Gastroenterol.* 2020;115(11):1849–56.
247. Miller A, Jalil S, Hussan H, et al. Apprising the choice of bariatric surgery in patients with compensated cirrhosis with morbid obesity: results of a national analysis. *Updates Surg.* 2021;73(5):1899–908.
248. Klebanoff MJ, Corey KE, Samur S, et al. Cost-effectiveness analysis of bariatric surgery for patients with nonalcoholic steatohepatitis cirrhosis. *JAMA Netw Open.* 2019;2(2):e190047.
249. Shimizu H, Phuong V, Maia M, et al. Bariatric surgery in patients with liver cirrhosis. *Surg Obes Relat Dis.* 2013;9(1):1–6.
250. Dallal RM, Mattar SG, Lord JL, et al. Results of laparoscopic gastric bypass in patients with cirrhosis. *Obes Surg.* 2004;14(1):47–53.
251. Mittal T, Ahuja A, Dey A, et al. Safety and efficacy of laparoscopic sleeve gastrectomy in patients with portal hypertension with liver function of Childs A. *Surg Endosc.* 2022;36(5):2942–8.
252. Minambres I, Rubio MA, de Hollanda A, et al. Outcomes of bariatric surgery in patients with cirrhosis. *Obes Surg.* 2019;29(2):585–92.
253. Vuppalanchi R, McCabe MET, Tandra SR, et al. Safety and efficacy of bariatric surgery in cirrhosis patients with extreme obesity. *Ann Surg.* 2022;275(1):e174–80.
254. Blumer V, Greene SJ, Ortiz M, et al. In-hospital outcomes after bariatric surgery in patients with heart failure. *Am Heart J.* 2020;230:59–62.
255. Sun D, Schaff HV, McKenzie TJ, et al. Safety of bariatric surgery in obese patients with hypertrophic cardiomyopathy. *Am J Cardiol.* 2022;15(167):93–7.
256. Balakumaran K, Jabri A, Haddadin F, et al. Bariatric weight loss surgery improves systolic function in obese patients with heart failure. *Curr Probl Cardiol.* 2023;48(8):101231.
257. Mentias A, Aminian A, Youssef D, et al. Long-term cardiovascular outcomes after bariatric surgery in the Medicare population. *J Am Coll Cardiol.* 2022;79(15):1429–37.
258. Val FRD, Cheon O, Menser T, et al. Bariatric surgery in end-stage heart failure: feasibility in successful attainment of a target body mass index. *J Card Fail.* 2020;26(11):944–7.
259. Alsabrook GD, Goodman HR, Alexander JW. Gastric bypass for morbidly obese patients with established cardiac disease. *Obes Surg.* 2006;16(10):1272–7.
260. Vest AR, Patel P, Schauer PR, et al. Clinical and echocardiographic outcomes after bariatric surgery in obese patients with left ventricular systolic dysfunction. *Circ Heart Fail.* 2016;9(3):e002260.
261. Naslund E, Stenberg E, Hofmann R, et al. Association of metabolic surgery with major adverse cardiovascular outcomes in patients with previous myocardial infarction and severe obesity: a nationwide cohort study. *Circulation.* 2021;143(15):1458–67.
262. Doumouras AG, Wong JA, Paterson JM, et al. Bariatric surgery and cardiovascular outcomes in patients with obesity and cardiovascular disease: a population-based retrospective cohort study. *Circulation.* 2021;143(15):1468–80.
263. Aminian A, Al-Kurd A, Wilson R, et al. Association of bariatric surgery with major adverse liver and cardiovascular outcomes in patients with biopsy-proven nonalcoholic steatohepatitis. *JAMA.* 2021;326(20):2031–42.
264. Lundberg CE, Jamaly S, Adiels M, et al. Surgical treatment of obesity and excess risk of developing heart failure in a controlled cohort study. *ESC Heart Fail.* 2022;9(3):1844–52.
265. Yuan H, Medina-Inojosa JR, Lopez-Jimenez F, et al. The long-term impact of bariatric surgery on development of atrial fibrillation and cardiovascular events in obese patients: an historical cohort study. *Front Cardiovasc Med.* 2021;8:647118.
266. Ng M, Rodgers B, Rehman S, et al. Left ventricular assist device support and longitudinal sleeve gastrectomy combined with diet in bridge to heart transplant. *Tex Heart Inst J.* 2022;49(1):e207521.
267. Benotti PN, Wood GC, Carey DJ, et al. Gastric bypass surgery produces a durable reduction in cardiovascular disease risk factors and reduces the long-term risks of congestive heart failure. *J Am Heart Assoc.* 2017;6(5):e005126.
268. Hoskuldsdottir G, Sattar N, Miftaraj M, et al. *J Am Heart Assoc.* 2021;10(7):e019323.
269. Yang TWW, Johari Y, Burton PR, et al. Bariatric surgery in patients with severe heart failure. *Obes Surg.* 2020;30(8):2863–9.
270. Strzelczyk J, Kalinowski P, Zieniewicz K, et al. The influence of surgical weight reduction on left atrial strain. *Obes Surg.* 2021;31(12):5243–50.
271. Stenberg E, Cao Y, Jernberg T, Naslund E. Safety of bariatric surgery in patients with previous acute coronary events or heart failure: nationwide cohort study. *BJS Open.* 2022;6(3):zrac083.

272. Busetto L, Dicker D, Azran C, et al. Practical Recommendations of the Obesity Management Task Force of the European Association for the Study of Obesity for the Post-Bariatric Surgery Medical Management. *Obes Facts*. 2017;10(6):597–632.
273. Aarts MA, Sivapalan N, Nikzad SE, et al. Optimizing bariatric surgery multidisciplinary follow-up: a focus on patient-centered care. *Obes Surg*. 2017;27(3):730–6.
274. Gould J, Ellsmere J, Fanelli R, et al. Panel report: best practices for the surgical treatment of obesity. *Surg Endosc*. 2011;25(6):1730–40.
275. Warman JL. The application of laparoscopic bariatric surgery for treatment of severe obesity in adolescents using a multidisciplinary adolescent bariatric program. *Crit Care Nurs Q*. 2005;28(3):276–87.
276. Collazo-Clavell ML, Clark MM, McAlpine DE, et al. Assessment and preparation of patients for bariatric surgery. *Mayo Clin Proc*. 2006;81(10 Suppl):S11–7.
277. Elte JW, Castro Cabezas M, Vrijland WW, et al. Proposal for a multidisciplinary approach to the patient with morbid obesity: the St. Franciscus Hospital morbid obesity program. *Eur J Intern Med*. 2008;19(2):92–8.
278. Apovian CM, Cummings S, Anderson W, et al. Best practice updates for multidisciplinary care in weight loss surgery. *Obesity (Silver Spring)*. 2009;17(5):871–9.
279. Di Lorenzo N, Antoniou SA, Batterham RL, et al. Clinical practice guidelines of the European Association for Endoscopic Surgery (EAES) on bariatric surgery: update 2020 endorsed by IFSO-EC. *EASO and ESPCOP Surg Endosc*. 2020;34(6):2332–58.
280. van Hout GC, Vreeswijk CM, van Heck GL. Bariatric surgery and bariatric psychology: evolution of the Dutch approach. *Obes Surg*. 2008;18(3):321–5.
281. Pories WJ, Aronne LJ. It's time for multidisciplinary obesity management centers. *Obesity (Silver Spring)*. 2019;27(4):534.
282. Gill RS, Karmali S, Hadi G, et al. Predictors of attrition in a multidisciplinary adult weight management clinic. *Can J Surg*. 2012;55(4):239–43.
283. Andalib A, Bouchard P, Bougie A, et al. Variability in bariatric surgical care among various centers: a survey of all bariatric surgeons in the province of Quebec Canada. *Obes Surg*. 2018;28(8):2327–32.
284. Aboueid S, Jasinska M, Bourgeault I, et al. Current weight management approaches used by primary care providers in six multidisciplinary healthcare settings in Ontario. *Can J Nurs Res*. 2018;50(4):169–78.
285. Vahibe A, Aizpuru MJ, Sarr MG, et al. Safety and efficacy of reversible surgery as a treatment for malnutrition after bariatric surgery. *J Am Coll Surg*. 2023;236(1):156–66.
286. Vanetta C, Dreifuss NH, Schlottmann F, et al. Bariatric surgery conversions in MBSAQIP centers: current indications and outcomes. *Obes Surg*. 2022;32(10):3248–56.
287. Major P, Zarzycki P, Rymarowicz J, et al. Revisional operations among patients after surgical treatment of obesity: a multicenter Polish Revision Obesity Surgery Study (PROSS). *Wideochir Inne Tech Maloinwazyjne*. 2022;17(2):372–9.
288. Xie J, Dreifuss NH, Schlottmann F, et al. Minimally invasive revisional bariatric surgery in a MBSAQIP accredited high-volume center. *Front Surg*. 2022;9:880044.
289. Hernandez LA, Guilbert L, Sepulveda EM, et al. Causes of revisional surgery, reoperations, and readmissions after bariatric surgery. *Rev Gastroenterol Mex (Engl Ed)*. 2023;88(3):232–7.
290. Gero D, Vannijvel M, Okkema S, et al. Defining global benchmarks in elective secondary bariatric surgery comprising conversational, revisional, and reversal procedures. *Ann Surg*. 2021;274(5):821–8.
291. Dreifuss NH, Mangano A, Hassan C, et al. Robotic revisional bariatric surgery: a high-volume center experience. *Obes Surg*. 2021;31(4):1656–63.
292. King K, Galvez A, Stoltzfus J, et al. Robotic-assisted surgery results in a shorter hospital stay following revisional bariatric surgery. *Obes Surg*. 2021;31(2):634–9.
293. Cheema F, Choi M, Moran-Atkin E, et al. Outcomes in revisional bariatric surgery: a high-volume single institution experience. *Surg Endosc*. 2021;35(7):3932–9.
294. El Chaar M, King K, Pastrana M, et al. Outcomes of robotic surgery in revisional bariatric cases: a propensity score-matched analysis of the MBSAQIP registry. *J Robot Surg*. 2021;15(2):235–9.
295. Mora Oliver I, Cassinello Fernandez N, Alfonso Ballester R, et al. Revisional bariatric surgery due to failure of the initial technique: 25 years of experience in a specialized Unit of Obesity Surgery in Spain. *Cir Esp (Engl Ed)*. 2019;97(10):568–74.
296. Keren D, Romano-Zelekha O, Rainis T, et al. Revisional bariatric surgery in Israel: findings from the Israeli Bariatric Surgery Registry. *Obes Surg*. 2019;29(11):3514.
297. Acevedo E, Mazzei M, Zhao H, et al. Outcomes in conventional laparoscopic versus robotic-assisted revisional bariatric surgery: a retrospective, case-controlled study of the MBSAQIP database. *Surg Endosc*. 2020;34(4):1573–84.
298. Clapp B, Liggett E, Jones R, et al. Comparison of robotic revisional weight loss surgery and laparoscopic revisional weight loss surgery using the MBSAQIP database. *Surg Obes Relat Dis*. 2019;15(6):909–19.
299. Aleassa EM, Hassan M, Hayes K, et al. Effect of revisional bariatric surgery on type 2 diabetes mellitus. *Surg Endosc*. 2019;33(8):2642–8.
300. Qiu J, Lundberg PW, Javier T, et al. Revisional bariatric surgery for weight regain and refractory complications in a single MBSAQIP accredited center: what are we dealing with? *Obes Surg*. 2018;28(9):2789–95.
301. Gray KD, Moore MD, Elmously A, et al. Perioperative outcomes of laparoscopic and robotic revisional bariatric surgery in a complex patient population. *Obes Surg*. 2018;28(7):1852–9.
302. Pires Souto K, Meinhardt NG, de Azevedo Dossin I, et al. Revisional malabsorptive bariatric surgery: 29-year follow-up in a Brazilian public hospital. *Obes Surg*. 2001;28(6):1504–10.
303. Fulton C, Sheppard C, Birch D, et al. A comparison of revisional and primary bariatric surgery. *Can J Surg*. 2017;60(3):205–11.
304. Daigle CR, Chaudhry R, Boules M, et al. Revisional bariatric surgery can improve refractory metabolic disease. *Surg Obes Relat Dis*. 2016;12(2):392–7.
305. Shimizu H, Annaberdyev S, Motamarry I, et al. Revisional bariatric surgery for unsuccessful weight loss and complications. *Obes Surg*. 2013;23(11):1766–73.
306. Kuesters S, Grueneberger JM, Baumann T, et al. Revisional bariatric surgery: indications and outcome of 100 consecutive operations at a single center. *Surg Endosc*. 2012;26(6):1718–23.
307. Fronza JS, Prystowsky JB, Hungness ES, et al. Revisional bariatric surgery at a single institution. *Am J Surg*. 2010;200(5):651–4.
308. Spyropoulos C, Kehagias I, Panagiotopoulos S, et al. Revisional bariatric surgery: 13-year experience from a tertiary institution. *Arch Surg*. 2010;145(2):173–7.
309. Lim CS, Liew V, Talbot ML, et al. Revisional bariatric surgery. *Obes Surg*. 2009;19(7):827–32.
310. Nettet EM, Kendrick ML, Houghton SG, et al. A two-decade spectrum of revisional bariatric surgery at a tertiary referral center. *Surg Obes Relat Dis*. 2007;3(1):25–30.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Maurizio De Luca¹ · Scott Shikora² · Dan Eisenberg³ · Luigi Angrisani⁴ · Chetan Parmar⁵ · Aayed Alqahtani⁶ · Ali Aminian⁷ · Edo Aarts⁸ · Wendy Brown⁹ · Ricardo V. Cohen¹⁰ · Nicola Di Lorenzo¹¹ · Silvia L. Faria¹² · Kasey P. S. Goodpaster¹³ · Ashraf Haddad¹⁴ · Miguel Herrera¹⁵ · Raul Rosenthal¹⁶ · Jacques Himpens¹⁷ · Angelo Iossa¹⁸ · Mohammad Kermansaravi¹⁹ · Lilian Kow²⁰ · Marina Kurian²¹ · Sonja Chiappetta²² · Teresa LaMasters²³ · Kamal Mahawar²⁴ · Giovanni Merola²⁵ · Abdelrahman Nimeri² · Mary O’Kane²⁶ · Pavlos Papasavas²⁷ · Giacomo Piatto²⁸ · Jaime Ponce²⁹ · Gerhard Prager³⁰ · Janey S. A. Pratt³ · Ann M. Rogers³¹ · Paulina Salminen³² · Kimberley E. Steele³³ · Michel Suter³⁴ · Salvatore Tolone³⁵ · Antonio Vitiello³⁶ · Marco Zappa³⁷ · Shanu N. Kothari³⁸

✉ Maurizio De Luca
nnwdel@tin.it

¹ Department of General Surgery, Rovigo Hospital, Rovigo, Italy

² Department of Surgery, Center for Metabolic and Bariatric Surgery, Brigham and Women’s Hospital, and Harvard Medical School, Boston, MA, USA

³ Department of Surgery, Stanford School of Medicine, VA Palo Alto Health Care System, 3801 Miranda Avenue, GS 112, Palo Alto, CA 94304, USA

⁴ Department of Public Health, Federico II University of Naples, Naples, Italy

⁵ Whittington Hospital, London, UK

⁶ New You Medical Center, King Saud University, Riyadh, Saudi Arabia

⁷ Department of General Surgery, Bariatric and Metabolic Institute, Cleveland Clinic, Cleveland, OH, USA

⁸ Weight Works Clinics and Allurion Clinics, Amersfoort, The Netherlands

⁹ Department of Surgery, Central Clinical School, Alfred Health, Monash University, Melbourne, Victoria, Australia

¹⁰ Center for the Treatment of Obesity and Diabetes, Hospital Alemão Oswaldo Cruz, São Paulo, Brazil

¹¹ Department of Surgical Sciences, University of Rome “Tor Vergata”, Rome, Italy

¹² Gastrocirurgia de Brasilia, University of Brasilia, Brasilia, Brazil

¹³ Bariatric and Metabolic Institute, Cleveland Clinic, Cleveland, OH, USA

¹⁴ Gastrointestinal Bariatric and Metabolic Center (GBMC), Jordan Hospital, Amman, Jordan

¹⁵ Endocrine and Bariatric Surgery, UNAM at INCMNSZ, Mexico City, Mexico

¹⁶ Cleveland Clinic Florida, The Bariatric Institute, 2950 Cleveland Clinic Blvd, Weston, FL 33331, USA

¹⁷ Bariatric Surgery Unit, Delta Chirec Hospital, Brussels, Belgium

¹⁸ Department of Medico Surgical Sciences and Biotechnologies Sapienza Polo Pontino, ICOT Hospital Latina, Latina, Italy

¹⁹ Department of Surgery, Minimally Invasive Surgery Research Center, Division of Minimally Invasive

and Bariatric Surgery, Hazrat-e Fatemeh Hospital, Iran University of Medical Sciences, Tehran, Iran

²⁰ Adelaide Bariatric Centre, Flinders University of South Australia, Adelaide, Australia

²¹ Department of Surgery, New York University Grossman School of Medicine, New York, NY, USA

²² Department of General and Laparoscopic Surgery, Obesity and Metabolic Surgery Unit, Ospedale Evangelico Betania, Naples, Italy

²³ Department of Surgery, University of Iowa, Iowa City, IA, USA

²⁴ South Tyneside and Sunderland Foundation NHS Trust, Sunderland, UK

²⁵ General and Laparoscopic Surgery, San Giovanni di Dio Hospital - Frattamaggiore, Naples, Italy

²⁶ Department of Nutrition and Dietetics, Leeds Teaching Hospitals NHS Trust, Leeds, UK

²⁷ Division of Metabolic and Bariatric Surgery, Hartford Hospital, Hartford, CT, USA

²⁸ UOC Chirurgia Generale e d’Urgenza, Ospedale di Montebelluna, Montebelluna, Italy

²⁹ Bariatric Surgery Program, CHI Memorial Hospital, Chattanooga, TN, USA

³⁰ Medical University of Vienna, Vienna, Austria

³¹ Department of Surgery, Penn State Health Milton S. Hershey Medical Center, Hershey, PA, USA

³² Division of Digestive Surgery and Urology, Department of Digestive Surgery, Turku University Hospital, Turku, Finland

³³ NIDDK Metabolic and Obesity Research Unit, National Institutes of Health, Bethesda, MD, USA

³⁴ Department of Visceral Surgery, University Hospital, Lausanne, Switzerland

³⁵ Seconda Università di Napoli, Naples, Italy

³⁶ Department of Advanced Biomedical Sciences, Università Degli Studi Di Napoli “Federico II”, Naples, Italy

³⁷ General Surgery Unit, Asst Fatebenefratelli-Sacco Milan, Milan, Italy

³⁸ Department of Surgery, Prisma Health, University of South Carolina School of Medicine, Greenville, SC, USA