

Purification and characterization of a 40.8-kDa cutinase in ungerminated conidia of *Botrytis cinerea* Pers.: Fr.

Katia Gindro^a *, Roger Pezet^b

^a University of Lausanne, Institute of Systematical Botany and Geobotany, Biology Building, CH-1015 Lausanne, Switzerland

^b Swiss Federal Research Station for Plant Production of Changins, Route de Duillier, CH-1260 Nyon, Switzerland

Received 4 December 1998; received in revised form 14 December 1998; accepted 15 December 1998

Abstract

Cytoplasmic soluble proteins from ungerminated conidia of *Botrytis cinerea* exhibited cutinase activity. A 40.8-kDa cutinase was purified to homogeneity from this crude conidial protein extract. This cutinase does not correspond either to constitutive or to induced lytic cutin enzymes already described by other authors. The possible role of this constitutive cutinase in the induction of other cutinolytic proteins in the early stages of infection of plants by *B. cinerea* is discussed. © 1999 Federation of European Microbiological Societies. Published by Elsevier Science B.V. All rights reserved.

Keywords: *Botrytis cinerea*; Esterase; Cutinase; [³H]Cutin

1. Introduction

Many plant pathogenic fungi penetrate the unwounded epidermal layer of their hosts by enzymatic hydrolysis of the cuticle [1]. Attachment of their spores to the plant surface is considered to be crucial for infection. Fungal esterases and particularly cutinases are involved in these phenomena [2]. A cutin-hydrolyzing enzyme was first described in *Fusarium solani* f. sp. *pisi* [3]. Cutinases were characterized and purified from several fungal pathogens [4–7], bacteria

[8], and pollen [9]. A culture filtrate of *Botrytis cinerea* Pers.: Fr. contained a cutinase induced by tomato cutin, which was the sole carbon source [10]. This extracellular cutinase was purified and the gene was cloned [11]. Recently it was demonstrated that ungerminated conidia of *B. cinerea* contained a constitutive cytoplasmic cutinase [12] which could initiate cutin degradation and release of cutin monomers. These fatty acids are known to induce cutinase gene transcription in germinated conidia of *F. solani* f. sp. *pisi* and enhance the pathogenicity of this fungus [13]. This paper describes the purification and characterization of a constitutive cutinase in ungerminated conidia of *B. cinerea*. The possible role of this enzyme in the early events of infection of grape berries is discussed.

* Corresponding author.

Tel.: +41 (22) 363 43 53; Fax: +41 (22) 362 13 25;
E-mail: katia.gindro@rac.admin.ch

2. Materials and methods

2.1. Organism and growth conditions

B. cinerea Pers.: Fr., isolate P69, was cultivated in Petri dishes on oatmeal agar (OMA, Difco). Cultures were placed at 21°C under alternating light and dark (12 h each) for 1 week. Conidia were harvested according to the method of Pezet and Pont [14] and stored dry at –80°C until required.

2.2. ³H-Labeled tomato cutin preparation

Pure tomato cutin was prepared according to Salinas et al. [10] and labeled with [³H]NaBH₄ (3.7 × 10⁹ Bq; 5.2 × 10¹¹ Bq mmol⁻¹) according to Köller et al. [15]. Cutinase activity was determined in purified fractions using [³H]tomato cutin according to Pascholati et al. [5].

2.3. Extraction and purification of a cytoplasmic constitutive cutinase

Cytoplasmic proteins were extracted from 15 g of ungerminated conidia with glass beads according to the method of Van Etten and Freer [16], slightly modified: buffer was replaced by distilled water. The crude protein extract, placed in dialysis tubing, was concentrated on polyethylene glycol (PEG 20000) to 20 ml, then filtered (Akrodisk Nalgen, 0.2 µm) and stored at –20°C until use. Crude extract, dialyzed against piperazine-HCl buffer (PHB, 10 mM, pH 5.5), was loaded on a Superdex Prep Grade column (Pharmacia, 1.5 × 30 cm) and eluted with the same buffer. Non-specific esterase activity was measured in the eluted fractions using *para*-nitrophenylbutyrate (PNB) according to Pascholati et al. [6]. Active fractions were pooled and concentrated to 10 ml on PEG 20000 as described before, then loaded on a DEAE Sepharose Cl-6B column (Pharmacia, 1.5 × 10 cm) suspended in PHB. A step gradient system of NaCl in the buffer (0, 0.05, 0.1, 0.15, 0.2, 0.25, 0.3 M) was used to elute eight fractions. The 0.2 M NaCl fraction contained non-specific esterase activity. This fraction was dialyzed against PHB, concentrated to 6 ml on PEG 20000 and 30% (w/v) of (NH₄)₂SO₄ was added. It was loaded on a butyl Sepharose 4B column (Pharmacia, 1 × 15

cm) in PHB added with 30% (w/v) of (NH₄)₂SO₄. A decreasing step gradient system of (NH₄)₂SO₄ (30% to 0% in 5% steps) was used to elute seven fractions. These fractions were extensively dialyzed against distilled water and concentrated to 4 ml on PEG 20000 before the PNB assay, where (NH₄)₂SO₄ interferes strongly. The active fractions (5% and 10% (NH₄)₂SO₄) were concentrated to 2 ml using microdialysis (Microsep, cut-off 10000). Protein concentration was determined using the method of Bradford [18] (protein determination kit, Bio-Rad).

2.4. Gel electrophoresis analysis

SDS-PAGE (4%/12.5%) was performed according to Hames [19]. Protein bands were revealed by the silver staining method (Bio-Rad Silver Stain Kit) and esterase activity was revealed on renatured SDS-PAGE by incubation of the gel for 1 h at room temperature in Tris-HCl buffer (50 mM, pH 7.0+2% (w/v) Triton X-100) followed by incubation in the same buffer with α,β-naphthyl acetate according to Shaw and Prasad [17]. The molecular mass of cutinase was determined by SDS-PAGE using low-range molecular mass prestained standards (111–20.5 kDa, Bio-Rad). The isoelectric point of the purified protein was determined by IEF analysis on precoated gels (Serva, 3–6 pH gradient, 300 µm, 125 × 125 cm, pI marker protein test mixture 9) according to the manufacturer.

2.5. Immunoblotting

Proteins separated on SDS-PAGE were transferred to PVDF membranes (Immobilon-P, Millipore) for 1.5 h at 100 V in electrotransfer buffer (Tris 25 mM, glycine 192 mM, methanol 20% v/v, pH 8.3). The membranes were then soaked overnight at 4°C in blocking buffer (Tris saline blocking buffer (TSBB): Tris-HCl 10 mM pH 7.5; NaCl 100 mM; Tween-20 0.1% (v/v); BSA 5% (w/v)). After three 5-min washes (TSBB without BSA) they were exposed for 3 h at room temperature to primary antibody (10 µg ml⁻¹ in TSBB containing 1% (w/v) BSA) raised against a constitutive esterase isolated from *B. cinerea* conidia [20] or an induced cutinase found in the culture filtrate [21] (mAb 14E5 and mAb 21C5 respectively, provided by Dr. A. Schots,




Fig. 1. Electrophoretic analysis of cutinase extracted from ungerminated conidia of *B. cinerea* Pers.: Fr. Lane 1: Molecular mass standards (silver stain coloration). Lane 2: SDS-PAGE of the 10% $(\text{NH}_4)_2\text{SO}_4$ fraction. Lane 3: SDS-PAGE of the pure cutinase (5% butyl $(\text{NH}_4)_2\text{SO}_4$ Sepharose fraction, silver stain coloration). Lane 4: Renaturated pure cutinase on SDS-PAGE (5% butyl $(\text{NH}_4)_2\text{SO}_4$ Sepharose fraction, α,β -naphthyl acetate coloration). Lane 5: IEF of the pure cutinase (5% butyl $(\text{NH}_4)_2\text{SO}_4$ Sepharose fraction, α,β -naphthyl acetate coloration). Lane 6: pI standards (Coomassie coloration).

Laboratory of Monoclonal Antibody, Wageningen, The Netherlands). Positive reactions were revealed with a goat anti-mouse IgG conjugated to alkaline phosphatase (Sigma, dilution 1/3000 in TSBB). Alkaline phosphatase activity was revealed with the fast red/naphthol method (Sigma Fast Red[®]/Naphthol AS-MX, N° F-4523) according to the manufacturer.

3. Results and discussion

Linskens and Haage [22] first provided evidence that penetration by *B. cinerea* in potato leaf cutin

in vitro was linked to a cutinolytic activity. An 18-kDa cutinase was isolated by Salinas [21] in culture filtrates of *B. cinerea*. More recently, Comménil et al. [23] described a lipase excreted by *B. cinerea* which exhibited strong cutinolytic activity. Purified polyclonal antibody raised against this 60-kDa protein suppressed *B. cinerea* lesion formation on tomato leaves.

We have demonstrated previously that ungerminated conidia of *B. cinerea* contained a constitutive cytoplasmic cutinase [12], and we have now described its purification to homogeneity using gel filtration, ion exchange and hydrophobic interaction chromatography (Table 1). The 5% $(\text{NH}_4)_2\text{SO}_4$ fraction separated on butyl Sepharose contained a single

Table 1

Purification steps of the constitutive cutinase in ungerminated conidia of *B. cinerea* Pers.: Fr

Purification step	Volume (ml)	Total protein ($\mu\text{g } \mu\text{l}^{-1}$)	Total enzyme ^a ($\mu\text{mol min}^{-1}$)	Specific activity ^a ($\mu\text{mol min}^{-1} \mu\text{g}^{-1}$)	Purification	Yield (%)
Crude extract	20	17.00	24.0	7.0	1.0	100
Superdex	10	0.40	7.9	98.7	14.1	32.5
DEAE (0.2 M NaCl)	6	0.04	3.6	446.0	63.7	14.9
Butyl Sepharose (5% $(\text{NH}_4)_2\text{SO}_4$)	2	0.01	1.2	640.0	91.4	5
Butyl Sepharose (10% $(\text{NH}_4)_2\text{SO}_4$)	2	0.03	0.7	466.0	66.5	2.9

^aDetermined with PNB as the substrate.

band with a molecular mass of 40.8 kDa revealed on SDS-PAGE by silver staining (Fig. 1) and with α , β -naphthyl acetate (Fig. 1). This fraction applied to ^3H -labeled cutin released ethyl acetate-soluble radioactive cutin monomers (Table 2). PNB hydrolytic activity (Table 1) and specific cutinase activity (Table 2) were also measured in the 10% $(\text{NH}_4)_2\text{SO}_4$ fraction (Table 2). Apart from the 40.8-kDa band, other unknown proteins were revealed in this latter fraction which will require further purification (Fig. 1). The constitutive 40.8-kDa cutinase does not correspond to other cutinases isolated from *B. cinerea*. Salinas [21] described a 110–111-kDa constitutive esterase isolated from *B. cinerea* conidia with an assumed cutinolytic activity and a 18-kDa cutinase purified from culture filtrates. mAb 14E5 raised against this constitutive esterase and mAb 21C5 raised against the excreted cutinase did not reveal the 40.8-kDa cutinase (Fig. 2). However, mAb 14E5 has revealed a 110-kDa protein present in a 2-week-old culture filtrate of *B. cinerea* (Fig. 2). Salinas [21] has described this 110-kDa constitutive conidial putative cutinase as a membrane-bound protein. The extraction protocols of conidial proteins used in this work did not allow the solubilization of such proteins. We can conclude that ungerminated conidia may contain two constitutive cutinolytic enzymes, one soluble (40.8 kDa) and another bound to membranes or cell walls (110 kDa), which are excreted at conidial germination and produced during mycelial growth. Further research will be necessary to elucidate this hypothesis.

Isoelectric focusing analysis of the purified 40.8-kDa cutinase has revealed seven isoforms with acidic properties (pI 3.5–4.2) (Fig. 1). This constitutive cutinase might be excreted during the early events of conidial germination. It could release, through cutin degradation, sufficient amounts of cutin monomers,




Fig. 2. Demonstration by immunoblotting that the 40.8-kDa cutinase does not react with mAb 14E5 and mAb 21C5 raised against a constitutive esterase in ungerminated conidia of *B. cinerea* and an induced 18-kDa cutinase purified from culture filtrates of *B. cinerea*. Lane 1: Prestained molecular mass standards. Lane 2: Detection of an 18-kDa cutinase with mAb 21C5 in 2-week-old cultures of *B. cinerea*. Lane 3: Detection of a 110-kDa constitutive conidial esterase with mAb 14E5 in 2-week-old cultures of *B. cinerea*. Lane 4: Pure 40.8-kDa cutinase (5% butyl $(\text{NH}_4)_2\text{SO}_4$ Sepharose fraction) is not detected by either mAb 21C5 or mAb 14E5.

such as 16-hydroxyhexadecanoic acid, to activate the *cutA* gene expression, described by Van der Vlugt-Bergmans [11], which is responsible for the synthesis of the 18-kDa induced cutinase. Both 40.8-kDa con-

Table 2

Radioactivity released from ^3H -labeled cutin by the pure 40.8-kDa constitutive cutinase contained in the butyl Sepharose (5% $(\text{NH}_4)_2\text{SO}_4$) fraction and in the partially purified 40.8-kDa cutinase (10% $(\text{NH}_4)_2\text{SO}_4$), compared to radioactivity released by cutinolytic activity in the crude extract

Extract	CPM ($\times 10^5$)	Specific activity (CPM $\times 10^4 \mu\text{g}^{-1}$ protein)
Control	12.0 \pm 0.4	0.00 \pm 0.00
Crude extract	38.0 \pm 0.7	0.21 \pm 0.02
Butyl Sepharose (5% $(\text{NH}_4)_2\text{SO}_4$)	52.0 \pm 0.6	436 \pm 2.09
Butyl Sepharose (10% $(\text{NH}_4)_2\text{SO}_4$)	41.0 \pm 0.4	273 \pm 1.21

stitutive and 18-kDa induced cutinase could be strongly implicated in early infection processes by *B. cinerea* preceding latent stages and final plant colonization. The role of the 110-kDa cutinase is still unknown. The raising of polyclonal antibodies against the 40.8-kDa cutinase is in progress and such antibodies should be useful to study its role during conidial germination, its development during mycelial growth, and to explain some aspects of the complex enzymatic events of the infection process of *B. cinerea*.

Acknowledgments

The authors thank Dr. A. Schots (Laboratory of Monoclonal Antibodies, Wageningen Agricultural University, Wageningen, The Netherlands) for providing 14E5 and 21C5 monoclonal antibodies, Drs. A. and T.C. Caesar and Dr. M. Cole for critically reading the manuscript, and Mrs. I. De Groote for valuable technical assistance.

References

- [1] Köller, W. (1990) Plant cuticles: the first barrier to be overcome by plant pathogens. In: *The Fungal Spore and Disease Initiation in Plants and Animals* (Cole, G.T. and Hoch, H.C., Eds.), pp. 219–240. Plenum, New York.
- [2] Schäfer, W. (1994) Molecular mechanisms of fungal pathogenicity to plants. *Annu. Rev. Phytopathol.* 32, 461–477.
- [3] Purdy, R.E. and Kolattukudy, P.E. (1975) Hydrolysis of plant cuticle by plant pathogens. Purification, amino acid composition, and molecular weight of two isozymes of cutinase and a non specific esterase from *Fusarium solani* f. *pisi*. *Biochemistry* 14, 2824–2831.
- [4] Köller, W. and Parker, D.M. (1989) Purification and characterization of cutinase from *Venturia inaequalis*. *Phytopathology* 79, 278–283.
- [5] Pascholati, S.F., Deising, H., Leite, B., Anderson, D. and Nicholson, R.L. (1993) Cutinase and non-specific esterase activities in the conidial mucilage of *Colletotrichum graminicola*. *Physiol. Mol. Plant Pathol.* 42, 37–51.
- [6] Pascholati, S.F., Yoshioka, H., Kunoh, H. and Nicholson, R.L. (1992) Preparation of the infection court by *Erysiphe graminis* f. sp. *hordei*: cutinase is a component of the conidial exudate. *Physiol. Mol. Plant Pathol.* 41, 53–59.
- [7] Trail, F. and Köller, W. (1993) Diversity of cutinases from plant pathogenic fungi: purification and characterization of two cutinases from *Alternaria brassicicola*. *Physiol. Mol. Plant Pathol.* 42, 205–220.
- [8] Sebastian, J. and Kolattukudy, P.E. (1988) Purification and characterization of cutinase from a fluorescent *Pseudomonas putida* bacterial strain isolated from phyllosphere. *Arch. Biochem. Biophys.* 263, 77–85.
- [9] Maiti, I.B., Kolattukudy, P.E. and Shaykh, M. (1979) Purification and characterization of a novel cutinase from *Nasturtium (Tropaeolum majus)* pollen. *Arch. Biochem. Biophys.* 196, 412–423.
- [10] Salinas, J., Warnaar, F. and Verhoeff, K. (1986) Production of cutin hydrolyzing enzymes by *Botrytis cinerea* in vitro. *J. Phytopathol.* 116, 299–307.
- [11] Van Der Vlugt-Bergmans, C.J.B., Wagemakers, C.A.M. and Van Kan, J.A.L. (1997) Cloning and expression of the cutinase A gene of *Botrytis cinerea*. *Mol. Plant-Microb. Interact.* 10, 21–29.
- [12] Gindro, K. and Pezet, R. (1997) Evidence for a constitutive cytoplasmic cutinase in ungerminated conidia of *Botrytis cinerea* Pers.: Fr. *FEMS Microbiol. Lett.* 149, 89–92.
- [13] Woloshuk, C.P. and Kolattukudy, P.E. (1986) Mechanisms by which contact with plant cuticle triggers cutinase gene expression in the spores of *Fusarium solani* f. sp. *pisi*. *Proc. Natl. Acad. Sci. USA* 83, 1704–1708.
- [14] Pezet, R. and Pont, V. (1990) Ultrastructural observations of pterostilbene fungitoxicity in dormant conidia of *Botrytis cinerea* Pers. *J. Phytopathol.* 129, 19–30.
- [15] Köller, W., Allan, C.R. and Kolattukudy, P.E. (1982) Role of cutinase and cell wall degrading enzymes in infection of *Pisum sativum* by *Fusarium solani* f. sp. *pisi*. *Physiol. Plant Pathol.* 20, 47–60.
- [16] Van Etten, J.L. and Freer, S.N. (1978) Simple procedure for disruption of fungal spores. *Appl. Environ. Microbiol.* 35, 622–623.
- [17] Shaw, C.R. and Prasad, R. (1970) Starch gel electrophoresis of enzymes – a compilation of recipes. *Biochem. Genet.* 4, 297–330.
- [18] Bradford, M.M. (1976) A rapid and sensitive method for the quantification of microgram quantities of protein utilising the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254.
- [19] Hames, B.D. (1990) One-dimensional polyacrylamide gel electrophoresis. In: *Gel Electrophoresis of Proteins. A Practical Approach* (Hames, B.D. and Rickwood, D., Eds.), pp. 30–38. Oxford University Press, Oxford.
- [20] Salinas, J. and Schots, A. (1994) Monoclonal antibodies-based immunofluorescence test for detection of conidia of *Botrytis cinerea* on cut flowers. *Phytopathology* 84, 351–356.
- [21] Salinas, J. (1992) Function of Cutinolytic Enzymes in the Infection of Gerbera Flowers by *Botrytis cinerea*. PhD Thesis, University of Utrecht, Utrecht.
- [22] Linskens, H.F. and Haage, P. (1963) Cutinase-Nachweis in phytopathogene Pilzen. *Phytopathol. Z.* 48, 306–311.
- [23] Comménil, P., Belingheri, L. and Dehorter, B. (1998) Anticlutinase antibodies prevent infection of tomato leaves by *Botrytis cinerea*. *Physiol. Mol. Plant Pathol.* 52, 1–14.