

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: Genetic studies of body mass index yield new insights for obesity biology.

Authors: Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, Powell C, Vedantam S, Buchkovich ML, Yang J, Croteau-Chonka DC, Esko T, Fall T, Ferreira T, Gustafsson S, Kutalik Z, Luan J, Mägi R, Randall JC, Winkler TW, Wood AR, Workalemahu T, Faul JD, Smith JA, Hua Zhao J, Zhao W, Chen J, Fehrmann R, Hedman ÅK, Karjalainen J, Schmidt EM, Absher D, Amin N, Anderson D, Beekman M, Bolton JL, Bragg-Gresham JL, Buyske S, Demirkan A, Deng G, Ehret GB, Feenstra B, Feitosa MF, Fischer K, Goel A, Gong J, Jackson AU, Kanoni S, Kleber ME, Kristiansson K, Lim U, Lotay V, Mangino M, Mateo Leach I, Medina-Gomez C, Medland SE, Nalls MA, Palmer CD, Pasko D, Pechlivanis S, Peters MJ, Prokopenko I, Shungin D, Stančáková A, Strawbridge RJ, Ju Sung Y, Tanaka T, Teumer A, Trompet S, van der Laan SW, van Setten J, Van Vliet-Ostaptchouk JV, Wang Z, Yengo L,

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.



Published in final edited form as:

Nature. 2015 February 12; 518(7538): 197–206. doi:10.1038/nature14177.

Genetic studies of body mass index yield new insights for obesity biology

A full list of authors and affiliations appears at the end of the article.

These authors contributed equally to this work.

Abstract

Obesity is heritable and predisposes to many diseases. To understand the genetic basis of obesity better, here we conduct a genome-wide association study and MetaboChip meta-analysis of body mass index (BMI), a measure commonly used to define obesity and assess adiposity, in up to 339,224 individuals. This analysis identifies 97 BMI-associated loci ($P < 5 \times 10^{-8}$), 56 of which are novel. Five loci demonstrate clear evidence of several independent association signals, and many loci have significant effects on other metabolic phenotypes. The 97 loci account for ~2.7% of BMI variation, and genome-wide estimates suggest that common variation accounts for >20% of BMI variation. Pathway analyses provide strong support for a role of the central nervous system in obesity susceptibility and implicate new genes and pathways, including those related to synaptic function, glutamate signalling, insulin secretion/action, energy metabolism, lipid biology and adipogenesis.

Obesity is a worldwide epidemic associated with increased morbidity and mortality that imposes an enormous burden on individual and public health. Around 40–70% of inter-individual variability in BMI, commonly used to assess obesity, has been attributed to genetic factors^{1–3}. At least 77 loci have previously been associated with an obesity measure⁴, 32 loci from our previous meta-analysis of BMI genome-wide association studies (GWAS)⁵. Nevertheless, most of the genetic variability in BMI remains unexplained. Moreover, although analyses of previous genetic association results have suggested intriguing biological processes underlying obesity susceptibility, few specific genes supported these pathways^{5,6}. For the vast majority of loci, the probable causal gene(s) and pathways remain unknown.

Correspondence and requests for materials should be addressed to E.K.S. (espeliot@med.umich.edu), R.J.F.L. (ruth.loos@mssm.edu), and J.N.H. (joelh@broadinstitute.org).

†Present address: Second Floor, B-dong, AICT Building, 145 Gwanggyo-ro, Yeongyong-gu, Suwon-si, Gyeonggi-do, 443-270, South Korea.

‡A list of authors and affiliations appears in the Supplementary Information.

§These authors jointly supervised this work.

Online Content Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.

Supplementary Information is available in the online version of the paper.

Author Contributions A full list of author contributions can be found in the Supplementary Information.

Author Information Reprints and permissions information is available at www.nature.com/reprints.

The authors declare competing financial interests: details are available in the online version of the paper. Readers are welcome to comment on the online version of the paper.

To expand the catalogue of BMI susceptibility loci and gain a better understanding of the genes and biological pathways influencing obesity, we performed the largest GWAS meta-analysis for BMI so far. This work doubles the number of individuals contributing GWAS results, incorporates results from >100,000 individuals genotyped with MetaboChip⁷, and nearly doubles the number of BMI-associated loci. Comprehensive assessment of meta-analysis results provides several lines of evidence supporting candidate genes at many loci and highlights pathways that reinforce and expand our understanding of biological processes underlying obesity.

Identification of 97 genome-wide significant loci

This BMI meta-analysis included association results for up to 339,224 individuals from 125 studies, 82 with GWAS results ($n = 236,231$) and 43 with results from MetaboChip ($n = 103,047$; Extended Data Table 1 and Supplementary Tables 1–3). After regression on age and sex and inverse normal transformation of the residuals, we carried out association analyses with genotypes or imputed genotype dosages. GWAS were meta-analysed together, as were MetaboChip studies, followed by a combined GWAS plus MetaboChip meta-analysis. In total, we analysed data from 322,154 individuals of European descent and 17,072 individuals of non-European descent (Extended Data Fig. 1).

Our primary meta-analysis of European-descent individuals from GWAS and MetaboChip studies ($n = 322,154$) identified 77 loci reaching genome-wide significance (GWS) and separated by at least 500 kilo-bases (kb) (Table 1, Extended Data Table 2 and Supplementary Figs 1 and 2). We carried out additional analyses to explore the effects of power and heterogeneity. The inclusion of 17,072 non-European-descent individuals (total $n = 339,224$) identified ten more loci, while secondary analyses identified another ten GWS loci (Table 2, Supplementary Tables 4–8 and Supplementary Figs 3–9). Of the 97 BMI-associated loci, 41 have previously been associated with one or more obesity measure^{5,8–12}. Thus, our current analyses identified 56 novel loci associated with BMI (Tables 1 and 2 and Extended Data Table 2).

Effects of associated loci on BMI

Newly identified loci generally have lower minor allele frequency and/or smaller effect size estimates than previously known loci (Extended Data Fig. 2a, b). On the basis of effect estimates in the discovery data set, which may be inflated owing to winner's curse, the 97 loci account for 2.7% of BMI phenotypic variance (Supplementary Table 4 and Extended Data Fig. 2a, b). We conservatively used only GWS single nucleotide polymorphisms (SNPs) after strict double genomic control correction, which probably over-corrects association statistics given the lack of evidence for population stratification in family-based analyses¹³ (Extended Data Fig. 3 and Extended Data Table 1). Polygene analyses suggest that SNPs with P values well below GWS add significantly to the phenotypic variance explained. For example, 2,346 SNPs selected from conditional and joint multiple-SNP analysis with $P < 5 \times 10^{-3}$ explained $6.6 \pm 1.1\%$ (mean \pm s.e.m.) of variance, compared to $21.6 \pm 2.2\%$ explained by all HapMap3 SNPs (31–54% of heritability; Fig. 1a). Furthermore, of 1,909 independent SNPs (pairwise distance >500 kb and $r^2 < 0.1$) included

on Metabochip for replication of suggestive BMI associations, 1,458 (76.4%) have directionally consistent effects with our previous GWAS meta-analysis⁵ and the non-overlapping samples in the current meta-analysis (Extended Data Fig. 2c). On the basis of the significant excess of these directionally consistent observations (sign test $P = 2.5 \times 10^{-123}$), we estimate ~1,007 of the 1,909 SNPs represent true BMI associations.

We compared the effects of our 97 BMI-associated SNPs between the sexes, between ethnicities, and across several cross-sections of our data (Supplementary Tables 4–11 and Extended Data Fig. 4). Two previously identified loci, near *SEC16B* ($P = 5.2 \times 10^{-5}$) and *ZFP64* ($P = 9.1 \times 10^{-5}$), showed evidence of heterogeneity between men and women. Both have stronger effects in women (Supplementary Table 10). Two SNPs, near *NEGR1* ($P = 9.1 \times 10^{-5}$) and *PRKDI* ($P = 1.9 \times 10^{-5}$), exhibited significant evidence for heterogeneity of effect between European- and African-descent samples, and one SNP, near *GBE1* ($P = 1.3 \times 10^{-4}$), exhibited evidence for heterogeneity between European and east Asian individuals (Supplementary Table 9). These findings may reflect true heterogeneity at these loci, but are most likely due to linkage disequilibrium (LD) differences across ancestries. Effect estimates for 79% of BMI-associated SNPs in African-descent samples ($P = 9.2 \times 10^{-9}$) and 91% in east Asian samples ($P = 1.8 \times 10^{-15}$) showed directional consistency with our European-only analyses. These results suggest that common BMI-associated SNPs have comparable effects across ancestries and between sexes. In additional heterogeneity analyses, we detected an influence of ascertainment at *TCF7L2* (stronger effects in type 2 diabetes case/control studies than in population-based studies); however, we saw no evidence of systematic ascertainment bias at other loci owing to inclusion of case/control studies (Supplementary Tables 10 and 11).

We also took advantage of LD differences across populations to fine-map association signals using Bayesian methods^{14,15}. At 10 of 27 loci fine-mapped for BMI on Metabochip, the addition of non-European individuals into the meta-analysis either narrowed the genomic region containing the 99% credible set, or decreased the number of SNPs in the credible set (Supplementary Table 12 and Supplementary Fig. 10). At the *SEC16B* and *FTO* loci, the all ancestries credible set includes a single SNP, although the SNP we highlight at *FTO* (rs1558902) differs from that identified by a recent fine-mapping effort in African-American cohorts¹⁶. Fine-mapping efforts using larger, more diverse study samples and more complete catalogues of variants will help to further narrow association signals.

We examined the combined effects of lead SNPs at the 97 loci in an independent sample of 8,164 European-descent individuals from the Health and Retirement Study¹⁷. We observed an average increase of 0.1 BMI units (kg per m^2) per BMI-increasing allele, equivalent to 260–320 g for an individual 160–180 cm in height. There was a 1.8 kg per m^2 difference in mean BMI between the 145 individuals (1.78%) carrying the most BMI-increasing alleles (>104) and those carrying the mean number of BMI-increasing alleles in the sample (91; Extended Data Fig. 2d), corresponding to a difference of 4.6–5.8 kg for an individual 160–180 cm in height, and a 1.5 kg m^{-2} difference (3.8–4.9 kg difference) in mean BMI between the 95 individuals (1.16%) carrying the least BMI-increasing alleles (<78) and those carrying the mean number. Such differences are medically significant in predisposing to development of metabolic disease¹⁸. For predicting obesity ($\text{BMI} \geq 30 \text{ kg per m}^2$), adding

genetic risk score to a model including age, age squared, sex and four genotype-based principal components slightly, but significantly increases the area under the receiver-operating characteristic curve from 0.576 to 0.601.

Additional associated variants at BMI loci

To identify additional SNPs with independent BMI associations at the 97 established loci, we used genome-wide complex trait analysis (GCTA)¹⁹ to perform approximate joint and conditional association analysis²⁰ using summary statistics from European sex-combined meta-analysis after removing family-based validation studies (TwinGene and QIMR). GCTA confirmed two signals at *MC4R* previously identified using exact conditional analyses⁵, and identified five loci with evidence of independent associations (Table 3): second signals near *LINC01122*, *NLRC3-ADCY9*, *GPRC5B-GP2* and *BDNF*, and a third signal near *MC4R* (rs9944545, Fig. 1b). Joint conditional analyses at two genomic regions separated by >500 kb (the *AGBL4-ELAVL4* regions on chr. 1, and the *ATP2A1-SBKI* regions on chr. 16), indicate that these pairs of signals may not be independent owing to extended LD.

Effects of BMI variants on other traits

We tested for associations between our 97 BMI-associated index SNPs and other metabolic phenotypes (Supplementary Tables 13–15 and Extended Data Figs 5 and 6). Thirteen of the twenty-three phenotypes tested had significantly more SNPs with effects in the anticipated direction than expected by chance (Supplementary Table 16). These results corroborate the epidemiological relationships of BMI with metabolic traits. Whether this reflects a common genetic aetiology or a causal relationship of BMI on these traits requires further investigation.

Interestingly, some loci showed significant association with traits in the opposite direction than expected based on their phenotypic correlation with BMI (Extended Data Fig. 5). For example, at *HHIP*, the BMI-increasing allele is associated with decreased type 2 diabetes risk and higher high-density lipoprotein cholesterol (HDL). At *LOC646736* and *IRSI*, the BMI-increasing allele is associated with reduced risk of coronary artery disease (CAD) and diabetic nephropathy, decreased triglyceride levels, increased HDL, higher adiponectin, and lower fasting insulin. This may be due to increased subcutaneous fat and possible production of metabolic mediators protective against the development of metabolic disease despite increased adiposity⁸. These unexpected associations may help us to understand better the complex pathophysiology underlying these traits, and may indicate benefits or side effects if these regions contain targets of therapeutic intervention. Furthermore, of our 97 GWS loci, 35 (binomial $P = 0.0019$) were in high LD ($r^2 > 0.7$) with one or more GWS SNPs in the National Human Genome Research Institute (NHGRI) GWAS catalogue ($P < 5 \times 10^{-8}$), even after removing anthropometric trait-associated SNPs. These SNPs were associated not only with cardiometabolic traits, but also with schizophrenia, smoking behaviour, irritable bowel syndrome, and Alzheimer's disease (Supplementary Table 17a, b).

BMI tissues, biological pathways and gene sets

We anticipated the expanded sample size would not only identify additional BMI-associated variants, but also more clearly highlight the biology implicated by genetic studies of BMI. By applying multiple complementary methods, we identified biologically relevant tissues, pathways and gene sets, and highlighted potentially causal genes at associated loci. These approaches included systematic methods incorporating diverse data types, including the novel approach, Data-driven Expression Prioritized Integration for Complex Traits (DEPICT)²¹, and extensive manual review of the literature.

DEPICT used 37,427 human gene expression microarray samples to identify tissues and cell types in which genes near BMI-associated SNPs are highly expressed, and then tested for enrichment of specific tissues by comparing results with randomly selected loci matched for gene density. In total, 27 out of 31 significantly enriched tissues were in the central nervous system (CNS) (out of 209 tested; Fig. 2a and Supplementary Table 18). Current results are not sufficient to isolate specific brain regions important in regulating BMI. However, we observe enrichment not only in the hypothalamus and pituitary gland—key sites of central appetite regulation—but even more strongly in the hippocampus and limbic system, tissues that have a role in learning, cognition, emotion and memory.

As a complementary approach, we examined overlap of associated variants at the 97 loci ($r^2 > 0.7$ with the lead SNP) with five regulatory marks found in most of the 14 selected cell types from brain, blood, liver, pancreatic islet and adipose tissue from the ENCODE Consortium²² and Roadmap Epigenomics Project²³ (Supplementary Table 19a–c). We found evidence of enrichment ($P < 1.2 \times 10^{-3}$) in 24 out of 41 data sets examined. The strongest enrichment was observed with promoter (histone 3 Lys 4 trimethylation (H3K4me3), histone 3 Lys 9 acetylation (H3K9ac)) and enhancer (H3K4me1, HeK27ac) marks detected in mid-frontal lobe, anterior caudate, astrocytes and substantia nigra, supporting neuronal tissues in BMI regulation.

To identify pathways or gene sets implicated by the BMI-associated loci, we first used Meta-Analysis Gene-set Enrichment of varia NT Associations (MAGENTA)²⁴, which takes as input pre-annotated gene sets, and then tests for overrepresentation of gene set genes at BMI-associated loci. We found enrichment (false discovery rate (FDR) < 0.05) of seven gene sets, including neurotrophin signalling. Other highlighted gene sets related to general growth and patterning: basal cell carcinoma, acute myeloid leukaemia, and hedgehog signalling (Supplementary Table 20a, b).

Second, we used DEPICT, that uses predefined gene sets reconstituted using coexpression data, to perform gene set enrichment analysis. After merging highly correlated gene sets, nearly 500 gene sets were significantly enriched (FDR < 0.05) for genes in BMI-associated loci (Fig. 2b and Supplementary Table 21a, b). The most strongly enriched gene sets highlight potentially novel pathways in the CNS. These include gene sets related to synaptic function, long-term potentiation and neurotransmitter signalling (glutamate signalling in particular, but also noradrenaline, dopamine and serotonin release cycles, and GABA (γ -aminobutyric acid) receptor activity; Fig. 2c). Potentially relevant mouse behavioural

phenotypes, such as physical activity and impaired coordination were also highly enriched (Fig. 2b and Supplementary Table 21a). Several gene sets previously linked to obesity, such as integration of energy metabolism, polyphagia, secretion and action of insulin and related hormones (for example, ‘regulation of insulin secretion by glucagon-like peptide 1’ and ‘glucagon signalling in metabolic regulation’), mTOR signalling (which affects cell growth in response to nutrient intake via insulin and growth factors²⁵), and gene sets overlapping the neurotrophin signalling pathway identified by MAGENTA were also enriched, although not as significantly as other CNS processes (Fig. 2d). DEPICT also identified significant enrichment for additional cellular components and processes: calcium channels, MAP kinase activity, chromatin organization and modification, and ubiquitin ligases.

Third, we manually reviewed literature related to all 405 genes within 500 kb and $r^2 > 0.2$ of the 97 index SNPs. We classified these genes into one or more biological categories, and observed 25 categories containing three or more genes (Supplementary Table 22). The largest category comprised genes involved in neuronal processes, including monogenic obesity genes involved in hypothalamic function and energy homeostasis, and genes involved in neuronal transmission and development. Other processes highlighted by the manual literature review included glucose and lipid homeostasis and limb development, which were less notable in the above methods, but may still be related to the underlying biology of BMI.

To identify specific genes that may account for BMI association, we considered each of the following to represent supportive evidence for a gene within a locus: (1) the gene nearest the index SNP²⁶; (2) genes containing missense, nonsense or copy number variants, or a *cis*-expression quantitative trait locus (eQTL) in LD with the index SNP; (3) genes prioritized by integrative methods implemented in DEPICT; (4) genes prioritized by connections in published abstracts by GRAIL (Gene Relationships Across Implicated Loci)²⁷; or (5) genes biologically related to obesity, related metabolic disease, or energy expenditure based on manual literature review (Tables 1 and 2, Extended Data Tables 2–4 and Supplementary Tables 23–25). We first focused on the 64 genes in associated loci with more than one consistent line of supporting evidence. As expected, many of these genes overlap with CNS processes, including synaptic function, cell–cell adhesion, and glutamate signalling (*ELAVL4*, *GRID1*, *CADM2*, *NRXN3*, *NEGR1* and *SCG3*), cause monogenic obesity syndromes (*MC4R*, *BDNF*, *BBS4* and *POMC*), or function in extreme/early onset obesity in humans and mouse models (*SH2B1* and *NEGR1*)^{6,28,29}. Other genes with several lines of supporting evidence are related to insulin secretion and action, energy metabolism, lipid biology, and/or adipogenesis (*TCF7L2*, *GIPR*, *IRS1*, *FOXO3*, *ASB4*, *RPTOR*, *NPC1*, *CREB1*, *FAM57B*, *APOBR* and *HSD17B12*), encode RNA binding/processing proteins (*PTBP2*, *ELAVL4*, *CELF1* and possibly *RALYL*), are in the MAP kinase signalling pathway (*MAP2K5* and *MAPK3*), or regulate cell proliferation or cell survival (*FAIM2*, *PARK2* and *OLFM4*). Although we cannot be certain that any individual gene is related to the association at a given locus, the strong enrichment of pathways among genes within associated loci argues for a causal role for these pathways, prioritizes specific genes for follow-up experiments, and provides the strongest genetic evidence so far for a role of particular biological and CNS processes in the regulation of human body mass.

Discussion

Our meta-analysis of nearly 340,000 individuals identified 97 GWS loci associated with BMI, 56 of which are novel. These loci account for 2.7% of the variation in BMI, and suggest that as much as 21% of BMI variation can be accounted for by common genetic variation. Our analyses provide robust evidence to implicate particular genes and pathways affecting BMI, including synaptic plasticity and glutamate receptor activity—pathways that respond to changes in feeding and fasting, are regulated by key obesity-related molecules such as BDNF and MC4R, and impinge on key hypothalamic circuits^{30–32}. These pathways also overlap with one of the several proposed mechanisms of action of topiramate, a component of one of two weight-loss drugs approved by the US Food and Drug Administration^{33,34}. This observation suggests that the relevant site of action for this drug may be glutamate receptor activity, supporting the idea that these genes and pathways could reveal more targets for weight-loss therapies. BMI-associated loci also overlap with genes and pathways implicated in neurodevelopment (Supplementary Tables 21 and 22). Finally, consistent with previous work and findings from monogenic obesity syndromes, we confirm a role for the CNS—particularly genes expressed in the hypothalamus—in the regulation of body mass.

Examining the genes at BMI-associated loci in the context of gene expression, molecular pathways, eQTL results, mutational evidence and genomic location provides several complementary avenues through which to prioritize genes for relevance in BMI biology. Genes such as *NPC1* and *ELAVL4* are implicated by many lines of evidence (literature, mutational, eQTL and DEPICT) and become strong candidate genes in their respective locations. It is important to recognize that pathway methods and literature reviews are limited by current data sets and knowledge, and thus provide only a working model of obesity biology. For example, little is known about host genetic factors that regulate the microbiome. Variation in immune-related genes such as *TLR4* could presumably exert an influence on obesity through the microbiome³⁵. Together, our results underscore the heterogeneous aetiology of obesity and its links with several related metabolic diseases and processes.

BMI variants are generally associated with related cardiometabolic traits in accord with established epidemiological relationships. This could be due to shared genetic effects or to other causes of cross-phenotypic correlations. However, some BMI-associated variants have effects on related traits counter to epidemiological expectations. Once better understood, these mechanisms may not only help to explain why not all obese individuals develop related metabolic diseases, but also suggest possible mechanisms to prevent development of metabolic disease in those who are already obese.

Larger studies of common genetic variation, studies of rare variation (including those based on imputation, exome chips and sequencing), and improved computational tools will continue to identify genetic variants associated with BMI and help to further refine the biology of obesity. The 97 loci identified here represent an important step in understanding the physiological mechanisms leading to obesity. These findings strengthen the connection between obesity and other metabolic diseases, enhance our appreciation of the tissues,

physiological processes, and molecular pathways that contribute to obesity, and will guide future research aimed at unravelling the complex biology of obesity.

METHODS

Study design

We conducted a two-stage meta-analysis to identify BMI-associated loci in European adults (Extended Data Fig. 1 and Extended Data Table 1). In stage 1 we performed meta-analysis of 80 GWAS ($n = 234,069$); and stage 2 incorporated data from 34 additional studies ($n = 88,137$) genotyped using Metabochip⁷ (Supplementary Tables 1–3). Secondary meta-analyses were also conducted for: (1) all ancestries, (2) European men, (3) European women, and (4) European population-based studies. The total number of subjects and SNPs included in each stage for all analyses is shown in Extended Data Table 1. No statistical methods were used to predetermine sample size.

Phenotype

BMI, measured or self-reported weight in kg per height in metres squared (Supplementary Tables 1 and 3) was adjusted for age, age squared, and any necessary study-specific covariates (for example, genotype-derived principal components) in a linear regression model. The resulting residuals were transformed to approximate normality using inverse normal scores. For studies with no known related individuals, residuals were calculated separately by sex and case/control status. For family-based studies, residuals were calculated with men and women together, adding sex as an additional covariate in the linear regression model. Relatedness was accounted for in a study-specific manner (Supplementary Table 2).

Sample quality control, imputation and association

Following study-specific quality control measures (Supplementary Table 2), all contributing GWAS common SNPs were imputed using the HapMap phase II CEU reference panel for European-descent studies³⁷, and CEU+YRI+CHB+JPT HapMap release 22 for the African-American and Hispanic GWAS. Directly genotyped (GWAS and Metabochip) and imputed variants (GWAS only) were then tested for association with the inverse normally transformed BMI residuals using linear regression assuming an additive genetic model. Quality control following study level analyses was conducted following procedures outlined elsewhere³⁸.

Meta-analysis

Fixed effects meta-analyses were conducted using the inverse variance-weighted method implemented in METAL³⁹. Study-specific GWAS results as well as GWAS meta-analysis results were corrected for genomic control using all SNPs⁴⁰. Study-specific Metabochip results as well as Metabochip meta-analysis results were genomic-control-corrected using 4,425 SNPs included on Metabochip for replication of associations with QT-interval, a phenotype not correlated with BMI, after pruning of SNPs within 500 kb of an anthropometry replication SNP. The final meta-analysis combined the genomic-control-corrected GWAS and Metabochip meta-analysis results.

Identification of novel loci

We used a distance criterion of ± 500 kb surrounding each GWS peak ($P < 5 \times 10^{-8}$) to define independent loci and to place our results in the context of previous studies, including our previous GIANT meta-analyses. Of several locus models tested, this definition most closely reflected the loci defined by approximate conditional analysis using GCTA (Tables 1 and 2, respectively). Current index SNPs falling within 500 kb of a SNP previously associated with BMI, weight, extreme obesity or body fat percentage^{5,8-11} were considered previously identified.

Characterization of BMI-associated SNP effects

To investigate potential sources of heterogeneity between groups we compared the effect estimates of our 97 GWS SNPs for men versus women of European ancestry and Europeans versus non-Europeans. To address the effects of studies ascertained on a specific disease or phenotype on our results we also compare the effect estimates of European ancestry studies of population-based studies with the following European-descent subsets of studies: (1) non-population-based studies (that is, those ascertained on a specific disease or phenotype); (2) type 2 diabetes cases; (3) type 2 diabetes controls; (4) combined type 2 diabetes cases and controls; (5) CAD cases; (6) CAD controls; and (7) combined CAD cases and controls (Supplementary Tables 10 and 11). We also tested for heterogeneity of effect estimates between our European sex-combined meta-analysis and results from recent GWAS meta-analyses for BMI in individuals of African or east Asian ancestry^{10,41} (Supplementary Table 9). Heterogeneity was assessed as described previously⁴². A Bonferroni-corrected $P < 5 \times 10^{-4}$ (corrected for 97 tests) was used to assess significance. For heterogeneity tests assessing effects of ascertainment, we also used a 5% FDR threshold to assess significance of heterogeneity statistics (Supplementary Table 11).

Fine-mapping

We compared the meta-analysis results and credible sets of SNPs likely to contain the causal variant, based on the method described previously¹⁴, across the European-only, non-European, and all ancestries sex-combined meta-analyses. For each index SNP falling within a Metachip fine-mapping region (27 for BMI), all SNPs available within 500 kb on either side of the index SNP were selected. Effect size estimates and standard errors for each SNP were converted to approximate Bayes' factors according to the method described previously¹⁵. All approximate Bayes' factors were then summed across the 1-megabase (Mb) region and the proportion of the posterior odds of being the causal variant was calculated for each variant (approximate Bayes' factor for SNP_{*i*}/sum of approximate Bayes' factors for the region). The set of SNPs that accounts for 99% of posterior odds of association in the region denotes the set most likely to contain the causal variant for that association region (Supplementary Table 12).

Cumulative effects, risk prediction and variance explained

We assessed the cumulative effects of the 97 GWS loci on mean BMI and on their ability to predict obesity (BMI ≥ 30 kg m⁻²) using the *c* statistic from logistic regression models in the Health and Retirement Study¹⁷, a longitudinal study of 26,000 European Americans 50

years or older. The variance explained (VarExp) by each SNP was calculated using the effect allele frequency (f) and beta (β) from the meta-analyses using the formula $\text{VarExp} = \beta^2(1 - f)2f$.

For polygene analyses, the approximate conditional analysis from GCTA^{19,20}, was used to select SNPs using a range of P value thresholds (that is, 5×10^{-8} , 5×10^{-7} , ..., 5×10^{-3}) based on summary data from the European sex-combined meta-analysis excluding TwinGene and QIMR studies. We performed a within-family prediction analysis using full-sib pairs selected from independent families (1,622 pairs from the QIMR cohort and 2,758 pairs from the TwinGene cohort) and then SNPs at each threshold were used to calculate the percentage of phenotypic variance explained and predict risk (Extended Data Figs 2 and 3). We then confirmed the results from population-based prediction and estimation analyses in an independent sample of unrelated individuals from the TwinGene ($n = 5,668$) and QIMR ($n = 3,953$) studies (Extended Data Fig. 3 and Fig. 1c). The SNP-derived predictor was calculated using the profile scoring approach implemented in PLINK and estimation analyses were performed using the all-SNP estimation approach implemented in GCTA.

Enrichment analysis of MetaboChip SNPs selected for replication

The 5,055 SNPs that were included for BMI replication on MetaboChip included 1,909 independent SNPs ($r^2 < 0.1$ and > 500 kb apart), of which 1,458 displayed directionally consistent effect estimates with those reported previously⁵. To estimate the number of MetaboChip SNPs truly associated with BMI, we counted the number of SNPs with directional consistency (DC) between ref. 5 and a meta-analysis of non-overlapping samples for these 1,909 SNPs. We then calculated DC in the presence of a mixture of associated and non-associated SNPs assuming $P(\text{DC} \mid \text{associated}) = 1$ and $P(\text{DC} \mid \text{not associated}) = 0.5$. In this formulation, $\text{DC} = R/2 + S$, meaning that $S = 2\text{DC} - T$, in which T equals the total number of SNPs, R equals the number of SNPs not associated with BMI, and S equals the number of SNPs associated with BMI. With $\text{DC} = 1,458$ and $T = 1,909$, we estimate S to be $2\text{DC} - T = 2 \times 1,458 - 1,909 = 1,007$.

Joint and conditional multiple SNP association analysis

To identify additional signals in regions of association, we used GCTA¹⁹, an approach that uses meta-analysis summary statistics and an LD matrix derived from a reference sample, to perform approximate joint and conditional SNP association analysis. We used 6,654 unrelated individuals of European ancestry from the ARIC cohort as the reference sample to approximate conditional P values.

Manual gene annotation and biological description

All genes within 500 kb of an index SNP were annotated for molecular function, cellular function, and for evidence of association with BMI-related traits in human or animal model experiments (Supplementary Table 22). We used several avenues for annotation, including Spotter (<http://csg.sph.umich.edu/boehnke/spotter/>), SNIPPER (<http://csg.sph.umich.edu/boehnke/snipper/>), PubMed (<http://www.ncbi.nlm.nih.gov/pubmed/>), OMIM (<http://www.omim.org>) and UNIPROT (<http://www.uniprot.org/>). When no genes mapped to this interval the nearest gene on each side of the index SNP was annotated. In examining

possible functions of genes in the region, we excluded any references to GWAS or other genetic association studies. We analysed 405 genes in the 97 GWS loci and manually curated them into 25 biological categories containing more than three genes.

Functional variants

All variants within 500 kb (HapMap release 22/1000 Genomes CEU) and in LD ($r^2 > 0.7$) with an index SNP were annotated for functional effects based on RefSeq transcripts using Annovar⁴³ (<http://www.openbioinformatics.org/annovar/>). PhastCon, Grantham, GERP, and PolyPhen⁴⁴ predictions were accessed via the Exome Variant Server⁴⁵ (<http://evs.gs.washington.edu/EVS>), and from SIFT⁴⁶ (<http://sift.jcvi.org/>) (Extended Data Table 4).

Copy number variations correlated with BMI index SNPs

To study common copy number variations, we used a list of copy number variations well-tagged by SNPs in high LD ($r^2 > 0.8$) with deletions in European populations from phase 1 release of the 1000 Genomes Project⁴⁷ (Supplementary Table 25).

eQTLs

We examined the *cis* associations between the 97 GWS SNPs and expression of nearby genes in whole blood, lymphocytes, skin, liver, omental fat, subcutaneous fat and brain tissue^{48–55} (Supplementary Table 23). Conditional analyses were performed by including both the BMI-associated SNP and the most significant *cis*-associated SNP for the given transcript. Conditional analyses were conducted for all data sets, except the brain tissue data set due to limited power. To minimize the potential for false-positives, only *cis* associations below a study-specific FDR of 5% (or 1% for some data sets), in LD with the peak SNP ($r^2 > 0.7$) for the transcript, and with conditional $P > 0.05$ for the peak SNP, are reported (Extended Data Table 2).

MAGENTA

We used the MAGENTA method to test predefined gene sets for enrichment at BMI-associated loci²⁴. We used the GWAS + Metabochip data as input and applied default settings.

GRAIL

We used GRAIL²⁷ to identify genes near BMI-associated loci having similarities in the published scientific text using PubMed abstracts as of December 2006. The BMI loci were queried against HapMap release 22 for the European panel, and we controlled for gene size.

DEPICT

We used DEPICT to identify the most likely causal gene at a given associated locus, reconstituted gene sets enriched for BMI associations, and tissues and cell types in which genes from associated loci are highly expressed²¹. To accomplish this, the method relies on publicly available gene sets (including molecular pathways) and uses gene expression data from 77,840 gene expression arrays⁷⁵ to predict which other genes are likely to be part of

these gene sets, thus combining known annotations with predicted annotations. For details and negative control analyses please see Supplementary Methods.

We first clumped the European-only GWAS-based meta-analysis summary statistics using 500 kb flanking regions, LD $r^2 > 0.1$ and excluded SNPs with $P < 5 \times 10^{-4}$; which resulted in a list of 590 independent SNPs. HapMap phase II CEU genotype data³⁷ was used to compute LD and genomic coordinates were defined by genome build GRCh38. Because the GWAS meta-analysis was based on both GWAS and MetaboChip studies, there were discrepancies in the index SNPs that are referenced in Table 1 of the paper and the ones used in DEPICT, which was run on the GWAS data only. Therefore we forced in GWS index SNPs from the GWAS plus MetaboChip GWA meta-analysis into the DEPICT GWAS-only based analysis. This enabled a more straightforward comparison of genes in DEPICT loci and genes in GWS loci highlighted by manual lookups, and did not lead to any significant bias towards SNPs on MetaboChip (data not shown). We forced in 62 of the GWS loci in Table 1, so all of the 97 SNPs were among the 590 SNPs. The 590 SNPs were further merged into 511 non-overlapping regions (FDR < 0.05) used in DEPICT analysis. For additional information on the analysis please refer to Supplementary Methods.

Cross-trait analyses

To explore the relationship between BMI and an array of cardiometabolic traits and diseases, association results for the 97 BMI index SNPs were requested from 13 GWAS meta-analysis consortia: DIAGRAM (type 2 diabetes)⁵⁶, CARDIoGRAM-C4D (CAD)⁵⁷, ICBP (systolic and diastolic blood pressure (SBP, DBP))⁵⁸, GIANT (waist-to-hip ratio, hip circumference, and waist circumference, each unadjusted and adjusted for BMI)^{13,59}, GLGC (HDL, low density lipoprotein cholesterol, triglycerides, and total cholesterol)⁶⁰, MAGIC (fasting glucose, fasting insulin, fasting insulin adjusted for BMI, and two-hour glucose)^{61–63}, ADIPOGen (BMI-adjusted adiponectin)⁶⁴, CKDgen (urine albumin-to-creatinine ratio (UACR), estimated glomerular filtration rate, and overall CKD)^{65,66}, ReproGen (age at menarche, age at menopause)^{67,68}, GENIE (diabetic nephropathy)^{69,70}. Proxies ($r^2 > 0.8$ in CEU) were used when an index SNP was unavailable.

Enrichment of concordant effects

We compared the effects for the 97 BMI index SNP across these related traits using a one-sided binomial test of the number of concordant effects versus a null expectation of $P = 0.5$. Concordant and nominally significant ($P < 0.05$) SNP effects were similarly tested using a one-sided binomial test with a null expectation of $P = 0.05$. We evaluated significance in either test with a Bonferroni-corrected threshold of $P = 0.002$ ($0.05/23$ traits tested).

Joint effects of cross-trait associations

To determine the joint effect of all 97 BMI loci on other cardiometabolic phenotypes, we used the meta-regression technique from ref. 64 to correlate the effect estimates of the BMI-increasing alleles with effect estimates from meta-analyses for each of the metabolic traits from other consortia (DIAGRAM, MAGIC, ICBP, GLGC, ADIPOGen, ReproGen and CARDIoGRAM).

Cross-traits heatmap

To explore observed concordance in effects of BMI loci on other cardiometabolic and anthropometric traits, we converted the effect estimates and standard errors (or P values) from meta-analysis to Z -scores oriented with respect to the BMI-increasing allele, for each of the 97 BMI index SNPs in the twenty-three traits. We then classified each Z -score as follows to generate a vector of the Z -score of each trait at each locus: 0 (not significant) if $-2 < Z < 2$; 1 (significant positive) if $Z > 2$; -1 (significant negative) if $Z < -2$.

Extended Data Fig. 5 displays these locus-trait relationships in a heatmap using Euclidean distance and complete linkage clustering to order both loci and traits.

Cross-traits bubble plot

We also represent the genetic overlap between other cardiometabolic traits and BMI susceptibility loci with a bubble plot in which the size of each bubble is proportional to the fraction of BMI-associated loci for which there was a significant association ($P < 5 \times 10^{-4}$). Each pair of bubbles is connected by a line proportional to the number of significant BMI-increasing loci overlapping between the traits.

NHGRI GWAS catalogue lookups


We extracted previously reported GWAS association within 500 kb of and $r^2 > 0.7$ with any BMI-index SNP from the NHGRI GWAS catalogue⁷¹ (<http://www.genome.gov/gwastudies>; Supplementary Table 17a, b). For studies reporting greater than 30 significant hits, additional SNP-trait associations were pulled from the literature and compared to BMI index SNPs the same as with other GWAS catalogue studies.

ENCODE/Roadmap

To identify global enrichment of data sets at the BMI-associated loci we performed permutation-based tests in a subset of 41 open chromatin (DNase-seq), histone modification (H3K27ac, H3K4me1, H3K4me3 and H3K9ac), and transcription factor binding data sets from the ENCODE Consortium²², Roadmap Epigenomics Project²³ and when available the ENCODE Integrative Analysis^{60,72} (Supplementary Table 19). We processed Roadmap Epigenomics sequencing data with multiple biological replicates using MACS2 (ref. 73) and then applied same Irreproducible Discovery Rate pipeline used in the ENCODE Integrative Analysis^{60,72}. Roadmap Epigenomics data with only a single replicate were analysed using MACS2 alone. We examined variants in LD with 97 BMI index SNPs based on $r^2 > 0.7$ from the 1000 Genomes phase 1 version 2 EUR samples⁷⁴. We matched the index SNP at each locus with 500 variants having no evidence of association ($P > 0.5$, ~1.2 million total variants) with a similar distance to the nearest gene ($\pm 11,655$ bp), number of variants in LD (± 8 variants), and minor allele frequency. Using these pools, we created 10,000 sets of control variants for each of the 97 loci and identified variants in LD ($r^2 > 0.7$) and within 1 Mb. For each SNP set, we calculated the number of loci with at least one variant located in a regulatory region under the assumption that one regulatory variant is responsible for each association signal. We estimated the P value assuming a sum of binomial distributions to represent the number of index SNPs (or their LD proxies; $r^2 > 0.7$) that overlap a regulatory


data set compared to the expectation observed in the 500 matched control sets. Data sets were considered significantly enriched if the P value was below a Bonferroni-corrected threshold of 1.2×10^{-3} , adjusting for 41 tests.

Extended Data




Extended Data Figure 1. Study design

*The SNP counts reflect sample size filter of $n = 50,000$. §Counts represent the primary European sex-combined analysis. Please see Extended Data Table 1 for counts for secondary analyses.




Extended Data Figure 2. Genetic characterization of BMI-associated variants

a, Plot of the cumulative phenotypic variance explained by each locus ordered by decreasing effect size. **b**, The relationship between effect size and allele frequency. Previously identified loci are blue circles and novel loci are red triangles. **c**, Quantile–quantile (Q–Q) plot of meta-analysis *P* values for all 1,909 BMI-replication SNPs (blue) and after removing SNPs near the 97 associated loci (green). **d**, Histogram of cumulative effect of BMI risk alleles. Mean BMI for each bin is shown by the black dots (with standard deviation) and corresponds to the right-hand y axis.




Extended Data Figure 3. Partitioning the variance in and risk prediction from SNP-derived predictor


a, b, The analyses were performed using 2,758 full sibling pairs from the TwinGene cohort (**a**) and 1,622 pairs from the QIMR cohort (**b**). The SNP-based predictor was adjusted for the first 20 principal components. The variance of the SNP-based predictor can be partitioned into four components (V_g , V_e , C_g and C_e) using the within-family prediction analysis, in which V_g is the variance explained by real SNP effects, C_g is the covariance between predictors attributed to the real effects of SNPs that are not in LD but correlated due to population stratification, V_e is the accumulated variance due to the errors in estimating SNP effects, and C_e is the covariance between predictors attributed to errors in estimating the effects of SNPs that are correlated due to population stratification. Error bars reflect s.e.m. of estimates. **c**, The prediction R^2 shown on the y axis is the squared correlation between phenotype and SNP-based genetic predictor in unrelated individuals from the TwinGene ($n = 5,668$) and QIMR ($n = 3,953$) studies. The number shown in each column is the number of SNPs selected from the GCTA joint and conditional analysis at a range of *P*-value thresholds. In each case, the predictor was adjusted by the first 20 principal components. The column in orange is the average prediction R^2 weighted by sample size over the two cohorts. The dashed grey line is the value inferred from the within-family prediction analyses using this equation $R^2 = (V_g + C_g)^2 / (V_g + V_e + C_g + C_e)$.



Extended Data Figure 4. Comparison of BMI-associated index SNPs across ethnicities
a, b, BMI effects observed in European ancestry individuals (*x* axes) compared to African ancestry (**a**) or Asian ancestry (**b**) individuals (*y* axes). **c, d,** Allele frequencies between ancestry groups, as in **a** and **b**. **e, f,** Comparison of the estimates of explained variance. In all plots, novel loci are in red and previously identified loci are in blue.



Extended Data Figure 5. Effects of BMI-associated loci on related metabolic traits
 Unsupervised hierarchical clustering of the 97 BMI-associated loci (y axis) on 23 related metabolic traits (x axis). The top row shows the a priori expected relationship with BMI (green is concordant effect direction, purple is opposite). Loci with statistically significant concordant direction of effect are highlighted in green, and significant but opposing effects are in purple. Grey indicates a non-significant relationship and those with no information are in white. The key in the top left corner also shows the count of gene-phenotype pairs in each category (cyan bars).



Extended Data Figure 6. Bubble chart representing the genetic overlap across traits at BMI susceptibility loci

Each bubble represents a trait for which association results were requested for the 97 GWS BMI loci. The size of the bubble is proportional to the number of BMI-increasing loci with a significant association. A line connects each pair of bubbles with thickness proportional to the number of significant loci shared between the traits. Traits tested include the current study BMI SNPs, African-American BMI (AA BMI), hip circumference (HIP), HIP adjusted for BMI (HIPadjBMI), waist circumference (WC), waist circumference adjusted for BMI (WCadjBMI), waist-to-hip ratio (WHR), waist-to-hip ratio adjusted for BMI (WHRadjBMI), height, adiponectin, coronary artery disease (CAD), diastolic blood pressure (DBP), systolic blood pressure (SBP), high-density lipoprotein (HDL), low-density lipoprotein (LDL), total cholesterol (TC), triglycerides (TG), type 2 diabetes (T2D), fasting glucose (FG), fasting insulin (FI), fasting insulin adjusted for BMI (FIadjBMI), two-hour glucose (Glu2hr), diabetic nephropathy (Diab_Neph), age at menopause (AgeMenopause), and age at menarche (AgeMenarche).

Extended Data Table 1

Descriptive characteristics of meta-analyses

Meta-analysis	Total number of studies	Maximum number of subjects	Number of SNPs*	λ_{GC}
<i>European sex-combined</i>				
GWAS	80	234,069	2,550,021	1.526
Metabochip	34	88,137	156,997	1.25
Joint GWAS+Metabochip	114	322,154	2,554,623	1.084
<i>European men</i>				
GWAS	72	104,666	2,473,152	1.279
Metabochip	34	48,274	152,326	1.121
Joint GWAS+Metabochip	106	152,893	2,477,617	1.006

Meta-analysis	Total number of studies	Maximum number of subjects	Number of SNPs*	λ_{GC}
European women				
GWAS	74	132,115	2,491,697	1.336
Metabochip	33	39,864	153,086	1.029
Joint GWAS+Metabochip	107	171,977	2,494,571	1.002
*European population-based				
GWAS	49	162,262	2,502,573	1.385
Metabochip	20	46,263	155,617	1.034
Joint GWAS+Metabochip	69	209,521	2,506,448	1.003
All ancestries				
GWAS	82	236,231	2,550,614	1.451
Metabochip	43	103,047	181,718	1.25
Joint GWAS+Metabochip	125	339,224	2,555,496	1.004

*For the GWAS and joint GWAS+Metabochip analyses, SNP count reflects $n = 50,000$.

Extended Data Table 2

Previously known GWS BMI loci in European meta-analysis

SNP	Chr:Position	*Notable gene(s)	Alleles	EAF	β	SE	<i>P</i> value
rs1558902	16:52,361,075	<i>FTO</i> (B,N)	A/T	0.415	0.082	0.003	7.51E-153
rs6567160	18:55,980,115	<i>MC4R</i> (B,N)	C/T	0.236	0.056	0.004	3.93E-53
rs13021737	2:622,348	<i>TMEM18</i> (N)	G/A	0.828	0.06	0.004	1.11E-50
rs10938397	4:44,877,284	<i>GNPDA2</i> (N); <i>GABRG1</i> (B)	G/A	0.434	0.04	0.003	3.21E-38
rs543874	1:176,156,103	<i>SEC16B</i> (N)	G/A	0.193	0.048	0.004	2.62E-35
rs2207139	6:50,953,449	<i>TFAP2B</i> (B,N)	G/A	0.177	0.045	0.004	4.13E-29
rs11030104	11:27,641,093	<i>BDAF</i> (B,M,N)	A/G	0.792	0.041	0.004	5.56E-28
rs3101336	1:72,523,773	<i>NEGR1</i> (B,C,D,N)	C/T	0.613	0.033	0.003	2.66E-26
rs7138803	12:48,533,735	<i>BCDIN3D</i> (N); <i>FAIM2</i> (D)	A/G	0.384	0.032	0.003	8.15E-24
rs10182181	2:25,003,800	<i>ADCY3</i> (B,M,N,Q); <i>POMC</i> (B,G); <i>NCOA1</i> (B) <i>SH2B1</i> (B,M,Q); <i>APOBR</i> (M,Q);	G/A	0.462	0.031	0.003	8.78E-24
rs3888190	16:28,796,987	<i>ATXN2L</i> (Q); <i>SBK1</i> (Q,D); <i>SULT1A2</i> (Q); <i>TUFM</i> (Q)	A/C	0.403	0.031	0.003	3.14E-23
rs1516725	3:187,306,698	<i>E7V5</i> (N)	C/T	0.872	0.045	0.005	1.89E-22
rs12446632	16:19,842,890	<i>GPRC5B</i> (C,N); <i>IQCK</i> (Q)	G/A	0.865	0.04	0.005	1.48E-18
rs2287019	19:50,894,012	<i>QPCTL</i> (N); <i>GIPR</i> (B,M)	C/T	0.804	0.036	0.004	4.59E-18
rs16951275	15:65,864,222	<i>M4P2K5</i> (B,D,N); <i>LBXCOR1</i> (M)	T/C	0.784	0.031	0.004	1.91E-17
rs3817334	11:47,607,569	<i>MTCH2</i> (M,Q); <i>C1QTNF4</i> (Q,I); <i>SPII</i> (Q); <i>CELF1</i> (D)	T/C	0.407	0.026	0.003	5.15E-17
rs2112347	5:75,050,998	<i>POC5</i> (M); <i>HMGCR</i> (B); <i>COL4A3BP</i> (B)	T/G	0.629	0.026	0.003	6.19E-17
rs12566985	1:74,774,781	<i>FPGT-TNNI3K</i> (N)	G/A	0.446	0.024	0.003	3.28E-15
rs3810291	19:52,260,843	<i>ZC3H4</i> (D,N,Q)	A/G	0.666	0.028	0.004	4.81E-15
rs7141420	14:78,969,207	<i>NRXN3</i> (D,N)	T/C	0.527	0.024	0.003	1.23E-14
rs13078960	3:85,890,280	<i>CADM2</i> (D,N)	G/T	0.196	0.03	0.004	1.74E-14
rs10968576	9:28,404,339	<i>LINGO2</i> (D,N)	G/A	0.32	0.025	0.003	6.61E-14

SNP	Chr:Position	*Notable gene(s)	Alleles	EAF	β	SE	P value
rs17024393	1:109,956,211	<i>GNAT2</i> (N); <i>AMPD2</i> (D)	C/T	0.04	0.066	0.009	7.03E-14
rs12429545	13:53,000,207	<i>OLFM4</i> (B,N)	A/G	0.133	0.033	0.005	1.09E-12
rs13107325	4:103,407,732	<i>SLC39A8</i> (M,N,Q)	T/C	0.072	0.048	0.007	1.83E-12
rs11165643	1:96,696,685	<i>PTBP2</i> (D,N)	T/C	0.583	0.022	0.003	2.07E-12
rs17405819	8:76,969,139	<i>HNF4G</i> (B,N)	T/C	0.7	0.022	0.003	2.07E-11
rs1016287	2:59,159,129	<i>LINC01122</i> (N)	T/C	0.287	0.023	0.003	2.25E-11
rs4256980	11:8,630,515	<i>TRIM66</i> (D,M,N); <i>TUB</i> (B)	G/C	0.646	0.021	0.003	2.90E-11
rs12401738	1:78,219,349	<i>FUBP1</i> (N); <i>USP33</i> (D)	A/G	0.352	0.021	0.003	1.15E-10
rs205262	6:34,671,142	<i>C6orf106</i> (N); <i>SNRPC</i> (Q)	G/A	0.273	0.022	0.004	1.75E-10
rs12016871	13:26,915,782	<i>MTIF3</i> (N); <i>GTF3A</i> (Q)	T/C	0.203	0.03	0.005	2.29E-10
rs12940622	17:76,230,166	<i>RPTOR</i> (B,N)	G/A	0.575	0.018	0.003	2.49E-09
rs11847697	14:29,584,863	<i>PRKD1</i> (N)	T/C	0.042	0.049	0.008	3.99E-09
rs2075650	19:50,087,459	<i>TOMM40</i> (B,N); <i>APOE</i> (B); <i>APOC1</i> (B)	A/G	0.848	0.026	0.005	1.25E-08
rs2121279	2:142,759,755	<i>LRP1B</i> (N)	T/C	0.152	0.025	0.004	2.31E-08
rs29941	19:39,001,372	<i>KCTD15</i> (N)	G/A	0.669	0.018	0.003	2.41E-08
rs1808579	18:19,358,886	<i>NPC1</i> (B,G,M,Q); <i>C18orf8</i> (N,Q)	C/T	0.534	0.017	0.003	4.17E-08

SNP positions are reported according to Build 36 and their alleles are coded based on the positive strand. Effect alleles, allele frequencies, betas (β), s.e.m., sample sizes (n), and P values are based on the meta-analysis of GWAS I + II + MetaboChip association data from the European sex-combined data set.

* Notable genes from biological relevance to obesity (B); GRAIL results (G); BMI-associated variant is in strong LD (r^2 0.7) with a missense variant in the indicated gene (M); gene nearest to Index SNP (N); association and eQTL data converge to affect gene expression (Q); DEPICT analyses (D); copy number variation (C).

Extended Data Table 3

Association of the GWS SNPs for BMI with *cis*-gene expression (*cis*-eQTLs)

SNP	Chr.	BMI increasing allele	Tissue	Gene	β for Giant SNP	<i>P</i> for GIANT SNP	<i>P</i> _{adj} for GIANT SNP	PeaksNP	<i>r</i> ²	<i>P</i> for peak SNP	<i>P</i> _{adj} for peak SNP	Reference
Novel loci												
rs11583200	1	c	Subcutaneous	<i>ELAVL4</i>	-0.066	1.90E-12	0.44	rs6588374	0.78	1.07E-12	0.36	Zhong et al.
rs492400	2	c	Liver	<i>PLCD4</i>	-0.054	4.64E-40	0.98	rs10187066	1.00	4.49E-40	0.98	Zhong et al.
rs492400	2	c	Lymphocyte	<i>RQCD1</i>	0.392	7.11E-22	0.94	rs526134	1.00	4.06E-22	0.21	Dixon et al.
rs492400	2	c	PBMC	<i>RQCD1</i>	-0.102	2.43E-06	0.98	rs526134	0.95	2.21E-06	0.96	PBMC meta-analysis
rs492400	2	c	Omental	<i>TTLA</i>	0.018	1.33E-10	0.82	rs12987009	0.73	2.82E-13	0.07	Zhong et al.
rs492400	2	c	Lymphocyte	<i>TTLA</i>	0.158	9.02E-06	1	rs492400	1.00	9.02E-06	1	Dixon et al.
rs17001654	4	G	Lymphocyte	<i>SCARB2</i>	0.248	5.57E-09	0.59	rs6835324	0.94	3.42E-09	0.25	Dixon et al.
rs9400239	6	C	Subcutaneous	<i>HSS00296402</i>	0.034	9.51E-22	0.97	rs2153960	0.94	1.93E-23	0.48	Zhong et al.
rs9400239	6	C	Omental	<i>HSS00296402</i>	0.015	1.34E-13	0.50	rs2153960	0.93	4.64E-17	0.22	Zhong et al.
rs1167827	7	G	Blood	<i>PMS2P3</i>	-0.595	4.20E-32	0.66	rs6963105	0.93	3.00E-32	0.39	Emilsson et al.
rs1167827	7	G	Omental	<i>PMS2P3</i>	-0.027	1.57E-11	0.95	rs6963105	0.98	6.94E-12	0.86	Zhong et al.
rs1167827	7	G	Subcutaneous	<i>PMS2P3</i>	-0.030	1.30E-10	0.71	rs1167796	0.73	1.04E-12	0.10	Zhong et al.
rs1167827	7	G	Adipose	<i>PMS2P3</i>	-0.346	3.40E-09	1	rs1167827	1.00	3.40E-09	1	Emilsson et al.
rs1167827	7	G	Blood	<i>PMS2P5</i>	-0.367	1.20E-11	0.47	rs6963105	0.93	5.00E-12	0.14	Emilsson et al.
rs1167827	7	G	Subcutaneous	<i>WBSCR16</i>	0.025	1.44E-10	1	rs1167827	1.00	1.44E-10	1	Zhong et al.
rs1167827	7	G	Omental	<i>WBSCR16</i>	0.017	1.75E-06	1	rs1167827	1.00	1.75E-06	1	Zhong et al.
rs9641123	7	C	Abdominal SAT	<i>hsa-miR-653</i>	-0.344	1.54E-04	0.23	rs16868443	0.71	1.38E-04	0.20	Partis et al.
rs11191560	10	C	Gluteal SAT	<i>SFXN2</i>	0.153	1.72E-05	0.20	rs71496550	NA	4.42E-06	0.41	Min et al.
rs11191560	10	C	Abdominal SAT	<i>SFXN2</i>	0.628	1.44E-04	0.02	rs71496550	NA	9.13E-05	0.94	Min et al.
rs7164727	15	T	Lymphocyte	<i>BBS4</i>	-0.163	3.14E-05	1	rs7164727	1.00	3.14E-05	1	Dixon et al.
rs9925964	16	A	Liver	<i>VKORC1</i>	0.122	4.41E-37	0.84	rs2303223	0.88	3.62E-44	0.05	Zhong et al.
rs9925964	16	A	Subcutaneous	<i>ZNF646</i>	0.017	2.55E-06	1	rs9925964	1.00	2.55E-06	1	Zhong et al.
rs9925964	16	A	Blood	<i>ZNF668</i>	-0.382	1.70E-12	0.48	rs10871454	0.93	1.10E-12	0.26	Emilsson et al.
rs9914578	17	G	Subcutaneous	<i>C17orf13</i>	-0.010	3.01E-06	0.99	rs7225843	0.99	2.86E-06	0.97	Zhong et al.
rs1808579	18	C	SKIN	<i>C18orf8</i>	-0.073	5.74E-10	0.86	rs1788781	0.90	1.67E-10	0.13	Grundberg et al.
rs1808579	18	C	Subcutaneous	<i>C18orf8</i>	-0.014	8.41E-08	1	rs1808579	1.00	8.41E-08	1	Zhong et al.

SNP	Chr.	BMI increasing allele	Tissue	Gene	β for Giant SNP	P for GIANT SNP	P_{adj} for GIANT SNP	PeakSNP	r^2	P for peak SNP	P_{adj} for peak SNP	Reference
rs17724992	19	A	Blood	<i>PGPEP1</i>	-0.825	1.60E-40	1	rs17724992	1.00	1.60E-40	1	Emilsson et al.
Previously reported loci												
rs10182181	2	G	Subcutaneous	<i>ADCY3</i>	0.022	7.57E-06	0.69	rs11684619	0.72	8.70E-09	0.05	Zhong et al.
rs2176040	2	A	Omental	<i>IRSI</i>	-0.036	3.74E-09	0.97	rs908252	0.87	3.98E-10	0.47	Zhong et al.
rs13107325	4	T	Liver	<i>SLC39A8</i>	-0.101	1.29E-17	1	rs13107325	1.00	1.29E-17	1	Zhong et al.
rs205262	6	G	Blood	<i>SNRPC</i>	-0.462	9.60E-15	0.58	rs6457792	0.96	9.40E-15	0.55	Emilsson et al.
rs205262	6	G	PBMC	<i>SNRPC</i>	-0.127	3.40E-09	0.03	rs2744943	0.73	3.15E-11	0.12	PBMC meta-analysis
rs205262	6	G	Omental	<i>SNRPC</i>	-0.012	6.64E-06	0.81	rs2814984	0.75	8.03E-07	0.30	Zhong et al.
rs3817334	11	T	SKIN	<i>C10TNF4</i>	-0.051	1.34E-09	0.82	rs7124681	1.00	9.42E-10	0.34	Grundberg et al.
rs3817334	11	T	Subcutaneous	<i>MTCH2</i>	0.044	7.64E-13	0.76	rs12794570	0.76	2.54E-15	0.10	Zhong et al.
rs3817334	11	T	Brain	<i>MTCH2</i>	28.255	7.51E-08	NA	NA	NA	NA	NA	Myers et al.
rs3817334	11	T	FAT	<i>SPI1</i>	-0.090	9.90E-07	0.90	rs10769262	0.70	1.15E-08	1	Grundberg et al.
rs12016871	13	T	PBMC	<i>GTF3A</i>	-0.258	6.68E-34	0.90	rs7988412	0.81	1.81E-36	0.29	PBMC meta-analysis
rs12016871	13	T	Lymphocyte	<i>GTF3A</i>	-0.375	3.89E-15	0.32	rs7988412	0.86	1.32E-15	0.06	Dixon et al.
rs12446632	16	G	Omental	<i>IQCK</i>	0.028	2.27E-10	0.83	rs11865578	0.83	4.14E-13	0.14	Zhong et al.
rs12446632	16	G	Liver	<i>IQCK</i>	0.031	5.39E-06	0.74	rs9921401	0.70	3.82E-07	0.20	Zhong et al.
rs3888190	16	A	Blood	<i>APOBR</i>	0.303	2.10E-08	0.68	rs2411453	0.83	1.10E-08	0.25	Emilsson et al.
rs3888190	16	A	PBMC	<i>ATXN2L</i>	0.084	1.04E-04	0.99	rs8049439	0.99	8.59E-05	0.88	PBMC meta-analysis
rs3888190	16	A	SKIN	<i>SBK1</i>	-0.063	1.63E-06	0.41	rs4788084	0.82	2.87E-07	0.10	Grundberg et al.
rs3888190	16	A	Adipose	<i>SH2B1</i>	-0.407	4.10E-13	0.67	rs12928404	0.92	2.40E-13	0.30	Emilsson et al.
rs3888190	16	A	Omental	<i>SH2B1</i>	-0.014	5.29E-07	0.87	rs12928404	0.93	4.65 E-07	0.83	Zhong et al.
rs3888190	16	A	Subcutaneous	<i>SULT1A2</i>	0.067	3.36E-21	0.52	rs1074631	0.80	3.93E-23	0.14	Zhong et al.
rs3888190	16	A	PBMC	<i>TUFM</i>	0.694	9.81E-198	0.94	rs8049439	0.99	9.81E-198	0.12	PBMC meta-analysis
rs1808579	18	C	Subcutaneous	<i>NPC1</i>	-0.027	2.52E-10	0.83	rs1805081	0.78	7.86E-14	0.06	Zhong et al.
rs3888190	16	A	SKIN	<i>TUFM</i>	0.074	7.90E-10	0.46	rs2411453	0.76	1.91E-10	0.09	Grundberg et al.
rs3810291	19	A	Adipose	<i>ZC3H4</i>	-0.386	3.70E-09	1	rs3810291	1.00	3.70E-09	1	Emilsson et al.

Extended Data Table 4

Putative coding variants in LD ($r^2 > 0.7$) with GWS BMI loci

BMI SNP	Chr.	Source	Putative Coding Variant	r^2	Gene	Protein Alteration	PhastCon Score	GERP Score	Grantham Score	PolyPhen	SIFT Prediction	SIFT Score
Novel genome-wide significant loci												
rs492400	2	1000G	rs3770213	0.89	ZNF142	L956H	0	-1.6	99	possibly damaging	Damaging	0
rs492400	2	1000G	rs3770214	0.89	ZNF142	S751G	0.2	1.4	56	benign	Tolerated	0.08
rs492400	2	1000G	rs2230115	0.963	ZNF142	A541S	0.5	5.1	99	benign	Tolerated	0.044
rs492400	2	1000G	rs1344642	0.963	STK36	R583Q	0	2.4	43	possibly damaging	Damaging	0
rs492400	2	1000G	rs1863704	0.89	STK36	G1003D	0	2	94	possibly damaging	Tolerated	0.41
rs492400	2	1000G	rs1863704	0.89	STK36	G982D	0	2	94	possibly damaging	-	-
rs492400	2	1000G	rs3731877	0.792	TLL4	E34Q	1	5.5	29	probably damaging	Unknown	Not scored
rs17001654	4	1000G	rs61750814	1	NUP54	N250S	1	5.5	46	benign	Damaging	0.05
rs4740619	9	1000G	rs4741510	0.901	CCDC171	S121T	1	2	58	benign	Damaging	0.05
rs4740619	9	1000G	rs1539172	0.74	CCDC171	K1069R	1	4.1	26	benign	Tolerated	1
rs2176598	11	1000G	rs11555762	0.774	HSD17B12	S280L	0	0.4	145	benign	Tolerated	0.74
rs3849570	3	1000G	rs2229519	0.771	GBE1	R190G	1	4.8	125	benign	Damaging	0.04
rs3736485	15	1000G	rs12102203	0.966	DMXL2	S1288P	0.7	1.7	74	benign	Tolerated	0.32
rs7164727	15	1000G	rs2277598	0.839	BBS4	I182T	0	-4.4	89	benign	Tolerated	0.47
rs9925964	16	1000G	rs749670	0.869	ZNF646	E327G	1	4.2	98	benign	Tolerated	0.44
Previously identified genome-wide significant loci												
rs10182181	2	HapMap	rs11676272	0.967	ADCY3	S107P	0	2.9	74	benign	Tolerated	0.28
rs13107325	4	1000G	rs13107325	1	SLC39A8	A324T	1	4.4	5.8	benign	Tolerated	0.09
rs13107325	4	1000G	rs13107325	1	SLC39A8	A391T	1	4.4	5.8	benign	Tolerated	0.09
rs2112347	5	1000G	rs2307111	0.862	POC5	H11R	0.9	5.8	29	benign	Unknown	Not scored
rs2112347	5	1000G	rs2307111	0.862	POC5	H36R	0.9	5.8	29	benign	Unknown	Not scored
rs4256980	11	HapMap	rs7935453	0.729	TRIM66	L630V	-	-	-	-	Tolerated	1
rs4256980	11	1000G	rs11042022	0.876	TRIM66	H466R	-	-	-	-	Tolerated	0.38
rs4256980	11	1000G	rs11042023	0.959	TRIM66	H324R	1	5.1	29	probably damaging	Damaging	0.03
rs11030104	11	1000G	rs6265	0.817	BDNF	V148M	1	5.2	21	probably damaging	Damaging	0
rs11030104	11	1000G	rs6265	0.817	BDNF	V66M	1	5.2	21	probably damaging	Damaging	0

BMI SNP	Chr.	Source	Putative Coding Variant	r^2	Gene	Protein Alteration	PhastCon Score	GERP Score	Grantham Score	PolyPhen	SIFT Prediction	SIFT Score
rs11030104	11	1000G	rs6265	0.817	<i>BDNF</i>	V74M	1	5.2	21	probably damaging	Damaging	0
rs11030104	11	1000G	rs6265	0.817	<i>BDNF</i>	V81M	1	5.2	21	probably damaging	Damaging	0
rs11030104	11	1000G	rs6265	0.817	<i>BDNF</i>	V95M	1	5.2	21	probably damaging	Damaging	0
rs3817334	11	1000G	rs1064608	0.809	<i>MTCH2</i>	P290A	1	5.1	27	probably damaging	Tolerated	0.12
rs3888190	16	1000G	rs180743	0.789	<i>APOBR</i>	P428A	0.1	0.5	27	benign	Unknown	Not scored
rs3888190	16	1000G	rs7498665	1	<i>SH2BI</i>	T484A	1	3.1	58	benign	Tolerated	0.25
rs16951275	15	1000G	rs7170185	1	<i>LBXCOR1</i>	W200R	-	-	-	-	-	-
rs1808579	18	1000G	rs1805082	0.935	<i>NPCI</i>	I858V	1	6.1	29	benign	Tolerated	0.24
rs1808579	18	1000G	rs1805081	0.905	<i>NPCI</i>	H215R	0	-1.1	29	benign	Tolerated	0.59
rs2287019	19	1000G	rs1800437	0.714	<i>GIPR</i>	E354Q	1	3.1	29	probably damaging	Tolerated	0.09

r^2 is the LD between the BMI index SNP and the putative coding variant.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Authors

Adam E. Locke^{#1}, Bratati Kahali^{#2}, Sonja I. Berndt^{#3}, Anne E. Justice^{#4}, Tune H. Pers^{#5,6,7,8}, Felix R. Day⁹, Corey Powell², Sailaja Vedantam^{5,6}, Martin L. Buchkovich¹⁰, Jian Yang^{11,12}, Damien C. Croteau-Chonka^{10,13}, Tonu Esko^{5,6,7,14}, Tove Fall^{15,16,17}, Teresa Ferreira¹⁸, Stefan Gustafsson^{16,17}, Zoltán Kutalik^{19,20,21}, Jian'an Luan⁹, Reedik Mägi^{14,18}, Joshua C. Randall^{18,22}, Thomas W. Winkler²³, Andrew R. Wood²⁴, Tsegaselassie Workalemahu²⁵, Jessica D. Faul²⁶, Jennifer A. Smith²⁷, Jing Hua Zhao⁹, Wei Zhao²⁷, Jin Chen²⁸, Rudolf Fehrmann²⁹, Åsa K. Hedman^{16,17,18}, Juha Karjalainen²⁹, Ellen M. Schmidt³⁰, Devin Absher³¹, Najaf Amin³², Denise Anderson³³, Marian Beekman^{34,35}, Jennifer L. Bolton³⁶, Jennifer L. Bragg-Gresham^{1,37}, Steven Buyske^{38,39}, Ayse Demirkan^{32,40}, Guohong Deng^{41,42,43}, Georg B. Ehret^{44,45}, Bjarke Feenstra⁴⁶, Mary F. Feitosa⁴⁷, Krista Fischer¹⁴, Anuj Goel^{18,48}, Jian Gong⁴⁹, Anne U. Jackson¹, Stavroula Kanoni⁵⁰, Marcus E. Kleber^{51,52}, Kati Kristiansson⁵³, Unhee Lim⁵⁴, Vaneet Lotay⁵⁵, Massimo Mangino⁵⁶, Irene Mateo Leach⁵⁷, Carolina Medina-Gomez^{58,59,60}, Sarah E. Medland⁶¹, Michael A. Nalls⁶², Cameron D. Palmer^{5,6}, Dorota Pasko²⁴, Sonali Pechlivanis⁶³, Marjolein J. Peters^{58,60}, Inga Prokopenko^{18,64,65}, Dmitry Shungin^{66,67,68}, Alena Stan áková⁶⁹, Rona J. Strawbridge⁷⁰, Yun Ju Sung⁷¹, Toshiko Tanaka⁷², Alexander Teumer⁷³, Stella Trompet^{74,75}, Sander W. van der Laan⁷⁶, Jessica van Setten⁷⁷, Jana V. Van Vliet-Ostaptchouk⁷⁸, Zhaoming Wang^{3,79}, Loïc Yengo^{80,81,82}, Weihua Zhang^{41,83}, Aaron Isaacs^{32,84}, Eva Albrecht⁸⁵, Johan Ärnlöv^{16,17,86}, Gillian M. Arscott⁸⁷, Antony P. Attwood^{88,89}, Stefania Bandinelli⁹⁰, Amy Barrett⁶⁴, Isabelita N. Bas⁹¹, Claire Bellis^{92,93}, Amanda J. Bennett⁶⁴, Christian Berne⁹⁴, Roza Blagieva⁹⁵, Matthias Blüher^{96,97}, Stefan Böhringer^{34,98}, Lori L. Bonnycastle⁹⁹, Yvonne Böttcher⁹⁶, Heather A. Boyd⁴⁶, Marcel Bruinenberg¹⁰⁰, Ida H. Caspersen¹⁰¹, Yii-Der Ida Chen^{102,103}, Robert Clarke¹⁰⁴, E. Warwick Daw⁴⁷, Anton J. M. de Craen⁷⁵, Graciela Delgado⁵¹, Maria Dimitriou¹⁰⁵, Alex S. F. Doney¹⁰⁶, Niina Eklund^{53,107}, Karol Estrada^{6,60,108}, Elodie Eury^{80,81,82}, Lasse Folkersen⁷⁰, Ross M. Fraser³⁶, Melissa E. Garcia¹⁰⁹, Frank Geller⁴⁶, Vilmantas Giedraitis¹¹⁰, Bruna Gigante¹¹¹, Alan S. Go¹¹², Alain Golay¹¹³, Alison H. Goodall^{114,115}, Scott D. Gordon⁶¹, Mathias Gorski^{23,116}, Hans-Jürgen Grabe^{117,118}, Harald Grallert^{85,119,120}, Tanja B. Grammer⁵¹, Jürgen Gräßler¹²¹, Henrik Grönberg¹⁵, Christopher J. Groves⁶⁴, Gaëlle Gusto¹²², Jeffrey Haessler⁴⁹, Per Hall¹⁵, Toomas Haller¹⁴, Goran Hallmans¹²³, Catharina A. Hartman¹²⁴, Maija Hassinen¹²⁵, Caroline Hayward¹²⁶, Nancy L. Heard-Costa^{127,128}, Quinta Helmer^{34,98,129}, Christian Hengstenberg^{130,131}, Oddgeir Holmen¹³², Jouke-Jan Hottenga¹³³, Alan L. James^{134,135}, Janina M. Jeff⁵⁵, Åsa Johansson¹³⁶, Jennifer Jolley^{88,89}, Thorhildur Juliusdottir¹⁸, Leena Kinnunen⁵³, Wolfgang Koenig⁵², Markku Koskenvuo¹³⁷, Wolfgang Kratzer¹³⁸, Jaana Laitinen¹³⁹, Claudia Lamina¹⁴⁰, Karin Leander¹¹¹, Nanette R. Lee⁹¹, Peter Lichtner¹⁴¹, Lars Lind¹⁴², Jaana Lindström⁵³, Ken Sin Lo¹⁴³, Stéphane Lobbens^{80,81,82}, Roberto Lorbeer¹⁴⁴, Yingchang Lu^{55,145},

François Mach⁴⁵, Patrik K. E. Magnusson¹⁵, Anubha Mahajan¹⁸, Wendy L. McArdle¹⁴⁶, Stela McLachlan³⁶, Cristina Menni⁵⁶, Sigrun Merger⁹⁵, Evelin Mihailov^{14,147}, Lili Milani¹⁴, Alireza Moayyeri^{56,148}, Keri L. Monda^{4,149}, Mario A. Morken⁹⁹, Antonella Mulas¹⁵⁰, Gabriele Müller¹⁵¹, Martina Müller-Nurasyid^{85,130,152,153}, Arthur W. Musk¹⁵⁴, Ramaiah Nagaraja¹⁵⁵, Markus M. Nöthen^{156,157}, Ilja M. Nolte¹⁵⁸, Stefan Pilz^{159,160}, Nigel W. Rayner^{18,22,64}, Frida Renstrom⁶⁶, Rainer Rettig¹⁶¹, Janina S. Ried⁸⁵, Stephan Ripke^{108,162}, Neil R. Robertson^{18,64}, Lynda M. Rose¹⁶³, Serena Sanna¹⁵⁰, Hubert Scharnagl¹⁶⁴, Salome Scholtens¹⁰⁰, Fredrick R. Schumacher¹⁶⁵, William R. Scott^{41,83}, Thomas Seufferlein¹³⁸, Jianxin Shi¹⁶⁶, Albert Vernon Smith^{167,168}, Joanna Smolonska^{29,169}, Alice V. Stanton¹⁷⁰, Valgerdur Steinthorsdottir¹⁷¹, Kathleen Stirrups^{22,50}, Heather M. Stringham¹, Johan Sundström¹⁴², Morris A. Swertz²⁹, Amy J. Swift⁹⁹, Ann-Christine Syvänen^{16,172}, Sian-Tsung Tan^{41,173}, Bamidele O. Tayo¹⁷⁴, Barbara Thorand^{120,175}, Gudmar Thorleifsson¹⁷¹, Jonathan P. Tyrer¹⁷⁶, Hae-Won Uh^{34,98}, Liesbeth Vandenput¹⁷⁷, Frank C. Verhulst¹⁷⁸, Sita H. Vermeulen^{179,180}, Niek Verweij⁵⁷, Judith M. Vonk¹⁶⁹, Lindsay L. Waite³¹, Helen R. Warren¹⁸¹, Dawn Waterworth¹⁸², Michael N. Weedon²⁴, Lynne R. Wilkens⁵⁴, Christina Willenborg^{183,184}, Tom Wilsgaard¹⁸⁵, Mary K. Wojczynski⁴⁷, Andrew Wong¹⁸⁶, Alan F. Wright¹²⁶, Qunyuan Zhang⁴⁷, The LifeLines Cohort Study[‡], Eoin P. Brennan¹⁸⁷, Murim Choi¹⁸⁸, Zari Dastani¹⁸⁹, Alexander W. Drong¹⁸, Per Eriksson⁷⁰, Anders Franco-Cereceda¹⁹⁰, Jesper R. Gådin⁷⁰, Ali G. Gharavi¹⁹¹, Michael E. Goddard^{192,193}, Robert E. Handsaker^{6,7}, Jinyan Huang^{194,195}, Fredrik Karpe^{64,196}, Sekar Kathiresan^{6,197}, Sarah Keildson¹⁸, Krzysztof Kiryluk¹⁹¹, Michiaki Kubo¹⁹⁸, Jong-Young Lee^{199,†}, Liming Liang^{194,200}, Richard P. Lifton²⁰¹, Baoshan Ma^{194,202}, Steven A. McCarroll^{6,7,162}, Amy J. McKnight²⁰³, Josine L. Min¹⁴⁶, Miriam F. Moffatt¹⁷³, Grant W. Montgomery⁶¹, Joanne M. Murabito^{127,204}, George Nicholson^{205,206}, Dale R. Nyholt^{61,207}, Yukinori Okada^{208,209}, John R. B. Perry^{18,24,56}, Rajkumar Dorajoo²¹⁰, Eva Reinmaa¹⁴, Rany M. Salem^{5,6,7}, Niina Sandholm^{211,212,213}, Robert A. Scott⁹, Lisette Stolk^{34,60}, Atsushi Takahashi²⁰⁸, Toshihiro Tanaka^{209,214,215}, Ferdinand M. van 't Hooft⁷⁰, Anna A. E. Vinkhuyzen¹¹, Harm-Jan Westra²⁹, Wei Zheng²¹⁶, Krina T. Zondervan^{18,217}, The ADIPOGen Consortium[‡], The AGEN-BMI Working Group[‡], The CARDIOGRAMplusC4D Consortium[‡], The CKDGen Consortium[‡], The GLGC[‡], The ICBP[‡], The MAGIC Investigators[‡], The MuTHER Consortium[‡], The MiGen Consortium[‡], The PAGE Consortium[‡], The ReproGen Consortium[‡], The GENIE Consortium[‡], The International Endogene Consortium[‡], Andrew C. Heath²¹⁸, Dominique Arveiler²¹⁹, Stephan J. L. Bakker²²⁰, John Beilby^{87,221}, Richard N. Bergman²²², John Blangero⁹², Pascal Bovet^{223,224}, Harry Campbell³⁶, Mark J. Caulfield¹⁸¹, Giancarlo Cesana²²⁵, Aravinda Chakravarti⁴⁴, Daniel I. Chasman^{163,226}, Peter S. Chines⁹⁹, Francis S. Collins⁹⁹, Dana C. Crawford^{227,228}, L. Adrienne Cupples^{127,229}, Daniele Cusi^{230,231}, John Danesh²³², Ulf de Faire¹¹¹, Hester M. den Ruijter^{76,233}, Anna F. Dominiczak²³⁴, Raimund Erbel²³⁵, Jeanette Erdmann^{183,184}, Johan G. Eriksson^{53,236,237}, Martin Farrall^{18,48}, Stephan B. Felix^{238,239}, Ele Ferrannini^{240,241}, Jean Ferrières²⁴², Ian Ford²⁴³, Nita G. Frouhi⁹, Terrence Forrester²⁴⁴, Oscar H. Franco^{58,59}, Ron T. Gansevoort²²⁰, Pablo V. Gejman²⁴⁵, Christian Gieger⁸⁵, Omri

Gottesman⁵⁵, Vilmondur Gudnason^{167,168}, Ulf Gyllensten¹³⁶, Alistair S. Hall²⁴⁶, Tamara B. Harris¹⁰⁹, Andrew T. Hattersley²⁴⁷, Andrew A. Hicks²⁴⁸, Lucia A. Hindorff²⁴⁹, Aroon D. Hingorani²⁵⁰, Albert Hofman^{58,59}, Georg Homuth⁷³, G. Kees Hovingh²⁵¹, Steve E. Humphries²⁵², Steven C. Hunt²⁵³, Elina Hyppönen^{254,255,256,257}, Thomas Illig^{119,258}, Kevin B. Jacobs^{3,79}, Marjo-Riitta Jarvelin^{83,259,260,261,263,263}, Karl-Heinz Jöckel⁶³, Berit Johansen¹⁰¹, Pekka Jousilahti⁵³, J. Wouter Jukema^{74,264,265}, Antti M. Jula⁵³, Jaakko Kaprio^{53,107,137}, John J. P. Kastelein²⁵¹, Sirkka M. Keinänen-Kiukaanniemi^{263,266}, Lambertus A. Kiemeny^{179,267}, Paul Knekt⁵³, Jaspal S. Kooner^{41,173,268}, Charles Kooperberg⁴⁹, Peter Kovacs^{96,97}, Aldi T. Kraja⁴⁷, Meena Kumari^{269,270}, Johanna Kuusisto²⁷¹, Timo A. Lakka^{125,272,273}, Claudia Langenberg^{9,269}, Loic Le Marchand⁵⁴, Terho Lehtimäki²⁷⁴, Valeriya Lyssenko^{275,276}, Satu Männistö⁵³, André Marette^{277,278}, Tara C. Matise³⁹, Colin A. McKenzie²⁴⁴, Barbara McKnight²⁷⁹, Frans L. Moll²⁸⁰, Andrew D. Morris¹⁰⁶, Andrew P. Morris^{14,18,281}, Jeffrey C. Murray²⁸², Mari Nelis¹⁴, Claes Ohlsson¹⁷⁷, Albertine J. Oldehinkel¹²⁴, Ken K. Ong^{9,186}, Pamela A. F. Madden²¹⁸, Gerard Pasterkamp⁷⁶, John F. Peden²⁸³, Annette Peters^{119,130,175}, Dirkje S. Postma²⁸⁴, Peter P. Pramstaller^{248,285}, Jackie F. Price³⁶, Lu Qi^{13,25}, Olli T. Raitakari^{286,287}, Tuomo Rankinen²⁸⁸, D. C. Rao^{47,71,218}, Treva K. Rice^{71,218}, Paul M. Ridker^{163,226}, John D. Rioux^{143,289}, Marylyn D. Ritchie²⁹⁰, Igor Rudan^{36,291}, Veikko Salomaa⁵³, Nilesh J. Samani^{114,115}, Jouko Saramies²⁹², Mark A. Sarzynski²⁸⁸, Heribert Schunkert^{130,131}, Peter E. H. Schwarz^{121,293}, Peter Sever²⁹⁴, Alan R. Shuldiner^{295,296,297}, Juha Sinisalo²⁹⁸, Ronald P. Stolk¹⁶⁹, Konstantin Strauch^{85,153}, Anke Tönjes^{96,97}, David-Alexandre Trégouët^{299,300,301}, Angelo Tremblay³⁰², Elena Tremoli³⁰³, Jarmo Virtamo⁵³, Marie-Claude Vohl^{278,304}, Uwe Völker^{73,239}, Gérard Waeber³⁰⁵, Gonneke Willemsen¹³³, Jacqueline C. Witteman⁵⁹, M. Carola Zillikens^{58,60}, Linda S. Adair³⁰⁶, Philippe Amouyel³⁰⁷, Folkert W. Asselbergs^{250,264,308}, Themistocles L. Assimes³⁰⁹, Murielle Bochud^{223,224}, Bernhard O. Boehm^{310,311}, Eric Boerwinkle³¹², Stefan R. Bornstein¹²¹, Erwin P. Bottinger⁵⁵, Claude Bouchard²⁸⁸, Stéphane Cauchi^{80,81,82}, John C. Chambers^{41,83,268}, Stephen J. Chanock³, Richard S. Cooper¹⁷⁴, Paul I. W. de Bakker^{77,313,314}, George Dedoussis¹⁰⁵, Luigi Ferrucci⁷², Paul W. Franks^{25,66,67}, Philippe Froguel^{65,80,81,82}, Leif C. Groop^{107,276}, Christopher A. Haiman¹⁶⁵, Anders Hamsten⁷⁰, Jennie Hui^{87,221,315}, David J. Hunter^{13,25,194}, Kristian Hveem¹³², Robert C. Kaplan³¹⁶, Mika Kivimäki²⁶⁹, Diana Kuh¹⁸⁶, Markku Laakso²⁷¹, Yongmei Liu³¹⁷, Nicholas G. Martin⁶¹, Winfried März^{51,164,318}, Mads Melbye^{309,319}, Andres Metspalu^{14,147}, Susanne Moebus⁶³, Patricia B. Munroe¹⁸¹, Inger Njølstad¹⁸⁵, Ben A. Oostra^{32,84,320}, Colin N. A. Palmer¹⁰⁶, Nancy L. Pedersen¹⁵, Markus Perola^{14,53,107}, Louis Pérusse^{278,302}, Ulrike Peters⁴⁹, Chris Power²⁵⁷, Thomas Quertermous³⁰⁹, Rainer Rauramaa^{125,273}, Fernando Rivadeneira^{58,59,60}, Timo E. Saaristo^{321,322}, Danish Saleheen^{232,323,324}, Naveed Sattar³²⁵, Eric E. Schadt³²⁶, David Schlessinger¹⁵⁵, P. Eline Slagboom^{34,35}, Harold Snieder¹⁶⁹, Tim D. Spector⁵⁶, Unnur Thorsteinsdottir^{171,327}, Michael Stumvoll^{96,97}, Jaakko Tuomilehto^{53,328,329,330}, André G. Uitterlinden^{58,59,60}, Matti Uusitupa^{331,332}, Pim van der Harst^{29,57,264}, Mark Walker³³³, Henri Wallaschofski^{239,334}, Nicholas J. Wareham⁹, Hugh Watkins^{18,48}, David R. Weir²⁶, H-Erich Wichmann^{335,336,337},

James F. Wilson³⁶, Pieter Zanen³³⁸, Ingrid B. Borecki⁴⁷, Panos Deloukas^{22,50,339}, Caroline S. Fox¹²⁷, Iris M. Heid^{23,85}, Jeffrey R. O'Connell^{295,296}, David P. Strachan³⁴⁰, Kari Stefansson^{171,327}, Cornelia M. van Duijn^{32,58,59,84}, Gonçalo R. Abecasis¹, Lude Franke²⁹, Timothy M. Frayling²⁴, Mark I. McCarthy^{18,64,341}, Peter M. Visscher^{11,12}, André Scherag^{63,342}, Cristen J. Willer^{28,30,343}, Michael Boehnke¹, Karen L. Mohlke¹⁰, Cecilia M. Lindgren^{6,18}, Jacques S. Beckmann^{20,21,344}, Inês Barroso^{22,345,346}, Kari E. North^{4,347,§}, Erik Ingelsson^{16,17,18,§}, Joel N. Hirschhorn^{5,6,7,§}, Ruth J. F. Loos^{9,55,145,348,§}, and Elizabeth K. Speliotes^{2,§}

Affiliations

¹Center for Statistical Genetics, Department of Biostatistics, University of Michigan, Ann Arbor, Michigan 48109, USA. ²Department of Internal Medicine, Division of Gastroenterology, and Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, Michigan 48109, USA. ³Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, Maryland 20892, USA. ⁴Department of Epidemiology, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA. ⁵Divisions of Endocrinology and Genetics and Center for Basic and Translational Obesity Research, Boston Children's Hospital, Boston, Massachusetts 02115, USA. ⁶Broad Institute of the Massachusetts Institute of Technology and Harvard University, Cambridge, Massachusetts 02142, USA. ⁷Department of Genetics, Harvard Medical School, Boston, Massachusetts 02115, USA. ⁸Center for Biological Sequence Analysis, Department of Systems Biology, Technical University of Denmark, Lyngby 2800, Denmark. ⁹MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Institute of Metabolic Science, Cambridge Biomedical Campus, Cambridge CB2 0QQ, UK. ¹⁰Department of Genetics, University of North Carolina, Chapel Hill, North Carolina 27599, USA. ¹¹Queensland Brain Institute, The University of Queensland, Brisbane 4072, Australia. ¹²The University of Queensland Diamantina Institute, The Translation Research Institute, Brisbane 4012, Australia. ¹³Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts 02115, USA. ¹⁴Estonian Genome Center, University of Tartu, Tartu 51010, Estonia. ¹⁵Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm 17177, Sweden. ¹⁶Science for Life Laboratory, Uppsala University, Uppsala 75185, Sweden. ¹⁷Department of Medical Sciences, Molecular Epidemiology, Uppsala University, Uppsala 75185, Sweden. ¹⁸Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford OX3 7BN, UK. ¹⁹Institute of Social and Preventive Medicine (IUMSP), Centre Hospitalier Universitaire Vaudois (CHUV), Lausanne 1010, Switzerland. ²⁰Swiss Institute of Bioinformatics, Lausanne 1015, Switzerland. ²¹Department of Medical Genetics, University of Lausanne, Lausanne 1005, Switzerland. ²²Wellcome Trust Sanger Institute, Hinxton, Cambridge CB10 1SA, UK. ²³Department of Genetic Epidemiology, Institute of Epidemiology and Preventive Medicine, University of Regensburg, D-93053 Regensburg, Germany. ²⁴Genetics of Complex Traits, University of Exeter Medical School, University of Exeter, Exeter EX1 2LU, UK. ²⁵Department of

Nutrition, Harvard School of Public Health, Boston, Massachusetts 02115, USA. ²⁶Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor, Michigan 48104, USA. ²⁷Department of Epidemiology, University of Michigan, Ann Arbor, Michigan 48109, USA. ²⁸Department of Internal Medicine, Division of Cardiovascular Medicine, University of Michigan, Ann Arbor, Michigan 48109, USA. ²⁹Department of Genetics, University Medical Center Groningen, University of Groningen, 9700 RB Groningen, The Netherlands. ³⁰Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, Michigan 48109, USA. ³¹HudsonAlpha Institute for Biotechnology, Huntsville, Alabama 35806, USA. ³²Genetic Epidemiology Unit, Department of Epidemiology, Erasmus MC University Medical Center, 3015 GE Rotterdam, The Netherlands. ³³Telethon Institute for Child Health Research, Centre for Child Health Research, The University of Western Australia, Perth, Western Australia 6008, Australia. ³⁴Netherlands Consortium for Healthy Aging (NCHA), Leiden University Medical Center, Leiden 2300 RC, The Netherlands. ³⁵Department of Molecular Epidemiology, Leiden University Medical Center, 2300 RC Leiden, The Netherlands. ³⁶Centre for Population Health Sciences, University of Edinburgh, Teviot Place, Edinburgh EH8 9AG, UK. ³⁷Kidney Epidemiology and Cost Center, University of Michigan, Ann Arbor, Michigan 48109, USA. ³⁸Department of Statistics & Biostatistics, Rutgers University, Piscataway, New Jersey 08854, USA. ³⁹Department of Genetics, Rutgers University, Piscataway, New Jersey 08854, USA. ⁴⁰Department of Human Genetics, Leiden University Medical Center, 2333 ZC Leiden, The Netherlands. ⁴¹Ealing Hospital NHS Trust, Middlesex UB1 3HW, UK. ⁴²Department of Gastroenterology and Hepatology, Imperial College London, London W2 1PG, UK. ⁴³Institute of infectious Diseases, Southwest Hospital, Third Military Medical University, Chongqing, China. ⁴⁴Center for Complex Disease Genomics, McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, USA. ⁴⁵Cardiology, Department of Specialties of Internal Medicine, Geneva University Hospital, Geneva 1211, Switzerland. ⁴⁶Department of Epidemiology Research, Statens Serum Institut, Copenhagen DK-2300, Denmark. ⁴⁷Department of Genetics, Washington University School of Medicine, St Louis, Missouri 63110, USA. ⁴⁸Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford OX3 9DU, UK. ⁴⁹Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, Washington 98109, USA. ⁵⁰William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London EC1M 6BQ, UK. ⁵¹Vth Department of Medicine (Nephrology, Hypertensiology, Endocrinology, Diabetology, Rheumatology), Medical Faculty of Mannheim, University of Heidelberg, D-68187 Mannheim, Germany. ⁵²Department of Internal Medicine II, Ulm University Medical Centre, D-89081 Ulm, Germany. ⁵³National Institute for Health and Welfare, FI-00271 Helsinki, Finland. ⁵⁴Epidemiology Program, University of Hawaii Cancer Center, Honolulu, Hawaii 96813, USA. ⁵⁵The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, New York 10029, USA. ⁵⁶Department

of Twin Research and Genetic Epidemiology, King's College London, London SE1 7EH, UK. ⁵⁷Department of Cardiology, University Medical Center Groningen, University of Groningen, 9700RB Groningen, The Netherlands. ⁵⁸Netherlands Consortium for Healthy Aging (NCHA), 3015GE Rotterdam, The Netherlands. ⁵⁹Department of Epidemiology, Erasmus MC University Medical Center, 3015GE Rotterdam, The Netherlands. ⁶⁰Department of Internal Medicine, Erasmus MC University Medical Center, 3015GE Rotterdam, The Netherlands. ⁶¹QIMR Berghofer Medical Research Institute, Brisbane, Queensland 4006, Australia. ⁶²Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, Maryland 20892, USA. ⁶³Institute for Medical Informatics, Biometry and Epidemiology (IMIBE), University Hospital Essen, 45147 Essen, Germany. ⁶⁴Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Oxford OX3 7LJ, UK. ⁶⁵Department of Genomics of Common Disease, School of Public Health, Imperial College London, Hammersmith Hospital, London W12 0NN, UK. ⁶⁶Department of Clinical Sciences, Genetic & Molecular Epidemiology Unit, Lund University Diabetes Center, Skåne University Hospital, Malmö 205 02, Sweden. ⁶⁷Department of Public Health and Clinical Medicine, Unit of Medicine, Umeå University, Umeå 901 87, Sweden. ⁶⁸Department of Odontology, Umeå University, Umeå 901 85, Sweden. ⁶⁹University of Eastern Finland, FI-70210 Kuopio, Finland. ⁷⁰Atherosclerosis Research Unit, Center for Molecular Medicine, Department of Medicine, Karolinska Institutet, Stockholm 17176, Sweden. ⁷¹Division of Biostatistics, Washington University School of Medicine, St Louis, Missouri 63110, USA. ⁷²Translational Gerontology Branch, National Institute on Aging, Baltimore, Maryland 21225, USA. ⁷³Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald, D-17475 Greifswald, Germany. ⁷⁴Department of Cardiology, Leiden University Medical Center, 2300 RC Leiden, The Netherlands. ⁷⁵Department of Gerontology and Geriatrics, Leiden University Medical Center, 2300 RC Leiden, The Netherlands. ⁷⁶Experimental Cardiology Laboratory, Division Heart and Lungs, University Medical Center Utrecht, 3584 CX Utrecht, The Netherlands. ⁷⁷Department of Medical Genetics, University Medical Center Utrecht, 3584 CX Utrecht, The Netherlands. ⁷⁸Department of Endocrinology, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, The Netherlands. ⁷⁹Core Genotyping Facility, SAIC-Frederick, Inc., NCI-Frederick, Frederick, Maryland 21702, USA. ⁸⁰CNRS UMR 8199, F-59019 Lille, France. ⁸¹European Genomic Institute for Diabetes, F-59000 Lille, France. ⁸²Université de Lille 2, F-59000 Lille, France. ⁸³Department of Epidemiology and Biostatistics, Imperial College London, London W2 1PG, UK. ⁸⁴Center for Medical Systems Biology, 2300 RC Leiden, The Netherlands. ⁸⁵Institute of Genetic Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental Health, D-85764 Neuherberg, Germany. ⁸⁶School of Health and Social Studies, Dalarna University, SE-791 88 Falun, Sweden. ⁸⁷PathWest Laboratory Medicine of Western Australia, Nedlands, Western Australia 6009, Australia. ⁸⁸Department of Haematology, University of Cambridge, Cambridge CB2 0PT, UK. ⁸⁹NHS Blood and Transplant, Cambridge CB2 0PT, UK. ⁹⁰Geriatric Unit, Azienda Sanitaria Firenze

(ASF), 50125 Florence, Italy. ⁹¹USC-Office of Population Studies Foundation, Inc., University of San Carlos, Cebu City 6000, Philippines. ⁹²Department of Genetics, Texas Biomedical Research Institute, San Antonio, Texas 78227, USA. ⁹³Genomics Research Centre, Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Queensland 4001, Australia. ⁹⁴Department of Medical Sciences, Endocrinology, Diabetes and Metabolism, Uppsala University, Uppsala 75185, Sweden. ⁹⁵Division of Endocrinology, Diabetes and Metabolism, Ulm University Medical Centre, D-89081 Ulm, Germany. ⁹⁶Integrated Research and Treatment Center (IFB) Adiposity Diseases, University of Leipzig, D-04103 Leipzig, Germany. ⁹⁷Department of Medicine, University of Leipzig, D-04103 Leipzig, Germany. ⁹⁸Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, 2300 RC Leiden, The Netherlands. ⁹⁹Medical Genomics and Metabolic Genetics Branch, National Human Genome Research Institute, NIH, Bethesda, Maryland 20892, USA. ¹⁰⁰LifeLines Cohort Study, University Medical Center Groningen, University of Groningen, 9700 RB Groningen, The Netherlands. ¹⁰¹Department of Biology, Norwegian University of Science and Technology, 7491 Trondheim, Norway. ¹⁰²Department of Pediatrics, University of California Los Angeles, Torrance, California 90502, USA. ¹⁰³Transgenomics Institute, Los Angeles Biomedical Research Institute, Torrance, California 90502, USA. ¹⁰⁴Clinical Trial Service Unit and Epidemiological Studies Unit, Nuffield Department of Population Health, University of Oxford, Oxford OX3 7LF, UK. ¹⁰⁵Department of Dietetics-Nutrition, Harokopio University, 17671 Athens, Greece. ¹⁰⁶Medical Research Institute, University of Dundee, Ninewells Hospital and Medical School, Dundee DD1 9SY, UK. ¹⁰⁷Institute for Molecular Medicine, University of Helsinki, FI-00014 Helsinki, Finland. ¹⁰⁸Analytic and Translational Genetics Unit, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts 02114, USA. ¹⁰⁹Laboratory of Epidemiology and Population Sciences, National Institute on Aging, NIH, Bethesda, Maryland 20892, USA. ¹¹⁰Department of Public Health and Caring Sciences, Geriatrics, Uppsala University, Uppsala 75185, Sweden. ¹¹¹Division of Cardiovascular Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden, Stockholm 17177, Sweden. ¹¹²Kaiser Permanente, Division of Research, Oakland, California 94612, USA. ¹¹³Service of Therapeutic Education for Diabetes, Obesity and Chronic Diseases, Geneva University Hospital, Geneva CH-1211, Switzerland. ¹¹⁴Department of Cardiovascular Sciences, University of Leicester, Glenfield Hospital, Leicester LE3 9QP, UK. ¹¹⁵National Institute for Health Research (NIHR) Leicester Cardiovascular Biomedical Research Unit, Glenfield Hospital, Leicester LE3 9QP, UK. ¹¹⁶Department of Nephrology, University Hospital Regensburg, D-93053 Regensburg, Germany. ¹¹⁷Department of Psychiatry and Psychotherapy, University Medicine Greifswald, HELIOS-Hospital Stralsund, D-17475 Greifswald, Germany. ¹¹⁸German Center for Neurodegenerative Diseases (DZNE), Rostock, Greifswald, D-17475 Greifswald, Germany. ¹¹⁹Research Unit of Molecular Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental Health, D-85764 Neuherberg, Germany. ¹²⁰German Center for Diabetes Research (DZD), 85764 Neuherberg,

Germany. ¹²¹Department of Medicine III, University Hospital Carl Gustav Carus, Technische Universität Dresden, D-01307 Dresden, Germany. ¹²²Institut inter Régional pour la Santé, Synergies, F-37520 La Riche, France. ¹²³Department of Public Health and Clinical Medicine, Unit of Nutritional Research, Umeå University, Umeå 90187, Sweden. ¹²⁴Department of Psychiatry, University of Groningen, University Medical Center Groningen, 9700RB Groningen, The Netherlands. ¹²⁵Kuopio Research Institute of Exercise Medicine, 70100 Kuopio, Finland. ¹²⁶MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Edinburgh EH4 2XU, UK. ¹²⁷National Heart, Lung, and Blood Institute, the Framingham Heart Study, Framingham, Massachusetts 01702, USA. ¹²⁸Department of Neurology, Boston University School of Medicine, Boston, Massachusetts 02118, USA. ¹²⁹Faculty of Psychology and Education, VU University Amsterdam, 1081BT Amsterdam, The Netherlands. ¹³⁰Deutsches Forschungszentrum für Herz-Kreislaferkrankungen (DZHK) (German Research Centre for Cardiovascular Research), Munich Heart Alliance, D-80636 Munich, Germany. ¹³¹Deutsches Herzzentrum München, Technische Universität München, D-80636 Munich, Germany. ¹³²Department of Public Health and General Practice, Norwegian University of Science and Technology, Trondheim 7489, Norway. ¹³³Biological Psychology, VU University Amsterdam, 1081BT Amsterdam, The Netherlands. ¹³⁴Department of Pulmonary Physiology and Sleep Medicine, Nedlands, Western Australia 6009, Australia. ¹³⁵School of Medicine and Pharmacology, University of Western Australia, Crawley 6009, Australia. ¹³⁶Uppsala University, Department of Immunology, Genetics, Pathology, SciLifeLab, Rudbeck Laboratory, SE-751 85 Uppsala, Sweden. ¹³⁷Hjelt Institute Department of Public Health, University of Helsinki, FI-00014 Helsinki, Finland. ¹³⁸Department of Internal Medicine I, Ulm University Medical Centre, D-89081 Ulm, Germany. ¹³⁹Finnish Institute of Occupational Health, FI-90100 Oulu, Finland. ¹⁴⁰Division of Genetic Epidemiology, Department of Medical Genetics, Molecular and Clinical Pharmacology, Innsbruck Medical University, 6020 Innsbruck, Austria. ¹⁴¹Institute of Human Genetics, Helmholtz Zentrum München - German Research Center for Environmental Health, D-85764 Neuherberg, Germany. ¹⁴²Department of Medical Sciences, Cardiovascular Epidemiology, Uppsala University, Uppsala 75185, Sweden. ¹⁴³Montreal Heart Institute, Montreal, Quebec H1T 1C8, Canada. ¹⁴⁴Institute for Community Medicine, University Medicine Greifswald, D-17475 Greifswald, Germany. ¹⁴⁵The Genetics of Obesity and Related Metabolic Traits Program, The Icahn School of Medicine at Mount Sinai, New York, New York 10029, USA. ¹⁴⁶School of Social and Community Medicine, University of Bristol, Bristol BS8 2BN, UK. ¹⁴⁷Institute of Molecular and Cell Biology, University of Tartu, Tartu 51010, Estonia. ¹⁴⁸Farr Institute of Health Informatics Research, University College London, London NW1 2DA, UK. ¹⁴⁹The Center for Observational Research, Amgen, Inc., Thousand Oaks, California 91320, USA. ¹⁵⁰Istituto di Ricerca Genetica e Biomedica (IRGB), Consiglio Nazionale delle Ricerche, Cagliari, Sardinia 09042, Italy. ¹⁵¹Center for Evidence-based Healthcare, University Hospital Carl Gustav Carus, Technische Universität Dresden, D-01307 Dresden, Germany.

¹⁵²Department of Medicine I, University Hospital Grosshadern, Ludwig-Maximilians-Universität, D-81377 Munich, Germany. ¹⁵³Institute of Medical Informatics, Biometry and Epidemiology, Chair of Genetic Epidemiology, Ludwig-Maximilians-Universität, D-81377 Munich, Germany. ¹⁵⁴Department of Respiratory Medicine, Sir Charles Gairdner Hospital, Nedlands, Western Australia 6009, Australia. ¹⁵⁵Laboratory of Genetics, National Institute on Aging, Baltimore, Maryland 21224, USA. ¹⁵⁶Department of Genomics, Life & Brain Center, University of Bonn, 53127 Bonn, Germany. ¹⁵⁷Institute of Human Genetics, University of Bonn, 53127 Bonn, Germany. ¹⁵⁸Department of Epidemiology, University Medical Center Groningen, University of Groningen, 9700 RB Groningen, The Netherlands. ¹⁵⁹Department of Epidemiology and Biostatistics, Institute for Research in Extramural Medicine, Institute for Health and Care Research, VU University Medical Center, 1081BT Amsterdam, The Netherlands. ¹⁶⁰Department of Internal Medicine, Division of Endocrinology and Metabolism, Medical University of Graz, 8036 Graz, Austria. ¹⁶¹Institute of Physiology, University Medicine Greifswald, D-17495 Karlsburg, Germany. ¹⁶²Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, Massachusetts 02142, USA. ¹⁶³Division of Preventive Medicine, Brigham and Women's Hospital, Boston, Massachusetts 02215, USA. ¹⁶⁴Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical University of Graz, Graz 8036, Austria. ¹⁶⁵Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California 90089, USA. ¹⁶⁶National Cancer Institute, Bethesda, Maryland 20892, USA. ¹⁶⁷Icelandic Heart Association, Kopavogur 201, Iceland. ¹⁶⁸University of Iceland, Reykjavik 101, Iceland. ¹⁶⁹Department of Epidemiology, University Medical Center Groningen, University of Groningen, 9700 RB Groningen, The Netherlands. ¹⁷⁰Molecular & Cellular Therapeutics, Royal College of Surgeons in Ireland, 123 St Stephen's Green, Dublin 2, Ireland. ¹⁷¹deCODE Genetics, Amgen Inc., Reykjavik 101, Iceland. ¹⁷²Department of Medical Sciences, Molecular Medicine, Uppsala University, Uppsala 75144, Sweden. ¹⁷³National Heart and Lung Institute, Imperial College London, London SW3 6LY, UK. ¹⁷⁴Department of Public Health Sciences, Stritch School of Medicine, Loyola University of Chicago, Maywood, Illinois 61053, USA. ¹⁷⁵Institute of Epidemiology II, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany, D-85764 Neuherberg, Germany. ¹⁷⁶Department of Oncology, University of Cambridge, Cambridge CB2 0QQ, UK. ¹⁷⁷Centre for Bone and Arthritis Research, Department of Internal Medicine and Clinical Nutrition, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg 413 45, Sweden. ¹⁷⁸Department of Child and Adolescent Psychiatry/Psychology, Erasmus MC University Medical Centre, 3000 CB Rotterdam, The Netherlands. ¹⁷⁹Department for Health Evidence, Radboud University Medical Centre, 6500 HB Nijmegen, The Netherlands. ¹⁸⁰Department of Genetics, Radboud University Medical Centre, 6500 HB Nijmegen, The Netherlands. ¹⁸¹Department of Clinical Pharmacology, William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London EC1M 6BQ, UK. ¹⁸²Genetics,

GlaxoSmithKline, King of Prussia, Pennsylvania 19406, USA. ¹⁸³German Center for Cardiovascular Research, partner site Hamburg/Lubeck/Kiel, 23562 Lubeck, Germany. ¹⁸⁴Institut für Integrative und Experimentelle Genomik, Universität zu Lübeck, D-23562 Lübeck, Germany. ¹⁸⁵Department of Community Medicine, Faculty of Health Sciences, UiT The Arctic University of Norway, 9037 Tromsø, Norway. ¹⁸⁶MRC Unit for Lifelong Health and Ageing at University College London, London WC1B 5JU, UK. ¹⁸⁷Diabetes Complications Research Centre, Conway Institute, School of Medicine and Medical Sciences, University College Dublin, Dublin 4, Ireland. ¹⁸⁸Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul, Korea. ¹⁸⁹Lady Davis Institute, Departments of Human Genetics, Epidemiology and Biostatistics, McGill University, Montréal, Québec H3T1E2, Canada. ¹⁹⁰Cardiothoracic Surgery Unit, Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm 17176, Sweden. ¹⁹¹Department of Medicine, Columbia University College of Physicians and Surgeons, New York 10032, USA. ¹⁹²Biosciences Research Division, Department of Primary Industries, Victoria 3083, Australia. ¹⁹³Department of Food and Agricultural Systems, University of Melbourne, Victoria 3010, Australia. ¹⁹⁴Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts 02115, USA. ¹⁹⁵State Key Laboratory of Medical Genomics, Shanghai Institute of Hematology, Rui Jin Hospital Affiliated with Shanghai Jiao Tong University School of Medicine, Shanghai, China. ¹⁹⁶NIHR Oxford Biomedical Research Centre, OUH Trust, Oxford OX3 7LE, UK. ¹⁹⁷Cardiovascular Research Center, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, USA. ¹⁹⁸Laboratory for Genotyping Development, RIKEN Center for Integrative Medical Sciences, Yokohama 230-0045, Japan. ¹⁹⁹Center for Genome Science, National Institute of Health, Chungcheongbuk-do, Chungbuk 363–951, Republic of Korea. ²⁰⁰Harvard School of Public Health, Department of Biostatistics, Harvard University, Boston, Massachusetts 2115, USA. ²⁰¹Department of Genetics, Howard Hughes Medical Institute, Yale University School of Medicine, New Haven, New Haven, Connecticut 06520, USA. ²⁰²College of Information Science and Technology, Dalian Maritime University, Dalian, Liaoning 116026, China. ²⁰³Nephrology Research, Centre for Public Health, Queen's University of Belfast, Belfast, County Down BT9 7AB, UK. ²⁰⁴Section of General Internal Medicine, Boston University School of Medicine, Boston, Massachusetts 02118, USA. ²⁰⁵Department of Statistics, University of Oxford, 1 South Parks Road, Oxford OX1 3TG, UK. ²⁰⁶MRC Harwell, Harwell Science and Innovation Campus, Harwell OX11 0QG, UK. ²⁰⁷Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Queensland 4059, Australia. ²⁰⁸Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences, Yokohama 230-0045, Japan. ²⁰⁹Department of Human Genetics and Disease Diversity, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, 113-8510 Tokyo, Japan. ²¹⁰Genome Institute of Singapore, Agency for Science, Technology and Research, 138672 Singapore. ²¹¹Department of Biomedical Engineering and Computational Science, Aalto University School of Science, Helsinki FI-00076, Finland.

²¹²Department of Medicine, Division of Nephrology, Helsinki University Central Hospital, FI-00290 Helsinki, Finland. ²¹³Folkhälsan Institute of Genetics, Folkhälsan Research Center, FI-00290 Helsinki, Finland. ²¹⁴Laboratory for Cardiovascular Diseases, RIKEN Center for Integrative Medical Sciences, Yokohama 230-0045, Japan. ²¹⁵Division of Disease Diversity, Bioresource Research Center, Tokyo Medical and Dental University, 113-8510 Tokyo, Japan. ²¹⁶Division of Epidemiology, Department of Medicine; Vanderbilt Epidemiology Center; and Vanderbilt-Ingram Cancer Center, Vanderbilt University Medical Center, Nashville, Tennessee 37075, USA. ²¹⁷Nuffield Department of Obstetrics & Gynaecology, University of Oxford, Oxford OX3 7BN, UK. ²¹⁸Department of Psychiatry, Washington University School of Medicine, St Louis, Missouri 63110, USA. ²¹⁹Department of Epidemiology and Public Health, EA3430, University of Strasbourg, Faculty of Medicine, Strasbourg, France. ²²⁰Department of Internal Medicine, University Medical Center Groningen, University of Groningen, 9700RB Groningen, The Netherlands. ²²¹Pathology and Laboratory Medicine, The University of Western Australia, Perth, Western Australia 6009, Australia. ²²²Cedars-Sinai Diabetes and Obesity Research Institute, Los Angeles, California 90048, USA. ²²³Institute of Social and Preventive Medicine (IUMSP), Centre Hospitalier Universitaire Vaudois and University of Lausanne, 1010 Lausanne, Switzerland. ²²⁴Ministry of Health, Victoria, Republic of Seychelles. ²²⁵University of Milano, Bicocca, 20126, Italy. ²²⁶Harvard Medical School, Boston, Massachusetts 02115, USA. ²²⁷Center for Human Genetics Research, Vanderbilt University Medical Center, Nashville, Tennessee 37203, USA. ²²⁸Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, Tennessee 37232, USA. ²²⁹Department of Biostatistics, Boston University School of Public Health, Boston, Massachusetts 02118, USA. ²³⁰Department of Health Sciences, University of Milano, I 20142, Italy. ²³¹Fondazione Filarete, Milano I 20139, Italy. ²³²Department of Public Health and Primary Care, University of Cambridge, Cambridge CB1 8RN, UK. ²³³Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, 3584 CX Utrecht, The Netherlands. ²³⁴Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow G12 8TA, UK. ²³⁵Clinic of Cardiology, West-German Heart Centre, University Hospital Essen, 45147 Essen, Germany. ²³⁶Department of General Practice and Primary Health Care, University of Helsinki, FI-00290 Helsinki, Finland. ²³⁷Unit of General Practice, Helsinki University Central Hospital, Helsinki 00290, Finland. ²³⁸Department of Internal Medicine B, University Medicine Greifswald, D-17475 Greifswald, Germany. ²³⁹DZHK (Deutsches Zentrum für Herz-Kreislaufforschung – German Centre for Cardiovascular Research), partner site Greifswald, D-17475 Greifswald, Germany. ²⁴⁰Department of Internal Medicine, University of Pisa, 56100 Pisa, Italy. ²⁴¹National Research Council Institute of Clinical Physiology, University of Pisa, 56124 Pisa, Italy. ²⁴²Department of Cardiology, Toulouse University School of Medicine, Rangueil Hospital, 31400 Toulouse, France. ²⁴³Robertson Center for Biostatistics, University of Glasgow, Glasgow G12 8QQ, UK. ²⁴⁴UWI Solutions for Developing Countries, The University of the West Indies, Mona, Kingston 7, Jamaica. ²⁴⁵NorthShore University

HealthSystem, Evanston, IL 60201, University of Chicago, Chicago, Illinois, USA. ²⁴⁶Leeds MRC Medical Bioinformatics Centre, University of Leeds, Leeds LS2 9LU, UK. ²⁴⁷Institute of Biomedical & Clinical Science, University of Exeter, Barrack Road, Exeter EX2 5DW, UK. ²⁴⁸Center for Biomedicine, European Academy Bozen, Bolzano (EURAC), Bolzano 39100, Italy (affiliated institute of the University of Lübeck, D-23562 Lübeck, Germany). ²⁴⁹Division of Genomic Medicine, National Human Genome Research Institute, National Institutes of Health, Bethesda, Maryland 20892, USA. ²⁵⁰Institute of Cardiovascular Science, University College London, London WC1E 6BT, UK. ²⁵¹Department of Vascular Medicine, Academic Medical Center, 1105 AZ Amsterdam, The Netherlands. ²⁵²Centre for Cardiovascular Genetics, Institute Cardiovascular Sciences, University College London, London WC1E 6JJ, UK. ²⁵³Cardiovascular Genetics Division, Department of Internal Medicine, University of Utah, Salt Lake City, Utah 84108, USA. ²⁵⁴Sansom Institute for Health Research, University of South Australia, Adelaide 5000, South Australia, Australia. ²⁵⁵School of Population Health, University of South Australia, Adelaide 5000, South Australia, Australia. ²⁵⁶South Australian Health and Medical Research Institute, Adelaide, South Australia 5000, Australia. ²⁵⁷Population, Policy, and Practice, University College London Institute of Child Health, London WC1N 1EH, UK. ²⁵⁸Hannover Unified Biobank, Hannover Medical School, Hannover, D-30625 Hannover, Germany. ²⁵⁹National Institute for Health and Welfare, FI-90101 Oulu, Finland. ²⁶⁰MRC Health Protection Agency (HPA) Centre for Environment and Health, School of Public Health, Imperial College London, London W2 1PG, UK. ²⁶¹Unit of Primary Care, Oulu University Hospital, FI-90220 Oulu, Finland. ²⁶²Biocenter Oulu, University of Oulu, FI-90014 Oulu, Finland. ²⁶³Institute of Health Sciences, University of Oulu, FI-90014 Oulu, Finland. ²⁶⁴Durrer Center for Cardiogenetic Research, Interuniversity Cardiology Institute Netherlands (ICIN), 3501 DG Utrecht, The Netherlands. ²⁶⁵Interuniversity Cardiology Institute of the Netherlands (ICIN), 3501 DG Utrecht, The Netherlands. ²⁶⁶Unit of Primary Health Care/General Practice, Oulu University Hospital, FI-90220 Oulu, Finland. ²⁶⁷Department of Urology, Radboud University Medical Centre, 6500 HB Nijmegen, The Netherlands. ²⁶⁸Imperial College Healthcare NHS Trust, London W12 0HS, UK. ²⁶⁹Department of Epidemiology and Public Health, University College London, London WC1E 6BT, UK. ²⁷⁰Department of Biological and Social Epidemiology, University of Essex, Wivenhoe Park, Colchester, Essex CO4 3SQ, UK. ²⁷¹Department of Medicine, Kuopio University Hospital and University of Eastern Finland, FI-70210 Kuopio, Finland. ²⁷²Department of Physiology, Institute of Biomedicine, University of Eastern Finland, Kuopio Campus, FI-70211 Kuopio, Finland. ²⁷³Department of Clinical Physiology and Nuclear Medicine, Kuopio University Hospital and University of Eastern Finland, FI-70210 Kuopio, Finland. ²⁷⁴Department of Clinical Chemistry, Fimlab Laboratories and School of Medicine University of Tampere, FI-33520 Tampere, Finland. ²⁷⁵Steno Diabetes Center A/S, Gentofte DK-2820, Denmark. ²⁷⁶Lund University Diabetes Centre and Department of Clinical Science, Diabetes & Endocrinology Unit, Lund University, Malmö 221 00, Sweden. ²⁷⁷Institut Universitaire de Cardiologie et de Pneumologie de Québec,

Faculty of Medicine, Laval University, Quebec, QC G1V 0A6, Canada. ²⁷⁸Institute of Nutrition and Functional Foods, Laval University, Quebec, QC G1V 0A6, Canada. ²⁷⁹Department of Biostatistics, University of Washington, Seattle, Washington 98195, USA. ²⁸⁰Department of Surgery, University Medical Center Utrecht, 3584 CX Utrecht, The Netherlands. ²⁸¹Department of Biostatistics, University of Liverpool, Liverpool L69 3GA, UK. ²⁸²Department of Pediatrics, University of Iowa, Iowa City, Iowa 52242, USA. ²⁸³Illumina, Inc, Little Chesterford, Cambridge CB10 1XL, UK. ²⁸⁴University of Groningen, University Medical Center Groningen, Department of Pulmonary Medicine and Tuberculosis, Groningen, The Netherlands. ²⁸⁵Department of Neurology, General Central Hospital, Bolzano 39100, Italy. ²⁸⁶Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, FI-20521 Turku, Finland. ²⁸⁷Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, FI-20521 Turku, Finland. ²⁸⁸Human Genomics Laboratory, Pennington Biomedical Research Center, Baton Rouge, Louisiana 70808, USA. ²⁸⁹Université de Montréal, Montreal, Quebec H1T 1C8, Canada. ²⁹⁰Center for Systems Genomics, The Pennsylvania State University, University Park, Pennsylvania 16802, USA. ²⁹¹Croatian Centre for Global Health, Faculty of Medicine, University of Split, 21000 Split, Croatia. ²⁹²South Carelia Central Hospital, 53130 Lappeenranta, Finland. ²⁹³Paul Langerhans Institute Dresden, German Center for Diabetes Research (DZD), 01307 Dresden, Germany. ²⁹⁴International Centre for Circulatory Health, Imperial College London, London W2 1PG, UK. ²⁹⁵Division of Endocrinology, Diabetes and Nutrition, University of Maryland School of Medicine, Baltimore, Maryland 21201, USA. ²⁹⁶Program for Personalized and Genomic Medicine, University of Maryland School of Medicine, Baltimore, Maryland 21201, USA. ²⁹⁷Geriatric Research and Education Clinical Center, Veterans Administration Medical Center, Baltimore, Maryland 21201, USA. ²⁹⁸Helsinki University Central Hospital Heart and Lung Center, Department of Medicine, Helsinki University Central Hospital, FI-00290 Helsinki, Finland. ²⁹⁹Sorbonne Universités, UPMC Univ Paris 06, UMR S 1166, F-75013 Paris, France. ³⁰⁰INSERM, UMR S 1166, Team Genomics and Physiopathology of Cardiovascular Diseases, F-75013 Paris, France. ³⁰¹Institute for Cardiometabolism And Nutrition (ICAN), F-75013 Paris, France. ³⁰²Department of Kinesiology, Laval University, Quebec QC G1V 0A6, Canada. ³⁰³Dipartimento di Scienze Farmacologiche e Biomolecolari, Università di Milano & Centro Cardiologico Monzino, Istituto di Ricovero e Cura a Carattere Scientifico, Milan 20133, Italy. ³⁰⁴Department of Food Science and Nutrition, Laval University, Quebec QC G1V 0A6, Canada. ³⁰⁵Department of Internal Medicine, University Hospital (CHUV) and University of Lausanne, Lausanne 1011, Switzerland. ³⁰⁶Department of Nutrition, University of North Carolina, Chapel Hill, North Carolina 27599, USA. ³⁰⁷Institut Pasteur de Lille; INSERM, U744; Université de Lille 2; F-59000 Lille, France. ³⁰⁸Department of Cardiology, Division Heart and Lungs, University Medical Center Utrecht, 3584 CX Utrecht, The Netherlands. ³⁰⁹Department of Medicine, Stanford University School of Medicine, Palo Alto, California 94304, USA. ³¹⁰Lee Kong Chian School of Medicine, Imperial College London and Nanyang Technological University, Singapore, 637553

Singapore, Singapore. ³¹¹Department of Internal Medicine I, Ulm University Medical Centre, D-89081 Ulm, Germany. ³¹²Health Science Center at Houston, University of Texas, Houston, Texas 77030, USA. ³¹³Department of Medicine, Division of Genetics, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115, USA. ³¹⁴Department of Epidemiology, University Medical Center Utrecht, 3584 CX Utrecht, The Netherlands. ³¹⁵School of Population Health, The University of Western Australia, Nedlands, Western Australia 6009, Australia. ³¹⁶Albert Einstein College of Medicine, Department of Epidemiology and Population Health, Belfer 1306, New York 10461, USA. ³¹⁷Center for Human Genetics, Division of Public Health Sciences, Wake Forest School of Medicine, Winston-Salem, North Carolina 27157, USA. ³¹⁸Synlab Academy, Synlab Services GmbH, 68163 Mannheim, Germany. ³¹⁹Department of Clinical Medicine, Copenhagen University, 2200 Copenhagen, Denmark. ³²⁰Department of Clinical Genetics, Erasmus MC University Medical Center, 3000 CA Rotterdam, The Netherlands. ³²¹Finnish Diabetes Association, Kirjoniementie 15, FI-33680 Tampere, Finland. ³²²Pirkanmaa Hospital District, FI-33521 Tampere, Finland. ³²³Center for Non-Communicable Diseases, Karachi, Pakistan. ³²⁴Department of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA. ³²⁵BHF Glasgow Cardiovascular Research Centre, Division of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow G12 8TA, UK. ³²⁶Icahn Institute for Genomics and Multiscale Biology, Icahn School of Medicine at Mount Sinai, New York, New York 10580, USA. ³²⁷Faculty of Medicine, University of Iceland, Reykjavik 101, Iceland. ³²⁸Institute for Health Research, University Hospital of La Paz (IdiPaz), 28046 Madrid, Spain. ³²⁹Diabetes Research Group, King Abdulaziz University, 21589 Jeddah, Saudi Arabia. ³³⁰Centre for Vascular Prevention, Danube-University Krems, 3500 Krems, Austria. ³³¹Department of Public Health and Clinical Nutrition, University of Eastern Finland, Finland. ³³²Research Unit, Kuopio University Hospital, FI-70210 Kuopio, Finland. ³³³Institute of Cellular Medicine, Newcastle University, Newcastle NE1 7RU, UK. ³³⁴Institute of Clinical Chemistry and Laboratory Medicine, University Medicine Greifswald, D-17475 Greifswald, Germany. ³³⁵Institute of Medical Informatics, Biometry and Epidemiology, Chair of Epidemiology, Ludwig-Maximilians-Universität, D-85764 Munich, Germany. ³³⁶Klinikum Grosshadern, D-81377 Munich, Germany. ³³⁷Institute of Epidemiology I, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany, D-85764 Neuherberg, Germany. ³³⁸Department of Pulmonology, University Medical Center Utrecht, 3584 CX Utrecht, The Netherlands. ³³⁹Princess Al-Jawhara Al-Brahim Centre of Excellence in Research of Hereditary Disorders (PACER-HD), King Abdulaziz University, 21589 Jeddah, Saudi Arabia. ³⁴⁰Division of Population Health Sciences & Education, St George's, University of London, London SW17 0RE, UK. ³⁴¹Oxford NIHR Biomedical Research Centre, Oxford University Hospitals NHS Trust, Oxford OX3 7LJ, UK. ³⁴²Clinical Epidemiology, Integrated Research and Treatment Center, Center for Sepsis Control and Care (CSCC), Jena University Hospital, 07743 Jena, Germany. ³⁴³Department of Human Genetics, University of Michigan, Ann Arbor, Michigan

48109, USA. ³⁴⁴Service of Medical Genetics, CHUV University Hospital, 1011 Lausanne, Switzerland. ³⁴⁵University of Cambridge Metabolic Research Laboratories, Institute of Metabolic Science, Addenbrooke's Hospital, Cambridge CB2 0QQ, UK. ³⁴⁶NIHR Cambridge Biomedical Research Centre, Institute of Metabolic Science, Addenbrooke's Hospital, Cambridge CB2 0QQ, UK. ³⁴⁷Carolina Center for Genome Sciences, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA. ³⁴⁸The Mindich Child Health and Development Institute, Icahn School of Medicine at Mount Sinai, New York, New York 10029, USA.

Acknowledgements

A full list of acknowledgments can be found in the Supplementary Information.

References

1. Maes HH, Neale MC, Eaves LJ. Genetic and environmental factors in relative body weight and human adiposity. *Behav. Genet.* 1997; 27:325–351. [PubMed: 9519560]
2. Visscher PM, Brown MA, McCarthy MI, Yang J. Five years of GWAS discovery. *Am. J. Hum. Genet.* 2012; 90:7–24. [PubMed: 22243964]
3. Zaitlen N, et al. Using extended genealogy to estimate components of heritability for 23 quantitative and dichotomous traits. *PLoS Genet.* 2013; 9:e1003520. [PubMed: 23737753]
4. Fall T, Ingelsson E. Genome-wide association studies of obesity and metabolic syndrome. *Mol. Cell. Endocrinol.* 2014; 382:740–757. [PubMed: 22963884]
5. Speliotes EK, et al. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature Genet.* 2010; 42:937–948. [PubMed: 20935630]
6. Willer CJ, et al. Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. *Nature Genet.* 2009; 41:25–34. [PubMed: 19079261]
7. Voight BF, et al. The metabochip, a custom genotyping array for genetic studies of metabolic, cardiovascular, and anthropometric traits. *PLoS Genet.* 2012; 8:e1002793. [PubMed: 22876189]
8. Kilpeläinen TO, et al. Genetic variation near *IRS1* associates with reduced adiposity and an impaired metabolic profile. *Nature Genet.* 2011; 43:753–760. [PubMed: 21706003]
9. Bradfield JP, et al. A genome-wide association meta-analysis identifies new childhood obesity loci. *Nature Genet.* 2012; 44:526–531. [PubMed: 22484627]
10. Monda KL, et al. A meta-analysis identifies new loci associated with body mass index in individuals of African ancestry. *Nature Genet.* 2013; 45:690–696. [PubMed: 23583978]
11. Berndt SI, et al. Genome-wide meta-analysis identifies 11 new loci for anthropometric traits and provides insights into genetic architecture. *Nature Genet.* 2013; 45:501–512. [PubMed: 23563607]
12. Guo Y, et al. Gene-centric meta-analyses of 108 912 individuals confirm known body mass index loci and reveal three novel signals. *Hum. Mol. Genet.* 2013; 22:184–201. [PubMed: 23001569]
13. Wood AR, et al. Defining the role of common variation in the genomic and biological architecture of adult human height. *Nature Genet.* 2014; 46:1173–1186. [PubMed: 25282103]
14. Maller JB, et al. Bayesian refinement of association signals for 14 loci in 3 common diseases. *Nature Genet.* 2012; 44:1294–1301. [PubMed: 23104008]
15. Wakefield J. A Bayesian measure of the probability of false discovery in genetic epidemiology studies. *Am. J. Hum. Genet.* 2007; 81:208–227. [PubMed: 17668372]
16. Peters U, et al. A systematic mapping approach of 16q12.2/*FTO* and BMI in more than 20,000 African Americans narrows in on the underlying functional variation: results from the Population Architecture using Genomics and Epidemiology (PAGE) study. *PLoS Genet.* 2013; 9:e1003171. [PubMed: 23341774]

17. Juster FT, Suzman R. An overview of the Health and Retirement Study. *J. Hum. Resour.* 1995; 30:S7–S56.
18. Bouchonville M, et al. Weight loss, exercise or both and cardiometabolic risk factors in obese older adults: results of a randomized controlled trial. *Int. J. Obes.* 2013; 38:423–431.
19. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* 2011; 88:76–82. [PubMed: 21167468]
20. Yang J, et al. Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nature Genet.* 2012; 44:369–375. [PubMed: 22426310]
21. Pers T, et al. Biological interpretation of genome-wide association studies using predicted gene functions. *Nat. Commun.* 2014; 5:5890.
22. The ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human genome. *Nature.* 2012; 489:57–74. [PubMed: 22955616]
23. Bernstein BE, et al. The NIH Roadmap Epigenomics Mapping Consortium. *Nature Biotechnol.* 2010; 28:1045–1048. [PubMed: 20944595]
24. Segrè AV, Groop L, Mootha VK, Daly MJ, Altshuler D. Common inherited variation in mitochondrial genes is not enriched for associations with type 2 diabetes or related glycaemic traits. *PLoS Genet.* 2010; 6:e1001058. [PubMed: 20714348]
25. Wullschleger S, Loewith R, Hall MN. TOR signaling in growth and metabolism. *Cell.* 2006; 124:471–484. [PubMed: 16469695]
26. Lango Allen H, et al. Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature.* 2010; 467:832–838. [PubMed: 20881960]
27. Raychaudhuri S, et al. Identifying relationships among genomic disease regions: predicting genes at pathogenic SNP associations and rare deletions. *PLoS Genet.* 2009; 5:e1000534. [PubMed: 19557189]
28. Mägi R, et al. Contribution of 32 GWAS-identified common variants to severe obesity in European adults referred for bariatric surgery. *PLoS ONE.* 2013; 8:e70735. [PubMed: 23950990]
29. Lee AW, et al. Functional inactivation of the genome-wide association study obesity gene neuronal growth regulator 1 in mice causes a body mass phenotype. *PLoS ONE.* 2012; 7:e41537. [PubMed: 22844493]
30. Yang Y, Atasoy D, Su HH, Sternson SM. Hunger states switch a flip-flop memory circuit via a synaptic AMPK-dependent positive feedback loop. *Cell.* 2011; 146:992–1003. [PubMed: 21925320]
31. Wu Q, Clark MS, Palmiter RD. Deciphering a neuronal circuit that mediates appetite. *Nature.* 2012; 483:594–597. [PubMed: 22419158]
32. Shen Y, Fu WY, Cheng EY, Fu AK, Ip NY. Melanocortin-4 receptor regulates hippocampal synaptic plasticity through a protein kinase A-dependent mechanism. *J. Neurosci.* 2013; 33:464–472. [PubMed: 23303927]
33. Gibbs JW III, Sombati S, DeLorenzo RJ, Coulter DA. Cellular actions of topiramate: blockade of kainate-evoked inward currents in cultured hippocampal neurons. *Epilepsia.* 2000; 41(suppl. 1):S10–S16. [PubMed: 10768293]
34. Poulsen CF, et al. Modulation by topiramate of AMPA and kainate mediated calcium influx in cultured cerebral cortical, hippocampal and cerebellar neurons. *Neurochem. Res.* 2004; 29:275–282. [PubMed: 14992287]
35. Henao-Mejia J, et al. Inflammation-mediated dysbiosis regulates progression of NAFLD and obesity. *Nature.* 2012; 482:179–185. [PubMed: 22297845]
36. Pruim RJ, et al. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics.* 2010; 26:2336–2337. [PubMed: 20634204]
37. Frazer KA, et al. A second generation human haplotype map of over 3.1 million SNPs. *Nature.* 2007; 449:851–861. [PubMed: 17943122]
38. Winkler TW, et al. Quality control and conduct of genome-wide association meta-analyses. *Nature Protocols.* 2014; 9:1192–1212.

39. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010; 26:2190–2191. [PubMed: 20616382]
40. Devlin B, Roeder K. Genomic control for association studies. *Biometrics*. 1999; 55:997–1004. [PubMed: 11315092]
41. Wen W, et al. Meta-analysis identifies common variants associated with body mass index in east Asians. *Nature Genet*. 2012; 44:307–311. [PubMed: 22344219]
42. Randall JC, et al. Sex-stratified genome-wide association studies including 270,000 individuals show sexual dimorphism in genetic loci for anthropometric traits. *PLoS Genet*. 2013; 9:e1003500. [PubMed: 23754948]
43. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res*. 2010; 38:e164. [PubMed: 20601685]
44. Adzhubei IA, et al. A method and server for predicting damaging missense mutations. *Nature Methods*. 2010; 7:248–249. [PubMed: 20354512]
45. NHLBI Exome Sequencing Project (ESP). Exome Variant Server. <http://evs.gs.washington.edu/EVS/>.
46. Ng PC, Henikoff, S. Predicting deleterious amino acid substitutions. *Genome Res*. 2001; 11:863–874.
47. Mills RE, et al. Mapping copy number variation by population-scale genome sequencing. *Nature*. 2011; 470:59–65. [PubMed: 21293372]
48. Emilsson V, et al. Genetics of gene expression and its effect on disease. *Nature*. 2008; 452:423–428. [PubMed: 18344981]
49. Zhong H, Yang X, Kaplan LM, Molony C, Schadt EE. Integrating pathway analysis and genetics of gene expression for genome-wide association studies. *Am. J. Hum. Genet*. 2010; 86:581–591. [PubMed: 20346437]
50. Grundberg E, et al. Mapping *cis*- and *trans*-regulatory effects across multiple tissues in twins. *Nature Genet*. 2012; 44:1084–1089. [PubMed: 22941192]
51. Dixon AL, et al. A genome-wide association study of global gene expression. *Nature Genet*. 2007; 39:1202–1207. [PubMed: 17873877]
52. Fehrmann RS, et al. Trans-eQTLs reveal that independent genetic variants associated with a complex phenotype converge on intermediate genes, with a major role for the HLA. *PLoS Genet*. 2011; 7:e1002197. [PubMed: 21829388]
53. Nelis M, et al. Genetic structure of Europeans: a view from the North-East. *PLoS ONE*. 2009; 4:e5472. [PubMed: 19424496]
54. Myers AJ, et al. A survey of genetic human cortical gene expression. *Nature Genet*. 2007; 39:1494–1499. [PubMed: 17982457]
55. Westra HJ, et al. Systematic identification of trans eQTLs as putative drivers of known disease associations. *Nature Genet*. 2013; 45:1238–1243. [PubMed: 24013639]
56. Morris AP, et al. Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nature Genet*. 2012; 44:981–990. [PubMed: 22885922]
57. Deloukas P, et al. Large-scale association analysis identifies new risk loci for coronary artery disease. *Nature Genet*. 2013; 45:25–33. [PubMed: 23202125]
58. Ehret GB, et al. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature*. 2011; 478:103–109. [PubMed: 21909115]
59. Shungin D, et al. New genetic loci link adipose and insulin biology to body fat distribution. *Nature*. (this issue) <http://dx.doi.org/nature14132>.
60. Willer C, et al. Discovery and refinement of loci associated with lipid levels. *Nature Genet*. 2013; 45:1274–1283. [PubMed: 24097068]
61. Scott RA, et al. Large-scale association analyses identify new loci influencing glycemic traits and provide insight into the underlying biological pathways. *Nature Genet*. 2012; 44:991–1005. [PubMed: 22885924]
62. Manning AK, et al. A genome-wide approach accounting for body mass index identifies genetic variants influencing fasting glycemic traits and insulin resistance. *Nature Genet*. 2012; 44:659–669. [PubMed: 22581228]

63. Saxena R, et al. Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge. *Nature Genet.* 2010; 42:142–148. [PubMed: 20081857]
64. Dastani Z, et al. Novel loci for adiponectin levels and their influence on type 2 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals. *PLoS Genet.* 2012; 8:e1002607. [PubMed: 22479202]
65. Pattaro C, et al. Genome-wide association and functional follow-up reveals new loci for kidney function. *PLoS Genet.* 2012; 8:e1002584. [PubMed: 22479191]
66. Böger CA, et al. CUBN is a gene locus for albuminuria. *J. Am. Soc. Nephrol.* 2011; 22:555–570. [PubMed: 21355061]
67. Stolk L, et al. Meta-analyses identify 13 loci associated with age at menopause and highlight DNA repair and immune pathways. *Nature Genet.* 2012; 44:260–268. [PubMed: 22267201]
68. Elks CE, et al. Thirty new loci for age at menarche identified by a meta-analysis of genome-wide association studies. *Nature Genet.* 2010; 42:1077–1085. [PubMed: 21102462]
69. Williams WW, et al. Association testing of previously reported variants in a large case-control meta-analysis of diabetic nephropathy. *Diabetes.* 2012; 61:2187–2194. [PubMed: 22721967]
70. Sandholm N, et al. New susceptibility loci associated with kidney disease in type 1 diabetes. *PLoS Genet.* 2012; 8:e1002921. [PubMed: 23028342]
71. Hindorff LA, et al. Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proc. Natl Acad. Sci. USA.* 2009; 106:9362–9367. [PubMed: 19474294]
72. Li Q, Brown JB, Huang H, Bickel PJ. Measuring reproducibility of high-throughput experiments. *Ann. Appl. Stat.* 2011; 5:1752–1779.
73. Feng J, Liu T, Qin B, Zhang Y, Liu XS. Identifying ChIP-seq enrichment using MACS. *Nature Protocols.* 2012; 7:1728–1740.
74. Abecasis GR, et al. An integrated map of genetic variation from 1,092 human genomes. *Nature.* 2012; 491:56–65. [PubMed: 23128226]
75. Fehrmann RS, et al. Gene expression analysis identifies global gene dosage sensitivity in cancer. *Nature Genet.* 2015; 47:115–125. [PubMed: 25581432]




Figure 1. Cumulative variance explained and example of secondary signals

a, The estimated variance in BMI explained by SNPs selected at a range of P values using unrelated individuals from the QIMR ($n = 3,924$; purple) and TwinGene ($n = 5,668$; gold), their weighted average (cyan), inferred from within-family prediction (red; Extended Data Fig. 2), and by all HapMap phase III SNPs in 16,275 unrelated individuals from the QIMR, TwinGene and ARIC studies (orange). **b**, Plot of the region surrounding *MC4R* (ref. 36). SNP associations from the European sex-combined meta-analysis are plotted with joint conditional P values (P_j) indicated for the three conditionally significant signals. SNPs are shaded and shaped based on the index SNP with which they are in strongest LD (rs6567160 in blue, rs994545 in yellow and rs17066842 in green).




Figure 2. Tissues and reconstituted gene sets significantly enriched for genes within BMI-associated loci

a, DEPICT predicts genes within BMI-associated loci ($P < 5 \times 10^{-4}$) are enriched for expression in the brain and central nervous system. Tissues are sorted by physiological system and significantly enriched tissues are in black; the dotted line represents statistically significant enrichment. **b**, The gene sets most significantly enriched for BMI-associated loci by DEPICT ($P < 10^{-6}$, FDR $< 4 \times 10^{-4}$). Nodes represent reconstituted gene sets and are colour-coded by P value. Edge thickness between nodes is proportional to degree of gene overlap as measured by the Jaccard index. Nodes with gene overlap greater than 25% were collapsed into a single ‘meta-node’ (blue border). **c**, The nodes contained within the most enriched meta-node, ‘clathrin-coated vesicle’, which shares genes with other gene sets relevant to glutamate signalling and synapse biology. **d**, The ‘generation of a signal involved in cell–cell signalling’ meta-node represents several overlapping gene sets relevant to obesity and energy metabolism (gene sets with $P < 4 \times 10^{-3}$, FDR < 0.05 shown). For the complete list of enriched gene sets refer to Supplementary Table 21a.

Table 1

Novel GWS BMI loci In European meta-analysis

SNP	Chr:position	Notable gene(s)*	Alleles	EAF	β	s.e.m.	P value
rs657452	1:49,362,434	<i>AGBL4</i> (N)	A/G	0.394	0.023	0.003	5.48×10^{-13}
rs12286929	11:114,527,614	<i>CADM1</i> (N)	G/A	0.523	0.022	0.003	1.31×10^{-12}
rs7903146	10:114,748,339	<i>TCF7L2</i> (B,N)	C/T	0.713	0.023	0.003	1.11×10^{-11}
rs10132280	14:24,998,019	<i>STXBP6</i> (N)	C/A	0.682	0.023	0.003	1.14×10^{-11}
rs17094222	10:102,385,430	<i>HIF1AN</i> (N)	C/T	0.211	0.025	0.004	5.94×10^{-11}
rs7599312	2:213,121,476	<i>ERBB4</i> (D,N)	G/A	0.724	0.022	0.003	1.17×10^{-10}
rs2365389	3:61,211,502	<i>FHIT</i> (N)	C/T	0.582	0.020	0.003	1.63×10^{-10}
rs2820292	1:200,050,910	<i>NAVI</i> (N)	C/A	0.555	0.020	0.003	1.83×10^{-10}
rs12885454	14:28,806,589	<i>PRKD1</i> (N)	C/A	0.642	0.021	0.003	1.94×10^{-10}
rs16851483	3:142,758,126	<i>RASA2</i> (N)	T/G	0.066	0.048	0.008	3.55×10^{-10}
rs1167827	7:75,001,105	<i>HIP1</i> (B,N); <i>PMS2L3</i> (B,Q); <i>PMS2P5</i> (Q); <i>WBSR16</i> (Q)	G/A	0.553	0.020	0.003	6.33×10^{-10}
rs758747	16:3,567,359	<i>NLRC3</i> (N)	T/C	0.265	0.023	0.004	7.47×10^{-10}
rs1928295	9:119,418,304	<i>TLR4</i> (B,N)	T/C	0.548	0.019	0.003	7.91×10^{-10}
rs9925964	16:31,037,396	<i>KAT8</i> (N); <i>ZNF646</i> (M,Q); <i>VKORC1</i> (Q); <i>ZNF668</i> (Q); <i>STX1B</i> (D); <i>FBXL19</i> (D)	A/G	0.620	0.019	0.003	8.11×10^{-10}
rs11126666	2:26,782,315	<i>KCNK3</i> (D,N)	A/G	0.283	0.021	0.003	1.33×10^{-9}
rs2650492	16:28,240,912	<i>SBK1</i> (D,N); <i>APOBR</i> (B)	A/G	0.303	0.021	0.004	1.92×10^{-9}
rs6804842	3:25,081,441	<i>RARB</i> (B)	G/A	0.575	0.019	0.003	2.48×10^{-9}
rs4740619	9:15,624,326	<i>C9orf93</i> (C,M,N)	T/C	0.542	0.018	0.003	4.56×10^{-9}
rs13191362	6:162,953,340	<i>PARK2</i> (B,D,N)	A/G	0.879	0.028	0.005	7.34×10^{-9}
rs3736485	15:49,535,902	<i>SCG3</i> (B,D); <i>DMXL2</i> (M,N)	A/G	0.454	0.018	0.003	7.41×10^{-9}
rs17001654	4:77,348,592	<i>NUP54</i> (M); <i>SCARB2</i> (Q,N)	G/C	0.153	0.031	0.005	7.76×10^{-9}
rs11191560	10:104,859,028	<i>NT5C2</i> (N); <i>CYP17A1</i> (B); <i>SFXN2</i> (Q)	C/T	0.089	0.031	0.005	8.45×10^{-9}
rs1528435	2:181,259,207	<i>UBE2E3</i> (N)	T/C	0.631	0.018	0.003	1.20×10^{-8}
rs1000940	17:5,223,976	<i>RABEP1</i> (N)	G/A	0.320	0.019	0.003	1.28×10^{-8}
rs2033529	6:40,456,631	<i>TDRG1</i> (N); <i>LRFN2</i> (D)	G/A	0.293	0.019	0.003	1.39×10^{-8}
rs11583200	1:50,332,407	<i>ELAVL4</i> (B,D,N,Q)	C/T	0.396	0.018	0.003	1.48×10^{-8}
rs9400239	6:109,084,356	<i>FOXO3</i> (B,N); <i>HSS00296402</i> (Q)	C/T	0.688	0.019	0.003	1.61×10^{-8}
rs10733682	9:128,500,735	<i>LMX1B</i> (B,N)	A/G	0.478	0.017	0.003	1.83×10^{-8}
rs11688816	2:62,906,552	<i>EHBPI</i> (B,N)	G/A	0.525	0.017	0.003	1.89×10^{-8}
rs11057405	12:121,347,850	<i>CLIP1</i> (N)	G/A	0.901	0.031	0.006	2.02×10^{-8}
rs11727676	4:145,878,514	<i>HHIP</i> (B,N)	T/C	0.910	0.036	0.006	2.55×10^{-8}
rs3849570	3:81,874,802	<i>GBE1</i> (B,M,N)	A/C	0.359	0.019	0.003	2.60×10^{-8}
rs6477694	9:110,972,163	<i>EPB41L4B</i> (N); <i>C9orf4</i> (D)	C/T	0.365	0.017	0.003	2.67×10^{-8}
rs7899106	10:87,400,884	<i>GRID1</i> (B,N)	G/A	0.052	0.040	0.007	2.96×10^{-8}
rs2176598	11:43,820,854	<i>HSD17B12</i> (B,M,N)	T/C	0.251	0.020	0.004	2.97×10^{-8}
rs2245368	7:76,446,079	<i>PMS2L11</i> (N)	C/T	0.180	0.032	0.006	3.19×10^{-8}

SNP	Chr:position	Notable gene(s)*	Alleles	EAF	β	s.e.m.	<i>P</i> value
rs17724992	19:18,315,825	<i>GDF15</i> (B); <i>PGPEP1</i> (Q,N)	A/G	0.746	0.019	0.004	3.42×10^{-8}
rs7243357	18:55,034,299	<i>GRP</i> (B,G,N)	T/G	0.812	0.022	0.004	3.86×10^{-8}
rs2033732	8:85,242,264	<i>RALYL</i> (D,N)	C/T	0.747	0.019	0.004	4.89×10^{-8}

GWS is defined as $P < 5 \times 10^{-8}$. SNP positions are reported according to Build 36 and their alleles are coded based on the positive strand. Alleles (effect/other), effect allele frequency (EAF), beta (β), standard error of the mean (s.e.m.) and *P* values are based on the meta-analysis of GWAS I + II + MetaboChip association data from the European sex-combined data set.

* Notable genes from biological relevance to obesity (B); copy number variation (C); DEPICT analyses (D); GRAIL results (G); BMI-associated variants in strong LD ($r^2 \geq 0.7$) with a missense variant in the indicated gene (M); gene nearest to index SNP (N); association and eQTL data converge to affect gene expression (Q).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

GWS BMI loci from secondary analyses

SNP	Chr:position	Notable gene(s)*	Alleles	EAF	β	s.e.m.	P value	Analysis
Novel loci								
rs9641123	7:93,035,668	<i>CALCR</i> (B,N); <i>hsa-miR-653</i> (Q)	C/G	0.430	0.029	0.005	2.08×10^{-10}	EPB
rs7164727	15:70,881,044	<i>LOC100287559</i> (N), <i>BBS4</i> (B,M,Q)	T/C	0.671	0.019	0.003	3.92×10^{-9}	All
rs492400	2:219,057,996	<i>PLCD4</i> (B,Q); <i>CYP27A1</i> (B); <i>USP37</i> (N); <i>TTLA4</i> (M,Q); <i>STK36</i> (B,M); <i>ZNF142</i> (M); <i>RQCD1</i> (Q)	C/T	0.424	0.024	0.004	6.78×10^{-9}	Men
rs2080454	16:47,620,091	<i>CBLN1</i> (N)	C/A	0.413	0.017	0.003	8.60×10^{-9}	All
rs7239883	18:38,401,669	<i>LOC284260</i> (N); <i>RIT2</i> (B,D)	G/A	0.391	0.023	0.004	1.51×10^{-8}	Women
rs2836754	21:39,213,610	<i>ETS2</i> (N)	C/T	0.599	0.017	0.003	1.61×10^{-8}	All
rs9914578	17:1,951,886	<i>SMG6</i> (D,N); <i>N29617</i> (Q)	G/C	0.229	0.020	0.004	2.07×10^{-8}	All
rs977747	1:47,457,264	<i>TALI</i> (N)	T/G	0.403	0.017	0.003	2.18×10^{-8}	All
rs9374842	6:120,227,364	<i>LOC285762</i> (N);	T/C	0.744	0.023	0.004	2.67×10^{-8}	EPB
rs4787491	16:29,922,838	<i>MAPK3</i> (D); <i>KCTD13</i> (D); <i>INO80E</i> (N); <i>TAOK2</i> (D); <i>YPEL3</i> (D); <i>DOC2A</i> (D); <i>FAM57B</i> (D)	G/A	0.510	0.022	0.004	2.70×10^{-8}	EPB
rs1441264	13:78,478,920	<i>MIR548A2</i> (N)	A/G	0.613	0.017	0.003	2.96×10^{-8}	All
rs17203016	2:207,963,763	<i>CREB1</i> (B,N); <i>KLF7</i> (B)	G/A	0.195	0.021	0.004	3.41×10^{-8}	All
rs16907751	8:81,538,012	<i>ZBTB10</i> (N)	C/T	0.913	0.047	0.009	3.89×10^{-8}	Men
rs13201877	6:137,717,234	<i>IFNGR1</i> (N); <i>OLIG3</i> (G)	G/A	0.140	0.024	0.004	4.29×10^{-8}	All
rs9540493	13:65,103,705	<i>MIR548X2</i> (N); <i>PCDH9</i> (D)	A/G	0.452	0.021	0.004	4.97×10^{-8}	EPB
rs1460676	2:164,275,935	<i>FIGN</i> (N)	C/T	0.179	0.021	0.004	4.98×10^{-8}	All
rs6465468	7:95,007,450	<i>ASB4</i> (B,N)	T/G	0.306	0.025	0.005	4.98×10^{-8}	Women
Previously identified loci								
rs6091540	20:50,521,269	<i>ZFP64</i> (N)	C/T	0.721	0.030	0.004	2.15×10^{-11}	Women
rs7715256	5:153,518,086	<i>GALNT10</i> (N)	G/T	0.422	0.017	0.003	8.85×10^{-9}	All
rs2176040	2:226,801,046	<i>LOC646736</i> (N); <i>IRS1</i> (B,Q)	A/G	0.365	0.024	0.004	9.99×10^{-9}	Men

SNP positions are reported according to Build 36 and their alleles are coded based on the positive strand. Alleles (effect/other), EAF, beta (β), s.e.m. and P values are based on the meta-analysis of GWAS I + II+ MetaboChip association data from the data set shown in the 'Analysis' column. EPB denotes European population-based studies, 'All' denotes all ancestries.

* Notable genes from biological relevance to obesity (B); copy number variation (C); DEPICT analyses (D); GRAIL results (G); BMI-associated variant is in strong LD ($r^2 > 0.7$) with a missense variant in the indicated gene (M); gene nearest to the index SNP (N); association and eQTL data converge to affect gene expression (Q).

Table 3

Secondary signals reaching GWS by conditional analysis

SNP	Chr: position	Nearest gene	Alleles	EAF	β	s.e.m.	Variance explained	<i>P</i> value
rs1016287	2:59159129	<i>LINC01122</i>	T/C	0.294	0.023	0.003	0.021%	2.62×10^{-11}
rs4671328	2:58788786	<i>LINC01122</i>	T/G	0.457	0.021	0.004	0.021%	2.73×10^{-8}
rs758747	16:3567359	<i>NLRC3</i>	T/C	0.241	0.022	0.004	0.018%	2.00×10^{-9}
rs879620	16:3955730	<i>ADCY9</i>	T/C	0.620	0.024	0.004	0.027%	2.17×10^{-9}
rs12446632	16:19842890	<i>GPRC5B</i>	G/A	0.860	0.036	0.005	0.031%	1.06×10^{-14}
rs11074446	16:20162624	<i>GP2</i>	T/C	0.867	0.029	0.005	0.019%	1.71×10^{-10}
rs6567160	18:55980115	<i>MC4R</i>	C/T	0.233	0.048	0.004	0.084%	3.52×10^{-38}
rs17066842	18:56191604	<i>MC4R</i>	G/A	0.960	0.051	0.008	0.020%	6.99×10^{-10}
rs9944545	18:56109224	<i>MC4R</i>	T/C	0.296	0.020	0.004	0.017%	1.01×10^{-8}
rs11030104	11:27641093	<i>BDNF</i>	A/G	0.791	0.051	0.004	0.087%	1.26×10^{-34}
rs10835210	11:27652486	<i>BDNF</i>	C/A	0.570	0.020	0.004	0.020%	1.25×10^{-8}

SNP positions are reported according to Build 36 and their alleles are coded based on the positive strand. Alleles (effect/other), EAF, estimated beta (β), s.e.m., explained variance, and *P* values from GCTA. First row at each locus represents lead signal, other row(s) represent secondary signals.