



Society position statements/white papers

Guidelines for postoperative care in gynecologic/oncology surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations – Part II



G. Nelson^{a,*}, A.D. Altman^b, A. Nick^c, L.A. Meyer^c, P.T. Ramirez^c, C. Ahtari^d, J. Antrobus^e, J. Huang^f, M. Scott^{g,h}, L. Wijkⁱ, N. Acheson^j, O. Ljungqvist^k, S.C. Dowdy^l

^a Department of Gynecologic Oncology, Tom Baker Cancer Centre, Calgary, Alberta, Canada

^b Department of Obstetrics, Gynecology and Reproductive Sciences, University of Manitoba, Winnipeg, Manitoba, Canada

^c Department of Gynecologic Oncology and Reproductive Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX, USA

^d Department of Obstetrics and Gynecology, Lausanne University Hospital, Lausanne, Switzerland

^e Department of Anesthesiology, Borders General Hospital, Melrose, Roxburghshire, United Kingdom

^f Anesthesiologists of Greater Orlando, Orlando, FL, USA

^g Department of Anaesthesia and Intensive Care Medicine, Royal Surrey County NHS Foundation Hospital, Egerton Road, Guildford, United Kingdom

^h Surrey Peri-operative Anaesthesia Critical Care Research group (SPACeR) Clinical Academic Group, FHMS, University of Surrey, United Kingdom

ⁱ Department of Obstetrics and Gynecology, Faculty of Medicine and Health, Örebro University, Örebro, Sweden

^j Department of Gynaecologic Oncology, Royal Devon & Exeter NHS Foundation Trust, Exeter, United Kingdom

^k Faculty of Medicine and Health, School of Health and Medical Sciences, Department of Surgery, Örebro University, Örebro, Sweden

^l Division of Gynecologic Surgery, Mayo Clinic College of Medicine, Rochester, MN, USA

HIGHLIGHTS

- We provide evidence supporting postoperative management of patients undergoing gynecologic/oncology surgery.
- This guideline will help integrate knowledge into practice, align perioperative care, and encourage future investigations.

ARTICLE INFO

Article history:

Received 24 September 2015

Received in revised form 14 December 2015

Accepted 21 December 2015

Available online 3 January 2016

Keywords:

Enhanced recovery after surgery

Gynecologic/oncology

Evidence based postoperative care

ABSTRACT

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The “Guidelines for Pre- and Intra-operative Care in Gynecologic/Oncology Surgery: Enhanced Recovery After Surgery (ERAS®) Society Recommendations – Part I,” [1] examined the evidence surrounding care elements including preoperative medical optimization, bowel preparation, carbohydrate loading, thromboembolism prophylaxis, skin preparation, standard anesthetic protocol and intraoperative fluid

management. The goal of this article is to critically review existing evidence and make recommendations for elements of postoperative care. This effort forms the basis of the ERAS® Guideline for postoperative care in gynecologic/oncology surgery.

2. Methods

2.1. Literature search

The authors convened in July 2014 to discuss topics for inclusion – the topic list was based on the ERAS® Colonic Surgery [2] and Rectal/

* Corresponding author at: Tom Baker Cancer Centre, 1331 29th St NW, Calgary, Alberta T2N 4N2, Canada.

E-mail address: gsnelson@ucalgary.ca (G. Nelson).

Pelvic [3] Guidelines which were used as templates. After the topics were agreed upon they were then allocated amongst the group according to expertise. The literature search (1966–2014) used Embase and PubMed to search medical subject headings including “gynecology”, “gynecologic oncology” and all postoperative ERAS® items (see Table 1). Reference lists of all eligible articles were crosschecked for other relevant studies.

2.2. Study selection

Titles and abstracts were screened by individual reviewers to identify potentially relevant articles. Discrepancies in judgment were resolved by the lead (GN) and senior authors (OL, SD). Meta-analyses, systematic reviews, randomized controlled studies, non-randomized controlled studies, reviews, and case series were considered for each individual topic.

2.3. Quality assessment and data analyses

The quality of evidence and recommendations were evaluated according to the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) system (see Tables 2a and 2b) [4] whereby recommendations are given as follows: Strong recommendations indicate that the panel is confident that the desirable effects of adherence to a recommendation outweigh the undesirable effects. Weak recommendations indicate that the desirable effects of adherence to a recommendation probably outweigh the undesirable effects, but the panel is less

Table 2a
GRADE system for rating quality of evidence.

Evidence level	Definition
High quality	Further research unlikely to change confidence in estimate of effect
Moderate quality	Further research likely to have important impact on confidence in estimate of effect and may change the estimate
Low quality	Further research very likely to have important impact on confidence in estimate of effect and likely to change the estimate
Very low quality	Any estimate of effect is very uncertain

Reference [4].

confident. Recommendations are based on quality of evidence: high, moderate, low and very low but also on the balance between desirable and undesirable effects; and on values and preferences. As such, consistent with other ERAS® Guideline Working groups [2,5], in some cases strong recommendations may be reached from low-quality data and vice versa. Of note, this would be considered a modified GRADE evaluation since we did not consider resource utilization when making our recommendations [6].

3. Results

The evidence base, recommendations, evidence level, and recommendation grade are provided for each individual ERAS® item below.

Table 1
Guidelines for postoperative care in gynecologic/oncology surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations.

Item	Recommendation	Evidence level	Recommendation grade
Prophylaxis against thromboembolism	Patients should wear well-fitting compression stockings and have intermittent pneumatic compression	High	Strong
	Extended prophylaxis (28 days) should be given to patients after laparotomy for abdominal or pelvic malignancies	High	Strong
Postoperative fluid therapy	Intravenous fluids should be terminated within 24 h after surgery; balanced crystalloid solutions are preferred to 0.9% normal saline	Moderate	Strong
Perioperative nutritional care	A regular diet within the first 24 h after gynecologic/oncology surgery is recommended	High	Strong
	The use of postoperative laxatives should be considered	Low	Weak
Prevention of postoperative ileus	The use of chewing gum should be considered	Moderate	Weak
Postoperative glucose control	ERAS elements that reduce metabolic stress should be employed to reduce insulin resistance and the development of hyperglycemia	High	Strong
	Perioperative maintenance of blood glucose levels (<180–200 mg/dL) results in improved perioperative outcomes; glucose levels above this range should be treated with insulin infusions and regular blood glucose monitoring to avoid the risk of hypoglycemia	High	Strong
Postoperative analgesia	A multimodal approach to analgesia should be adopted including use of NSAIDs/acetaminophen, gabapentin and dexamethasone (unless contraindications exist)	Multimodal: high NSAIDs/acetaminophen: high Gabapentin: moderate Dexamethasone: low	Strong
	Vaginal hysterectomy		
	Paracervical nerve block or intrathecal morphine can be used to reduce pain and opioid consumption	Low	Weak
	Open general gynecologic surgery		
	Spinal anesthesia with intrathecal morphine is recommended	Moderate	Strong
	Alternatively, thoracic epidural analgesia (TEA) with low concentration local anesthetic solutions with the addition of opiates for 24–48 h can be considered	High	Strong
	Truncal nerve blocks (TAP or ilioinguinal) can be recommended where patients have undergone general anesthesia without neuraxial blockade	Moderate	Strong
	Continuous wound infiltration (CWI) of local anesthetic can be considered	Moderate	Strong
	Major oncologic surgery		
	TEA may be considered but patients frequently require additional IV opioids in addition to TEA to achieve adequate analgesia	Low	Weak
Laparoscopic gynecologic/oncology surgery			
Lack of evidence makes it difficult to recommend one analgesic intervention over another, however a multimodal approach should be employed	Low	Weak	
Peritoneal drainage	Peritoneal drainage is not recommended routinely in gynecologic/oncology surgery including for patients undergoing lymphadenectomy or bowel surgery	Moderate	Strong
Urinary drainage	Urinary catheters should be used for postoperative bladder drainage for a short period preferably <24 h postop	Low	Strong
Early mobilization	Patients should be encouraged to mobilize within 24 h of surgery	Low	Strong

Table 2b
GRADE system for rating strength of recommendations.

Recommendation strength	Definition
Strong	When desirable effects of intervention clearly outweigh the undesirable effects, or clearly do not
Weak	When trade-offs are less certain – either because of low quality evidence or because evidence suggests desirable and undesirable effects are closely balanced

Reference [4].

4. Postoperative thromboembolism prophylaxis

4.1. Immediate postoperative prophylaxis

Pneumatic compression stockings reduce the rate of VTE (venous thromboembolism) when compared to observation [7]. The risk reduction is equivalent when compared to heparin [8] and improved when combined with heparin [9] in gynecologic oncology patients. Graduated compression stockings decrease the rate of DVT in hospitalized patients, especially when combined with another method [10].

4.2. Extended postoperative prophylaxis

A large prospective cohort trial showed an increased rate of VTE within 30 days of surgery in cancer patients [11], and extended prophylaxis (28 days) is now considered a common practice within major gynecologic oncology surgery [12]. A Cochrane review of 4 randomized controlled trials examining extended prophylaxis has shown a decrease in VTE (14.3% vs. 6.1%; $p < 0.0005$) and a decrease in symptomatic VTE (1.7% vs. 0.2%; $p = 0.02$) [13]. The role of extended prophylaxis in minimally invasive surgery is likely not necessary without other high-risk features (elevated BMI, previous VTE, coagulopathy, decreased mobility) [14].

4.2.1. Summary and recommendations

Patients should wear well-fitting compression stockings and have intermittent pneumatic compression. Extended prophylaxis (28 days) should be given to patients after laparotomy for abdominal or pelvic malignancies.

4.2.2. Evidence level

High.

4.2.3. Recommendation grade

Strong.

5. Postoperative fluid therapy

Oral intake of fluid and food should be started the day of surgery whenever possible. With the commencement of oral diet and oral analgesia as soon as tolerated after surgery the need for postoperative intravenous fluids beyond 12–24 h is rarely needed in an uncomplicated recovery. Patients can drink immediately after surgery. Flavored high energy protein drinks prescribed three times a day are safe and can bridge the postoperative period of building back up to a normal diet to ensure some protein and calorie intake early in the recovery process. They are usually 200–250 ml in volume with around 150 kJ/100 ml of carbohydrate and 3–6 g/100 mL of protein with the addition of vitamins, mineral and trace elements. If intravenous fluids must be maintained then a total hourly volume of no more than 1.2 mL/kg (including drugs, approximately 90 mL/h for a 75 kg female) should be given [15]. Balanced crystalloid solutions are preferred to 0.9% normal saline due to the cumulative risk of hyper-chloremic acidosis. The use of starch solutions during the perioperative period should be limited by dose and duration to avoid the adverse effects seen in studies on intensive care

patients such as bleeding and renal dysfunction [16]. Oliguria as low as 20cm³/h is a normal response to surgery, and the need for further intravenous fluid boluses should be assessed within clinical context. A small proportion of patients undergoing major surgery will develop SIRS (Systemic Inflammatory Response Syndrome) causing marked vasodilation and hypotension without sepsis. These patients will require vasopressor therapy such as a noradrenaline infusion during surgery and postoperatively until resolution.

5.1. Summary and recommendations

Intravenous fluids should be terminated within 24 h after surgery. Balanced crystalloid solutions are preferred to 0.9% normal saline.

5.2. Evidence level

Moderate.

5.3. Recommendation grade

Strong.

6. Perioperative nutritional care

A number of randomized trials on the subject of early feeding (defined as having oral intake of fluids or food within the first 24 h after surgery) have been performed in gynecologic oncology [17–20]. Effects include accelerated return of bowel activity, reduced length of stay, with no evidence of higher complication rates related to wound healing, anastomotic leaks, or pulmonary complications. A randomized study in patients with ovarian cancer showed a significantly lower rate of complications for patients receiving early feeding. However, complication rates were not different between groups when the analysis was limited to a smaller cohort of patients undergoing intestinal resections [18,19]. It is important to note that early feeding is associated with a higher rate of nausea, but not vomiting, abdominal distension, or nasogastric tube use. Patient satisfaction with control of vomiting in one series was over 90% with early feeding despite a higher incidence of nausea in the enhanced recovery group [21].

6.1. Summary and recommendation

A regular diet within the first 24 h after gynecologic/oncology surgery is recommended.

6.2. Evidence level

High.

6.3. Recommendation grade

Strong.

7. Prevention of postoperative ileus

Laxatives are commonly used within enhanced recovery protocols to hasten the return of bowel function, but no high quality data is available in gynecologic oncology. In one prospective, but nonrandomized trial of 20 patients undergoing open radical hysterectomy, milk of magnesia and biscolic suppositories were well tolerated and associated with a reduction in hospital stay compared with historical controls [22]. In 68 patients undergoing hepatic resection via laparotomy, patients randomized to magnesium hydroxide experienced a median one-day reduction in time to passage of stool [23]. Although data are limited and effects appear modest, continued use of laxatives is reasonable given the low cost and side effect profile.

In patients undergoing hysterectomy and colonic resection, randomized trials have shown improved recovery when a peripheral mu antagonist was administered [24]. Its use in patients undergoing planned enteric resections is reasonable, but we cannot provide a recommendation for its use at the present time as cost-effectiveness and efficacy data in patients with gynecologic malignancies continues to be collected. Perioperative use of chewing gum had a positive effect on the incidence of postoperative ileus (36% vs. 15%) and length of stay (1 day reduction) in a randomized trial of patients undergoing staging for gynecologic malignancies [25]. A meta-analysis of randomized trials investigating prokinetics such as erythromycin, the cholecystokinin-like drugs, cisapride, dopamine-antagonists, propranolol, vasopressin, and intravenous lidocaine [26] or neostigmine [27] failed to demonstrate benefit.

7.1. Summary and recommendations

The use of postoperative laxatives and chewing gum should be considered.

7.2. Evidence level

Laxatives: Low.

Chewing gum: Moderate.

7.3. Recommendation grade

Weak.

8. Postoperative control of glucose

Perioperative hyperglycemia, classically defined as blood glucose levels greater than 180 to 200 mg/dL is associated with poor clinical outcomes including increased perioperative mortality, hospital length of stay, ICU length of stay and postoperative infection [28,29]. Most clinicians would agree that prevention of perioperative hyperglycemia is a desirable intervention, the optimal blood glucose range remains controversial due to the potential adverse events related to iatrogenic hypoglycaemia [30]. Clinical trials in adult surgical patients illustrate this paradigm of outcomes with intensive insulin therapy (IIT) defined as 140 to 200 mg/dL by the American College of Physicians. The Leuven surgical trial randomly assigned patients to IIT or conventional glucose management with a decrease in mean blood glucose levels and ICU mortality in the IIT group [31]. However, hypoglycemia was more frequent in the IIT group. These findings were supported by meta-analytical data from 5 randomized trials, which compared IIT to less stringent glycemic control and demonstrated significantly lower mortality [32]. In contrast, the Normoglycemic in Intensive Care Evaluation Survival Glucose Algorithm Regulation (NICE-SUGAR) trial demonstrated a higher rate of severe hypoglycemia and higher 90-day mortality in those patients who received IIT compared to conventional glucose management [33]. Hypoglycemia is the most common adverse effect of IIT and can lead to unwanted morbidity such as seizures, brain damage, and cardiac arrhythmias. As a result, more liberal blood glucose targets of 180 to 200 mg/dL are typically recommended in effort to prevent significant hyperglycemia whilst avoiding iatrogenic hypoglycemia [34–36].

The surgical stress response triggers a cascade of sympathetic nervous system and endocrine responses that include activation of the HPA axis and increased cortisol secretion, which leads to a net increase in peripheral insulin resistance [37]. Traditional perioperative interventions such as mechanical bowel preparation, pre-operative fasting, and slow resumption of normal diet all contribute to the relative insulin resistant state noted perioperatively and have been shown to correlate with perioperative complications and increased length of hospital stay [38]. Several elements of enhanced recovery protocols abrogate postoperative insulin resistance and thereby result in lower perioperative

glucose levels without resulting hypoglycemia. Examples of key elements include avoidance of oral mechanical preoperative bowel preparation and avoidance of preoperative fasting until 2 h prior to surgery, pre-operative carbohydrate loading along with stimulation of gut function by early resumption of postoperative oral intake and optimal fluid balance [1].

8.1. Summary and recommendation

ERAS elements that reduce metabolic stress should be employed to reduce insulin resistance and the development of hyperglycemia. Perioperative maintenance of blood glucose levels (<180–200 mg/dL) results in improved perioperative outcomes. Glucose levels above this range should be treated with insulin infusions and regular blood glucose monitoring to avoid the risk of hypoglycemia.

8.2. Evidence level

Use of stress reducing elements: High.

Treating hyperglycemia above 180–200 mg/dL: High.

8.3. Recommendation grade

Strong.

9. Postoperative analgesia

Pain following gynecological abdominal surgery can be severe [39]. Uncontrolled acute post-operative pain is associated with dissatisfaction [40], post-operative complications, and is a strong risk factor for development of chronic pain [41]. Morphine is commonly used to control post-operative pain but is associated with nausea, sedation [42], fatigue [43] and poorer quality of recovery [44] and may prolong time to mobilization. Opioid analgesics also contribute to the development of ileus. Therefore an enhanced recovery pathway for gynecological surgery must employ a strategy to effectively control post-operative pain and allow attainment of other ERAS targets such as early mobilization and return to oral diet whilst reducing the need for opiates. Many RCTs in the last 20 years in open surgery have focused on epidural analgesia, which can offer excellent analgesia, reduction in the surgical stress response, and earlier return of gut function. However the role of epidural analgesia is now a matter of debate. With the increasing uptake of laparoscopic and robotic assisted surgery the magnitude and duration of visceral and wound pain have been markedly reduced such that good post-operative pain control is achievable by many different analgesic techniques, often used in combination to tackle both the visceral and wound elements. The literature base is developing rapidly and may well impact on future recommendations.

9.1. Multimodal analgesia

The concept of achieving analgesia through the additive or synergistic effects of different types of analgesics is not new [45]. Non-steroidal anti-inflammatory drugs (NSAIDs) have been extensively investigated, both as part of a multi-modal analgesic regime as well as for gynecologic surgery, and are effective at reducing pain and opioid consumption and improving patient satisfaction [46,47], and a combination of NSAID and acetaminophen is more effective than either drug alone [48]. Both should be administered regularly unless contraindication exists. Gabapentin has recently become popular for treatment of post-operative pain. A recent systematic review found that pre-emptive administration of gabapentin for abdominal hysterectomy was effective in reducing post-operative pain, opioid consumption and side effects [49] and has been used in one gynecologic enhanced recovery program [21]. However, studies have not yet identified the optimal dose, or timing of administration. Dexamethasone appears to have analgesic

effects [50,51], as well as preventing post-operative nausea and vomiting, so may be useful as part of an ERP for gynecologic surgery. However its analgesic effects are yet to be fully investigated and it may cause transient post-operative hyperglycemia. Chronic administration of steroids are known to impair wound healing, although this has not been demonstrated following administration of a course <10 days duration [52]. Intravenous lidocaine is gaining popularity as an analgesic adjunct in abdominal surgery. A Cochrane analysis concluded that there was low quality evidence of an early reduction in pain, and opioid consumption and time to bowel recovery were reduced, though the effect on these outcomes was small [26].

9.1.1. Summary and recommendations

A multimodal analgesia strategy should be employed with the aim of reducing post-operative opioid requirement. Post-operatively, opioids should be given orally to patients who can tolerate diet. For patients unable to tolerate diet following surgery, then an opioid IV PCA can be used until resumption of GI function, but the oral route should be used as soon as possible.

Acetaminophen and NSAIDs in combination should be administered regularly to all patients unless contraindication exists.

Dexamethasone may be administered to prevent PONV and reduce pain, but should be used with caution in diabetic patients.

Gabapentin may reduce pain and side effects and may be considered, although the optimal dose is not known.

9.1.2. Level of evidence

Use of multimodal analgesia: High.

Combination of acetaminophen and NSAIDs: High.

Gabapentin: Moderate.

Dexamethasone as an analgesic: Low.

9.1.3. Recommendation grade

Strong.

9.2. Analgesia for vaginal hysterectomy

Few rigorous studies have been performed investigating analgesia in patients undergoing vaginal hysterectomy. In one study, intra-operative paracervical nerve block appeared to reduce post-operative pain and morphine consumption, and although the analgesic benefit appears to be limited to the first few hours after surgery, patients mobilized more quickly [53]. However a Cochrane review concluded that paracervical nerve block was ineffective for cervical dilatation [54]. One study investigated high-volume local anesthetic infiltration of the surrounding tissues and found that although the analgesic benefit was limited to the first four hours post-operatively, patients used less opioid analgesics and mobilized earlier [55]. In another study, spinal anesthesia with intrathecal morphine and clonidine also reduced early post-operative pain and morphine consumption, though the effect was modest [56]. Both spinal morphine and paracervical nerve block have been used to facilitate early discharge in enhanced recovery vaginal hysterectomy pathways [57,58].

9.2.1. Summary and recommendations

Local anesthetic infiltration may be effective at reducing early post-operative pain and opioid consumption, and facilitating early mobilization. Either paracervical nerve block or intrathecal morphine may reduce pain and opioid consumption after vaginal hysterectomy. However, the effect is small.

9.2.2. Evidence level

Low.

9.2.3. Recommendation grade

Weak.

9.3. Analgesia for open general gynecologic surgery

The optimal analgesic regimen for open gynecologic surgery is currently a subject of debate. Thoracic epidural analgesia (TEA) has gained widespread acceptance in providing post-operative analgesia for major abdominal surgery [2], and has been shown to be superior to intravenous PCA [59]. TEA is effective in attenuating the surgical stress response and reducing pain and opioid consumption for up to 72 h [59, 60] following abdominal incisions, and also has an impact on complications following abdominal surgery, reducing the time to bowel recovery by up to 36 h and may reduce cardiac and respiratory complications in high risk patients. TEA has been shown to effectively reduce pain following abdominal hysterectomy [61] and gynecologic cancer surgery [62], and reduces time to return of gut function. However the role of TEA in enhanced recovery surgery is now less clear: whilst analgesia and recovery seem to be better with TEA than IV PCA, epidural failure rates may be as high as 30% [63], and many of these patients will require supplemental opiates. Even if patients are normovolemic the sympathetic block that results from TEA may result in hypotension that may require treatment with vasopressors [64]. Patients who undergo abdominal hysterectomy in ERAS protocols can target a length of hospital stay of 1–2 days, in which case TEA may hinder achievement of other ERAS goals such as mobilization [65] and removal of urinary catheter, and TEA has been shown to increase hospital stay and complication rates in gynaecologic cancer surgery [66].

Where TEA is to be used local anesthetic should be of low concentration, and should be combined with an opioid such as fentanyl. Post-operative hypotension may require treatment with vasopressors. Some consideration must be made to the impact on early ERAS goals and how they will be achieved, and expert post-operative input may be required to ensure reliable analgesia. Although epidurals sited in the lumbar spine have an evidence base for analgesic benefit in gynecologic surgery, thoracic epidurals are preferred: epidurals should be sited at the level appropriate for innervation of the surgical area, and thoracic epidurals are likely to cause less hypotension [67] and motor block [68] than those in the lumbar segments.

An alternative to TEA is spinal anesthesia with low-dose intrathecal morphine (ITM). As a single injection, this has benefits over TEA in allowing early mobilization and removal of urinary catheter as well as facilitating early discharge from hospital [69]. When compared to a general anesthetic without neuraxial block, spinal anesthesia with ITM significantly reduces pain and morphine consumption both for hysterectomy [39,44,70] and this analgesic benefit may persist for up to 48 h post-operatively [39,44]. The added benefit of reduced morphine consumption is the reduced risk of post-operative ileus. Additionally ITM appears to reduce peri-operative stress hormone release [71], improve post-operative recovery [44,72] and reduce post-operative drowsiness and fatigue, though at the expense of increased pruritus [73]. Most studies do not indicate an increase in vomiting with low-dose ITM when compared with IV PCA [39,74]. Dose-finding studies appear to show a ceiling of effect at 200 mcg [74] and doses of ITM within this range do not appear to increase the risk of respiratory depression [75]. Spinal anesthesia without long-acting opioids does not improve post-operative pain when compared to general anesthesia [76].

Experience dictates that to improve patient acceptability of spinal anesthesia with ITM, general anesthesia may need to be offered in addition, in which case the dose of intrathecal local anesthetic should be reduced to avoid intra-operative hypotension and intraoperative narcotics should be minimized to reduce side effects.

Where patients have undergone general anesthesia without neuraxial blockade, truncal nerve blocks may serve to reduce pain and reduce post-operative morphine requirement. Transversus abdominis plane (TAP) blocks involve the injection of a large volume of local anesthetic in between the muscle layers of the trunk, and may now be performed under ultrasound guidance to ensure accurate delivery of local anesthetic. This technique has been shown to be efficacious for

abdominal incisions [77], including abdominal hysterectomy [78], and a meta-analysis concluded that TAP blocks reduce pain and morphine requirement up to 24 h after open gynecologic surgery [79]. In patients undergoing Cesarean section, TAP blocks appear to be less effective than intrathecal morphine at controlling post-operative pain, although side effects were fewer than ITM [80] and TAP blocks do not appear to add any analgesic benefit when used in combination with ITM [81]. Bilateral ilioinguinal nerve blocks also appear to reduce post-operative morphine requirement, however may not reduce morphine-related side effects [82].

Wound infiltration with local anesthetic is safe and easy to perform, though any effect on post-operative pain and opioid consumption is modest and short-lived [83]. Prolongation of this analgesic effect may be achieved through insertion of sub-cutaneous wound catheters [84]. A meta-analysis concluded that continuous wound infiltration (CWI) reduced pain and opioid consumption and improved recovery after major abdominal surgery [85], and may provide analgesia equivalent to TEA for abdominal surgery [86]. Another study found that, when compared to TEA for open colorectal surgery, CWI reduced opioid usage, vomiting and time to bowel recovery, and improved patient satisfaction [87]. However for gynecologic surgery the data is less clear, and although CWI has been shown to improve analgesia, reduce opioid requirements and reduce time to return of gut function [88] a number of studies have either only demonstrated benefit in the first few hours after surgery [84], or failed to demonstrate benefit at all [89–91]. There is lack of agreement concerning ideal catheter placement [92, 93], though in most studies the infusion catheter was placed below the abdominal fascia. The impact of continuous wound infiltration on wound healing has not been fully studied, though existing data has not shown an increase in wound complication rates. More research with this technique is required in this patient group before any conclusions may be drawn.

Intraperitoneal local anesthetic (IPLA) has been utilized to reduce post-operative pain, and one trial demonstrated reduced opioid consumption and improved surgical recovery score when used alongside TEA following colorectal surgery [94]. A systematic review of other trials concluded that IPLA reduces post-operative pain but not opioid consumption, and recovery parameters were unchanged [95]. IPLA has also been tested for open hysterectomy and has been found to reduce post-operative pain [96] and morphine consumption [97], however the benefit was limited to the first few hours after surgery, and analgesia from IPLA does not seem to be dose-responsive [98].

9.3.1. Summary and recommendations

For open surgery a multimodal, opiate sparing analgesic strategy should be utilized. TEA or spinal anesthesia with intrathecal morphine may improve recovery parameters and are recommended. However TEA may increase time to mobilization and removal of urinary catheter, and may potentially impact on hospital stay.

Where patients have undergone general anesthesia without neuraxial blockade, a truncal block, such as TAP blocks, may reduce pain and opioid consumption for up to 24 h and should be employed. Continuous wound infiltration or intraperitoneal instillation of local anesthetic may improve recovery for colorectal surgery and may be considered as an alternative to TAP blocks or TEA, however the evidence of benefit in gynecologic surgery is lacking.

Post-operatively, multimodal analgesia should be used. Systemic opioids may be given either orally or by intravenous PCA. The IV PCA should be discontinued when normal gut function resumes.

9.3.2. Evidence level

Intrathecal morphine: Moderate.
Thoracic epidural analgesia: High.
TAP blocks: Moderate.
CWI: Moderate.

9.3.3. Recommendation grade

Strong.

9.4. Analgesia for major oncologic surgery

In patients undergoing cytoreductive surgery, the large surgical area and complex patient pain history means that post-operative pain is often severe. TEA is widely used, and was associated with superior pain control at rest and on movement for the first 3 post-operative days in one observational study [99], and a randomized controlled study [100] found improved pain control on coughing for the first 3 post-operative days. However other investigators found no benefit in pain, bowel recovery or length of stay in patients with peri-operative TEA, and an increase in vasopressor requirement [101]. In patients undergoing heated intraperitoneal chemotherapy (HIPEC), the use of TEA is controversial. TEA may reduce opioid consumption and reduce time to extubation [102] although an IV PCA is often required in addition to TEA to achieve adequate analgesia [103]. HIPEC may be associated with a post-operative coagulopathy that may complicate removal of the epidural catheter, however in one study removal of epidural catheter was delayed in only 0.5% of cases [104]. Some centers use intravenous PCA in preference to TEA, citing adequate analgesia with fewer hemodynamic effects [105]. TAP blocks were examined in one retrospective study and appeared to reduce opioid consumption on post-operative day 1 only [106].

9.4.1. Summary and recommendations

TEA is effective in reducing post-operative pain after gynecologic laparotomy. However TEA may not improve other post-operative outcomes and patients may require additional IV opioids in addition to TEA to achieve adequate analgesia. TEA may compound hypotension that requires vasopressor support. Intravenous PCA appears to be a suitable alternative.

9.4.2. Evidence level

Low.

9.4.3. Recommendation grade

Weak.

9.5. Analgesia for laparoscopic gynecologic/oncology surgery

A meta-analysis examining TAP blocks for laparoscopic surgery across a range of abdominal procedures [107] found only pain at rest, and not dynamic pain, was reduced. For laparoscopic hysterectomy, one trial showed that TAP blocks improved post-operative quality of recovery (QoR40) score [108] however 3 further trials did not show benefit [109–111]. Intra-peritoneally administered local anesthetic has been used successfully for minor gynecologic laparoscopic procedures but this technique does not seem to be effective for major gynecologic laparoscopic surgery, either by single instillation or continuous infusion [112–116]. Additionally, a Cochrane analysis of intraperitoneal local anesthetic for laparoscopic cholecystectomy found low quality evidence of benefit, though the effect was likely to be clinically insignificant [117]. ITM showed a small benefit in robotic surgery [118]. TEA has been investigated for laparoscopic colorectal surgery and appeared to prolong hospital stay without improving patient outcomes [69].

9.5.1. Summary and recommendations

For laparoscopic gynecologic/oncology surgery, neither TAP blocks nor intraperitoneal instillation of local anesthetic are recommended on the current level of evidence. For laparoscopic abdominal surgery, TEA may prolong hospital stay without improving outcomes. Multimodal analgesia should be employed, and post-operative opioids may be given either orally or by IV PCA depending on magnitude of surgery and predicted post-operative gut function.

9.5.2. Level of evidence

Low.

9.5.3. Recommendation grade

Weak.

10. Peritoneal drainage

Peritoneal drainage has traditionally been used to prevent accumulation of fluid in the bed of dissection, to evacuate blood, serous collections, or infection, and in colorectal surgery it has been thought to prevent anastomotic leakage. However, peritoneal drainage has not been shown to prevent anastomotic leaks or improve overall outcome, and is not recommended routinely after either colonic or rectal surgery [119–121]. There is little research regarding drains after colonic or rectal anastomosis in gynecologic oncology surgery [122]. It is difficult to extrapolate the results from the colorectal literature directly to all gynecological surgery. For patients with metastatic ovarian cancer, the scope of surgery is larger, encompassing other organ resections, most of the peritoneal surfaces, and the risk factor profile for postoperative morbidity is elevated with poor nutritional status, ascites, peritoneal carcinomatosis, extended operative times, and cytotoxic therapy. Regardless, the rate of anastomotic leakage in ovarian cancer surgery in the literature ranges from 1 to 7%, in the range found in colorectal surgery [122–124]. In summary, we did not find evidence that drainage gives better outcomes after gynecological surgery. Furthermore, a Cochrane systematic review including 4 studies with 571 participants [125] concluded that drains did not prevent lymphocysts, but were rather associated with a higher risk of cyst formation after pelvic lymphadenectomy. Fewer studies have investigated para-aortic lymphadenectomy, but no evidence exists to recommend drainage [126]. Urological surgical techniques are frequently employed during major gynecologic oncology cases. Historically drains have been placed at the site of bladder resection/reconstruction, ureteral reimplantation, and urinary diversion (ileal conduit, continent reservoir) with the aim of identifying early urine leaks. Nevertheless, there are no specific studies that have evaluated the use of drains in such surgeries in our patient population. Looking to the urological literature, it should be noted that the ERAS Radical Cystectomy guideline found no evidence to support or refute the use of drains in this setting and as such further research is required in this area [5].

10.1.1. Summary and recommendation

Peritoneal drainage is not recommended routinely in gynecologic/oncology surgery including for patients undergoing lymphadenectomy or bowel surgery.

10.1.2. Evidence level

Moderate.

10.1.3. Recommendation grade

Strong.

11. Urinary drainage

The primary indications for postoperative bladder drainage are to monitor urine output and prevent urinary retention. However, there is considerable variation in the method and duration of bladder drainage following surgery for gynecological cancers. In addition, there is a high incidence of bladder related morbidity to the lower genital tract following such surgery, which may include effects on urinary voiding and bladder capacity [127].

A review of policies for removal of short-term urinary catheters identified only a small number of studies including patients undergoing

gynecologic surgery [128]. When comparing the timing of removal of the catheter, time to first voiding was longer, but larger volumes of urine were passed following midnight removal compared to early morning. In one study, midnight removal of catheters was also associated with significantly shorter length of stay [129]. A recent single center study following uncomplicated total abdominal hysterectomy compared removal of urethral catheters immediately after surgery, 6 h, or 24 h postoperatively. The intermediate group had fewer re-catheterizations compared to the immediate removal group, and lower rates of urinary tract infection than the prolonged users [130]. These findings are supported by a recent review [131]. In the same review, a greater number of patients required re-catheterization following a urethral compared to a suprapubic catheter. Two small studies focusing on patients undergoing radical hysterectomy for cervical cancer showed the suprapubic route to be associated with fewer bladder infections [132,133]. In one of these studies intermittent self-catheterization was associated with a higher infection rate but patients found the technique to be catheterization [133].

11.1.1. Summary and recommendation

Urinary catheters should be used for postoperative bladder drainage for a short period preferably <24 h postoperatively.

11.1.2. Evidence level

Low.

11.1.3. Recommendation grade

Strong.

12. Early mobilization

There are multiple hypothesized benefits to early mobilization, including a reduction in pulmonary complications, decreased insulin resistance, less muscle atrophy, and reduced length of hospital stay [134, 135]. Early mobilization has been shown to be an integral part of systematic efforts to reduce venous thromboembolic complications in the surgical patient [136]. Foley catheters, poor pain control, and IV poles, have been identified by gynecologic surgical patients as barriers to ambulation [137]. Therefore, compliance with other aspects of enhanced recovery protocols may improve early mobilization by limiting these barriers [138]. A care plan listing daily mobilization goals and patient engagement with an activity diary may be helpful [135,139].

12.1.1. Summary and recommendation

Patients should be encouraged to mobilize within 24 h of surgery.

12.1.2. Evidence level

Low.

12.1.3. Recommendation grade

Strong.

13. Discussion

This guideline outlines the recommendations of the ERAS® Group for the postoperative management of patients undergoing gynecologic/oncology surgery, and is based on the best available evidence. As was the case in Part I [1], in some instances good quality data was not available. This was particularly true for the evidence surrounding urinary drainage, early mobilization and postoperative analgesia in which the

optimal analgesic regimen for vaginal surgery/MIS and open gynecologic surgery is currently a subject of debate. In some instances recommendations were made based on findings from other surgical disciplines in which major abdominal surgery is routinely utilized.

We are hopeful that these gynecologic/oncology ERAS® guidelines will help integrate existing knowledge into practice, align perioperative care, and encourage future investigations to address existing knowledge gaps. Measuring compliance has proven to be a key factor required for success and sustainability of ERAS® protocols [140]. A process is currently underway whereby the gynecologic/oncology guidelines are being translated into their corresponding audit system (ERAS Interactive Audit System, EIAS) which will help to ensure compliance [141] and allow surgeons/clinicians to improve the care delivered to our patient population.

Conflict of interest statement

Dr. Acheson reports personal fees from Baxter UK Ltd., outside the submitted work. In addition, Dr. Acheson has a commercial (future royalties on a medical device in development) relationship with Mediplus Ltd. He has also held the following appointments: Joint National Clinical Advisor (Gynaecology) to the Enhanced Recovery Partnership Programme, Department of Health (2010–2011), and continued under NHS Improvement (2011–2013); Member of Steering Board, Enhanced Recovery ERAS (UK) (2011–2013).

Dr. Scott received honoraria for lecturing and travel expenses from Baxter Healthcare, Merck, and Deltex. He is an Executive Committee member of the ERAS Society.

Dr. Ljungqvist has an appointment with Nutricia Advisory Board, has received speakers honoraria from Nutricia, MSD, B Braun and Fresenius-Kabi. He is the current Chairman of the ERAS Society (www.erasociety.org). He founded, serves on the Board and owns stock in Encare AB that runs the ERAS Society Interactive Audit System (EIAS).

References

- G. Nelson, A. Altman, A. Nick, L. Meyer, P.T. Ramirez, C. Achari, et al., Guidelines for pre- and intra-operative care in gynecologic/oncology surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations, *Gynecol. Oncol.* (2015) <http://dx.doi.org/10.1016/j.ygyno.2015.11.015>.
- U.O. Gustafsson, M.J. Scott, W. Schwenk, N. Demartines, D. Roulin, N. Francis, et al., Guidelines for perioperative care in elective colonic surgery: Enhanced Recovery After Surgery (ERAS) Society recommendations, *World J. Surg.* 37 (2) (2012) 259–284.
- J. Nygren, J. Thacker, F. Carli, K.C. Fearon, S. Norderval, D.N. Lobo, et al., Guidelines for perioperative care in elective rectal/pelvic surgery: Enhanced Recovery After Surgery (ERAS) Society recommendations, *World J. Surg.* 37 (2) (Feb 2013) 285–305.
- G.H. Guyatt, A.D. Oxman, G.E. Vist, R. Kunz, Y. Falck-Ytter, P. Alonso-Coello, et al., GRADE: an emerging consensus on rating quality of evidence and strength of recommendations, *BMJ* 336 (7650) (2008) 924–926.
- Y. Cerantola, M. Valerio, B. Persson, P. Jichlinski, O. Ljungqvist, M. Hubner, et al., Guidelines for perioperative care after radical cystectomy for bladder cancer: Enhanced Recovery After Surgery (ERAS®) Society recommendations, *Clin. Nutr.* 32 (6) (Dec 2013) 879–887.
- M. Brunetti, I. Shemilt, S. Pregno, L. Vale, A.D. Oxman, J. Lord, et al., GRADE guidelines: 10. Considering resource use and rating the quality of economic evidence, *J. Clin. Epidemiol.* 66 (2) (Feb 2013) 140–150.
- D.L. Clarke-Pearson, L.S. Synan, R. Dodge, J.T. Soper, A. Berchuck, R.E. Coleman, A randomized trial of low-dose heparin and intermittent pneumatic calf compression for the prevention of deep venous thrombosis after gynecologic oncology surgery, *Am. J. Obstet. Gynecol.* 168 (4) (1993) 1146–1153 (discussion 1153–4).
- G.L. Maxwell, L. Synan, R. Dodge, B. Carroll, D.L. Clarke-Pearson, Pneumatic compression versus low molecular weight heparin in gynecologic oncology surgery: a randomized trial, *Obstet. Gynecol.* 98 (6) (2001) 989–995.
- M.H. Einstein, D.M. Kushner, J.P. Connor, A.A. Bohl, T.J. Best, M.D. Evans, et al., A protocol of dual prophylaxis for venous thromboembolism prevention in gynecologic cancer patients, *Obstet. Gynecol.* 112 (5) (2008) 1091–1097.
- A. Sachdeva, M. Dalton, S.V. Amaragiri, T. Lees, Graduated compression stockings for prevention of deep vein thrombosis, *Cochrane Database Syst. Rev.* 12 (2014), CD001484.
- G. Agnelli, G. Bolis, L. Capussotti, R.M. Scarpa, F. Tonelli, E. Bonizzoni, et al., A clinical outcome-based prospective study on venous thromboembolism after cancer surgery: the @RISTOS project, *Ann. Surg.* 243 (1) (2006) 89–95.
- M.K. Gould, D.A. Garcia, S.M. Wren, P.J. Karanicolas, J.I. Arcelus, J.A. Heit, et al., Prevention of VTE in nonorthopedic surgical patients: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines, *Chest* 141 (Suppl. 2) (2012) e227S–e277S.
- M.S. Rasmussen, L.N. Jorgensen, P. Wille-Jorgensen, Prolonged thromboprophylaxis with low molecular weight heparin for abdominal or pelvic surgery, *Cochrane Database Syst. Rev.* (1) (2009) CD004318 (doi:10.1002/CD004318).
- P.T. Ramirez, A.M. Nick, M. Frumovitz, K.M. Schmeler, Venous thromboembolic events in minimally invasive gynecologic surgery, *J. Minim. Invasive Gynecol.* 20 (6) (2013) 766–769.
- National Clinical Guideline Centre (UK), *Intravenous Fluid Therapy: Intravenous Fluid Therapy in Adults in Hospital*, Royal College of Physicians (UK), London, 2013.
- A. Perner, N. Haase, A.B. Guttormsen, J. Tenhunen, G. Klemenzson, A. Aneman, et al., Hydroxyethyl starch 130/0.42 versus Ringer's acetate in severe sepsis, *N. Engl. J. Med.* 367 (2) (2012 Jul 12) 124–134.
- K. Charoenkwan, G. Phillipson, T. Vutyavanich, Early versus delayed (traditional) oral fluids and food for reducing complications after major abdominal gynaecologic surgery, *Cochrane Database Syst. Rev.* (2007), CD004508.
- L. Minig, R. Biffi, V. Zanagnolo, A. Attanasio, C. Beltrami, L. Bocciolone, et al., Early oral versus “traditional” postoperative feeding in gynecologic oncology patients undergoing intestinal resection: a randomized controlled trial, *Ann. Surg. Oncol.* 16 (2009) 1660–1668.
- L. Minig, R. Biffi, V. Zanagnolo, A. Attanasio, C. Beltrami, L. Bocciolone, et al., Reduction of postoperative complication rate with the use of early oral feeding in gynecologic oncology patients undergoing a major surgery: a randomized controlled trial, *Ann. Surg. Oncol.* 16 (2009) 3101–3110.
- J.M. Schilder, J.A. Hurteau, K.Y. Look, D.H. Moore, G. Raff, F.B. Stehman, et al., A prospective controlled trial of early postoperative oral intake following major abdominal gynecologic surgery, *Gynecol. Oncol.* 67 (1997) 235–240.
- E. Kalogera, J.N. Bakkum-Gamez, C.J. Jankowski, E. Trabuco, J.K. Lovely, S. Dhanorker, et al., Enhanced recovery in gynecologic surgery, *Obstet. Gynecol.* 122 (2 Pt 1) (2013) 319–328.
- J. Fanning, S. Yu-Brekke, Prospective trial of aggressive postoperative bowel stimulation following radical hysterectomy, *Gynecol. Oncol.* 73 (1999) 412–414.
- P.O. Hendry, R.M. van Dam, S.F. Bukkems, D.W. McKeown, R.W. Parks, T. Preston, et al., Randomized clinical trial of laxatives and oral nutritional supplements within an enhanced recovery after surgery protocol following liver resection, *Br. J. Surg.* 97 (2010) 1198–1206.
- U. Traut, L. Brugger, R. Kunz, C. Pauli-Magnus, K. Haug, H.C. Bucher, et al., Systemic prokinetic pharmacologic treatment for postoperative adynamic ileus following abdominal surgery in adults, *Cochrane Database Syst. Rev.* (2008), CD004930.
- I.E. Ertas, K. Gungorduk, A. Ozdemir, U. Solmaz, A. Dogan, Y. Yildirim, Influence of gum chewing on postoperative bowel activity after complete staging surgery for gynecological malignancies: a randomized controlled trial, *Gynecol. Oncol.* 131 (1) (2013 Oct) 118–122.
- P. Kranke, J. Jokinen, N.L. Pace, A. Schnabel, M.W. Hollmann, K. Hahnenkamp, et al., Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery, *Cochrane Database Syst. Rev.* 7 (Jul 16 2015) CD009642.
- P.J. Karanicolas, S.E. Smith, B. Kanbur, E. Davies, G.H. Guyatt, The impact of prophylactic dexamethasone on nausea and vomiting after laparoscopic cholecystectomy: a systematic review and meta-analysis, *Ann. Surg.* 248 (2008) 751–762.
- R.P. Kiran, M. Turina, J. Hammel, V. Fazio, The clinical significance of an elevated postoperative glucose value in nondiabetic patients after colorectal surgery: evidence for the need for tight glucose control? *Ann. Surg.* 258 (4) (2013) 599–604.
- M. Ramos, Z. Khalpey, S. Lipsitz, J. Steinberg, M.T. Panizales, M. Zinner, et al., Relationship of perioperative hyperglycemia and postoperative infections in patients who undergo general and vascular surgery, *Ann. Surg.* 248 (4) (2008) 585–591.
- A. Qaseem, L.L. Humphrey, R. Chou, V. Snow, S. P. Clinical Guidelines Committee of the American College of Physicians. Use of intensive insulin therapy for the management of glycemic control in hospitalized patients: a clinical practice guideline from the American College of Physicians, *Ann. Intern. Med.* 154 (4) (2011) 260–267.
- G. van den Berghe, P. Wouters, F. Weekers, C. Verwaest, F. Bruyninckx, M. Schetz, et al., Intensive insulin therapy in critically ill patients, *N. Engl. J. Med.* 345 (19) (2001) 1359–1367.
- D.E. Griesdale, R.J. de Souza, R.M. van Dam, D.K. Heyland, D.J. Cook, A. Malhotra, et al., Intensive insulin therapy and mortality among critically ill patients: a meta-analysis including NICE-SUGAR study data, *CMAJ* 180 (8) (2009) 821–827.
- NICE-SUGAR Study Investigators, S. Finfer, D.R. Chittock, S.Y. Su, D. Blair, D. Foster, V. Dhingra, et al., Intensive versus conventional glucose control in critically ill patients, *N. Engl. J. Med.* 360 (13) (2009) 1283–1297.
- J.C. Preiser, P. Devos, S. Ruiz-Santana, C. Mélot, D. Annane, J. Groeneveld, et al., A prospective randomised multi-centre controlled trial on tight glucose control by intensive insulin therapy in adult intensive care units: the Glucontrol study, *Intensive Care Med.* 35 (10) (2009) 1738–1748.
- I. Vanhorebeek, L. Langouche, G. Van den Berghe, Tight blood glucose control with insulin in the ICU: facts and controversies, *Chest* 132 (1) (2007) 268–278.
- J.S. Kinsley, A. Grover, Severe hypoglycemia in critically ill patients: risk factors and outcomes, *Crit. Care Med.* 35 (10) (2007) 2262–2267.
- J.P. Desborough, The stress response to trauma and surgery, *Br. J. Anaesth.* 85 (1) (2000) 109–117.
- J. Nygren, The metabolic effects of fasting and surgery, *Best Pract. Res. Clin. Anaesthesiol.* 20 (3) (2006) 429–438.
- L. Massicotte, K.D. Chalaoui, D. Beaulieu, J.D. Roy, F. Bissonnette, Comparison of spinal anesthesia with general anesthesia on morphine requirement after abdominal hysterectomy, *Acta Anaesthesiol. Scand.* 53 (5) (2009) 641–647.
- P.S. Myles, B. Weitkamp, K. Jones, J. Melick, S. Hensen, Validity and reliability of a postoperative quality of recovery score: the QoR-40, *Br. J. Anaesth.* 84 (1) (2000) 11–15.
- W.A. Macrae, Chronic post-surgical pain: 10 years on, *Br. J. Anaesth.* 101 (1) (2008) 77–86.
- A. Woodhouse, L.E. Mather, The effect of duration of dose delivery with patient-controlled analgesia on the incidence of nausea and vomiting after hysterectomy, *Br. J. Clin. Pharmacol.* 45 (1) (1998) 57–62.

- [43] S.J. Dolin, J.N. Cashman, Tolerability of acute postoperative pain management: nausea, vomiting, sedation, pruritis, and urinary retention. Evidence from published data, *Br. J. Anaesth.* 95 (5) (2005) 584–591.
- [44] L.J. Catro-Alves, V.L. De Azevedo, T.F. De Freitas Braga, A.C. Goncalves, G.S. De Oliveira Jr., The effect of neuraxial versus general anesthesia techniques on postoperative quality of recovery and analgesia after abdominal hysterectomy, *Anesth. Analg.* 113 (6) (2011) 1480–1486.
- [45] A. Buvanendran, J.S. Kroin, Multimodal analgesia for controlling acute postoperative pain, *Curr. Opin. Anaesthesiol.* 22 (5) (2009) 588–593.
- [46] S. Niruthisard, T. Werawataganon, P. Bunburaphong, M. Ussawanophakiat, C. Wongsakornchaikul, K. Toleb, Improving the analgesic efficacy of intrathecal morphine with parecoxib after total abdominal hysterectomy, *Anesth. Analg.* 105 (3) (2007) 822–824.
- [47] A. Blackburn, J.D. Stevens, R.G. Wheatley, T.H. Madej, D. Hunter, Balanced analgesia with intravenous ketorolac and patient-controlled morphine following lower abdominal surgery, *J. Clin. Anesth.* 7 (2) (1995) 103–108.
- [48] C.K. Ong, R.A. Seymour, P. Lirk, A.F. Merry, Combining paracetamol (acetaminophen) with nonsteroidal antiinflammatory drugs: a qualitative systematic review of analgesic efficacy for acute postoperative pain, *Anesth. Analg.* 110 (4) (2010) 1170–1179.
- [49] N. Alayed, N. Alghanaim, X. Tan, T. Tulandi, Preemptive use of gabapentin in abdominal hysterectomy: a systematic review and meta-analysis, *Obstet. Gynecol.* 123 (6) (2014) 1221–1229.
- [50] T.H. Lunn, B.B. Kristensen, L.Ø. Andersen, H. Husted, K.S. Otte, L. Gaarn-Larsen, et al., Effect of high-dose preoperative methylprednisolone on pain and recovery after total knee arthroplasty: a randomized, placebo-controlled trial, *Br. J. Anaesth.* 106 (2) (2011) 230–238.
- [51] G.S. De Oliveira Jr., S. Ahmad, P.C. Fitzgerald, R.J. Marcus, C.S. Altman, A.S. Panjwani, et al., Dose ranging study on the effect of preoperative dexamethasone on postoperative quality of recovery and opioid consumption after ambulatory gynaecological surgery, *Br. J. Anaesth.* 107 (3) (2011) 362–371.
- [52] A.S. Wang, E.J. Armstrong, A.W. Armstrong, Corticosteroids and wound healing: clinical considerations in the perioperative period, *Am. J. Surg.* 206 (3) (2013 Sep) 410–417.
- [53] A.M. Hristovska, B.B. Kristensen, M.A. Rasmussen, Y.H. Rasmussen, L.B. Elving, C.V. Nielsen, et al., Effect of systematic local infiltration analgesia on postoperative pain in vaginal hysterectomy: a randomized, placebo-controlled trial, *Acta Obstet. Gynecol. Scand.* 93 (3) (2014) 233–238.
- [54] T. Tangsirawatthana, U.S. Sangkomkhamhang, P. Lumbiganon, M. Laopaiboon, Paracervical local anaesthesia for cervical dilatation and uterine intervention, *Cochrane Database Syst. Rev.* 9 (Sep 30 2013) CD005056.
- [55] B.B. Kristensen, Y.H. Rasmussen, M. Agerlin, M.W. Topp, M.O. Weincke, H. Kehlet, Local infiltration analgesia in urogenital prolapse surgery: a prospective randomized, double-blind, placebo-controlled study, *Acta Obstet. Gynecol. Scand.* 90 (10) (Oct 2011) 1121–1125.
- [56] J. Sprung, M.S. Sanders, M.E. Warner, J.B. Gebhart, C.R. Stanhope, C.J. Jankowski, et al., Pain relief and functional status after vaginal hysterectomy: intrathecal versus general anesthesia, *Can. J. Anesth.* 53 (7) (2006) 690–700.
- [57] R. Penketh, A. Griffiths, S. Chawathe, A prospective observational study of the safety and acceptability of vaginal hysterectomy performed in a 24-hour day case surgery setting, *BJOG* 114 (4) (2007) 430–436.
- [58] M. Ottesen, M. Sørensen, Y. Rasmussen, S. Smidt-Jensen, H. Kehlet, B. Ottesen, Fast track vaginal surgery, *Acta Obstet. Gynecol. Scand.* 81 (2) (2001) 138–146.
- [59] C.L. Wu, S.R. Cohen, J.M. Richman, A.J. Rowlingson, G.E. Courpas, K. Cheung, et al., Efficacy of postoperative patient-controlled and continuous infusion epidural analgesia versus intravenous patient-controlled analgesia with opioids: a meta-analysis, *Anesthesiology* 103 (5) (2005) 1079–1088.
- [60] F. Carli, N. Mayo, K. Klubien, T. Schrickler, J. Trudel, P. Belliveau, Epidural analgesia enhances functional exercise capacity and health-related quality of life after colonic surgery: results of a randomized trial, *Anesthesiology* 97 (3) (2002) 540–549.
- [61] H. Jørgensen, J.S. Fomsgaard, J. Dirks, J. Wetterslev, B. Andreasson, J.B. Dahl, Effect of peri- and postoperative epidural anaesthesia on pain and gastrointestinal function after abdominal hysterectomy, *Br. J. Anaesth.* 87 (4) (Oct 2001) 577–583.
- [62] S.E.I. Ferguson, T. Malhotra, V.E. Seshan, D.A. Levine, Y. Sonoda, C. DS, et al., A prospective randomized trial comparing patient-controlled epidural analgesia to patient-controlled intravenous analgesia on postoperative pain control and recovery after major open gynecologic cancer surgery, *Gynecol. Oncol.* 114 (1) (2009) 111–116.
- [63] L.B. Ready, Acute pain: lessons learned from 25,000 patients, *Reg. Anesth. Pain Med.* 24 (6) (Nov–Dec 1999) 499–505.
- [64] M. Hübner, C. Blanc, D. Roulin, M. Winiker, S. Gander, N. Demartines, Randomized clinical trial on epidural versus patient-controlled analgesia for laparoscopic colorectal surgery within an enhanced recovery pathway, *Ann. Surg.* 261 (4) (Apr 2015) 648–653.
- [65] L.M. Chen, V.K. Weinberg, C. Chen, C.B. Powell, L.L. Chen, J.K. Chan, et al., Perioperative outcomes comparing patient controlled epidural versus intravenous analgesia in gynecologic oncology surgery, *Gynecol. Oncol.* 115 (3) (2009) 357–361.
- [66] D. Belavy, M. Janda, J. Baker, A. Obermair, Epidural analgesia is associated with an increased incidence of postoperative complications in patients requiring an abdominal hysterectomy for early stage endometrial cancer, *Gynecol. Oncol.* 131 (2) (2013) 423–429.
- [67] G. Sagirolgu, B. Meydan, E. Copuroglu, A. Baysal, Y. Yoruk, Y. Altemur Karamustafaoglu, et al., A comparison of thoracic or lumbar patient-controlled epidural analgesia methods after thoracic surgery, *World J. Surg. Oncol.* 12 (May 4 2014) 96.
- [68] W.A. Visser, R.A. Lee, M.J. Gielen, Factors affecting the distribution of neural blockade by local anesthetics in epidural anesthesia and a comparison of lumbar versus thoracic epidural anesthesia, *Anesth. Analg.* 107 (2) (Aug 2008) 708–721.
- [69] B.F. Levy, M.J. Scott, W. Fawcett, C. Fry, T.A. Rockall, Randomized clinical trial of epidural, spinal or patient-controlled analgesia for patients undergoing laparoscopic colorectal surgery, *Br. J. Surg.* 98 (8) (2011) 1068–1078.
- [70] N. Borendal Wodlin, L. Nilsson, P. Kjølhede, for the 'GASPI' Study Group, The impact of mode of anaesthesia on postoperative recovery from fast-track abdominal hysterectomy: a randomised clinical trial, *BJOG* 118 (3) (2010) 299–308.
- [71] S. Karaman, S. Kocabas, M. Uyar, C. Zincircioglu, V. Firat, Intrathecal morphine: effects on perioperative hemodynamics, postoperative analgesia, and stress response for total abdominal hysterectomy, *Adv. Ther.* 23 (2) (2006) 295–306.
- [72] N.B. Wodlin, L. Nilsson, P. Kjølhede, Health-related quality of life and postoperative recovery in fast-track hysterectomy, *Acta Obstet. Gynecol. Scand.* 90 (4) (2011) 362–368.
- [73] N.B. Wodlin, L. Nilsson, K. Årestedt, P. Kjølhede, for the 'GASPI' Study Group, Mode of anesthesia and postoperative symptoms following abdominal hysterectomy in a fast-track setting, *Acta Obstet. Gynecol. Scand.* 90 (4) (2011) 369–379.
- [74] A. Hein, P. Rösblad, C. Gillis-Haegerstrand, K. Schedvins, J. Jakobsson, G. Dahlgren, Low dose intrathecal morphine effects on post-hysterectomy pain: a randomized placebo-controlled study, *Acta Anaesthesiol. Scand.* 56 (1) (2011) 102–109.
- [75] M. Gehling, M. Tryba, Risks and side-effects of intrathecal morphine combined with spinal anaesthesia: a meta-analysis, *Anaesthesia* 64 (6) (2009) 643–651.
- [76] M.J. Dakin, O.Y. Osinubi, F. Carli, Preoperative spinal bupivacaine does not reduce postoperative morphine requirement in women undergoing total abdominal hysterectomy, *Reg. Anesth. Pain Med.* 21 (2) (1996) 99–102.
- [77] P.L. Petersen, O. Mathiesen, H. Torup, J.B. Dahl, The transversus abdominis plane block: a valuable option for postoperative analgesia? A topical review, *Acta Anaesthesiol. Scand.* 54 (5) (May 2010) 529–535.
- [78] J. Carney, J.G. McDonnell, A. Ochana, R. Bhinder, J.G. Laffey, The transversus abdominis plane block provides effective postoperative analgesia in patients undergoing total abdominal hysterectomy, *Anesth. Analg.* 107 (6) (2008) 2056–2060.
- [79] R. Champaneria, L. Shah, J. Geoghegan, J.K. Gupta, J.P. Daniels, Analgesic effectiveness of transversus abdominis plane blocks after hysterectomy: a meta-analysis, *Eur. J. Obstet. Gynecol. Reprod. Biol.* 166 (1) (2013) 1–9.
- [80] B.M. Mishriky, R.B. George, H. AS, Transversus abdominis plane block for analgesia after Cesarean delivery: a systematic review and meta-analysis, *Can. J. Anaesth.* 59 (8) (Aug 2012) 766–778.
- [81] F.W. Abdallah, S.H. Halpern, C.B. Margarido, Transversus abdominis plane block for postoperative analgesia after Caesarean delivery performed under spinal anaesthesia? A systematic review and meta-analysis, *Br. J. Anaesth.* 109 (5) (2012) 679–687.
- [82] F. Oriola, Y. Toque, A. Mary, O. Gagneur, S. Beloucif, H. Dupont, Bilateral ilioinguinal nerve block decreases morphine consumption in female patients undergoing nonlaparoscopic gynecologic surgery, *Anesth. Analg.* 104 (3) (2007) 731–734.
- [83] L. Lowenstein, E.Z. Zimmer, M. Deutsch, Y. Paz, D. Yaniv, P. Jakobi, Preoperative analgesia with local lidocaine infiltration for abdominal hysterectomy pain management, *Eur. J. Obstet. Gynecol. Reprod. Biol.* 136 (2) (2008) 239–242.
- [84] E. Zohar, B. Fredman, A. Phillipov, R. Jedeikin, A. Shapiro, The analgesic efficacy of patient-controlled bupivacaine wound instillation after total abdominal hysterectomy with bilateral salpingo-oophorectomy, *Anesth. Analg.* 93 (2) (2001) 482–487.
- [85] S.S. Liu, J.M. Richman, R.C. Thirby, C.L. Wu, Efficacy of continuous wound catheters delivering local anesthetic for postoperative analgesia: a quantitative and qualitative systematic review of randomized controlled trials, *J. Am. Coll. Surg.* 203 (6) (Dec 2006) 914–932.
- [86] N.T. Ventham, M. Hughes, S. O'Neill, N. Johns, R.R. Brady, S.J. Wigmore, Systematic review and meta-analysis of continuous local anaesthetic wound infiltration versus epidural analgesia for postoperative pain following abdominal surgery, *Br. J. Surg.* 100 (10) (Sep 2013) 1280–1289.
- [87] S. Bertoglio, F. Fabiani, P.D. Negri, A. Corcione, D.F. Merlo, F. Cafiero, et al., The postoperative analgesic efficacy of preperitoneal continuous wound infusion compared to epidural continuous infusion with local anesthetics after colorectal cancer surgery: a randomized controlled multicenter study, *Anesth. Analg.* 115 (6) (Dec 2012) 1442–1450.
- [88] S. Maric, M. Banovic, K.S. Zdravcevic, Continuous wound infusion of levobupivacaine after total abdominal hysterectomy with bilateral salpingo-oophorectomy, *Period. Biol.* 111 (2) (2009) 299–302.
- [89] B.B. Kristensen, D.S. Christensen, M. Ostergaard, K. Skjelsager, D. Nielsen, T.S. Mogensen, Lack of postoperative pain relief after hysterectomy using preperitoneally administered bupivacaine, *Reg. Anesth. Pain Med.* 24 (6) (Nov–Dec 1999) 576–580.
- [90] W.M. Leong, W.K. Lo, J.W. Chiu, Analgesic efficacy of continuous delivery of bupivacaine by an elastomeric balloon infuser after abdominal hysterectomy: a prospective randomised controlled trial, *Aust N Z J Obstet Gynaecol.* 42 (5) (2002) 515–518.
- [91] D.M. Kushner, R. LaGalbo, J.P. Connor, R. Chappell, S.L. Stewart, E.M. Hartenbach, Use of a bupivacaine continuous wound infusion system in gynecologic oncology: a randomized trial, *Obstet. Gynecol.* 106 (2) (2005) 227–233.
- [92] T. Rackelboom, S. Le Strat, S. Silvera, T. Schmitz, A. Bassot, F. Goffinet, et al., Improving continuous wound infusion effectiveness for postoperative analgesia after cesarean delivery: a randomized controlled trial, *Obstet. Gynecol.* 116 (4) (Oct 2010) 893–900.
- [93] M.C. Hafizoglu, K. Katircioglu, M.Y. Ozkalkanli, S. Savaci, Bupivacaine infusion above or below the fascia for postoperative pain treatment after abdominal hysterectomy, *Anesth. Analg.* 107 (6) (2008) 2068–2072.

- [94] A. Kahohehr, T. Sammour, K. Zargar Shoshtari, M. Taylor, A.G. Hill, Intraperitoneal local anesthetic improves recovery after colon resection: a double-blinded randomized controlled trial, *Ann. Surg.* 254 (1) (Jul 2011) 28–38.
- [95] A. Kahohehr, T. Sammour, M. Soop, A.G. Hill, Intraperitoneal local anaesthetic in abdominal surgery – a systematic review, *ANZ J. Surg.* 81 (4) (Apr 2011) 237–245.
- [96] A. Gupta, A. Perniola, K. Axelsson, S.E. Thörn, K. Crafoord, N. Rawal, Postoperative pain after abdominal hysterectomy: a double-blind comparison between placebo and local anesthetic infused intraperitoneally, *Anesth. Analg.* (2004) 1173–1179.
- [97] A. Ng, A. Swami, G. Smith, A.C. Davidson, J. Emembolu, The analgesic effects of intraperitoneal and incisional bupivacaine with epinephrine after total abdominal hysterectomy, *Anesth. Analg.* 95 (1) (Jul 2002) 158–162.
- [98] A. Perniola, A. Gupta, K. Crafoord, B. Darvish, A. Magnuson, K. Axelsson, Intraabdominal local anaesthetics for postoperative pain relief following abdominal hysterectomy: a randomized, double-blind, dose-finding study, *Eur. J. Anaesthesiol.* 26 (5) (2002) 421–429.
- [99] M. Courtney-Brooks, K.C. Tanner Kurtz, E.B. Pelkofski, J. Nakayama, L.R. Duska, Continuous epidural infusion anesthesia and analgesia in gynecologic oncology patients: less pain, more gain? *Gynecol. Oncol.* 136 (1) (Jan 2015) 77–81.
- [100] S.E. Ferguson, T. Malhotra, V.E. Seshan, D.A. Levine, Y. Sonoda, D.S. Chi, et al., A prospective randomized trial comparing patient-controlled epidural analgesia to patient-controlled intravenous analgesia on postoperative pain control and recovery after major open gynecologic cancer surgery, *Gynecol. Oncol.* 114 (1) (Jul 2009) 111–116.
- [101] L.M. Chen, V.K. Weinberg, C. Chen, C.B. Powell, L.L. Chen, J.K. Chan, et al., Perioperative outcomes comparing patient controlled epidural versus intravenous analgesia in gynecologic oncology surgery, *Gynecol. Oncol.* 115 (3) (Dec 2009) 357–361.
- [102] C. Schmidt, T. Steinke, S. Moritz, M. Bucher, Thoracic epidural anesthesia in patients with cytoreductive surgery and HIPEC, *J. Surg. Oncol.* 102 (2010) 545–546.
- [103] J.C. Bell, B.G. Rylah, R.W. Chambers, H. Peet, F. Mohamed, B.J. Moran, Perioperative management of patients undergoing cytoreductive surgery combined with heated intraperitoneal chemotherapy for peritoneal surface malignancy: a multi-institutional experience, *Ann. Surg. Oncol.* 19 (13) (Dec 2012) 4244–4251.
- [104] P. Owusu-Agyemang, J. Soliz, A. Hayes-Jordan, N. Harun, V. Gottumukkala, Safety of epidural analgesia in the perioperative care of patients undergoing cytoreductive surgery with hyperthermic intraperitoneal chemotherapy, *Ann. Surg. Oncol.* 21 (5) (May 2014) 1487–1493.
- [105] G. Nelson, E. Kalogera, S.C. Dowdy, Enhanced recovery pathways in gynecologic oncology, *Gynecol. Oncol.* 135 (3) (Dec 2014) 586–594.
- [106] C. Rivard, E.L. Dickson, R.I. Vogel, P.A. Argenta, D. Teoh, The effect of anesthesia choice on post-operative outcomes in women undergoing exploratory laparotomy for a suspected gynecologic malignancy, *Gynecol. Oncol.* 133 (2) (May 2014) 278–282.
- [107] G.S. De Oliveira Jr., L.J. Castro-Alves, A. Nader, M.C. Kendall, R.J. McCarthy, Transversus abdominis plane block to ameliorate postoperative pain outcomes after laparoscopic surgery: a meta-analysis of randomized controlled trials, *Anesth. Analg.* 118 (2) (2014) 454–463.
- [108] G.S. De Oliveira Jr., M.P. Milad, P. Fitzgerald, R. Rahmani, R.J. McCarthy, Transversus abdominis plane infiltration and quality of recovery after laparoscopic hysterectomy: a randomized controlled trial, *Obstet. Gynecol.* 118 (6) (Dec 2011) 1230–1237.
- [109] G.A. Calle, C.C. López, E. Sánchez, J.F. De Los Ríos, E.M. Vázquez, E. Serna, et al., Transversus abdominis plane block after ambulatory total laparoscopic hysterectomy: randomized controlled trial, *Acta Obstet. Gynecol. Scand.* 93 (4) (2014) 345–350.
- [110] S.M. Kane, V. Garcia-Tomas, M. Alejandro-Rodriguez, B. Astley, R.R. Pollard, Randomized trial of transversus abdominis plane block at total laparoscopic hysterectomy: effect of regional analgesia on quality of recovery, *Am. J. Obstet. Gynecol.* 207 (5) (2012) 419 (e1–5).
- [111] L. El Hachem, E. Small, P. Chung, E.L. Moshier, K. Friedman, S.S. Fenske, et al., Randomized controlled double-blind trial of transversus abdominis plane block versus trocar site infiltration in gynecologic laparoscopy, *Am. J. Obstet. Gynecol.* 212 (2) (Feb 2015) 182 (e1–9).
- [112] V. Andrews, J.T. Wright, F. Zakaria, S. Banerjee, K. Ballard, Continuous infusion of local anaesthetic following laparoscopic hysterectomy – a randomised controlled trial, *BJOG* 121 (6) (2014) 754–760 (discussion 761).
- [113] H. Keita, J.L. Benifla, V. Le Bouar, R. Porcher, B. Wachowska, K. Bedairia, et al., Prophylactic ip injection of bupivacaine and/or morphine does not improve post-operative analgesia after laparoscopic gynecologic surgery, *Can. J. Anaesth.* 50 (4) (Apr 2003) 362–367.
- [114] I.C. Shaw, J. Stevens, S. Krishnamurthy, The influence of intraperitoneal bupivacaine on pain following major laparoscopic gynaecological procedures, *Anaesthesia* 56 (11) (Nov 2001) 1041–1044.
- [115] Y. Kaufman, I. Hirsch, L. Ostrovsky, O. Klein, I. Shnaider, E. Khoury, et al., Pain relief by continuous intraperitoneal nebulization of ropivacaine during gynecologic laparoscopic surgery – a randomized study and review of the literature, *J. Minim. Invasive Gynecol.* 15 (5) (Sep-Oct 2008) 554–558.
- [116] D. Arden, E. Seifert, N. Donnellan, R. Guido, T. Lee, S. Mansuria, Intraperitoneal instillation of bupivacaine for reduction of postoperative pain after laparoscopic hysterectomy: a double-blind randomized controlled trial, *J. Minim. Invasive Gynecol.* 20 (5) (Sep-Oct 2013) 620–626.
- [117] K.S. Gurusamy, M. Nagendran, G.P. Guerrini, C.D. Toon, M. Zinnuroglu, B.R. Davidson, Intraperitoneal local anaesthetic instillation versus no intraperitoneal local anaesthetic instillation for laparoscopic cholecystectomy, *Cochrane Database Syst. Rev.* 3 (Mar 13 2014), CD007337.
- [118] D. Segal, N. Awad, H. Nasir, S. Mustafa, L. Lowenstein, Combined spinal and general anesthesia vs general anesthesia for robotic sacrocrvicopexy: a randomized controlled trial, *Int. Urogynecol. J.* 25 (3) (2014) 369–374.
- [119] A. Karliczek, E.C. Jesus, D. Matos, A.A. Castro, A.N. Atallah, T. Wiggers, Drainage or nondrainage in elective colorectal anastomosis: a systematic review and meta-analysis, *Color. Dis.* 8 (4) (May 2006) 259–265.
- [120] E.C. Jesus, A. Karliczek, D. Matos, A.A. Castro, A.N. Atallah, Prophylactic anastomotic drainage for colorectal surgery, *Cochrane Database Syst. Rev.* 4 (Oct 18 2004), CD002100.
- [121] H. Petrowsky, N. Demartines, V. Rousson, P.A. Clavien, Evidence-based value of prophylactic drainage in gastrointestinal surgery: a systematic review and meta-analysis, *Ann. Surg.* 240 (6) (2004) 1074–1084 (discussion 1084–5).
- [122] E. Kalogera, S.C. Dowdy, A. Mariani, G. Aletti, J.N. Bakkum-Gamez, W.A. Cliby, Utility of closed suction pelvic drains at time of large bowel resection for ovarian cancer, *Gynecol. Oncol.* 126 (3) (Sep 2012) 391–396.
- [123] E. Kalogera, S.C. Dowdy, A. Mariani, A.L. Weaver, G. Aletti, J.N. Bakkum-Gamez, et al., Multiple large bowel resections: potential risk factor for anastomotic leak, *Gynecol. Oncol.* 130 (1) (Jul 2013) 213–218.
- [124] M. Jurado, J.L. Alcazar, J. Baixauli, J.L. Hernandez-Lizoain, Low colorectal anastomosis after pelvic exenteration for gynecologic malignancies: risk factors analysis for leakage, *Int. J. Gynecol. Cancer* 21 (2011) 397–402.
- [125] K. Charoenkwan, C. Kietpeerakool, Retroperitoneal drainage versus no drainage after pelvic lymphadenectomy for the prevention of lymphocyst formation in patients with gynaecological malignancies, *Cochrane Database Syst. Rev.* 6 (Jun 4 2014), CD007387.
- [126] P. Morice, N. Lassau, P. Pautier, C. Haie-Meder, C. Lhomme, D. Castaigne, Retroperitoneal drainage after complete para-aortic lymphadenectomy for gynecologic cancer: a randomized trial, *Obstet. Gynecol.* 97 (2) (Feb 2001) 243–247.
- [127] R.A. Brooks, J.D. Wright, M.A. Powell, J.S. Rader, F. Gao, D.G. Mutch, et al., Long-term assessment of bladder and bowel dysfunction after radical hysterectomy, *Gynecol. Oncol.* 114 (1) (Jul 2009) 75–79.
- [128] R. Griffiths, R. Fernandez, Policies for the removal of short-term indwelling urethral catheters, *Cochrane Database Syst. Rev.* 1 (2005), CD004011.
- [129] T.E.J. Ind, R. Brown, V.M. Pyneandee, M. Swanne, G. Taylor, Midnight removal of urinary catheters – improved outcome following gynaecological surgery, *Int. Urogynecol. J.* 4 (1993) 342–345.
- [130] M.R. Ahmed, W.A. Sayed Ahmed, K.A. Atwa, L. Metwally, Timing of urinary catheter removal after uncomplicated total abdominal hysterectomy, *Eur. J. Obstet. Gynecol. Reprod. Biol.* 176 (2014) 60–63.
- [131] S. Phipps, Y.N. Lim, S. McClinton, C. Barry, A. Rane, J. N'Dow, Short term urinary catheter policies following urogenital surgery in adults, *Cochrane Database Syst. Rev.* (2) (2006) (Art no. CD 004374).
- [132] T.H. Wells, H. Steed, V. Capstick, A. Schepanksy, M. Hiltz, W. Faught, What is the optimal method of bladder drainage after radical hysterectomy? *J. Obstet. Gynaecol. Can.* 30 (11) (2008) 1034–1038.
- [133] R. Naik, K. Maughan, A. Nordin, A. Lopes, K.A. Godfrey, M.H. Hatem, A prospective randomised controlled trial of intermittent self catheterisation vs suprapubic catheterisation for postoperative bladder care following radical hysterectomy, *Gynecol. Oncol.* 99 (2) (2005) 437–442.
- [134] H. Kehlet, D.W. Wilmore, Multimodal strategies to improve surgical outcome, *Am. J. Surg.* 183 (6) (Jun 2002) 630–641.
- [135] M. Van der Leeden, R. Huijsmans, E. Geleijn, d.L.-d.K. ES, J. Dekker, H.J. Bonjer, et al., Early enforced mobilisation following surgery for gastrointestinal cancer: feasibility and outcomes, *Physiotherapy* (May 7 2015) (pii: S0031-9406(15)03780-3).
- [136] M.R. Cassidy, P. Rosenkranz, D. McAneny, Reducing postoperative venous thromboembolism complications with a standardized risk-stratified prophylaxis protocol and mobilization program, *J. Am. Coll. Surg.* 218 (6) (Jun 2014) 1095–1104.
- [137] M. Liebermann, M. Awad, M. Dejong, C. Rivard, J. Sinacore, L. Brubaker, Ambulation of hospitalized gynecologic surgical patients: a randomized controlled trial, *Obstet. Gynecol.* 121 (3) (Mar 2013) 533–537.
- [138] M.R. Ahmed, W.A. Sayed Ahmed, K.A. Atwa, L. Metwally, Timing of urinary catheter removal after uncomplicated total abdominal hysterectomy: a prospective randomized trial, *Eur. J. Obstet. Gynecol. Reprod. Biol.* 176 (May 2014) 60–63.
- [139] K. Lassen, M. Soop, J. Nygren, P.B. Cox, P.O. Hendry, C. Spies, et al., Consensus review of optimal perioperative care in colorectal surgery: Enhanced Recovery After Surgery (ERAS) group recommendations, *Arch. Surg.* 144 (10) (Oct 2009) 961–969.
- [140] U.O. Gustafsson, J. Hausel, A. Thorell, O. Ljungqvist, M. Soop, J. Nygren, Adherence to the enhanced recovery after surgery protocol and outcomes after colorectal cancer surgery, *Arch. Surg.* 146 (2011) 571–577.
- [141] ERAS Interactive Audit System (EIAS), <http://www.erasociety.org/index.php/eras-care-system/eras-interactive-audit-system>, 2015.