



Recurrent bridgehead effects accelerate global alien ant spread

Cleo Bertelsmeier^{a,1}, Sébastien Ollier^b, Andrew M. Liebhold^c, Ekehard G. Brockerhoff^d, Darren Ward^{e,f}, and Laurent Keller^{a,1}

^aDepartment of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland; ^bEcologie Systématique Evolution, Université Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, 91400 Orsay, France; ^cNorthern Research Station, US Forest Service, Morgantown, WV 26505; ^dForest Protection, Scion (New Zealand Forest Research Institute), Riccarton, Christchurch, 8440 New Zealand; ^eSystematics, Landcare Research, Auckland, New Zealand; and ^fSchool of Biological Sciences, University of Auckland, Auckland, New Zealand

Edited by Joan E. Strassmann, Washington University in St. Louis, St. Louis, MO, and approved April 6, 2018 (received for review February 8, 2018)

Biological invasions are a major threat to biological diversity, agriculture, and human health. To predict and prevent new invasions, it is crucial to develop a better understanding of the drivers of the invasion process. The analysis of 4,533 border interception events revealed that at least 51 different alien ant species were intercepted at US ports over a period of 70 years (1914–1984), and 45 alien species were intercepted entering New Zealand over a period of 68 years (1955–2013). Most of the interceptions did not originate from species' native ranges but instead came from invaded areas. In the United States, 75.7% of the interceptions came from a country where the intercepted ant species had been previously introduced. In New Zealand, this value was even higher, at 87.8%. There was an overrepresentation of interceptions from nearby locations (Latin America for species intercepted in the United States and Oceania for species intercepted in New Zealand). The probability of a species' successful establishment in both the United States and New Zealand was positively related to the number of interceptions of the species in these countries. Moreover, species that have spread to more continents are also more likely to be intercepted and to make secondary introductions. This creates a positive feedback loop between the introduction and establishment stages of the invasion process, in which initial establishments promote secondary introductions. Overall, these results reveal that secondary introductions act as a critical driver of increasing global rates of invasions.

Previous studies have identified bridgehead effects in several well-studied invasive species (13–19). However, all of these studies reconstructed the invasion histories of individual species, and it remains unknown how frequent secondary introductions are compared with primary introductions overall. To quantify the frequency of secondary introductions, we analyzed border interceptions of ants at air and maritime ports in the United States and New Zealand, which we used as a proxy for accidental species introductions (20). Many introductions of species occur via air and maritime ports; the analysis of such data therefore constitutes a powerful means to investigate the first step of the invasion process and to determine the proportion of primary and secondary introductions (21). We focused on ants, which make up a particularly prominent group of widespread invasive species (22) that can rapidly disassemble native communities (23). The variety of their lifestyles, colony sizes, habitat requirements, and diets also allows them to colonize almost all terrestrial habitats on all continents except Antarctica (8, 22). Their complex social structure has contributed to them becoming highly successful invasive species. There are currently more than 200 known ant species that have established populations outside of their native range (hereafter referred to as alien species; Table 1), and 19 are classified as invasive by the International Union for the Conservation of Nature because of their effects on biodiversity,

biological invasions | globalization | propagule pressure | invasive ants | secondary introductions

The globalization of trade and travel is facilitating the accidental transport of animal and plant species around the world (1–3). Although many of these species fail to overcome biotic and abiotic barriers to establishment outside of their native range, those that do succeed are among the greatest threats to global biodiversity, agriculture, and human health (4, 5). A cross-taxonomic comparison has shown that there has been no saturation in the global accumulation of introduced species (6), and that the rates of new biological invasions may continue to rise in the future (7). In particular, the second wave of trade globalization after World War II provided opportunities for transport of a new set of species establishing ranges spanning trans-continental scales (8) and increased invasions to countries with historically low levels of trade (9).

It has been suggested that biological invasions are a self-reinforcing phenomenon (“invasion begets invasion”) (10), via the so-called bridgehead effect (11). Under a bridgehead scenario, a successfully established invasive (Table 1) population serves as a source of colonists for new invasions and thereby gives rise to secondary introductions. A recent horizon scan of emerging challenges and opportunities in invasion science highlighted the bridgehead effect as one of the most important ecological issues likely to influence how biological invasions are studied and managed in the near future (12).

Significance

Because of the globalization of trade and travel, worldwide invasion rates are high. A potential driver of the global acceleration of new invasions is the so-called bridgehead effect, in which initial invasive populations serve as the source of additional invasions via secondary introductions. However, the frequency and overall importance of secondary introductions remain largely unknown. Using a remarkable dataset, spanning nearly 100 years (1914–2013), of ant interceptions at air and maritime ports in the United States and New Zealand, we found that most ant introductions arise via secondary transport via intermediate regions. Our analyses also reveal positive feedback between the introduction and establishment stages of the invasion process via secondary introductions acting as a critical driver of increasing global invasion rates.

Author contributions: C.B., S.O., A.M.L., and L.K. designed research; C.B., S.O., and L.K. performed research; A.M.L., E.G.B., and D.W. contributed new reagents/analytic tools; C.B. and S.O. analyzed data; and C.B., S.O., A.M.L., E.G.B., D.W., and L.K. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

¹To whom correspondence may be addressed. Email: cleo.bertelsmeier@unil.ch or Laurent.Keller@unil.ch.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1801990115/-DCSupplemental.

Published online May 7, 2018.

Table 1. Glossary

Status	Definition
Native species	Species with no established population outside of the native range.
Alien species	Species with at least one established (self-sustaining) population outside of the native range, at outdoor locations.
Invasive species	Alien species with documented effects on biodiversity, agriculture, health or ecosystem functioning; listed according to this criterion by the Invasive Species Specialist Group of the International Union for the Conservation of Nature

ecosystem functioning, agriculture, infrastructure, and human or animal health (8), causing important economic losses (24, 25). Five species are even among the “100 worst invasive species” (26).

We defined primary interceptions as those originating from any country within the native range of a given species, and secondary interceptions as those originating from any country outside the native range of a species. Our rich dataset of 4,533 ant

interception events provides an exceptional opportunity not only to quantify the relative importance of introductions arising from the native versus introduced range of intercepted species but also to identify potential sources of secondary transport and to test whether there is positive feedback among the introduction, establishment, and worldwide spread of species (i.e., if bridgehead effects have the potential to contribute to rising global invasion rates).




Fig. 1. Percentage of primary vs. secondary introductions of the most frequently intercepted species in the United States and New Zealand. The proportion of interceptions from the species' native countries is shown above the x-axis (in gray) and the proportion of interceptions from countries in the alien range below the x-axis (in color). The color code indicates the origin of the secondarily intercepted species. Species were visually grouped on the x-axis according to their native range (Dataset S1). Af, Africa; As, Asia; Eu, Europe; N. Am, North America; L. Am, Latin America; Oc, Oceania.




Fig. 4. Feedback loop between interceptions and successful establishment. (A) Link between the number of border interceptions per species and establishment probability \pm SD, modeled using a binomial GLM (red line). (B) Link between the number of continents where a species has established and the number of times it has been intercepted, modeled using a GLM with Poisson link function (red line). The size of the gray points is proportional to the number of cases.

2D). Thus, Latin America acted as a bridgehead to the United States, and Oceania as a bridgehead to New Zealand (Fig. 3). Our findings that a high frequency of secondary introduction reflects the topology of human transport does not exclude that some species evolved a greater invasiveness in some introduced populations.

The second prediction is that secondary introductions should generate a positive feedback loop whereby invasion begets invasion, similar to the way in which an infectious disease can increase the rate of spread when the number of infected hosts increases (35). To test this prediction, we first investigated whether the probability of establishment (i.e., attaining a self-sustaining population) was positively associated with the number of interceptions and, second, if successful establishments on a higher number of continents lead, in turn, to a higher number of interceptions. In line with these predictions, species that were more frequently intercepted at ports of entry were also more likely to become established in both the United States [generalized linear model (GLM), $\chi^2 = 4.6$; $P = 0.047$] and New Zealand (GLM, $\chi^2 = 7.1$; $P < 0.01$; Fig. 4A). Conversely, species that were successfully established in a greater number of continents were more likely to be intercepted at ports of entry in both the United States (GLM, $\chi^2 = 539$; $P < 0.0001$) and New Zealand (GLM, $\chi^2 = 879$; $P < 0.0001$; Fig. 4B). Species established in a greater number of continents also had a greater proportion of secondary interceptions in both the United States (GLM, $\chi^2 = 320$; $P < 0.0001$) and New Zealand (GLM, $\chi^2 = 618$; $P < 0.0001$). These data thus confirmed a positive effect of propagule pressure on invasion success (20, 37, 38) and demonstrated a positive feedback of the rate of successful establishment on the rate of interceptions. Moreover, some species-level morphological and life-history traits can also influence the probability that a species is transported, is established successfully, and will be transported further (20). It has been shown that nesting habits can interact with the effect of propagule

pressure on establishment probability (20). Additional traits such as the type of colony founding, queen number, and habitat generalism also influence colonization ability after transcontinental transport (8). Therefore, species with these life-history traits may be particularly likely to benefit from the positive feedback loop between introductions and new establishments once they have started to spread through human transport.

This study provides a quantitative global assessment of secondary introductions and reveals that a surprisingly large number of interceptions of alien ant species comes from invaded ranges. This has important implications for our understanding of invasion processes because bridgeheads appear to play a key role in increasing invasion rates when species reach regions that are highly connected. Our analyses also reveal a positive feedback among global introduction, spread, and subsequent invasion, which is likely to lead to further acceleration of the worldwide spread of invasive species in the future.

Materials and Methods

Interceptions. For the United States, we used records of ant interceptions made by inspectors of the US Department of Agriculture (USDA) and the Department of Homeland Security at ports of entry (maritime, land, and air) from 1914 to 1984, published by the US Department of Agriculture (39). For New Zealand, we used ant interceptions compiled by the New Zealand Ministry for Primary Industries (or predecessors to this ministry previously recorded under different names) responsible for examining cargo, goods, mail, and baggage that arrived from 1955 to 2013. Parts of these data have been described in an earlier paper on New Zealand interception records (40). Data in both interception databases included ant species names and the country of origin. Species names were checked for synonymy using the authoritative AntWeb v. 6.0.13 (antweb.org), which contains 15,961 valid species names based on the Bolton World catalog (41) and the taxonomic history of senior synonyms that have become outdated because of taxonomic revisions. Because countries have also changed names over the course of the last century, we revised the list of origins using ISO code 3166, which provides an international standard for country codes and their subdivisions.

We used the Antmaps database, a web interface of the Global Ant Biodiversity Informatics project (42), to obtain a list of alien species. To be conservative and consider only species with known established populations outdoors, we excluded all species listed as “needing verification,” “dubious,” or “indoors” (see [Dataset S1](#) for interceptions of alien ants and [Dataset S2](#) for other intercepted ant species without documented alien populations). The US dataset contained 1,428 alien ant interceptions with information on the country of origin and a specimen identified at the species level (51 species from 29 genera were recorded). The New Zealand dataset contained 3,105 alien ant interceptions with information on the country of origin and a specimen identified at the species level (45 species from 26 genera). In total, our dataset contains 69 alien ant species, 27 of which were intercepted in both countries. To test whether species classified as invasive exhibit a higher rate of secondary introductions than other alien species, we distinguished between alien and invasive alien species, according to the Invasive Species Specialist Group of the International Union for the Conservation of Nature, which lists as invasive species those with documented effects on biodiversity and/or human activities (www.iucngisd.org/gisd/).

Distribution and Establishment. To determine the global country-level distribution of species and to confirm the presence/absence of all intercepted species in the United States and New Zealand, we used georeferenced AntWeb v. 6.0.13 specimen data; species lists from the literature assembled by AntWiki (an authoritative database maintained by ant experts, which contains regional ant fauna lists from different countries over the world and is interlinked with the Encyclopedia of Life); the New Zealand Landcare Research database (43), which provides occurrence data of alien and invasive ant species; a dataset of recorded alien ants in the United States (44, 45); and a dataset of alien ant species worldwide (46). To determine the native and introduced range of the alien ant species, we combined information on the introduced ranges of alien ant species from the literature (46) and the Antmaps database. The alien range of a species was defined as all the countries comprising at least one introduced population (recorded by one or both of these datasets). Conversely, the native range was defined as all countries containing native but no introduced populations.

Secondary Introductions. For all species intercepted more than 20 times in the United States or New Zealand, we calculated the proportion of primary

introductions (interceptions originating from countries within the native range) and secondary introductions (interceptions originating from countries where the species has established an introduced population). To test whether the proportion of secondary introductions varied across species, we used a chi-square test. To visually represent the origin of secondary interceptions for the most frequently intercepted species (Fig. 1), we grouped species according to their native range (Asia, Africa, Europe, Oceania, Latin America, and North America; hereafter called continents). When the native range covered several continents, we selected the continent containing the larger number of native countries (Dataset S1).

To identify countries and regions acting as major source of interceptions in the United States and New Zealand, we analyzed the total number of ant interceptions originating from each country. We mapped these export countries, calculating an interception index, based on the log number of interceptions. To visually identify countries that are important exporters, we divided the number of interceptions by the log of land area of each country because the number of interceptions should be influenced by the size of the countries (Fig. 2). To map global flows of ant interceptions, we calculated the proportion of primary and secondary interceptions from each country, represented as a pie chart, where we subdivided secondary interceptions according to the native range of the species, using a color code (Fig. 3). When a species was native from more than one continent, we allocated the number of secondary interceptions to the respective continents proportional to the number of countries comprising native populations.

Propagule Pressure and Positive Feedback. To test for an association between propagule pressure and establishment likelihood, we modeled the estab-

lishment probability at the species-level with a binomial GLM (logit-link), including all alien ant species intercepted at least once. The interception counts for each species were log-transformed. To test for a positive feedback (i.e., if the establishment in a higher number of continents elsewhere is linked to a higher number of interceptions in the United States and New Zealand), we used a GLM (Poisson link) of the number of interceptions per species against the number of continents where it has established. To test whether the number of continents in which a species has established was linked to the proportion of secondary introductions in the United States and New Zealand, we used a GLM (binomial link), where we used the total number of interceptions per species as weights. This GLM parameter is used to specify the number of cases when the response variable is a proportion.

ACKNOWLEDGMENTS. We thank P. Christe, A. Guisan, T. Schwander, and four reviewers for their comments on the manuscript; Stefan Cover, Brian Fisher, and Ben Hoffmann for advice on native ranges of alien ants; and US Department of Agriculture Animal and Plant Health Inspection Service, and the New Zealand Ministry for Primary Industries for providing summaries of ant interception records from the United States and New Zealand, respectively. C.B. and L.K. were supported by the Swiss National Science Foundation and two European Research Council advanced grants. D.W. was supported from Ministry of Business, Innovation & Employment (MBIE) core backbone funding to Landcare Research within the Managing Invasives Portfolio, and the Better Border Biosecurity collaboration. E.G.B. acknowledges MBIE core funding (C04X1104) to Scion and the Better Border Biosecurity collaboration (www.b3nz.org).

- Nentwig W (2015) Introduction, establishment rate, pathways and impact of spiders alien to Europe. *Biol Invasions* 17:2757–2778.
- Rouget M, et al. (2015) Invasion debt—Quantifying future biological invasions. *Divers Distrib* 22:445–456.
- Essl F, et al. (2011) Socioeconomic legacy yields an invasion debt. *Proc Natl Acad Sci USA* 108:203–207.
- Clavero M, Garcia-Berthou E (2005) Invasive species are a leading cause of animal extinctions. *Trends Ecol Evol* 20:110.
- Mack R, et al. (2000) Biotic invasions: Causes, epidemiology, global consequences, and control. *Ecol Appl* 10:689–710.
- Seebens H, et al. (2017) No saturation in the accumulation of alien species worldwide. *Nat Commun* 8:14435.
- Seebens H, et al. (2015) Global trade will accelerate plant invasions in emerging economies under climate change. *Glob Change Biol* 21:4128–4140.
- Bertelsmeier C, Ollier S, Liebhold A, Keller L (2017) Recent human history governs global ant invasion dynamics. *Nat Ecol Evol* 1:0184.
- Early R, et al. (2016) Global threats from invasive alien species in the twenty-first century and national response capacities. *Nat Commun* 7:12485.
- Garnas JR, et al. (2016) Complex patterns of global spread in invasive insects: Evolutionary and management consequences. *Biol Invasions* 18:935–952.
- Lombaert E, et al. (2010) Bridgehead effect in the worldwide invasion of the bio-control harlequin ladybird. *PLoS One* 5:e9743.
- Ricciardi A, et al. (2017) Invasion science: A horizon scan of emerging challenges and opportunities. *Trends Ecol Evol* 32:464–474.
- Ascunce MS, et al. (2011) Global invasion history of the fire ant *Solenopsis invicta*. *Science* 331:1066–1068.
- Vogel V, Pedersen JS, Giraud T, Krieger MJB, Keller L (2010) The worldwide expansion of the Argentine ant. *Divers Distrib* 16:170–186.
- Miller N, et al. (2005) Multiple transatlantic introductions of the western corn rootworm. *Science* 310:992.
- Boubou A, et al. (2012) Test of colonisation scenarios reveals complex invasion history of the red tomato spider mite *Tetranychus evansi*. *PLoS One* 7:e35601.
- Darling JA, Bagley MJ, Roman J, Tepolt CK, Geller JB (2008) Genetic patterns across multiple introductions of the globally invasive crab genus *Carcinus*. *Mol Ecol* 17:4992–5007.
- Barker BS, Andonian K, Swope SM, Luster DG, Dlugosch KM (2017) Population genomic analyses reveal a history of range expansion and trait evolution across the native and invaded range of yellow starthistle (*Centaurea solstitialis*). *Mol Ecol* 26:1131–1147.
- Leduc A, et al. (2015) Bridgehead invasion of a monomorphic plant pathogenic bacterium: *Xanthomonas citri* pv. *citri*, an emerging citrus pathogen in Mali and Burkina Faso. *Environ Microbiol* 17:4429–4442.
- Suarez AV, Holway DA, Ward PS (2005) The role of opportunity in the unintentional introduction of nonnative ants. *Proc Natl Acad Sci USA* 102:17032–17035.
- Blackburn TM, et al. (2011) A proposed unified framework for biological invasions. *Trends Ecol Evol* 26:333–339.
- Holway DA, et al. (2002) The causes and consequences of ant invasions. *Annu Rev Ecol Syst* 33:181–233.
- Cerdá X, Angulo E, Caut S, Courchamp F (2012) Ant community structure on a small Pacific island: Only one native species living with the invaders. *Biol Invasions* 14:323–339.
- Pimentel D (2011) *Biological Invasions—Economic and Environmental Costs of Alien Plant, Animal, and Microbe Species* (CRC Press, Boca Raton, FL).
- Bradshaw CJA, et al. (2016) Massive yet grossly underestimated global costs of invasive insects. *Nat Commun* 7:12986.
- Lowe S, Browne M, Boudjelas S, De Poorter M (2000) *100 of the World's Worst Invasive Alien Species - A selection from the Global Invasive Species Database* (The Invasive Species Specialist Group, Auckland, New Zealand), pp 12.
- Keller SR, Gilbert KJ, Fields PD, Taylor DR (2012) Bayesian inference of a complex invasion history revealed by nuclear and chloroplast genetic diversity in the colonizing plant, *Silene latifolia*. *Mol Ecol* 21:4721–4734.
- De Kort H, Mergeay J, Jacquemyn H, Honnay O (2016) Transatlantic invasion routes and adaptive potential in North American populations of the invasive glossy buckthorn, *Frangula alnus*. *Ann Bot* 118:1089–1099.
- Foucaud J, et al. (2010) Worldwide invasion by the little fire ant: Routes of introduction and eco-evolutionary pathways. *Evol Appl* 3:363–374.
- Gotzke D, Axen HJ, Suarez AV, Helms Cahan S, Shoemaker D (2015) Global invasion history of the tropical fire ant: A stowaway on the first global trade routes. *Mol Ecol* 24:374–388.
- Ugelvig LV, et al. (2008) The introduction history of invasive garden ants in Europe: Integrating genetic, chemical and behavioural approaches. *BMC Biol* 6:11.
- Colautti RI, Lau JA (2015) Contemporary evolution during invasion: Evidence for differentiation, natural selection, and local adaptation. *Mol Ecol* 24:1999–2017.
- Colautti RI, Barrett SCH (2013) Rapid adaptation to climate facilitates range expansion of an invasive plant. *Science* 342:364–366.
- Banks NC, Paini DR, Bayliss KL, Hodda M (2015) The role of global trade and transport network topology in the human-mediated dispersal of alien species. *Ecol Lett* 18:188–199.
- Floerl O, Inglis GJ, Dey K, Smith A (2009) The importance of transport hubs in stepping-stone invasions. *J Appl Ecol* 46:37–45.
- Carlton JT (1996) Pattern, process, and prediction in marine invasion ecology. *Biol Conserv* 78:97–106.
- Lockwood JL, Cassey P, Blackburn T (2005) The role of propagule pressure in explaining species invasions. *Trends Ecol Evol* 20:223–228.
- Brockerhoff EG, Kimberley M, Liebhold AM, Haack RA, Cavey JF (2014) Predicting how altering propagule pressure changes establishment rates of biological invaders across species pools. *Ecology* 95:594–601.
- Magarey RD, Colunga-García M, Fieselman DA (2009) Plant biosecurity in the United States: Roles, responsibilities, and information needs. *Bioscience* 59:875–884.
- Ward DF, Beggs JR, Clout MN, Harris RJ, O'Connor S (2006) The diversity and origin of exotic ants arriving in New Zealand via human-mediated dispersal. *Divers Distrib* 12:601–609.
- Bolton B, Alpert G, Ward PS, Naskrecki P (2007) *Bolton's Catalogue of the Ants of the World* (Harvard Univ Press, Cambridge, MA).
- Janicki J, Narula N, Ziegler M, Guénard B, Economo EP (2016) Visualizing and interacting with large-volume biodiversity data using client-server web-mapping applications: The design and implementation of antmaps.org. *Ecol Inform* 32:185–193.
- Ward DF, Rees J, Harris RJ, Stanley MC (2009) New Zealand Ant Distribution Database. v2.0. Available at <https://www.landcareresearch.co.nz/science/plants-animals-fungi/animals/invertebrates/invasive-invertebrates/antsnz/ant-distribution-data>. Accessed May 15, 2017.
- Wittenborn D, Jeschke JM (2011) Characteristics of exotic ants in North America. *Neobiota (Berlin)* 64:47–64.
- Deyrup M, Davis L, Cover S (2000) Exotic ants in Florida. *Trans Am Entomol Soc* 126:293–326.
- McGlynn TP (1999) The worldwide transfer of ants: Geographical distribution and ecological invasions. *J Biogeogr* 26:535–548.