

Serveur Académique Lausannois SERVAL serval.unil.ch

Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

Title: The Effect of Cigarette Smoking on Diabetic Peripheral Neuropathy: A Systematic Review and Meta-Analysis.

Authors: Clair C, Cohen MJ, Eichler F, Selby KJ, Rigotti NA

Journal: Journal of general internal medicine

Year: 2015 Aug

Volume: 30

Issue: 8

Pages: 1193-203

DOI: 10.1007/s11606-015-3354-y

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

The Effect of Cigarette Smoking on Diabetic Peripheral Neuropathy: a Systematic Review and Meta-Analysis

Carole Clair, MD, MSc¹, Marya J Cohen, MD, MPH², Florian Eichler, MD³, Kevin J. Selby, MD¹, Nancy A. Rigotti, MD⁴

¹Department of Ambulatory Care and Community Medicine, Lausanne University Hospital, Lausanne, Switzerland, ²Division of General Internal Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, United States, ³Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, United States ⁴Massachusetts General Hospital, Tobacco Research and Treatment Center, Division of General Internal Medicine, Harvard Medical School, Boston, Massachusetts, United States

Authors' emails:

Carole Clair: carole.clair@hospvd.ch

Marya Cohen: mjcohen@partners.orgmicro

Florian Eichler: feichler@partners.org

Kevin J. Selby: kevin.selby@hospvd.ch

Nancy Rigotti: nrigotti@partners.org

Corresponding author

Carole Clair

Policlinique Médicale Universitaire

Rue du Bugnon 44

1011 Lausanne, Switzerland

Phone: +41 79 556 81 46

Fax: +41 21 314 61 01

Email address: Carole.Clair@hospvd.ch

Word count

Abstract: 286 words

Manuscript: 3257 words

References: 71

3 tables, 4 figures, 4 supplementary files

1 **Abstract**

2 **Objective:** Studies suggest that smoking might be a risk factor for the development of
3 microvascular complications such as diabetic peripheral neuropathy (DPN). The objective of
4 this study is to assess the relationship between smoking and DPN in people with type 1 or
5 type 2 diabetes.

6 **Research Design and Methods:** A systematic review of PubMed, EMBASE, and Cochrane
7 Clinical Trials databases was conducted from 1966 to November 2014 for cohort, cross-
8 sectional and case-controlled studies that assessed the relationship between smoking and
9 DPN. Separate meta-analyses for prospective cohort studies and case control or cross-
10 sectional studies were performed using random effects models.

11 **Results:** Thirty-eight studies (10 prospective cohort and 28 cross-sectional) were included.
12 The prospective cohort studies included 5,558 participants without DPN at baseline. During a
13 follow up ranging from 2 to 10 years, 1,550 cases of DPN occurred. The pooled unadjusted
14 odds ratio (OR) of developing DPN associated with smoking was 1.26 (95% CI 0.86-1.85;
15 $I^2=74.3\%$; Evidence grade: moderate strength). Overall prospective studies were of moderate
16 to high quality according to the Newcastle-Ottawa scale. Including only the six studies of
17 moderate quality, we found a higher and significant association, with an adjusted OR of 1.73
18 (95% CI 1.48-2.03, $I^2=0\%$, Evidence grade: moderate strength). The cross sectional studies
19 included 27,594 participants. The pooled OR of DPN associated with smoking was 1.42
20 (95% CI 1.21-1.65; $I^2=64.5\%$; Evidence grade: moderate strength). Overall cross-sectional
21 studies were of low to moderate quality, and after exclusion of studies of low quality (n=4),
22 the OR was 1.29 (95% CI 1.13-1.48; $I^2=50.5\%$). There was no evidence of publication bias.

23 **Conclusions:** Smoking may be associated with an increased risk of DPN in people with
24 diabetes. Future studies are needed to test whether smoking cessation reduces the risk of DPN

25 in adult with diabetes.

26 Diabetic peripheral neuropathy, also known as distal symmetrical polyneuropathy or
27 sensorimotor neuropathy, is part of a wider spectrum of microvascular complications of
28 diabetes; other microvascular complications include ulcer/amputations, erectile dysfunction
29 and autonomic dysfunction. Diabetic peripheral neuropathy is the most common
30 microvascular complication of diabetes, affecting approximately 30% of people with
31 diabetes¹⁻³. Symptoms include numbness, tingling or burning sensation in the legs and hands,
32 typically in a “stocking and glove” distribution¹. Ultimately, muscle weakness, loss of
33 reflexes, and foot deformities can result, leading to end clinical sequelae of ulcers, potential
34 infection, and amputation for some patients with poorly-controlled disease.

35 There is a complex interaction between metabolic and vascular factors in the pathogenesis of
36 diabetic peripheral neuropathy^{1,4}. Hyperglycemia has been the primary factor described and
37 leads to nerve-cell damage through several mechanisms, including oxidative stress or polyol
38 accumulation³. Reduced nerve perfusion, endoneurial hypoxia and endothelial dysfunction
39 also contribute to neuropathy development¹.

40 Previous studies have investigated potential risk factors for diabetic peripheral neuropathy,
41 including hypertension, microalbuminuria, dyslipidemia, and of particular interest, cigarette
42 smoking⁵⁻⁷. There appears to be an increased likelihood of neuropathy in people with diabetes
43 who smoke; however, prior studies investigating this relationship only included a small
44 number of participants⁷.

45 In order to better assess the relationship between smoking and diabetic neuropathy, we
46 conducted a systematic review and meta-analysis, analyzing cross-sectional, case control,
47 prospective and retrospective cohort studies.

48

49

50

51 **Research Design and Methods**

52 **Search Strategy and Selection Criteria**

53 We searched PubMed (1966 to November 2014), Embase (1980 to November 2014) and
54 Cochrane Clinical Trials (until November 2014). We also searched the references of the
55 relevant retrieved articles. Studies that assessed the effect of cigarette smoking on the risk of
56 peripheral neuropathy among patients with type 1 or type 2 diabetes were included
57 (population of interest). Only participants with diabetes at baseline were included, as we were
58 interested primarily in the effect of smoking on diabetic complications. The exposure of
59 interest was cigarette smoking. In order to be considered for inclusion in the systematic
60 review, all studies had to include a control or comparison group of participants with diabetes
61 who did not smoke. The outcome of interest was diabetic peripheral neuropathy.

62 Cohort studies as well as cross-sectional studies and case control studies were included based
63 on our search results. For cohort studies, we included studies with at least one year of follow-
64 up because we assumed that there should be a latency period of at least one year for smoking
65 to impact the development of diabetic neuropathy. We considered studies published in all
66 languages and did not restrict our search to published studies only.

67 For our search, we combined 3 search themes: 1) diabetes, 2) smoking and 3) neuropathy.

68 The full electronic search is available in the **online appendix 1**.

69 **Study Selection**

70 A first screen of retrieved citations was performed based on titles and abstracts; each citation
71 was screened by 2 different co-authors (CC, MJC, FE or KJS). The inclusion criteria for this
72 first screen were the following: population with diabetes (type 1 or type 2) at baseline,
73 neuropathy as one recorded outcome (not necessarily the primary outcome), and
74 identification as prospective, cohort or cross-sectional studies. We included studies even if
75 they did not mention smoking exposure in the title or abstract (although it was preferable).

76 Exclusion criteria included gestational diabetes, animal studies, or non-original study design
77 (such as reviews, editorials, and case reports/case series). A second screen was then
78 performed based on full-text review of retained citations. Exclusion criteria were the same as
79 for the first screen, with the addition of the following criteria: 1) Smoking-neuropathy
80 relationship was not assessed and/or data did not allow calculating it by hand; 2) Peripheral
81 neuropathy was not one of the outcomes; or 3) People without diabetes were included.

82 Two reviewers (CC, MJC, FE or KJS) independently reviewed the articles and any
83 disagreement was resolved by consensus.

84 **Data Extraction and Quality Assessment**

85 Two authors independently extracted the data from selected studies. To evaluate the risk of
86 bias in individual studies, and assess overall quality, we considered several criteria based on
87 the Newcastle-Ottawa scale⁸. The Newcastle-Ottawa scale for cohort studies has 3 categories:
88 1) selection (representativeness of the exposed cohort, selection of the non-exposed cohort,
89 ascertainment of exposure and demonstration that outcome of interest was not present at start
90 of study) (0-4 points), 2) comparability (comparability of cohorts on the basis of design or
91 analyses) (0-2 points) and 3) outcome (assessment of outcome, was follow-up long enough
92 for outcomes to occur, adequacy of follow-up of cohorts) (0-3 points). The Newcastle-Ottawa
93 scale exists for case control studies and we used a modified version to evaluate the quality of
94 cross-sectional studies. In the modified version we deleted the question on selection of
95 controls (in the “selection” category, yielding to a maximum of 3 points) and the questions on
96 methods of ascertainment for cases and controls and non-response rate (in the “exposure”
97 section, yielding to a maximum of 2 points). We reported the score for each subcategory in
98 the extraction form. We defined Additionally, we evaluated the strength of evidence of the
99 studies using the Cochrane GRADE criteria⁹. By definition, observational studies are
100 considered “Low quality” with the GRADE approach. We upgraded studies to “Moderate

101 quality” if they met at least one of the following factors: 1. “Large magnitude of effect”; 2.
102 “All plausible confounding would reduce a demonstrated effect or suggest a spurious effect
103 when results show no effect”; or 3. “Dose-response gradient”. Studies were downgraded to
104 “Very low quality” if at least one of the following factors was present: 1. “Limitations in the
105 design and implementation of available studies suggesting high likelihood of bias”; 2.
106 “Indirectness of evidence”; 3. “Unexplained heterogeneity or inconsistency of results”; 4.
107 “Imprecision of results”; or 5. “High probability of publication bias”.

108 Studies reported risk ratios (RR), odds ratios (OR) or absolute numbers when describing the
109 relationship between smoking and diabetic peripheral neuropathy. As most prospective and
110 cross-sectional studies reported ORs and not all studies provided information to convert OR
111 into RR, we used OR in our meta-analyses. For studies that did not provide OR or RR, we
112 calculated unadjusted ORs and Confidence intervals (CI) manually.

113 **Data Synthesis and Analysis**

114 We pooled our results using the DerSimonian and Laird random effect model¹⁰ because we
115 expected to have heterogeneity between studies. Anticipated sources of heterogeneity
116 included study population (people with type 1 versus type 2 diabetes), definition of smoking
117 and definition of neuropathy and were defined *a priori*. We explored other sources of
118 heterogeneity for 3 variables that were added post-hoc: level of adjustment, mean duration of
119 follow-up (for prospective study only) and level of quality assessed with the Newcastle-
120 Ottawa scale⁸. We then performed stratified analyses to assess/explore potential sources of
121 heterogeneity linked to a priori and post hoc variables. In parallel we performed univariate
122 metaregression analyses to quantify potential source of heterogeneity.

123 We performed separate meta-analyses stratified by type of design. To assess heterogeneity,
124 the Q statistic and I squared statistic were calculated^{11,12}.

125 The possibility of publication bias was assessed using the Begg test and visual inspection of
126 the funnel plot^{13,14}. STATA 13 (StataCorp, College Station, Texas) was used for statistical
127 analyses.

128 **Results**

129 **Study Selection**

130 In terms of study selection, the initial search included 2006 citations from PubMed, Embase
131 and Cochrane Clinical Trials databases. After excluding duplicates, 1554 unique citations
132 were available (see **Figure 1**). After the first screen, 126 citations were considered for further
133 review. After a second screen, 88 studies were excluded based on full text review. Agreement
134 between reviewers at this stage was good with a Kappa of 0.78. Reasons for exclusion at this
135 point included no estimate (or numbers to allow manual calculation) of the smoking-
136 neuropathy relationship (n=54), outcome other than peripheral neuropathy (n = 30), or
137 inclusion of participants without diabetes (n = 4). Finally, 38 studies were selected for final
138 inclusion in the systematic review and we performed separate meta-analyses for the 10
139 prospective studies^{5,15-23} and the 28 cross-sectional studies^{6,7,24-49}.

140 **Smoking and incidence of diabetic peripheral neuropathy in prospective cohort studies**

141 The main individual characteristics of the prospective studies are shown in **Table 1**. They
142 include 5,558 participants in total; 3 studies include participants with type 2 diabetes, 6
143 studies included participants with type 1 diabetes and 1 study included both participants with
144 type 1 and type 2 diabetes. Participants were from different settings including inpatient,
145 outpatient and the community; mean age of participants ranged from 25 to 66 years old and
146 mean diabetes duration ranged from 0 to 17 years. All studies excluded participants with
147 neuropathy at baseline and participants were followed for 2 to 10 years. Peripheral
148 neuropathy screening was done by neurological history and examination in most studies<sup>5,17-
149 20,23</sup>, by electromyography to measure nerve conduction velocities in one study¹⁵, through
150 measure of vibration perception with biothesiometers in one study²², and by monofilament
151 examinations in two studies^{16,21}. The definition of smoking exposure varied between studies;
152 six studies compared ever smokers (i.e. current and former smokers) to never smokers, one

153 study compared current to non smokers (i.e. former and never smokers) and 3 studies did not
154 clearly specify the smoking comparison groups. Most studies provided OR, two RR and one
155 gave numbers of smokers and non smokers and of participants who developed peripheral
156 neuropathy in each category. All studies except one performed multivariable-adjusted
157 analyses; five controlled for at least A1C and diabetes duration and 4 adjusted for either A1C
158 or diabetes duration and several other confounders - see **Online appendix 2**. The quality of
159 studies varied. Most were considered good quality with maximum points for selection and
160 exposure criteria on the Newcastle-Ottawa scale; however, two were classified as suboptimal
161 for quality with lower scores^{15,16}. Using the GRADE criteria, two studies were rated as
162 “Low quality”^{21,23}, two were downgraded as “Very low”^{15,16} because of a poorly defined
163 outcome and risk of selection bias, and six were upgraded to “Moderate quality”^{5,17-20,22}
164 mainly due to adjustment for confounding factors and a dose-response gradient. In terms of
165 the incidence of diabetic peripheral neuropathy, 7 studies showed a positive association with
166 smoking and 3 showed a negative association, OR ranged from 0.22 to 10.16. When we
167 pooled the data using a random effects model, the pooled OR was 1.26 (95% CI of 0.86 –
168 1.85) - see **Figure 2**. There was evidence of high heterogeneity across studies as suggested by
169 the *I-squared* statistic ($I^2 = 74.3\%$). Visual inspection of the funnel plot (**Online appendix 3**)
170 and the Begg test (p-value = 0.72) did not suggest publication bias (i.e. no evidence of small
171 negative unpublished studies) but showed a cluster of medium to large negative studies.
172 Trying to correct for eventual small unpublished negative studies using the “trim and fill”
173 method in STATA⁵⁰ did not significantly change the results (OR 1.26, 95% CI 0.86-1.83).
174 However, when restricting the analysis to studies of highest quality (i.e with “Moderate
175 quality” using the GRADE criteria for prospective studies) the pooled OR was 1.73 (95% CI
176 1.48-2.03) with no evidence of heterogeneity ($I^2 = 0\%$) (**Figure 3**). In stratified analyses,
177 studies with higher quality, better level of adjustment and longer follow-up showed a stronger

178 positive association between smoking and diabetic peripheral neuropathy (**Table 2**). Studies
179 including people with type 1 diabetes showed increased risks of diabetic peripheral
180 neuropathy for smokers compared with non-smokers whereas studies with people with type 2
181 diabetes did not show a statistically significant association between diabetic peripheral
182 neuropathy and smoking.

183

184 **Smoking and prevalence of diabetic peripheral neuropathy in cross sectional studies**

185 The main individual characteristics of the cross-sectional studies are shown in **Table 3**. They
186 include 27,594 participants in total; 21 studies include people with type 2 diabetes, 3 with
187 type 1 diabetes and 4 with both type 1 and type 2 diabetes. Mean age of participants ranged
188 from 19 to 68 years old and mean diabetes duration ranged from 0 to 20 years. There was a
189 high heterogeneity in the definition of exposure: 7 studies compared current smokers to non
190 smokers (i.e. former and never smokers), 4 studies compared ever smokers (i.e. current and
191 former smokers) to never smokers, 6 studies compared current vs. never smokers, 2 studies
192 compared smokers of 30 or more pack-years to smokers of less than 30 pack-years, one study
193 compared smokers of < 20 pack years to never smokers and 8 studies did not specify the
194 comparison groups. The majority of studies expressed the estimate in OR, two used RR and 9
195 used number or proportions, allowing us to calculate unadjusted OR and 95% CI manually.
196 Seven studies controlled for at least A1C and diabetes duration, one adjusted for either A1C
197 or diabetes duration, 4 adjusted for some confounders but not A1C and diabetes duration and
198 16 did not adjust for potential confounders.- see **Online appendix 4**. Using the GRADE
199 criteria, most studies were rated as “Low quality” and 4 studies were downgraded as “Very
200 low”^{25,28-30} mainly because of selection bias, lack of adjustment for confounders and poorly
201 defined exposure and/or outcome. The majority of studies showed increased odds of
202 neuropathy for smokers compared with non-smokers and ORs ranged from 0.68 to 8.20. The

203 pooled OR using a random effects model was 1.42 (95% CI of 1.21 – 1.65) - see **Figure 4**.
204 There was evidence of some heterogeneity between studies ($I^2 = 64.5\%$). There was no
205 evidence of publication bias as suggested by both visual inspection of the funnel plots
206 (**Online appendix 5**) and the Begg test (p-value = 0.17). In stratified analyses, studies with
207 higher level of adjustment, studies which included participants with type 1 diabetes and those
208 comparing ever vs. never smokers showed a higher and stronger association between
209 smoking and diabetic peripheral neuropathy. (**Table 2**) When analyses were restricted to
210 studies with the highest available level of evidence (i.e. “Low quality” using the GRADE
211 criteria) the pooled OR was 1.29 (95% CI 1.13-1.48; $I^2=50.5\%$). When analyzing only studies
212 which provided adjusted estimates OR was 1.59 (95% CI 1.28-1.97, $I^2=43.4\%$).

213

214

215 **Conclusions**

216 In summary, we found a positive association between smoking and diabetic peripheral
217 neuropathy prevalence and incidence. Although we performed two different meta-analyses,
218 both had similar findings and suggest that smoking is associated with an increased risk of
219 peripheral neuropathy among people with diabetes. There was substantial heterogeneity for
220 both prospective and cross-sectional meta-analyses. However, in stratified analyses, studies
221 with higher level of adjustment, longer follow-up (prospective studies only) and good level of
222 quality showed a stronger positive association between smoking and diabetic peripheral
223 neuropathy with less heterogeneity. Prospective studies comparing ever smokers (current and
224 former smokers) vs. never smokers as well as those including participants with type 1
225 diabetes showed a stronger positive association between smoking and diabetic peripheral
226 neuropathy. However these studies were of higher quality and might not necessarily reflect a
227 real effect modification.

228

229 In people without diabetes, cigarette smoking has been shown to be positively associated
230 with A1C, a surrogate for metabolic control which reflects average glycemia over the past
231 months⁵¹. A previous meta-analysis has shown that smokers have a 44% increased risk of
232 developing type 2 diabetes compared with non smokers⁵². Among people with diabetes prior
233 studies suggest that smoking is also associated with insulin resistance⁵³, higher insulin
234 needs^{54,55} and thus poor metabolic control⁵⁶⁻⁶¹. As microvascular complications in people
235 with type 1 or type 2 diabetes are highly linked to metabolic control^{62,63}, A1C probably acts
236 as a mediator in the relationship between smoking and diabetic peripheral neuropathy.
237 However, the fact that the association remains positive after adjustment for A1C suggests that
238 hyperglycemia may not entirely mediate this relationship. Furthermore, smoking is associated
239 with oxidative stress, systemic inflammation and endothelial dysfunction independent of
240 diabetes⁶⁴⁻⁶⁶. It might increase the risk of nerve damage through these pathways in parallel to
241 metabolic factors. Smoking may also have direct toxic effects and induce diabetic peripheral
242 neuropathy via hypoxemia and microvascular insufficiency. Similar to what occurs with
243 larger vessels (coronary arteries), smaller arteries, including vasa nervorum, might be
244 damaged by smoking which, in turn, leads to development and progression of diabetic
245 peripheral neuropathy. Smoking has been found to be a causal variable in other microvascular
246 complications such as retinopathy or nephropathy, and similar mechanisms might occur for
247 diabetic peripheral neuropathy to damage those target organs⁶⁷. Finally, confounding factors
248 might also contribute to the association between smoking and diabetic peripheral neuropathy.
249 Smokers might have poorer adherence to recommended self-care compared with non-
250 smokers⁶⁸. Smokers also tend to accumulate unhealthy behaviors such as alcohol abuse, lack
251 of physical activity or a diet rich in fat and poor in fruits and vegetables⁶⁹. Though these

252 factors might contribute to diabetes complications through poorer diabetes control, they do
253 not entirely explain the association, which remains after adjustment for diabetes control.

254

255 Our study has several strengths. We retrieved and pooled a substantial number of studies,
256 assessing the association between smoking and diabetic peripheral neuropathy. Contrary to
257 other microvascular complications such as nephropathy or retinopathy, few studies had to
258 date shown a clear positive association between smoking and diabetic peripheral neuropathy.
259 Indeed, few studies have been directly designed to measure the impact of smoking on
260 diabetic peripheral neuropathy and the complex, multifactorial pathogenesis of diabetic
261 peripheral neuropathy makes it difficult to measure the effect of smoking on a unique
262 outcome. Many prospective studies and some cross-sectional studies, included in our meta-
263 analysis, provided adjusted estimates which permitted control for some potential confounders
264 and exploration of mediating factors.

265

266 Our study has several limitations including the relatively small number of prospective studies
267 and the heterogeneity between studies. Stratified analyses allowed us to address the source of
268 heterogeneity but due to limited number of prospective studies, some conclusions cannot be
269 drawn. For example, studies including participants with type 1 diabetes were of higher
270 quality, and it is thus difficult to conclude that the association between smoking and diabetic
271 peripheral neuropathy is significant only among people with type 1 diabetes and not in those
272 with type 2 diabetes. Another limitation is that the cross-sectional studies were of medium to
273 poor quality. Some did not adjust for the main confounders, some did not assess the outcome
274 clinically, and the smoking exposure was very variable between studies. Finally, we cannot
275 prove that the association we observed is causal because of the design and limitations of our
276 selected studies.

277

278 Few studies have prospectively assessed the impact of smoking cessation on diabetes control
279 and complications. We identified only one study which prospectively assessed the impact of
280 smoking cessation on diabetic peripheral neuropathy⁷⁰. Among 193 participants newly
281 diagnosed with type 2 diabetes and microalbuminuria, 62% had quit smoking at 12 months.
282 In this population, the prevalence of diabetic peripheral neuropathy decreased significantly
283 more in participants who quit smoking compared to those who continued ($p < 0.04$), but no
284 absolute numbers were given. This was also the case for microalbuminuria, peripheral
285 vascular disease, blood pressure and dyslipidemia. This unique study of suboptimal quality
286 suggests that the effect of smoking on diabetic peripheral neuropathy might be reversible, but
287 more research is needed to assess the effect of smoking cessation on diabetes control and
288 micro-vascular complications.

289 In conclusion, smoking might be associated with an increased risk of developing diabetic
290 peripheral neuropathy. This is an important finding as this exposure is a modifiable behavior
291 to be targeted in clinical practice, as recommended in diabetes guidelines⁷¹. Even though we
292 cannot exclude the possibility of confounding, it seems reasonable based on the consistency
293 of results and plausibility of biological hypothesis that the association might be causal or at
294 least of concern for clinicians. Future research should be focused on evaluating the impact of
295 smoking cessation on improvement of diabetic neuropathy, helping to establish a causal link
296 between exposure and outcome.

Author's contributions:

CC contributed to the study conception and design, the data research, extraction, led the analyses, and drafted the manuscript, MJC and FE contributed to the study conception and design, the data research, extraction, the interpretation of data and the drafting of the manuscript as well as its review, KJS contributed to the data research, extraction, the interpretation of data and reviewed and edited the manuscript. NAR contributed to the study conception, interpretation of data and critically reviewed the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Parts of the data of this manuscript have been presented as an oral presentation at the 35th annual meeting of the Society of General Internal Medicine (SGIM) in May 2012.

Financial support:

CC was supported by a grant from the Swiss National Science Foundation PBLAP3-127728/1 and by a grant from the SICPA foundation.

Conflict of interest:

NR has been an unpaid consultant to Pfizer and Alere Wellbeing and receives royalties from UpToDate for chapters on smoking cessation. All other authors declare that they do not have a conflict of interest

References

1. Cameron NE, Eaton SE, Cotter MA, Tesfaye S. Vascular factors and metabolic interactions in the pathogenesis of diabetic neuropathy. *Diabetologia* 2001;44:1973-88.
2. Tesfaye S, Stevens LK, Stephenson JM, et al. Prevalence of diabetic peripheral neuropathy and its relation to glycaemic control and potential risk factors: the EURODIAB IDDM Complications Study. *Diabetologia* 1996;39:1377-84.
3. Vinik AI, Park TS, Stansberry KB, Pittenger GL. Diabetic neuropathies. *Diabetologia* 2000;43:957-73.
4. Vincent AM, Calabek B, Roberts L, Feldman EL. Biology of diabetic neuropathy. *Handbook of clinical neurology* 2013;115:591-606.
5. Tesfaye S, Chaturvedi N, Eaton SE, et al. Vascular risk factors and diabetic neuropathy. *N Engl J Med* 2005;352:341-50.
6. Maser RE, Steenkiste AR, Dorman JS, et al. Epidemiological correlates of diabetic neuropathy. Report from Pittsburgh Epidemiology of Diabetes Complications Study. *Diabetes* 1989;38:1456-61.
7. Mitchell BD, Hawthorne VM, Vinik AI. Cigarette smoking and neuropathy in diabetic patients. *Diabetes Care* 1990;13:434-7.
8. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Ottawa Hospital Research Institute. (Accessed February 16, 2015, at http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp.)
9. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924-6.
10. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-88.
11. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539-58.
12. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557-60.
13. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50:1088-101.
14. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629-34.
15. Lehtinen JM, Niskanen L, Hyvonen K, Siitonen O, Uusitupa M. Nerve function and its determinants in patients with newly-diagnosed type 2 (non-insulin-dependent) diabetes mellitus and in control subjects--a 5-year follow-up. *Diabetologia* 1993;36:68-72.
16. Adler AI, Boyko EJ, Ahroni JH, Stensel V, Forsberg RC, Smith DG. Risk factors for diabetic peripheral sensory neuropathy. Results of the Seattle Prospective Diabetic Foot Study. *Diabetes Care* 1997;20:1162-7.
17. Forrest KY, Maser RE, Pambianco G, Becker DJ, Orchard TJ. Hypertension as a risk factor for diabetic neuropathy: a prospective study. *Diabetes* 1997;46:665-70.
18. Sands ML, Shetterly SM, Franklin GM, Hamman RF. Incidence of distal symmetric (sensory) neuropathy in NIDDM. The San Luis Valley Diabetes Study. *Diabetes Care* 1997;20:322-9.
19. Christen WG, Manson JE, Bubes V, Glynn RJ. Risk factors for progression of distal symmetric polyneuropathy in type 1 diabetes mellitus. Sorbinil Retinopathy Trial Research Group. *Am J Epidemiol* 1999;150:1142-51.
20. Sibal L, Law HN, Gebbie J, Home P. Cardiovascular risk factors predicting the development of distal symmetrical polyneuropathy in people with type 1 diabetes: A 9-year follow-up study. *Ann N Y Acad Sci* 2006;1084:304-18.
21. Gerrits EG, Lutgers HL, Kleefstra N, et al. Skin autofluorescence: a tool to identify type 2 diabetic patients at risk for developing microvascular complications. *Diabetes Care* 2008;31:517-21.
22. Elliott J, Tesfaye S, Chaturvedi N, et al. Large-fiber dysfunction in diabetic peripheral neuropathy is predicted by cardiovascular risk factors. *Diabetes Care* 2009;32:1896-900.

23. Uruska A, Araszkiwicz A, Uruski P, Zozulinska-Ziolkiewicz D. Higher risk of microvascular complications in smokers with type 1 diabetes despite intensive insulin therapy. *Microvasc Res* 2014;92:79-84.
24. Franklin GM, Shetterly SM, Cohen JA, Baxter J, Hamman RF. Risk factors for distal symmetric neuropathy in NIDDM. The San Luis Valley Diabetes Study. *Diabetes Care* 1994;17:1172-7.
25. Gregory R, Tattersall RB, Allison SP. Peripheral neuropathy as a presenting feature of type 2 diabetes: a case-controlled study. *Diabet Med* 1994;11:407-9.
26. Matsumoto T, Ohashi Y, Yamada N, Kikuchi M. Hyperglycemia as a major determinant of distal polyneuropathy independent of age and diabetes duration in patients with recently diagnosed diabetes. *Diabetes Res Clin Pract* 1994;26:109-13.
27. Zafra Mezcuca JA, Mendez Segovia JC, Novalbos Ruiz JP, Costa Alonso MJ, Failde Martinez I. [Chronic complications in patients with type 2 diabetes mellitus cared for at a health center]. *Aten Primaria* 2000;25:529-35.
28. Barbosa AP, Medina JL, Ramos EP, Barros HP. Prevalence and risk factors of clinical diabetic polyneuropathy in a Portuguese primary health care population. *Diabetes Metab* 2001;27:496-502.
29. Gomez-Viera N, Soto-Lavastida A, Rosello-Silva H, Gomez de Molina-Iglesias M. [Risk factors involved in symmetrical distal diabetic neuropathy]. *Rev Neurol* 2001;32:806-12.
30. Tapp RJ, Shaw JE, de Courten MP, Dunstan DW, Welborn TA, Zimmet PZ. Foot complications in Type 2 diabetes: an Australian population-based study. *Diabet Med* 2003;20:105-13.
31. Boru UT, Alp R, Sargin H, et al. Prevalence of peripheral neuropathy in type 2 diabetic patients attending a diabetes center in Turkey. *Endocr J* 2004;51:563-7.
32. Tamer A, Yildiz S, Yildiz N, et al. The prevalence of neuropathy and relationship with risk factors in diabetic patients: a single-center experience. *Medical principles and practice : international journal of the Kuwait University, Health Science Centre* 2006;15:190-4.
33. Al-Mahroos F, Al-Roomi K. Diabetic neuropathy, foot ulceration, peripheral vascular disease and potential risk factors among patients with diabetes in Bahrain: a nationwide primary care diabetes clinic-based study. *Annals of Saudi medicine* 2007;27:25-31.
34. Cho HC. The Association between Serum GGT Concentration and Diabetic Peripheral Polyneuropathy in Type 2 Diabetic Patients. *Korean diabetes journal* 2010;34:111-8.
35. Jianbo L, Yuche C, Ming S, et al. Association of homocysteine with peripheral neuropathy in Chinese patients with type 2 diabetes. *Diabetes Res Clin Pract* 2011;93:38-42.
36. Spallone V, Morganti R, D'Amato C, et al. Clinical correlates of painful diabetic neuropathy and relationship of neuropathic pain with sensorimotor and autonomic nerve function. *European Journal of Pain* 2011;15:153-60.
37. Wang W, Balamurugan A, Biddle J, Rollins KM. Diabetic neuropathy status and the concerns in underserved rural communities: challenges and opportunities for diabetes educators. *Diabetes Educ* 2011;37:536-48.
38. Abougalambou SS, Abougalambou AS. Explorative study on diabetes neuropathy among type II diabetic patients in Universiti Sains Malaysia Hospital. *Diabetes & metabolic syndrome* 2012;6:167-72.
39. Ji N, Zhang N, Ren ZJ, et al. Risk factors and pain status due to diabetic neuropathy in chronic long-term diabetic patients in a Chinese urban population. *Chin Med J (Engl)* 2012;125:4190-6.
40. Katulanda P, Ranasinghe P, Jayawardena R, Constantine GR, Sheriff MH, Matthews DR. The prevalence, patterns and predictors of diabetic peripheral neuropathy in a developing country. *Diabetology & metabolic syndrome* 2012;4:21.
41. Rasul S, Ilhan A, Wagner L, Luger A, Kautzky-Willer A. Diabetic polyneuropathy relates to bone metabolism and markers of bone turnover in elderly patients with type 2 diabetes: greater effects in male patients. *Gender medicine* 2012;9:187-96.
42. Eleftheriadou I, Grigoropoulou P, Mourouzis I, et al. Serum osteoprotegerin levels are associated with peripheral neuropathy in patients with type 2 diabetes. *Diabetologia* 2013;56:S496.

43. Molina M, Gonzalez R, Folgado J, et al. [Correlation between plasma concentrations of homocysteine and diabetic polyneuropathy evaluated with the Semmes-Weinstein monofilament test in patients with type 2 diabetes mellitus]. *Med Clin (Barc)* 2013;141:382-6.
44. Aubert CE, Michel PL, Gillery P, et al. Association of peripheral neuropathy with circulating advanced glycation end products, soluble receptor for advanced glycation end products and other risk factors in patients with type 2 diabetes. *Diabetes Metab Res Rev* 2014;30:679-85.
45. Bener A, Al-Laftah F, Al-Hamaq AO, Daghsh M, Abdullatef WK. A study of diabetes complications in an endogamous population: an emerging public health burden. *Diabetes & metabolic syndrome* 2014;8:108-14.
46. Brownrigg JR, de Lusignan S, McGovern A, et al. Peripheral neuropathy and the risk of cardiovascular events in type 2 diabetes mellitus. *Heart* 2014;100:1837-43.
47. Hu Y, Liu F, Shen J, et al. Association between serum cystatin C and diabetic peripheral neuropathy: a cross-sectional study of a Chinese type 2 diabetic population. *Eur J Endocrinol* 2014;171:641-8.
48. Jaiswal M, Divers J, Isom S, et al. Prevalence and clinical correlates of diabetic peripheral neuropathy among youth with type 1 diabetes: Search for diabetes in youth cohort study. *Diabetes* 2014;63:A148-A9.
49. Wang DD, Bakhotmah BA, Hu FB, Ali Alzahrani H. Prevalence and correlates of diabetic peripheral neuropathy in a Saudi Arabic population: A cross-sectional study. *PloS one* 2014;9.
50. Duval S, Tweedie R. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000;56:455-63.
51. Clair C, Bitton A, Meigs JB, Rigotti NA. Relationships of cotinine and self-reported cigarette smoking with hemoglobin A1c in the U.S.: results from the National Health and Nutrition Examination Survey, 1999-2008. *Diabetes Care* 2011;34:2250-5.
52. Willi C, Bodenmann P, Ghali WA, Faris PD, Cornuz J. Active smoking and the risk of type 2 diabetes: a systematic review and meta-analysis. *JAMA* 2007;298:2654-64.
53. Targher G, Alberiche M, Zenere MB, Bonadonna RC, Muggeo M, Bonora E. Cigarette smoking and insulin resistance in patients with noninsulin-dependent diabetes mellitus. *J Clin Endocrinol Metab* 1997;82:3619-24.
54. Madsbad S, McNair P, Christensen MS, et al. Influence of smoking on insulin requirement and metabolic status in diabetes mellitus. *Diabetes Care* 1980;3:41-3.
55. Bott U, Jorgens V, Grusser M, Bender R, Muhlhauser I, Berger M. Predictors of glycaemic control in type 1 diabetic patients after participation in an intensified treatment and teaching programme. *Diabet Med* 1994;11:362-71.
56. Melin EO, Thunander M, Svensson R, Landin-Olsson M, Thulesius HO. Depression, obesity, and smoking were independently associated with inadequate glycemic control in patients with type 1 diabetes. *Eur J Endocrinol* 2013;168:861-9.
57. Gunton JE, Davies L, Wilmshurst E, Fulcher G, McElduff A. Cigarette smoking affects glycemic control in diabetes. *Diabetes Care* 2002;25:796-7.
58. Kapoor D, Jones TH. Smoking and hormones in health and endocrine disorders. *Eur J Endocrinol* 2005;152:491-9.
59. Nilsson PM, Gudbjornsdottir S, Eliasson B, Cederholm J. Smoking is associated with increased HbA1c values and microalbuminuria in patients with diabetes--data from the National Diabetes Register in Sweden. *Diabetes Metab* 2004;30:261-8.
60. Sargeant LA, Khaw KT, Bingham S, et al. Cigarette smoking and glycaemia: the EPIC-Norfolk Study. *European Prospective Investigation into Cancer. Int J Epidemiol* 2001;30:547-54.
61. Lundman BM, Asplund K, Norberg A. Smoking and metabolic control in patients with insulin-dependent diabetes mellitus. *J Intern Med* 1990;227:101-6.
62. Fullerton B, Jeitler K, Seitz M, Horvath K, Berghold A, Siebenhofer A. Intensive glucose control versus conventional glucose control for type 1 diabetes mellitus. *Cochrane Database Syst Rev* 2014;2:CD009122.

63. Hemmingsen B, Lund SS, Gluud C, et al. Targeting intensive glycaemic control versus targeting conventional glycaemic control for type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2013;11:CD008143.
64. Winkelmann BR, Boehm BO, Nauck M, et al. Cigarette smoking is independently associated with markers of endothelial dysfunction and hyperinsulinaemia in non-diabetic individuals with coronary artery disease. *Curr Med Res Opin* 2001;17:132-41.
65. Burke A, Fitzgerald GA. Oxidative stress and smoking-induced vascular injury. *Prog Cardiovasc Dis* 2003;46:79-90.
66. Benowitz NL. Cigarette smoking and cardiovascular disease: pathophysiology and implications for treatment. *Prog Cardiovasc Dis* 2003;46:91-111.
67. Eliasson B. Cigarette smoking and diabetes. *Prog Cardiovasc Dis* 2003;45:405-13.
68. Yu MK, Lyles CR, Bent-Shaw LA, Young BA. Sex disparities in diabetes process of care measures and self-care in high-risk patients. *Journal of diabetes research* 2013;2013:575814.
69. Chiolerio A, Wietlisbach V, Ruffieux C, Paccaud F, Cornuz J. Clustering of risk behaviors with cigarette consumption: A population-based survey. *Prev Med* 2006;42:348-53.
70. Voulgari C, Katsilambros N, Tentolouris N. Smoking cessation predicts amelioration of microalbuminuria in newly diagnosed type 2 diabetes mellitus: a 1-year prospective study. *Metabolism* 2011;60:1456-64.
71. Standards of medical care in diabetes--2014. *Diabetes Care* 2014;37 Suppl 1:S14-80.

Tables and figures legends:

Table 1: Characteristics of prospective studies included in the Meta-analysis

Table 2: Stratified analyses for prospective studies and cross-sectional studies

Table 3: Characteristics of cross-sectional studies included in the Meta-analysis

Figure 1: Study flow diagram

Figure 2: Forest plot for prospective studies: Adjusted Odds ratios of Neuropathy for Smokers Compared with Nonsmokers

Figure 3: Forest plot for prospective studies of highest quality (“moderate quality” using the GRADE criteria): Adjusted Odds ratios of Neuropathy for Smokers Compared with Nonsmokers

Figure 4: Forest plot for cross-sectional studies: Adjusted Odds Ratios of Neuropathy for Smokers Compared with Nonsmokers

Online appendix files

Online appendix 1: Electronic search strategy

Online appendix 1: Level of adjustment and quality of prospective studies

Online appendix 2: Funnel plot for prospective studies

Online appendix 3: Level of adjustment and quality of cross-sectional studies

Online appendix 4: Funnel plot for cross-sectional studies

Table 1: Characteristics of prospective studies included in the Meta-analysis

Author, year	Country/ region	Sample size	Population	Type of DM	% men	Mean age	Mean DM duration	Neuropathy screening	Smoking comparison	FUP y	Estimate
Lehtinen et al., 1993	Finland	113	Subjects with newly diagnosed DM from the community	2	51	56.4	0	Nerve conduction velocities	NS	5	N
Adler et al., 1997	USA	387	US veterans followed in an outpatient clinic	Both	96	61.7	9.8	Monofilament examination	Current vs. former + never smokers	2.6	OR
Forrest et al., 1997	USA	453	Subjects with childhood-onset DM	1	49	25.1	16.9	Neurological examination	Current + former vs. never smokers	5.3	RR
Sands et al., 1997	USA	231	Biethnic population in Colorado	2	NS	NS	NS	Neurological examination and history	Current + former vs. never smokers	4.7	OR
Christen et al., 1999	USA	407	Participants of a drug (Sorbinil) multi-center trial	1	75	31.4	6.5	Neurological examination and history	Current + former vs. never smokers*	2	RR
Tesfaye et al., 2005	Europe	1172	Subjects randomly selected from 31 diabetes clinics	1	51	30.7	12.4	Neurological examination	Current + former vs. never smokers	7.3	OR
Sibal et al., 2006	UK	334	Outpatients who attended diabetes services	1	54	39	20	Neurological examination and history	NS	9	OR
Gerrits et al.,	Netherlands	973	Subjects from	2	46	66	4	Monofilament	NS	3.1	OR

2008			primary care					examination			
Elliott et al., 2009	Europe	1407	Subjects randomly selected from 31 diabetes clinics	1	48	31.5	13.1	Vibration perception threshold measured by biothesiometers	Current + former vs. never smokers	7.3	OR
Uruska et al., 2014	Poland	81	Patients treated with intensive Insulin from onset of disease	1	63	34	10	Neurological examination (monofilament, vibration, temperature and ankle reflex)	Current + former vs. never smokers	10	OR

DM = diabetes mellitus, NS = non specified, FUP = follow-up, OR=Odds ratio, RR=Relative risk, N=number or proportion

* In the age-adjusted analyses, compared current smokers to never smokers, in the multivariable-adjusted model compares ever to never smokers

Table 2: Stratified analyses for prospective and cross-sectional studies

Stratified analysis	Total number of trials	OR	(95% CI)	<i>I-squared</i>	P value from metareg ⁺
Adjustment for confounding factors					
Prospective studies					0.71
Adjusted for at least HbA1c and DM duration	5	1.47	(1.01-2.13)	71.8%	
Not adjusted for HbA1c and DM duration	5	1.03	(0.34-3.09)	79.2%	
Cross sectional studies					0.31
Adjusted for at least HbA1c and DM duration	7	1.59	(1.23-2.06)	43.6%	
Not adjusted for HbA1c and DM duration	21	1.36	(1.11-1.66)	69.2%	
Type of diabetes					
Prospective studies					0.02
Type 1	6	1.74	(1.48-2.04)	0%	
Type 2	3	0.65	(0.16-2.71)	83.2%	
Both	1	0.22	(0.07-0.66)	--	
Cross sectional studies					0.19
Type 1	3	3.02	(2.03-4.47)	11.7%	
Type 2	21	1.24	(1.08-1.44)	50.5%	
Both	4	1.55	(0.94-2.57)	63.2%	
Smoking exposure					
Prospective studies					0.007
Ever (current + former) vs. never smoker	6	1.77	(1.51-2.08)	0%	
Current vs. never smoker	1	0.22	(0.07-0.66)	--	
Non specified	3	0.47	(0.21-1.06)	31.6%	
Cross sectional studies					0.79
Ever (current + former) vs. never smoker	4	1.78	(1.39-2.29)	10.6%	
Current vs. non smokers (former + never)	7	1.38	(0.87-2.20)	70.0%	

Current vs. never smoker	6	1.58	(1.00-2.48)	74.2%	
Non specified or other definition	11	1.28	(1.03-1.60)	57.2%	
Mean follow-up					
Prospective studies					0.322
< 5 years	4	0.77	(0.25-2.31)	86.4%	
≥ 5 years	6	1.63	(1.21-2.21)	47.7%	
Level of quality (GRADE criteria)					
Prospective studies					0.007
Moderate	6	1.73	(1.48-2.03)	0%	
Low	2	1.57	(0.05-50.83)	91%	
Very low	2	0.27	(0.11-0.65)	0%	
Cross sectional studies					0.75
Low	22	1.29	(1.13-1.48)	50.5%	
Very low	4	1.30	(0.61-2.75)	68.3%	

+ P-value for metaregression using the “metareg” Stata command

Table 3: Characteristics of cross-sectional studies included in the Meta-analysis

Author	Country	Sample size	Population	Type of DM	% men	Mean age	Mean DM duration	Neuropathy screening	Smoking measure	Estimate
Maser et al., 1989 ⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶⁶	USA	363	Cohort of patients with recent diagnosis	1	50	28.4	19.9	Neurological examination and history	Current + former vs. never smokers	OR
Mitchell et al., 1990	USA	214	Patients admitted to the inpatient diabetic clinic of a University hospital	1	37	46	14.7	Neurological examination and history	Smoking \geq 30 vs. < 30 pack-years	OR
Franklin et al., 1994	USA	277	Biethnic population in Colorado	2	43	59.5	9.7	Neurological examination and history	< 20 pack-years vs. never smokers	OR
Gregory et al., 1994	UK	136	Newly diagnosed patients attending a hospital	2	50	68	0	Neurological examination and history	Smoking \geq 30 vs. < 30 pack-years	N
Matsumoto et al., 1994	Japan	742	Outpatients who visited the diabetic unit of a department of internal medicine	2	54	49	1.3	Information from patient's charts and neurological examination t	NS	OR
Zafra Mezcua et al., 2000	Spain	504	Patients attending a medical outpatient clinic	2	42	63.9	8.6	Medical chart review	NS	RR
Barbosa et al., 2001	Portugal	93	Patients from primary health cares	2	40	65.4	10.1	Neurological examination	Current vs. never smokers	N
Gomez-Viera et al.,	Cuba	200	Patients diagnosed in clinic	Both	326	.	.	Clinical diagnosis with	NS	RR

2001								neuroinduction exam corroboration		
Tapp et al., 2003	Australia	821	Population-based survey	2	51	63.1	0.2	Neurological examination and history	Current vs. former + never smokers	OR
Boru et al., 2004	Turkey	866	Patients who attended a diabetic clinic	2	40	57.2	8.5	Neurological examination and history	NS	OR
Tamer et al., 2006	Turkey	191	Patients with type 2 DM recruited	2	43	58.7	.	Neurological examination and electromyography	Current + former vs. never smokers	OR
Al-Mahroos et al., 2007	Bahrain	1477	Patients from specialized clinics	2	43	57.3	9.5	Neurological examination and history	NS	OR
Cho et al., 2010	Korea	90	Patients who underwent work-ups for peripheral polyneuropathy	2	51	59	8.7	Neurological examination and history	NS	OR
Jianbo et al., 2011	China	227	Inpatients and outpatients	2	.	64.5	9.3	Neurologic exam and electromyography	Current vs. never smoker	N
Spallone et al., 2011	Italy	191	Diabetic patients with suspected neuropathic pain referred to a center	Both	56.5	58.6	16.7	History + electrodiagnostic studies in selected cases	NS	OR
Wang et al., 2011	USA	816	Patients referred to a diabetes education program	2	45.2	57	.	Questionnaires and review of medical records	Current + former vs. never	OR

									smokers	
Abougalambou et al., 2012	Malaysia	1077	Patient followed in an outpatient diabetic clinic	2	45.2	.	.	Neurologic exam	Current vs. former+ never smokers	OR
Ji et al., 2012	China	565	Mostly inpatients	2	47.8	66.6	16.2	Medical history and/or symptoms and/or neurological exam	Current vs. never smokers	N
Katulanda et al., 2012	Sri Lanka	337	Non institutionalized adults from the community	2	37.1	56.8	6.3	Symptoms and neurologic exam	Current vs. former + never smokers	OR
Rasul et al., 2012	Austria	120	Patients from an outpatient clinic	2	59.2	62.9	12.7	Neurological exam and nerve conduction velocity	Current vs. never smokers	N
Eleftheradiou et al., 2013 (Abstract)	Greece	71	Patients from an outpatient clinic	2	63.4	67.7	15	Neuropathy symptom and neuropathy disability scores, vibration perception threshold	Current vs. former + never smokers	OR
Molina et al., 2013	Spain	405	Patients from a diabetes clinic and primary care clinic	2	58.3	66	12.7	Semmes-Weinstein monofilament test	Current vs. former + never smokers	N
Aubert et al., 2014	France	198		2	79.8	65	13	Neuropathy disability score or inability to perceive	Current vs. former + never smokers	N

								monofilament		
Bener et al., 2014	Qatar	1633		Both	51.6	45.3	7.3	Not specified	Current vs. former + never smokers	OR
Brownrigg et al., 2014	UK	13043		2	51.8	63.8	.	Semmes-Weinstein monofilament test	Current vs. never smokers	N
Hu et al., 2014	China	937	Diabetic inpatients at a clinical medical center of diabetes	2	57.7	59.6	9.8	Neurological examination and nerve conduction tests	NS	N
Jaiswal et al., 2014 (Abstract)	USA	1448	Participants to the SEARCH for diabetes in youth study	1	50	19	8	Symptoms and neurologic exam	Current vs. never smokers	OR
Wang et al., 2014	Saudi Arabia	552	People with diabetes from the community	Both	62.7	53.4	.		Current + former vs. never smokers	OR

DM = diabetes mellitus, NS = non specified, FUP = follow-up, OR=Odds ratio, RR=Relative risk, N=number or proportion

Figure 1:






Figure 2




Dashed vertical line represents the estimated pooled effect size estimate; points in grey squares with lines represent odds ratios and 95% CIs of individual studies; the open diamond represent a visual summary of the overall 95% CI of the effect estimate of smoking on the incidence of DPN. Studies on the right of the 1 vertical line indicate a positive association between smoking and DPN, studies on the left a negative association.

Figure 3



Dashed vertical line represents the estimated pooled effect size estimate; points in grey squares with lines represent odds ratios and 95% CIs of individual studies; the open diamond represent a visual summary of the overall 95% CI of the effect estimate of smoking on the incidence of DPN. Studies on the right of the 1 vertical line indicate a positive association between smoking and DPN, studies on the left a negative association.

Figure 4



Dashed vertical line represents the estimated pooled effect size estimate; points in grey squares with lines represent odds ratios and 95% CIs of individual studies; the open diamond represent a visual summary of the overall 95% CI of the effect estimate of smoking on the prevalence of DPN. Studies on the right of the 1 vertical line indicate a positive association between smoking and DPN, studies on the left a negative association.

Online appendix 1

A) Electronic search strategy for PubMed (performed on November 17, 2014):

((polyneuropath*[tiab] OR neuropath*[tiab]) OR ("Peripheral Nervous System Diseases"[Mesh]) OR (((("Diabetic Neuropathies"[Mesh]) OR "Polyneuropathies"[Mesh]) OR "Diabetic Foot"[Mesh])) AND (("Diabetes Mellitus, Type 2"[Mesh] OR "Diabetes Mellitus, Type 1"[Mesh]) OR (diabetes[tiab] OR diabetic*[tiab])) AND (((("Smoking"[Mesh] OR "Smoking Cessation"[Mesh]) OR ("Tobacco Use Cessation"[Mesh] OR "Tobacco Use Disorder"[Mesh] OR "Tobacco, Smokeless"[Mesh] OR "Tobacco"[Mesh])) OR "Nicotine"[Mesh]) OR (smoking[tiab] OR smoker*[tiab] OR cigarett*[tiab] OR tobacco[tiab] OR nicotine[tiab])).

B) Electronic search strategy for Embase (performed on November 17, 2014):

```
'diabetic neuropathy'/exp OR 'peripheral neuropathy'/exp OR 'polyneuropathy'/exp AND [embase]/lim
```

```
'diabetes mellitus'/exp OR 'non insulin dependent diabetes mellitus'/exp OR 'insulin dependent diabetes mellitus'/exp AND [embase]/lim
```

```
'smoking'/exp OR 'smoking cessation'/exp OR 'smokeless tobacco'/exp OR 'tobacco dependence'/exp OR 'tobacco'/exp OR 'nicotine'/exp AND [embase]/lim
```

Online appendix 2: Level of adjustment and quality of prospective studies


Author, year	Adjustment	Quality score (NOS scale)		
		Selection ¹	Comparability ²	Exposure ³
Lehtinen et al., 1993	-	***	-	**
Adler et al., 1997	HbA1c, height, history of ulcer, age, alcohol, albumin, creatinin	*	*	*
Forrest et al., 1997	HbA1c, DM duration, height, hypertension	****	**	***
Sands et al., 1997	HbA1c, DM duration, age, insulin treatment, ethnicity, gender, history of myocardial infarct, angina	****	**	***
Christen et al., 1999	HbA1c, age, drug (Sorbinil vs plabebo), height, gender	****	*	***
Tesfaye et al., 2005	HbA1c, DM duration	****	*	***
Sibal et al., 2006	DM duration, systolic blood pressure, diastolic blood pressure, total cholesterol, glomerular filtration rate, triglycerides, microalbuminuria, retinopathy, foot complications	****	*	***
Gerrits et al., 2008	HbA1c, DM duration, sex, systolic blood pressure, LDL cholesterol, HDL cholesterol, triglycerides, body-mass index, skin autofluorescence	****	**	***
Elliott et al., 2009	HbA1C, DM duration, hypertension, body-mass index, retinopathy, cardiovascular disease history, cholesterol, triglycerides, albumin	****	*	***
Uruska et al., 2014	HbA1c, age, sexe, hypertension	***	*	*

HbA1c= glycated heamoglobin, DM=diabetes mellitus

NOS scale = Newcastle Ottawa scale

¹ Selection : minimum = -, maximum = ****, ² Comparability: minimum = -, maximum = **, ³ Exposure: minimum = -, maximum = ***

Online appendix 3



Cho et al., 2010	Retinopathy, systolic blood pressure, diastolic blood pressure, gamma-glutamyl transferase, Aspartate amino transferase, urine albumine/creatinie, C-reactive protein	*	*	*
Jianbo et al., 2011	-	***	-	*
Spallone et al., 2011	HbA1c, DM duration, gender, age, systolic blood pressure, triglycerides, peripheral arterial disease, Valsalva ratio, type of DM, body-mass index, waist circumference , Michigan Diabetic Neuropathy Score	*	**	*
Wang et al., 2011	DM duration, sex, race, education, self foot exam, foot exam by medical doctor	*	*	*
Abougambou et al., 2012	-	***	-	*
Ji et al., 2012	-	***	-	*
Katulanda et al., 2012	-	***	-	*
Rasul et al., 2012	-	***	-	*
Eleftheradiou et al., 2013 (abstract)	Sex, age, body-mass index			
Molina et al., 2013	-	**	-	**
Aubert et al., 2014	-	***	-	**
Bener et al., 2014	-	**	-	*
Brownrigg et al., 2014	-	****	-	**
Jaiswal et al., 2014 (abstract)	-			
Hu et al., 2014	-	**	-	**

Wang et al., 2014	Sex, age, nationality	***	*	*
----------------------	-----------------------	-----	---	---

HbA1c=
heamoglobin, DM=diabetes mellitus, NA= not applicable

glycated

NOS scale = Newcastle Ottawa scale

¹ Selection : minimum = -, maximum = ***, ² Comparability: minimum = -, maximum = **, ³ Exposure: minimum = -, maximum = **

