

Editorial Manager(tm) for Anesthesiology
Manuscript Draft

Manuscript Number: 200802049R3

Title: Activation of p38 Mitogen-activated Protein Kinase in Spinal Microglia Contributes to Incision-induced Mechanical Allodynia

Article Type: Laboratory Investigation

Section/Category:

Corresponding Author: Dr. Chia-Chuan Wang, Ph.D.

Corresponding Author's Institution: Fu Jen Catholic Universtiy

First Author: Yeong-Ray Wen, M.D.

Order of Authors: Yeong-Ray Wen, M.D.; Marc R Suter, M.D.; Ru-Rong Ji, Ph.D.; Geng-Chang Yeh, M.D., Ph.D.; Yen-Sheng Wu, B.S.; Kuo-Ching Wang, M.D.; Tatsuro Kohno, M.D.; Wei-Zen Sun, M.D.; Chia-Chuan Wang, Ph.D.

Manuscript Region of Origin:

Abstract: ABSTRACT:

Background: Recent studies have implicated the activation of stress-activated mitogen-activated protein kinase (MAPK) p38 in spinal microglial cells for development of neuropathic and inflammatory pain. The aim of the present study was to investigate whether phosphorylation of p38 (p-p38) also mediates mechanical allodynia and thermal hyperalgesia induced by plantar incision.

Methods: After rats received a plantar incision surgery, mechanical allodynia and thermal hyperalgesia were determined by von Frey filaments and radiant heat, respectively, and the number of p-p38 immunoreactive cells in the dorsal horn was quantified to determine p38 activation

at different time points after incision. The p38 inhibitor FR167653 was administered intrathecally 30 min before hindpaw plantar incision to determine the role of p38 in postoperative pain.

Results: A significant increase in number of p-p38 immunoreactive cells was observed in the ipsilateral L4-5 spinal dorsal horn from 1 h to 3 days after the incision. P-p38 was found predominantly in microglia. However, microglial activation (assessed by OX-42 upregulation) was not evident until 3 days after plantar incision. Intrathecal pretreatment of FR167653 attenuated incision-induced mechanical allodynia from 1 h to day 2 and significantly reduced activation of p38 in the dorsal horn 1 day after plantar incision. However, FR167653 only inhibited heat hyperalgesia at an early time point.

Conclusions: Plantar incision-induced mechanical allodynia can be prevented by the p38 inhibitor. Our results suggest that p38 activation in spinal microglia play a role in incision-induced mechanical allodynia in rats. Therefore, p38 inhibition may be useful in treating post-surgical pain.

*** Response to Reviewers (RevisionsOnly)**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

This piece of the submission is being sent via mail.

1
2
3
4
5 **Activation of p38 Mitogen-Activated Protein Kinase in Spinal**
6
7
8 **Microglia Contributes to Incision-Induced Mechanical Allodynia**
9

10
11
12 Yeong-Ray Wen, M.D.,*†‡ Marc R. Suter, M.D.,§ Ru-Rong Ji, Ph.D.,|| Geng-Chang Yeh, M.D.,
13
14
15 Ph.D.,# Yen-Sheng Wu, B.S.,** Kuo-Ching Wang, M.D.,††. Tatsuro Kohno, M.D.,‡‡ Wei-Zen
16
17
18 Sun, M.D.,§§ and Chia-Chuan Wang, Ph.D.##
19
20
21
22
23

24 * Student, # Professor, Graduate Institute of Medical Sciences, † Assistant Professor, School of
25
26
27 Medicine, Taipei Medical University, Taipei, Taiwan; ‡ †† Staff Anesthesiologist, Department
28
29
30 of Anesthesiology, Shin-Kong Wu Ho-Su Memorial Hospital, Taipei, Taiwan; § Research
31
32
33 Fellow, || Associate Professor, Pain Research Center, Department of Anesthesiology, Brigham
34
35
36 and Women’s Hospital and Harvard Medical School, Boston, Massachusetts, USA; ‡‡
37
38
39 Associate Professor, Department of Anesthesiology, Medical School of Niigata University,
40
41
42 Japan; §§ Professor, Department of Anesthesiology, National Taiwan University Hospital and
43
44
45 College of Medicine, National Taiwan University, Taipei, Taiwan; ** Research Assistant, ##
46
47
48 Assistant Professor, School of Medicine, Fu Jen Catholic University, Taipei County, Taiwan
49
50
51
52
53

54 Dr. Sun and Wang had equal contributions as corresponding authors in this study.
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 Corresponding author: Chia-Chuan Wang, Ph.D.
6

7 School of Medicine, Fu Jen Catholic University
8

9
10 510 Chung-Cheng Rd., Hsin-Chuang
11

12
13 Taipei County 24205, Taiwan.
14

15
16 Tel: +886-2-29053466
17

18
19 Fax: +886-2-29052096
20

21
22 E-mail: 050814@mail.fju.edu.tw
23
24
25
26

27 **Attributed Department and Institution:**
28

29
30 Graduate Institute of Medical Sciences, School of Medicine, Taipei Medical University, Taipei,
31

32
33 Shin-Kong Wu Ho-Hsu Memorial Hospital, Taipei, and College of Medicine, Fu Jen Catholic
34

35
36 University, Taipei County, Taiwan.
37
38
39
40
41

42 **Acknowledgements**
43

44
45 The authors wish to thank Chih-Chun Ma, B.S. (Department of Anesthesiology, Shin-Kong Wu
46

47
48 Ho-Su Memorial Hospital, Taipei, Taiwan) for technical assistance, Chyi-Huey Bai, Ph.D.
49

50
51 (Central Laboratory, Shin-Kong Wu Ho-Su Memorial Hospital, Taipei, Taiwan) for assistance
52

53
54 in the statistical analysis, and Robert W. Gereau IV, Associate Professor (Washington
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 University Pain Center and Department of Anesthesiology, Washington University School of
6
7
8 Medicine, St. Louis, Missouri) for critically reading this manuscript.
9

10
11
12
13
14 **Support Statement**

15
16 This study is supported by Research Grant Nos. SKH-FJU-95-22 (to Y.-R. Wen) and
17
18
19 SKH-FJU-95-17 (to C.-C. Wang) from Shin-Kong Wu Ho-Su Memorial Hospital and Fu Jen
20
21
22 Catholic University, Taipei County, Taiwan and NSC95-2320-B-030-009 (to C.-C. Wang) from
23
24
25 National Science Council, Taipei, Taiwan.
26
27
28
29
30

31
32 **Abbreviated Title**

33
34 p38 activation and postsurgical pain
35
36
37
38
39

40
41 **Summary Statement**

42
43 Incision-induced mechanical allodynia is partially attributed to p38 activation in spinal
44
45
46 microglia and can be prevented by p38 inhibitor. These suggest p38 activation
47
48
49 participates in the development of incision-induced mechanical allodynia in rats.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7 **Abstract**
8

9 *Background:* Recent studies have implicated the activation of stress-activated
10 mitogen-activated protein kinase (MAPK) p38 in spinal microglial cells for development of
11 neuropathic and inflammatory pain. The aim of the present study was to investigate whether
12 phosphorylation of p38 (p-p38) also mediates mechanical allodynia and thermal hyperalgesia
13 induced by plantar incision.
14
15
16
17
18
19
20
21
22

23 *Methods:* After rats received a plantar incision surgery, mechanical allodynia and
24 thermal hyperalgesia were determined by von Frey filaments and radiant heat, respectively, and
25 the number of p-p38 immunoreactive cells in the dorsal horn was quantified to determine p38
26 activation at different time points after incision. The p38 inhibitor FR167653 was administered
27 intrathecally 30 min before hindpaw plantar incision to determine the role of p38 in
28 postoperative pain.
29
30
31
32
33
34
35
36
37
38
39
40
41

42 *Results:* A significant increase in number of p-p38 immunoreactive cells was observed in
43 the ipsilateral L4-5 spinal dorsal horn from 1 h to 3 days after the incision. P-p38 was found
44 predominantly in microglia. However, microglial activation (assessed by OX-42 upregulation)
45 was not evident until 3 days after plantar incision. Intrathecal pretreatment of FR167653
46 attenuated incision-induced mechanical allodynia from 1 h to day 2 and significantly reduced
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

activation of p38 in the dorsal horn 1 day after plantar incision. However, FR167653 only inhibited heat hyperalgesia at an early time point.

Conclusions: Plantar incision-induced mechanical allodynia can be prevented by the p38 inhibitor. Our results suggest that p38 activation in spinal microglia play a role in incision-induced mechanical allodynia in rats. Therefore, p38 inhibition may be useful in treating post-surgical pain.

1
2
3
4
5 **Introduction**
6

7
8 Postoperative pain results from surgical injuries in skin, fascia, muscle, and small nerves
9
10 innervating these tissues and is a common form of acute pain in clinics. Despite increased basic
11
12 and clinical research and improved understanding of pathological mechanisms, optimal
13
14 postsurgical pain therapy remains a challenge for physicians. To investigate the underlying
15
16 mechanisms of postsurgical pain, Brennan et al¹ have developed a rat model of incisional pain
17
18 which is characterized by increased mechanical sensitivity after a surgical incision made in the
19
20 plantar aspect of rat hindpaw. This model has been shown to possess a battery of nociceptive
21
22 responses that parallel the time course of postoperative pain in human^{1;2}.
23
24
25
26
27
28
29
30

31
32 Several studies based on this surgical pain model have suggested discrepancies between
33
34 surgical incision and other pathological pain models³⁻⁵. For example, surgical incision-induced
35
36 central sensitization is more likely mediated by activation of non-*N*-methyl-D-aspartate,
37
38 metabotropic glutamate receptor and neurokinin-1 receptors in spinal dorsal horn^{4;6;7}, but not
39
40 by *N*-methyl-D-aspartate receptors^{5;8} which play an important role in inflammatory
41
42 hyperalgesia⁹. Therefore, our knowledge of inflammatory or neuropathic pain may not apply to
43
44 postsurgical pain conditions.
45
46
47
48
49
50

51
52 Mitogen-activated protein kinases (MAPKs) are activated through various upstream
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 protein kinases and transduce diverse extracellular stimuli into intracellular post-translational
6
7 and transcriptional responses¹⁰. The members of the MAPK family, including extracellular
8
9 signal-regulated kinase, c-Jun N-terminal kinase, and p38, have been shown to be critical for
10
11 the development and maintenance of inflammation- and nerve injury-induced pain
12
13 hypersensitivity^{11;12}. p38 is a stress-activated protein kinase and growing evidence suggests that
14
15 activation of p38 MAPK in the spinal cord and dorsal root ganglion contributes to the
16
17 development and maintenance of inflammatory¹³⁻¹⁶ and neuropathic pain^{14;17-20}. However, the
18
19 role of p38 MAPK in incision-induced pain hypersensitivity remains unclear.
20
21
22
23
24
25
26
27

28
29 The present study was designed to investigate whether activation (phosphorylation) of p38
30
31 MAPK in the spinal cord is involved in pain hypersensitivity after a plantar incision in the rat
32
33 hindpaw. Using immunohistochemical and behavioral approaches, we examined the temporal
34
35 correlation between the incision-induced nocifensive behaviors and p38 activation in the spinal
36
37 cord dorsal horn. We also investigated the preventive effects of intrathecal injection of p38
38
39 inhibitor, FR167653, before plantar incision on post-surgical mechanical allodynia and thermal
40
41 hyperalgesia.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 **Materials and Methods**
6

7
8
9 **Animals**

10 Male Sprague-Dawley rats (250-300gm, CD® (SD) IGS, National Laboratory Animal
11 Center, Taiwan) were used. The animals were housed in groups of three to four per cage and
12 acclimatized to the laboratory conditions (12-h light/dark cycle, 22 ± 1 °C room temperature) 1
13 week before experiments. Animals had free access to food and water. All experiments were
14 carried out in accordance with the guidelines established by the Institutional Animal Care and
15 Utilization Committee, Fu Jen Catholic University, Taipei County, Taiwan. Efforts were made
16 to minimize animal suffering and the number of animals.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32
33 **Surgery**

34
35 *Plantar incision*

36 The plantar surgery was performed as previously described¹. All animals were
37 anesthetized with isoflurane (4% to induce, 1.5-2% to maintain) in air delivered via a nose cone.
38
39 The plantar aspect of the left hindpaw was sterilized with a 10% povidone-iodine solution
40 before and after surgeries and was placed through a hole in a sterile drape. A 1-cm longitudinal
41 incision was made through skin and fascia of the plantar aspect of the foot, starting 0.5 cm from
42 the proximal edge of the heel and extending toward the toes. The plantaris muscle was elevated
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 and longitudinally incised through, leaving muscle origin and insertion intact. After hemostasis
6
7
8 with gentle pressure, the skin was apposed with 2 mattress sutures of 5-0 nylon. The animals
9
10 were allowed to recover. The incision was checked daily and any sign of wound infection or
11
12
13
14 dehiscence excluded the animal from the study.

15 16 *Intrathecal cannulation*

17
18
19 Animals were anesthetized with chloral hydrate (intraperitoneal injection, 400 mg/kg,
20
21
22 Kanto Chemical Co. Inc., Tokyo, Japan) and a thoracic laminectomy was performed. A midline
23
24
25 incision was made at the lower thoracic spines. The dura was cut and a PE5 polyethylene tubing
26
27
28 (OD = 0.355 mm, ID = 0.2 mm) adhering to PE50 tubing (OD = 0.956 mm, ID = 0.58 mm) was
29
30
31 inserted into the subarachnoid space and the tip of the catheter was located near the lumbar
32
33
34 enlargement of the spinal cord. The catheter was secured to the paravertebral muscles with
35
36
37 suture and the external end of the catheter was sealed by heat. The wound was then closed in
38
39
40 layers and animals were returned to their home cages for recovery at least 3 days before plantar
41
42
43 incision. The patency of catheter was confirmed by paralysis of hindlimbs following 10 μ l 2%
44
45
46 lidocaine injection the next day after catheterization. Rats with any neurological deficits were
47
48
49 excluded from the experiment.

50 51 52 **Spinal Drug Delivery**

1
2
3
4
5 The p38 inhibitor FR167653 (a gift from Fujisawa Pharmaceutical Company, Osaka,
6
7
8 Japan) was dissolved in 100% dimethyl sulfoxide (DMSO, Sigma-Aldrich, St. Louis, MO) to
9
10 obtain a concentration of 50 mg/ml as a stocking solution, then diluted to 10 µg/µl (20%
11
12 DMSO), 5 µg /µl (10% DMSO) and 2 µg/µl (4% DMSO) in sterile saline on the injection day.
13
14
15 FR167653 (100 µg, 50 µg, and 20 µg) or diluted vehicle (20%, 10%, and 4% DMSO) was
16
17
18 injected intrathecally (10 µl) followed by flush with 20 µl of saline 30 min before plantar
19
20
21 incision.
22
23

24 25 **Behavioral experiments**

26
27
28 All animals were habituated to the testing environment from 2 days before baseline testing.
29
30
31 For mechanical stimulation, animals were individually placed in a chamber (10 x 10 x 20 cm) of
32
33
34 a Plexiglas cage on an elevated iron mesh floor and allowed to acclimate for 20 min before
35
36
37 testing. The von Frey filaments were applied from underneath the mesh openings to stimulate
38
39
40 the plantar surface at an area adjacent to the wound near the medial heel. A series of von Frey
41
42
43 filaments with incremental stiffness (0.4, 0.6, 1, 2, 4, 6, 8, and 15 g; Stoelting, Wood Dale, IL),
44
45
46 always started from the 2.0 g filament, were presented perpendicular to the plantar surface for
47
48
49 5-6 seconds for each filament. Stimulation was conducted in a consecutive fashion, whether
50
51
52 ascending or descending using up-down method as previously described²¹. In the absence of
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 paw withdrawal response to the initial selected filament, a stronger stimulus was presented; in
6
7
8 the event of paw withdrawal, the next weaker stimulus was chosen. In cases in which threshold
9
10
11 fell outside the range of detection, 0.4 and 15.0 g were assigned for the minimal and maximal
12
13
14 value, respectively. Threshold value was an average of two test measurements at each time
15
16
17 point, and was used for calculation of the fifty percent withdrawal threshold²². The animals
18
19
20 were tested daily from 2 days before surgery, and then 1, 3, 5 h and 1, 2, 3, 5, and 7 d after
21
22
23 surgical incisions.

24
25
26 For thermal hyperalgesia, rats were placed individually in the Plexiglas cage on a glass
27
28
29 floor which had been pre-heated at constant 30°C (Plantar Test Apparatus, IITC, Woodland
30
31
32 Hills, CA). After animal accommodation, a focused radiant heat source underneath the glass
33
34
35 floor was applied at the incision midpoint of the left paw. The intensity of the emitting heat was
36
37
38 adjusted to produce withdrawal latency at about 20-25 s in normal rats. Cut-off latency was
39
40
41 pre-determined at 30 s to avoid paw injury. Each rat was tested three times, at an interval of 5
42
43
44 min, and the average of the three measurements was recorded as withdrawal latency at the time
45
46
47 point. The data were obtained before surgery, then 1, 3, and 5 h and 1, 2, 3, 5, and 7 d after
48
49
50 surgery. Preoperative baselines of both mechanical and thermal tests were tested daily for 2 d
51
52
53 and averaged.

1
2
3
4
5 Because DMSO was reported to have anti-inflammatory²³ and analgesic properties²⁴, its
6
7 use as a solvent for FR167653 may be of concern. Therefore, two control groups, saline and
8
9 DMSO, are included in the studies of p38 inhibitor administration. Specifically, 10% and 20%
10
11 DMSO were given as vehicle controls of 50 µg and 100 µg FR167653, respectively. At least 8
12
13
14
15 rats were included for each group in the behavioral experiment with FR167653 and vehicle
16
17 treatment. The same experimenter who did the mechanical and thermal stimulation was blinded
18
19
20
21 to the group allocation of the rats.
22
23
24

25 **Immunohistochemistry**

26
27
28 At 1 h, 3 h and 1 d, 3 d and 7 d post-incisional time points, animals were sacrificed with an
29
30 overdose of intraperitoneal injection of chloral hydrate (650mg/kg) and perfused transcardially
31
32 with saline (room temperature), followed by 4°C 4% paraformaldehyde in 0.1 M phosphate
33
34 buffer. The L4-5 spinal segments were carefully removed, post-fixed in the same fixative for
35
36 overnight, and then cryoprotected in cold 30% sucrose/phosphate buffer overnight. The tissues
37
38 were frozen at -20°C in a cryostat and transverse free-floating spinal sections (30 µm) were cut
39
40 and collected in 0.1 M phosphate buffer. After blocking with 2% normal goat serum containing
41
42 0.3% Triton X-100 for 1 hour at room temperature, all the sections were incubated with either a
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 Technology, Beverly, MA) or a mouse monoclonal anti-OX-42 (CD11b, microglia marker,
6
7
8 1:2000; Serotec, Indianapolis, IN) at 4°C for two nights. Then, sections were incubated with
9
10 either goat biotinylated anti-rabbit or goat biotinylated anti-mouse secondary antibody (1:200,
11
12
13 Vector Laboratories, Burlingame, CA) for 2 hr, and subsequently with ExtraAvidin peroxidase
14
15 (1:1000, Sigma-Aldrich) for 1 hr at room temperature. After rinsing in 0.1 M phosphate buffer
16
17
18 for 20 min, sections were reacted with 0.25% 3,3'-diaminobenzidine tetrahydrochloride
19
20
21 solution in phosphate buffer containing 0.0025% hydrogen peroxide (Sigma-Aldrich) for 6 min.
22
23
24 Finally, sections were mounted onto gelatin-coated glass slides, air-dried, dehydrated, cleared
25
26
27 with xylene and coverslipped with Entellan mounting medium (Merck, Darmstadt, Germany).
28
29
30

31
32 For double immunofluorescence, sections were incubated with a mixture of primary
33
34 antibodies rabbit polyclonal antibody against p-p38 (1:500; Cell Signaling Technology) and
35
36 mouse monoclonal antibody against neuronal specific nuclear protein (NeuN, a neuronal
37
38 marker) (1:4000, Chemicon, Temecula, CA), glial fibrillary acidic protein (GFAP; an astrocyte
39
40 marker) (1:5000; Chemicon), or OX-42 (1:4000, Serotec) for 36-48 hrs at 4°C, followed by a
41
42
43 mixture of goat anti-mouse fluorescein isothiocyanate- and goat anti-rabbit
44
45
46 indocarbocyanine-conjugated secondary antibodies (1:200, Jackson ImmunoResearch Inc.,
47
48
49 West Grove, PA) for 1 hour at room temperature. Then, the double-stained sections were
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 observed with a Nikon fluorescence microscopy (Tokyo, Japan), and images were captured
6
7
8 with a CCD camera (SPOT, Diagnostic Instruments, Sterling Heights, MI).
9

10 **Quantification and Statistical Analysis**

11
12
13
14 All the results were expressed as mean \pm SEM. To assess the incision-induced mechanical
15
16 and thermal thresholds before and after plantar incision, the main effects of different time points
17
18 and treatments as well as interaction of treatment \times control effect were analyzed by two-way
19
20 analysis of variance (ANOVA). Then, repeated-measures of ANOVA were applied to assess the
21
22 incision-induced mechanical and thermal threshold before and after plantar incision by using
23
24 the pre-incisional baselines as control and one-way ANOVA was applied to compare group
25
26 differences after FR167653 treatment at each time point. All tests were followed by Bonferroni
27
28 test as the multiple comparison analysis. Analysis was completed via software SPSS (14.0;
29
30 SPSS Inc., Chicago, IL).
31
32
33
34
35
36
37
38
39

40 For immunohistochemistry, phospho-p38-immunoreactive (p-p38-ir) cells in L4-5 spinal
41
42 dorsal horns (laminae I-V) were counted on six randomly chosen sections from each animal and
43
44 averaged for each animal. In all cases, the number of p-p38-ir cells was quantified at a
45
46 magnification of 100x. The density of OX-42 labeled microglia ipsilateral to the incision paw
47
48 was measured microscopically under 100x magnification in 8 to 10 randomly selected sections
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 from the L4 and L5 spinal cord. Images were digitally captured with a computer-assisted image
6
7
8 analysis system (Image J, NIH Image, National Institute of Health, Bethesda, MD). The images
9
10
11 with 2 pre-defined rectangles located at the medial one-half of laminae I-II and 3 rectangles at
12
13
14 the white matter ipsilateral to the operated paw were computerized-automatically adjusted with
15
16
17 gray scales ranging from 0 to 255, and the pixel numbers of each areas was counted, following
18
19
20 the method described by Lin et al²⁵. The pixel ratio of the average of the medial superficial
21
22
23 dorsal horn to the average of the white matter was calculated. Group differences in the number
24
25
26 of p-p38-ir cells and the pixel ratio of OX-42 staining were compared with either one-way
27
28
29 ANOVA followed by *post hoc* Bonferroni test, or student t-test when appropriate for two-group
30
31
32 comparison. $P < 0.05$ was considered statistically significant.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Results

Activation of p38 in the spinal cord after plantar incision

To investigate whether p38 MAPK is activated after surgery, we used immunohistochemistry to examine changes of phosphorylated p38 immunoreactive (p-p38-ir) cells in the spinal dorsal horn at different postoperative time points. A very low constitutive expression of p-p38 was found in the L4-5 spinal dorsal horn of naïve rats (n = 6; Fig. 1A). Incisions induced an increase in the number of p-p38-ir cells evident in the ipsilateral L4-5 spinal segments (Fig. 1B and C). Quantification of p-p38-ir cells in the L4-5 dorsal horn revealed an increase in p38 activation at 1 h (n = 5), 3 h (n = 7), 1 d (n = 7), 3 d (n = 6), and 7 d (n = 7) following plantar incisions. p-p38 level began to increase at 1 h, reached the peak at day 1, and then maintained for 3 days. Analysis of data revealed a significant difference in the number of p-p38-ir cells in the dorsal horn of the ipsilateral lumbar spinal cord [F(5,32) = 3.914; one-way ANOVA; $P < 0.01$] and significant increases between the naïve and the incision group were found at 1 h ($P < 0.05$), 3 h ($P < 0.01$), 1 d ($P < 0.001$) and 3 d ($P < 0.01$) postoperatively (Fig. 1D). Then, p38 activation decreased at day 7 and did not show statistical significance ($P = 0.062$) compared to the naïve group.

Postoperative nociceptive behavior

1
2
3
4
5 As reported previously¹, plantar incision produced a rapid mechanical allodynia beginning
6
7
8 at 1 h postoperatively. Analysis revealed that plantar incision resulted in a significant reduction
9
10 of mechanical withdrawal threshold to von Frey stimulation at 1 h (2.08 ± 0.68 g; $P < 0.001$), 3
11
12 h (3.62 ± 1.19 g; $P < 0.01$), 5 h (4.34 ± 1.19 g; $P < 0.01$), 1 d (5.08 ± 0.84 g; $P < 0.01$), 2 d (7.04
13
14 ± 1.31 g; $P < 0.05$), and 3 d (7.69 ± 1.12 g; $P < 0.05$) when compared with the baseline value
15
16
17 (13.5 ± 1.19 g; $n = 6$; Fig. 1E). This reduction in thresholds recovered at day 5 (11.26 ± 1.16 g;
18
19
20 $P > 0.05$).

21
22
23
24
25 In addition, plantar incision caused a rapid decrease in the paw withdrawal latency to
26
27
28 noxious thermal stimuli (Fig. 1F). The mean withdrawal latency to radiant heat was $25.33 \pm$
29
30
31 1.01 sec ($n = 8$) before incision. After incision, a significant decrease in withdrawal latency was
32
33
34 shown at 1 h (5.68 ± 0.98 s; $P < 0.01$), 3 h (4.83 ± 0.34 s; $P < 0.01$), 5 h (5.94 ± 0.58 s; $P < 0.01$),
35
36
37 1 d (8.36 ± 0.55 s; $P < 0.01$), 2 d (9.4 ± 0.58 s; $P < 0.01$), and 3 d (14.59 ± 0.57 s; $P < 0.01$)
38
39
40 postoperatively. On postoperative day 5, the withdrawal latency returned to preoperative
41
42
43 baseline level (21.85 ± 1.93 s; $P > 0.05$).

44 45 46 ***Plantar incision-induced p-p38 is localized primarily in microglia, and in some neurons***

47
48
49 Previous reports have demonstrated colocalization of p-p38 with OX-42, a marker for
50
51
52 microglial cells, in spinal cord in several pain models^{17;26}. Here, to identify whether p-p38-ir
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 was expressed in the microglial cells in the plantar incision model, we performed double
6
7
8 immunofluorescence staining of p-p38 with several cell-specific markers: NeuN (neurons),
9
10 GFAP (astrocytes), and OX-42 (microglia) at 1 h, 3 h and 1 d (Fig. 2; n = 2 for each time point)
11
12
13 postoperatively. p-p38 staining was found in a limited number of the spinal dorsal horn neurons
14
15
16 at 1 h (Fig. 2A and 2B) and 3 h (Fig. 2G and 2H). Neurons have close contacts with p-p38
17
18
19 immunoreactive cells at the time point of 1 d, but no colocalization was observed between
20
21
22 p-p38 and NeuN (Fig. 2M and 2N). The majority of p-p38 was localized in OX-42-positive
23
24
25 microglia at 1 h (Fig. 2D-2F), 3 h (Fig. 2J-2L) and 1 d (Fig. 2P-2R). The morphology of
26
27
28 p-p38-ir cells, as shown in Figure 2, shows small cell bodies with ramified processes. However,
29
30
31 no p-p38 was localized in GFAP-positive astrocytes at all the time points examined (Fig. 2C, 2I,
32
33
34 and 2O).

35 36 37 ***Plantar incision induced spinal microglial activation***

38
39
40 The OX-42 immunoreactivity in the L4-5 spinal dorsal horn at different time point after
41
42
43 plantar incision was shown in Figure 3. In tissue from rats in naïve group and 3 h post-operative
44
45
46 time, microglia showed small soma with thin branched processes (Fig. 3A). The topographic
47
48
49 expression of OX-42 developed gradually from the superficial dorsal horn (at 1 d; Fig. 3B) to
50
51
52 the whole dorsal horns (at 3 d; Fig. 3C) following surgery. Activated microglia were markedly
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 found in the laminae I-II of spinal dorsal horn 3 d after plantar incision, especially at the medial
6
7
8 half of the dorsal horn, with marked morphological changes such as increased cell size and
9
10 thickened branches (Fig. 3C). Thereafter, the microglial activation was decreased to basal level
11
12
13 at 7 d after surgery (Fig. 3D). Statistical analysis showed that there is a significant increase in
14
15
16 the pixel ratio of OX-42 immunoreactivity of the ipsilateral superficial dorsal horn to the white
17
18
19 matter in sections from 3 d (1.41 ± 0.05 ; $P < 0.001$, *post hoc* comparison; Fig. 3F)
20
21
22 postoperatively, but not 3 h (1.17 ± 0.05), 1 d (1.24 ± 0.06), and 7 d (1.21 ± 0.02) when
23
24
25 compared with those from naïve rats (1.12 ± 0.04 ; Fig. 3E). These results implicated the most
26
27
28 remarkable microglial activation was observed on day 3 following plantar incision.
29
30

31 ***Inhibition of p38 prevented mechanical hypersensitivity***

32
33
34 To investigate the functional roles of activation of p38 in incision-induced mechanical
35
36
37 hyperalgesia, we tested whether intrathecal administration of FR167653, a specific p38
38
39
40 inhibitor²⁷⁻²⁹, attenuated mechanical allodynia. First, to examine whether blocking p38
41
42
43 activation in the spinal cord has an effect on baseline withdrawal threshold, we injected saline,
44
45
46 vehicle (10% DMSO in saline), 20 μ g or 50 μ g FR167653 intrathecally 30 min before plantar
47
48
49 incision. Basal withdrawal thresholds to von Frey filaments were not different after FR167653
50
51
52 injection (Fig. 4A). The baseline values of paw withdrawal thresholds for rats in saline, vehicle,
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 20 μg and 50 μg FR167653-treatment group were 14.19 ± 0.45 g (n = 10), 13.93 ± 0.75 g (n =
6
7
8 9), 13.95 ± 0.38 g (n = 8), and 12.54 ± 0.72 g (n = 8), respectively ($P > 0.05$).
9

10
11 Then we tested the effects of pretreatment of FR167653 (20 μg or 50 μg) and the vehicle
12
13 on mechanical hypersensitivity following plantar incision. An initial analysis of two-way
14
15 ANOVA showed that the treatment effect, the time effect and the interaction of time \times treatment
16
17 were all significantly different (all $P < 0.001$), indicating different time trends in the four groups.
18
19 Then, the analysis of one-way ANOVA revealed that inhibition of spinal p38 activation by 50
20
21 μg FR167653 significantly prevented incision-induced mechanical allodynia at 1 h ($P < 0.001$),
22
23 3 h ($P < 0.001$), 5 h ($P < 0.001$), 1 d ($P < 0.001$) and 2 d ($P < 0.001$) (Fig. 4A) when compared
24
25 with animals intrathecally treated with saline. Intrathecal injection of 10% DMSO seemingly
26
27 produced a mild anti-allodynic effect lasting for about 1 day, however, the effects were not
28
29 statistically significant at any time point. FR167653 50 μg still showed significant differences
30
31 compared to the vehicle group at 1 h ($P < 0.001$), 3 h ($P < 0.01$), 5 h ($P < 0.01$), 1 d ($P < 0.05$),
32
33 and 2 d ($P < 0.05$). Moreover, treatment with 50 μg FR167653 results in significant increase of
34
35 withdrawal threshold in a dose-dependent manner when compared with 20 μg treatment at 3 h
36
37 ($P < 0.05$), 1 d ($P < 0.01$), and 2 d ($P < 0.05$). Intrathecal 20 μg FR167653 could only inhibit
38
39 postoperative allodynia for 1 h ($P < 0.01$ versus saline and vehicle) and soon lost its
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 anti-allodynic effect.
6

7
8 ***Inhibition of p38 only inhibited thermal hyperalgesia at an early time point***
9

10 To elucidate whether inhibition of p38 activation altered withdrawal responses to thermal
11 stimulation, the FR167653 was injected intrathecally 30 min before plantar incision.
12
13 Pretreatment of saline, vehicle (20% DMSO), 100 µg FR167653 (intrathecal injection) had no
14 effect on the baseline values of withdrawal latency (Fig. 4B). The baseline values of paw
15 withdrawal latencies for rats in saline, vehicle, and 100 µg FR167653-treatment group were
16 23.99 ± 1.03 sec (n = 12), 25.35 ± 1.21 sec (n = 8), and 24.69 ± 1.27 sec (n = 9), respectively (*P*
17 > 0.05; two-way ANOVA). Moreover, it is surprisingly that a very high dose of FR167653 (100
18 µg) could only produced a transient reduction of thermal hyperalgesia at 1 h after surgery (*P* <
19 0.05; *post hoc* comparison; Fig. 4B) when compared with saline or vehicle (20% DMSO in
20 saline) group, and had no effect at any other time points. Lower dose of FR167653 (50 µg) was
21 tested in a small number of rats and showed no effect (data not shown). Unlike von Frey test,
22 post-incision thermal hyperalgesia was not affected by intrathecal DMSO, though a high
23 concentration of 20% was used.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48
49 ***FR167653 reduced incision-induced number of p-p38-ir cells***
50

51 Because there was a maximal increase of p-p38-ir cells at 1 day after plantar incision, we
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 examined whether intrathecal pretreatment of p38 MAPK inhibitor reduced incision-induced
6
7
8 activation of p38 in the spinal dorsal horn at this time point. Results from this study showed that
9
10
11 pretreatment with 20 and 50 μg of FR167653 produced a significant reduction in the number of
12
13
14 p-p38-ir cell in lumbar dorsal horn (37.5 ± 16.8 at the 50 μg group and 47.0 ± 4.7 at the 20 μg
15
16
17 group vs. 157.0 ± 9.7 at the vehicle group, $P < 0.001$; $F(2,11) = 40.551$; one-way ANOVA) (Fig.
18
19
20 5).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Discussion

In this study, we have shown that in the L4-5 spinal dorsal horn, the plantar incision induced an increase in phosphorylation of p38 starting immediately after surgery and the activation of p38 is coincident with the development of incisional pain. Activation of p38 was found mainly in microglia and in a limited number of spinal dorsal horn neurons at 1h and 3h, but exclusively in microglia at 1 d after incision. Plantar incision produced a delayed microglial activation (OX-42 upregulation) at day 3. Pretreatment of p38 inhibitor FR167653 significantly prevented incision-induced mechanical allodynia and reduced p38 activation in the lumbar spinal dorsal horn, but only attenuated thermal hyperalgesia at an early time point. These results suggest that p38 activation in spinal microglial cells plays a role in the development of postoperative mechanical hypersensitivity.

p38 activation and pain hypersensitivity

The present study disclosed that activation of p38 MAPK plays a role in incision-induced mechanical hypersensitivity because blocking its activation with pretreatment of p38 inhibitor FR167653 effectively prevented postoperative tactile allodynia. These results are in agreement with previous observations in various experimental pain models, including models of nerve injury^{14;17;19}, inflammation^{14;15}, diabetes³⁰ and capsaicin³¹. All these studies identified the

1
2
3
4
5 colocalization of p-p38 in microglia after injury. However, the appearance of p-p38 in different
6
7 spinal cells is also observed in a adjuvant immunized rats²⁶, venom-induced inflammation³²,
8
9 and neuropathic pain model³³. Our study supports these studies indicating that involvement of
10
11 p38 MAPK cascade in neurons/microglia follows a time-sequence and is not exclusively in one
12
13
14
15
16
17 cell type in incisional pain.

18
19
20 This study investigated the temporal relationship between the p38 activation and
21
22 incision-induced mechanical hypersensitivity. The immunoreactivity of p-p38 reaches the
23
24 maximum at 1 d after paw incision; however, the mechanical allodynia is most obvious at 1 h.
25
26 Pretreatment of FR167653 at two different doses dose-dependently prevented post-incisional
27
28 allodynia, although they showed similar inhibition on p-p38. Because we treated the animals
29
30 with single dose of p38 inhibitor before plantar incision, the p-p38-IR existing on the day 1
31
32 represents a direct result of an extinct effect of FR167653. In contrast, the differential
33
34 anti-allodynic effects of two inhibitor doses reflect an integrated consequence of inhibition on
35
36 nociceptive development, or broadly on complicated neuron-glia interactions. These results
37
38 indicate that the activation of spinal p38 is required for incision-induced mechanical allodynia
39
40 but the quantitative p-p38 expression is only partially correlated with the nociceptive responses.
41
42
43
44
45
46
47
48
49
50 The p38 activation later than the appearance of the maximal allodynia suggests p-p38
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 contributes to the nociceptive development rather than initiation. In addition, previous reports
6
7
8 have demonstrated that the other members of MAPK, including extracellular signal-regulated
9
10
11 kinase 1 and 2, and c-Jun N-terminal kinase, play distinct roles in the development and
12
13
14 maintenance of inflammation- and nerve injury-induced pain hypersensitivity^{30;34;35}. However,
15
16
17 their functional roles in incisional pain remain to be elucidated.

18
19
20 Surprisingly, with one single intrathecal injection, 50 µg FR167653 produced a long
21
22
23 inhibition on tactile allodynia lasting for 2 days after incision, whereas a higher dose of
24
25
26 FR167653 only resulted in a very brief inhibition on heat hyperalgesia. It suggests that spinal
27
28
29 p38 activation is involved more in the pathophysiology of mechanical allodynia, than thermal
30
31
32 hyperalgesia in surgical pain. In clinics, the most common pain complaint after surgery is
33
34
35 mechanically evoked such as dressing change, coughing and ambulation³⁶. Furthermore,
36
37
38 different pain modalities may have discrepant drug responses. Single injection of 50 µg
39
40
41 FR167653 prevented incision-induced mechanical allodynia for 2 days, whereas in the spared
42
43
44 nerve injury, after the pain related behavior was established, it could reverse mechanical
45
46
47 allodynia for only 6 hours³⁷. This supports the idea that different pain models are underlined by
48
49
50 various mechanisms and may require a different specific treatment at a given timepoint. These
51
52
53 results implicate p38 inhibitor FR167653 could be a candidate in prevention of postsurgical
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 pain.
6
7

8 In experimental studies, drugs such as FR167653, SB203580, and CNI1493 were used to
9
10 test the functional roles of p38 MAPK in inflammation- and nerve injury-induced hyperalgesia
11
12 and allodynia. They all diminish p38 activity to reduce inflammation- and nerve injury-induced
13
14 pain hypersensitivity^{15;18;37;38}. Among them, FR167653 has a potent anti-inflammatory action
15
16 by inhibiting interleukin-1 β (IL-1 β) and tumor necrosis factor- α (TNF- α) production^{39;40}, and
17
18 these pro-inflammatory cytokines can modulate the synaptic transmission activity and promote
19
20 pain⁴¹. *In vivo* studies showed that FR167653 can effectively prevent and attenuate
21
22 ischemia/reperfusion injury through inhibiting p38 MAPK phosphorylation²⁷⁻²⁹, an action
23
24 different from SB203580 which binds to the adenosine triphosphate pocket in the enzyme to
25
26 block p38 activity⁴². In consequence, we confirmed that FR167653 prevented the tactile
27
28 allodynia via markedly inhibition of spinal p38 phosphorylation.
29
30
31
32
33
34
35
36
37
38
39

40 In contrast, our results did not show blocking of heating pain hypersensitivity by
41
42 FR167653 except a brief inhibition at the first post-surgical hour. Although thermal
43
44 hyperalgesia is less noteworthy in the postoperative patients, there is growing evidence that the
45
46 excitation of dorsal root ganglion (DRG) neurons plays an important role in heat
47
48 hypersensitivity⁴³⁻⁴⁵. Tissue injury induced thermal hyperalgesia by sensitization in DRG
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 neurons and activation of a candidate nociceptor, the cation-selective ion-channel transient
6
7 receptor potential vanilloid 1⁴³. Behavioral study has suggested transient receptor potential
8
9 vanilloid 1 is important in generating heat hyperalgesia by incision in a transient receptor
10
11 potential vanilloid 1 knockout model⁴⁶. Furthermore, Mizushima et al.⁴⁷ proved that capsaicin
12
13 injection into the hindpaw induced phosphorylation of p38 MAPK in transient receptor
14
15 potential vanilloid 1-positive neurons in the DRG immediately after capsaicin injection, and
16
17 capsaicin-induced thermal hyperalgesia was reversed by intrathecal FR167653 administration
18
19 when the intrathecal catheter tip is placed close to the L4-5 DRG level.
20
21
22
23
24
25
26
27

28
29 A site-dependent expression of p-p38 MAPK should be considered. Peripheral
30
31 inflammatory pain by complete Freund's adjuvant caused p-p38 expression in DRG neurons of
32
33 C fibers for several days, but did not activate p38 phosphorylation in the spinal dorsal horn¹³.
34
35 Because the dural membrane in rats extends over the capsule of the DRG^{17;48;49} and intrathecal
36
37 FR167653 might diffuse to affect neuronal cells in the DRG, we suggest that in this study
38
39 FR167653 could inhibit the p-p38 activation initiated in the DRG and nerves⁴⁷ at the immediate
40
41 postsurgical period, but was unable to prevent heat hyperalgesia. This hypothesis provides a
42
43 logical explanation that the immediate blocking of mechanical allodynia is mediated majorly
44
45 via p-p38 inhibition in the DRG but few in the spinal cord.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 Importantly, blocking of p38 activation with FR167653 did not result in change of baseline
6
7
8 nociceptive responses to tactile and thermal stimuli before plantar incision, indicating that this
9
10
11 agent does not alter the physiological responses to phasic pain. These results are consistent with
12
13
14 previous pharmacological studies demonstrating that spinal p38 inactivation does not modulate
15
16
17 baseline nociception^{15;17;19;31;47}. This pharmacological character adds to the merit of this agent
18
19
20 for the potential use in clinic.

21 22 ***Microglia activation and postoperative pain***

23
24
25 Compared with temporal correlation of an early activation of p38 and mechanical
26
27
28 allodynia in behavior, the increased microglial activation was observed much delayed, i.e. not
29
30
31 at the early (3 h and 1 d) but the late period (3 d). This finding is not surprising because previous
32
33
34 studies indicated that there is a dissociation between microglial activation and tactile allodynia
35
36
37 in plantar incision⁵⁰ and nerve injury models⁵¹. Our results correspond to current concepts
38
39
40 which regard the emergence of “activated” microglia, by evidence of upregulation of CD11b
41
42
43 expression, as a result of proliferation of “reactive” microglia and invading monocytes from
44
45
46 vessels in the vicinity, rather than an initiating sign of pathological cascades leading to pain^{52;53}.
47
48

49 The relationship between p38 phosphorylation and microglial activation in the nociceptive
50
51
52 hypersensitization has not been well established or understood. Most of pain modalities
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 including formalin^{15;54}, complete Freund's adjuvant²⁶, capsaicin injection¹³, nerve injury^{17;19;37},
5
6
7 spinal cord injury⁵⁵, and incision (this study) proved to activate the spinal p38 MAPK
8
9
10 phosphorylation in the microglia. However, the detection of microglia activation depending on
11
12
13 immunohistological up-regulation of the surface marker CD11b (OX-42) would reflect rather a
14
15
16 morphological change than a functional alteration. Furthermore, pain states may not be always
17
18
19 associated with the upregulation of this surface marker^{50;51;56} and other characteristics of
20
21
22 activation should be sought in the future. Another recent study supported our finding by using
23
24
25 the expression of ionized calcium-binding adapter molecule 1 (Iba-1), another microglia marker.
26
27
28 When measuring the whole spinal dorsal horn, Iba-1 expression 1 day after incision is
29
30
31 significantly increased when compared with that of the contralateral side and the naïve rats, but
32
33
34 when the measuring area is limited to laminae I-II, the increase of Iba-1 expression is
35
36
37 insignificant⁵⁷. Therefore, p38 MAPK phosphorylation may be a better marker than OX-42
38
39
40 expression in incision-induced pain model.
41
42
43
44
45

46 ***Conclusions***

47
48
49 In conclusion, activation (phosphorylation) of p38 MAPK in spinal microglial cells is
50
51
52 partially coincident with the development of the tactile allodynia induced by surgical incision.
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Most importantly, intrathecal administration of p38 MAPK inhibitor FR167653 before surgery can effectively prevent incision-induced mechanical pain hypersensitivity, a common problem in clinics. Accordingly, we suggest p38 inhibitor could be used as an effective therapeutic agent to relieve post-operative pain and microglia inhibition may be a new strategy for postoperative pain treatment.

1
2
3
4
5 Reference List
6
7

- 8 1. Brennan TJ, Vandermeulen EP, Gebhart GF: Characterization of a rat model of incisional
9
10 pain. *Pain* 1996; 64: 493-501
11
12
13
14 2. Kawamata M, Takahashi T, Kozuka Y, Nawa Y, Nishikawa K, Narimatsu E, Watanabe H,
15
16 Namiki A: Experimental incision-induced pain in human skin: effects of systemic
17
18 lidocaine on flare formation and hyperalgesia. *Pain* 2002; 100: 77-89
19
20
21
22
23 3. Honore P, Wade CL, Zhong C, Harris RR, Wu C, Ghayur T, Iwakura Y, Decker MW,
24
25 Faltynek C, Sullivan J, Jarvis MF: Interleukin-1 α gene-deficient mice show
26
27 reduced nociceptive sensitivity in models of inflammatory and neuropathic pain but not
28
29 reduced nociceptive sensitivity in models of inflammatory and neuropathic pain but not
30
31 post-operative pain. *Behav. Brain Res.* 2006; 167: 355-64
32
33
34
35 4. Zahn PK, Umali E, Brennan TJ: Intrathecal non-NMDA excitatory amino acid receptor
36
37 antagonists inhibit pain behaviors in a rat model of postoperative pain. *Pain* 1998; 74:
38
39 213-23
40
41
42
43
44 5. Zahn PK, Brennan TJ: Lack of effect of intrathecally administered N-methyl-D-aspartate
45
46 receptor antagonists in a rat model for postoperative pain. *Anesthesiology* 1998; 88:
47
48 143-56
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5 6. Yamamoto T, Sakashita Y: The role of the spinal opioid receptor like1 receptor, the NK-1
6
7
8 receptor, and cyclooxygenase-2 in maintaining postoperative pain in the rat.
9
10 Anesth.Analg. 1999; 89: 1203-8
11
12
13
- 14 7. Zhu CZ, Hsieh G, Ei-Kouhen O, Wilson SG, Mikusa JP, Hollingsworth PR, Chang R,
15
16
17 Moreland RB, Brioni J, Decker MW, Honore P: Role of central and peripheral mGluR5
18
19
20 receptors in post-operative pain in rats. Pain 2005; 114: 195-202
21
22
23
- 24 8. Zahn PK, Pogatzki-Zahn EM, Brennan TJ: Spinal administration of MK-801 and NBQX
25
26
27 demonstrates NMDA-independent dorsal horn sensitization in incisional pain. Pain 2005;
28
29
30 114: 499-510
31
32
33
- 34 9. Yamamoto T, Shimoyama N, Mizuguchi T: The effects of morphine, MK-801, an NMDA
35
36
37 antagonist, and CP-96,345, an NK1 antagonist, on the hyperesthesia evoked by
38
39
40 carageenan injection in the rat paw. Anesthesiology 1993; 78: 124-33
41
42
43
- 44 10. Widmann C, Gibson S, Jarpe MB, Johnson GL: Mitogen-activated protein kinase:
45
46
47 conservation of a three-kinase module from yeast to human. Physiol Rev. 1999; 79:
48
49
50 143-80
51
52
53
- 54 11. Ji RR, Strichartz G: Cell signaling and the genesis of neuropathic pain. Sci.STKE. 2004;
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 2004: reE14
6
7
8

- 9 12. Ji RR, Kawasaki Y, Zhuang ZY, Wen YR, Zhang YQ: Protein kinases as potential targets
10 for the treatment of pathological pain. *Handb.Exp.Pharmacol.* 2007; 359-89
11
12
13
14
15 13. Ji RR, Samad TA, Jin SX, Schmoll R, Woolf CJ: p38 MAPK activation by NGF in
16 primary sensory neurons after inflammation increases TRPV1 levels and maintains heat
17 hyperalgesia. *Neuron* 2002; 36: 57-68
18
19
20
21
22
23
24
25 14. Kim SY, Bae JC, Kim JY, Lee HL, Lee KM, Kim DS, Cho HJ: Activation of p38 MAP
26 kinase in the rat dorsal root ganglia and spinal cord following peripheral inflammation
27 and nerve injury. *Neuroreport* 2002; 13: 2483-6
28
29
30
31
32
33
34
35 15. Svensson CI, Marsala M, Westerlund A, Calcutt NA, Campana WM, Freshwater JD,
36 Catalano R, Feng Y, Protter AA, Scott B, Yaksh TL: Activation of p38 mitogen-activated
37 protein kinase in spinal microglia is a critical link in inflammation-induced spinal pain
38 processing. *J.Neurochem.* 2003; 86: 1534-44
39
40
41
42
43
44
45
46
47
48 16. Hua XY, Svensson CI, Matsui T, Fitzsimmons B, Yaksh TL, Webb M: Intrathecal
49 minocycline attenuates peripheral inflammation-induced hyperalgesia by inhibiting p38
50 MAPK in spinal microglia. *Eur.J.Neurosci.* 2005; 22: 2431-40
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5 17. Jin SX, Zhuang ZY, Woolf CJ, Ji RR: p38 mitogen-activated protein kinase is activated
6
7 after a spinal nerve ligation in spinal cord microglia and dorsal root ganglion neurons and
8
9 contributes to the generation of neuropathic pain. *J.Neurosci.* 2003; 23: 4017-22
10
11
12
13
14 18. Obata K, Yamanaka H, Kobayashi K, Dai Y, Mizushima T, Katsura H, Fukuoka T,
15
16 Tokunaga A, Noguchi K: Role of mitogen-activated protein kinase activation in injured
17
18 and intact primary afferent neurons for mechanical and heat hypersensitivity after spinal
19
20 nerve ligation. *J.Neurosci.* 2004; 24: 10211-22
21
22
23
24
25 19. Tsuda M, Mizokoshi A, Shigemoto-Mogami Y, Koizumi S, Inoue K: Activation of p38
26
27 mitogen-activated protein kinase in spinal hyperactive microglia contributes to pain
28
29 hypersensitivity following peripheral nerve injury. *Glia* 2004; 45: 89-95
30
31
32
33
34 20. Tsuda M, Inoue K, Salter MW: Neuropathic pain and spinal microglia: a big problem
35
36 from molecules in "small" glia. *Trends Neurosci.* 2005; 28: 101-7
37
38
39
40
41 21. Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL: Quantitative assessment of
42
43 tactile allodynia in the rat paw. *J.Neurosci.Methods* 1994; 53: 55-63
44
45
46
47
48 22. Dixon WJ: Efficient analysis of experimental observations.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5 23. Gorog P, Kovacs IB: Effect of dimethyl sulfoxide (DMSO) on various experimental
6
7 inflammations. *Curr.Ther.Res.Clin.Exp.* 1968; 10: 486-92
8
9
10
11 24. Evans MS, Reid KH, Sharp JB, Jr.: Dimethylsulfoxide (DMSO) blocks conduction in
12
13 peripheral nerve C fibers: a possible mechanism of analgesia. *Neurosci.Lett.* 1993; 150:
14
15 145-8
16
17
18
19
20
21 25. Lin CS, Tsaur ML, Chen CC, Wang TY, Lin CF, Lai YL, Hsu TC, Pan YY, Yang CH,
22
23 Cheng JK: Chronic intrathecal infusion of minocycline prevents the development of
24
25 spinal-nerve ligation-induced pain in rats. *Reg Anesth.Pain Med.* 2007; 32: 209-16
26
27
28
29
30
31 26. Boyle DL, Jones TL, Hammaker D, Svensson CI, Rosengren S, Albani S, Sorkin L,
32
33 Firestein GS: Regulation of peripheral inflammation by spinal p38 MAP kinase in rats.
34
35 *PLoS.Med.* 2006; 3: e338
36
37
38
39
40
41 27. Kobayashi M, Takeyoshi I, Yoshinari D, Matsumoto K, Morishita Y: P38
42
43 mitogen-activated protein kinase inhibition attenuates ischemia-reperfusion injury of the
44
45 rat liver. *Surgery* 2002; 131: 344-9
46
47
48
49
50
51 28. Furuichi K, Wada T, Iwata Y, Sakai N, Yoshimoto K, Kobayashi KK, Mukaida N,
52
53 Matsushima K, Yokoyama H: Administration of FR167653, a new anti-inflammatory
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 compound, prevents renal ischaemia/reperfusion injury in mice. *Nephrol.Dial.Transplant.*
6
7
8 2002; 17: 399-407
9

10
11 29. Kawashima Y, Takeyoshi I, Otani Y, Koibuchi Y, Yoshinari D, Koyama T, Kobayashi M,
12
13
14
15 Matsumoto K, Morishita Y: FR167653 attenuates ischemia and reperfusion injury of the
16
17
18 rat lung with suppressing p38 mitogen-activated protein kinase. *J.Heart Lung Transplant.*
19
20
21 2001; 20: 568-74
22

23
24 30. Daulhac L, Mallet C, Courteix C, Etienne M, Duroux E, Privat AM, Eschalier A, Fialip J:
25
26
27
28 Diabetes-induced mechanical hyperalgesia involves spinal mitogen-activated protein
29
30
31 kinase activation in neurons and microglia via N-methyl-D-aspartate-dependent
32
33
34 mechanisms. *Mol.Pharmacol.* 2006; 70: 1246-54
35

36
37 31. Sweitzer SM, Peters MC, Ma JY, Kerr I, Mangadu R, Chakravarty S, Dugar S, Medicherla
38
39
40
41 S, Protter AA, Yeomans DC: Peripheral and central p38 MAPK mediates
42
43
44 capsaicin-induced hyperalgesia. *Pain* 2004; 111: 278-85
45

46
47 32. Cui XY, Dai Y, Wang SL, Yamanaka H, Kobayashi K, Obata K, Chen J, Noguchi K:
48
49
50
51 Differential activation of p38 and extracellular signal-regulated kinase in spinal cord in a
52
53
54 model of bee venom-induced inflammation and hyperalgesia. *Mol.Pain* 2008; 4: 17
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

33. Xu M, Bruchas MR, Ippolito DL, Gendron L, Chavkin C: Sciatic nerve ligation-induced proliferation of spinal cord astrocytes is mediated by kappa opioid activation of p38 mitogen-activated protein kinase. *J.Neurosci.* 2007; 27: 2570-81

34. Zhuang ZY, Gerner P, Woolf CJ, Ji RR: ERK is sequentially activated in neurons, microglia, and astrocytes by spinal nerve ligation and contributes to mechanical allodynia in this neuropathic pain model. *Pain* 2005; 114: 149-59

35. Ma W, Quirion R: Partial sciatic nerve ligation induces increase in the phosphorylation of extracellular signal-regulated kinase (ERK) and c-Jun N-terminal kinase (JNK) in astrocytes in the lumbar spinal dorsal horn and the gracile nucleus. *Pain* 2002; 99: 175-84

36. Mogensen T, Eliassen K, Ejlersen E, Vegger P, Nielsen IK, Kehlet H: Epidural clonidine enhances postoperative analgesia from a combined low-dose epidural bupivacaine and morphine regimen. *Anesth.Analg.* 1992; 75: 607-10

37. Wen YR, Suter MR, Kawasaki Y, Huang J, Pertin M, Kohno T, Berde CB, Decosterd I, Ji RR: Nerve conduction blockade in the sciatic nerve prevents but does not reverse the activation of p38 mitogen-activated protein kinase in spinal microglia in the rat spared nerve injury model. *Anesthesiology* 2007; 107: 312-21

- 1
2
3
4
5 38. Milligan ED, Twining C, Chacur M, Biedenkapp J, O'Connor K, Poole S, Tracey K,
6
7
8 Martin D, Maier SF, Watkins LR: Spinal glia and proinflammatory cytokines mediate
9
10 mirror-image neuropathic pain in rats. *J.Neurosci.* 2003; 23: 1026-40
11
12
13
14
15 39. Yamamoto N, Sakai F, Yamazaki H, Nakahara K, Okuhara M: Effect of FR167653, a
16
17
18 cytokine suppressive agent, on endotoxin-induced disseminated intravascular coagulation.
19
20
21 *Eur.J.Pharmacol.* 1996; 314: 137-42
22
23
24
25 40. Yamamoto N, Sakai F, Yamazaki H, Sato N, Nakahara K, Okuhara M: FR167653, a dual
26
27
28 inhibitor of interleukin-1 and tumor necrosis factor-alpha, ameliorates endotoxin-induced
29
30
31 shock. *Eur.J.Pharmacol.* 1997; 327: 169-74
32
33
34
35 41. Scholz J, Woolf CJ: The neuropathic pain triad: neurons, immune cells and glia.
36
37
38 *Nat.Neurosci.* 2007; 10: 1361-8
39
40
41
42 42. Koistinaho M, Koistinaho J: Role of p38 and p44/42 mitogen-activated protein kinases in
43
44
45 microglia. *Glia* 2002; 40: 175-83
46
47
48
49 43. Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, Julius D: The
50
51
52 capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 1997; 389:
53
54
55 816-24
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

44. Caterina MJ, Leffler A, Malmberg AB, Martin WJ, Trafton J, Petersen-Zeitz KR, Koltzenburg M, Basbaum AI, Julius D: Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 2000; 288: 306-13

45. Davis JB, Gray J, Gunthorpe MJ, Hatcher JP, Davey PT, Overend P, Harries MH, Latcham J, Clapham C, Atkinson K, Hughes SA, Rance K, Grau E, Harper AJ, Pugh PL, Rogers DC, Bingham S, Randall A, Sheardown SA: Vanilloid receptor-1 is essential for inflammatory thermal hyperalgesia. *Nature* 2000; 405: 183-7

46. Pogatzki-Zahn EM, Shimizu I, Caterina M, Raja SN: Heat hyperalgesia after incision requires TRPV1 and is distinct from pure inflammatory pain. *Pain* 2005; 115: 296-307

47. Mizushima T, Obata K, Yamanaka H, Dai Y, Fukuoka T, Tokunaga A, Mashimo T, Noguchi K: Activation of p38 MAPK in primary afferent neurons by noxious stimulation and its involvement in the development of thermal hyperalgesia. *Pain* 2005; 113: 51-60

48. Michael GJ, Averill S, Nitkunan A, Rattray M, Bennett DL, Yan Q, Priestley JV: Nerve growth factor treatment increases brain-derived neurotrophic factor selectively in TrkA-expressing dorsal root ganglion cells and in their central terminations within the spinal cord. *J.Neurosci.* 1997; 17: 8476-90

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

49. Porreca F, Lai J, Bian D, Wegert S, Ossipov MH, Eglen RM, Kassotakis L, Novakovic S, Rabert DK, Sangameswaran L, Hunter JC: A comparison of the potential role of the tetrodotoxin-insensitive sodium channels, PN3/SNS and NaN/SNS2, in rat models of chronic pain. *Proc.Natl.Acad.Sci.U.S.A* 1999; 96: 7640-4

50. Obata H, Eisenach JC, Hussain H, Bynum T, Vincler M: Spinal glial activation contributes to postoperative mechanical hypersensitivity in the rat. *J.Pain* 2006; 7: 816-22

51. Colburn RW, DeLeo JA, Rickman AJ, Yeager MP, Kwon P, Hickey WF: Dissociation of microglial activation and neuropathic pain behaviors following peripheral nerve injury in the rat. *J.Neuroimmunol.* 1997; 79: 163-75

52. Suter MR, Wen YR, Decosterd I, Ji RR: Do glial cells control pain? *Neuron Glia Biol.* 2007; 3: 255-68

53. Echeverry S, Shi XQ, Zhang J: Characterization of cell proliferation in rat spinal cord following peripheral nerve injury and the relationship with neuropathic pain. *Pain* 2008; 135: 37-47

54. Svensson CI, Fitzsimmons B, Azizi S, Powell HC, Hua XY, Yaksh TL: Spinal p38beta isoform mediates tissue injury-induced hyperalgesia and spinal sensitization.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

J.Neurochem. 2005; 92: 1508-20

55. Crown ED, Ye Z, Johnson KM, Xu GY, McAdoo DJ, Hulsebosch CE: Increases in the activated forms of ERK 1/2, p38 MAPK, and CREB are correlated with the expression of at-level mechanical allodynia following spinal cord injury. *Exp.Neurol.* 2006; 199: 397-407
56. Winkelstein BA, DeLeo JA: Nerve root injury severity differentially modulates spinal glial activation in a rat lumbar radiculopathy model: considerations for persistent pain. *Brain Res.* 2002; 956: 294-301
57. Romero-Sandoval A, Eisenach JC: Spinal cannabinoid receptor type 2 activation reduces hypersensitivity and spinal cord glial activation after paw incision. *Anesthesiology* 2007; 106: 787-94

Figure Legends

Figure 1. Plantar incision induces p38 activation in the ipsilateral L4-5 spinal dorsal horns from (A) naïve rats and (B) rats at 1 day postoperatively. Higher magnification of p-p38-immunoreactive cells with long cytoplasmic processes (arrowhead) shown in (C) from the square marked in (B). (D) Quantification of the number of the p-p38 immunoreactive cells per section in the spinal dorsal horn after plantar incision. A significant increase is observed at 1 h, 3 h, day 1 and day 3 postoperative time. Plantar incision reduces paw withdrawal threshold to (E) von Frey filaments ($n = 6$) and to (F) radiant heat ($n = 8$). Data represent the mean \pm S.E.M. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ (repeated measurement ANOVA; $n = 6$) versus before incision (Pre-op). Scale bar = 100 μm for A and B, 20 μm for C.

Figure 2. Double-labeled immunofluorescence of p-p38, neuronal specific nuclear protein (NeuN, a neuronal marker), glial fibrillary acidic protein (GFAP, an astrocyte marker), and OX-42 (a microglia marker) in the spinal dorsal horn ipsilateral to paw incision at 1h (A-F), 3h (G-L), and 1d (M-R) postoperatively. (A, G, and H) Double staining demonstrates colocalization (arrows) of p-p38 (red) and NeuN (green) in neurons from spinal cord of 1h and 3h group, but not in that of 1d. B, H, and N show higher magnification of colocalization from A, G, and H, respectively. (C, I, and O)

Double labeling of p-p38 (red) with GFAP (green) showed no colocalization in the spinal dorsal horn at every time point. **(D-F, J-L, and P-R)** Double staining for p-p38 (red) and OX-42 (green) reveals a heavy colocalization Scale bar for A, C-G, I-M, and O-R 200 μm ; for B, H, and N 50 μm .

Figure 3. Representative photomicrographs illustrating OX-42 immunoreactivity in ipsilateral spinal dorsal horn **(A)** 3 h, **(B)** 1 day, **(C)** 3 days, **(D)** 7 days after hind paw incision. **(E)** Predefined rectangles at the superficial dorsal horn of a naïve rat were shown. The images were digitalized captured to calculate the pixel numbers. **(F)** Spinal microglial activation after a unilateral hind paw incision was indicated by the pixel ratio of the L4-5 superficial dorsal horn to the white matter. The pixel ratio was increased and reached the peak 3 days after incision. Scale bar = 100 μm for A-D and E. * $P < 0.05$ vs 3 h, repeated measure ANOVA with Bonferroni *post hoc* test; n = 4-5 for each time point.

Figure 4. Effects of p38 inhibitor FR167653 pretreatment on withdrawal threshold to von Frey stimulation and withdrawal threshold to thermal stimulation before and after plantar incision. **(A)** FR167653 attenuates plantar incision-induced mechanical hypersensitivity. Rats were pretreated with intrathecal bolus of saline (n = 10), vehicle

(10% DMSO in saline; n = 9), 20 μ g (n = 8), or 50 μ g (n = 8) FR167653. **(B)** FR167653 only prevents incision-induced thermal hyperalgesia at 1h. Rat from saline (n = 12), vehicle (20% DMSO in saline; n = 8) or 100 μ g (n = 9) FR167653. Data represented as the mean \pm S.E.M. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ versus saline; + $P < 0.05$, ++ $P < 0.01$ versus vehicle; # $P < 0.05$, ## $P < 0.01$ versus 20 μ g FR167653; one-way ANOVA with Bonferroni *post hoc* test.

Figure 5. Intrathecal injection of p38 inhibitor FR167653 attenuates plantar incision-induced p38 activation in the ipsilateral spinal dorsal horn. Rats were pretreated with either **(A)** saline or **(B)** FR167653 (50 μ g). **(C)** A significant decrease of p-p38-immunoreactive cells was observed both at dosage of 20 μ g and 50 μ g. Data represent as the mean \pm SEM. *** $P < 0.001$ versus saline-treated rats; ANOVA (n = 5 for each group). Scale bar is 200 μ m for A and B

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)




Figure 4
[Click here to download high resolution image](#)





Figure 5
[Click here to download high resolution image](#)

