

LETTER TO THE EDITOR

A POSSIBLE WAY TO MOTIVATE AMBIVALENT PATIENTS TO UNDERGO DETOXIFICATION

NICOLAS BERTHOLET and JEAN-BERNARD DAEPPEN*

Alcohol Treatment Centre, Lausanne, Switzerland

(Received 12 July 2005; first review notified 15 August 2005; in final revised form 19 September 2005; accepted 22 October 2005; advance access publication 18 November 2005)

Health-related quality-of-life and well-being are two important outcome measures for inpatients attending alcohol treatment programs. Rapid improvement in well-being and health related quality-of-life may be a possible way to motivate ambivalent patients to undergo detoxification.

The present letter is an ancillary report of a randomized controlled trial on pharmacological management of alcohol withdrawal. It focuses on changes in patients' perception of well-being and health-related quality-of-life over the first 3 days of inpatient detoxification.

During a 12-month period, consecutive patients admitted to an alcohol treatment inpatient unit were considered for inclusion. The selection process was described in the principal report (Daeppen *et al.*, 2002). Subjects' well-being and perceived health-related quality of life were assessed at intake using four questions from the well-being schedule (McDowell and Newell, 1996) and three dimensions from the MOS-SF-36 (physical functioning, vitality, and mental health) (Ware and Sherbourne, 1992). Perceived well-being and health-related quality of life were measured again at withdrawal day 4 using similar measures adapted to reflect the three first detoxification days. Wilcoxon tests were used to compare health-related measures between withdrawal days 1 and 4. Among the 126 included individuals, 4 left the programme within the first 3 days, 3 had exclusion criteria apparent only after inclusion, 1 had somnolence and 1 had several falls, resulting in a 117 (92.9 %) patient sample. The mean \pm SD age was 46.6 ± 9.52 years; 76.9% were men ($n = 90$); 94.9% were white ($n = 111$).

Compared to withdrawal day 1, all mean scores were statistically improved at withdrawal day 4 toward a self-perception of a better functioning and health: health concerns score

(3.72 vs 5.97, $Z = -5.68$, $P < 0.001$), anxiety score (3.84 vs 6.9, $Z = -7.46$, $P < 0.001$), energy score (5.43 vs 3.93, $Z = -5.20$, $P < 0.001$), feeling good vs. depressed score (6.12 vs 3.9, $Z = -7.06$, $P < 0.001$), physical functioning (MOS-SF-36) (87.91 vs 84.30, $Z = -4.08$, $P < 0.001$), vitality (MOS-SF-36) (57.61 vs 42.87, $Z = -6.85$, $P < 0.001$), and mental health (MOS-SF-36) (66.99 vs 46.26, $Z = -7.91$, $P < 0.001$).

This letter suggests rapid improvement in all examined dimensions of perceived well-being and health-related quality of life over the first 3 days of alcohol withdrawal among alcohol-dependent individuals. The data reported are consistent with observations on perceived health-related quality-of-life associated with abstinence published in the literature (Foster *et al.*, 1999). These findings may be used as a motivational tool to encourage alcohol-dependent patients to undergo detoxification and as positive feedback for patients' during early alcohol withdrawal.

REFERENCES

- Daeppen, J. B., Gache, P., Landry, U. *et al.* (2002) Symptom-triggered vs. fixed-schedule doses of benzodiazepine for alcohol withdrawal: a randomised treatment trial. *Archives of Internal Medicine* **162**, 1117–1121.
- Foster, J. H., Powell, J. E., Marshall, E. J. *et al.* (1999) Quality of life in alcohol-dependent subjects: a review. *Quality of Life Research* **8**, 255–261.
- McDowell, I. and Newell, C. (1996) The general well-being schedule. In *Measuring Health: A Guide to Rating Scales and Questionnaires*, 2nd edn, pp. 206–213. Oxford University Press, Oxford.
- Ware, J.E., Jr and Sherbourne, C. D. (1992) The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Medical Care* **30**, 473–483.

*Author to whom correspondence should be addressed at: Alcohol Treatment Centre, Mont-Paisible 16, 1011 Lausanne CHUV, Switzerland. Tel.: +4121 314 08 75; Fax: +4121 314 05 62; E-mail: Jean-Bernard.Daeppen@chuv.ch